



UvA-DARE (Digital Academic Repository)

The transient IGR J17361-4441 in the globular cluster NGC 6388

an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole

Wijnands, R. ; Yang, Y.J.; Degenaar, N. ; Altamirano, D.; Tudose, V.; Heinke, C.O.

Publication date

2011

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Wijnands, R., Yang, Y. J., Degenaar, N., Altamirano, D., Tudose, V., & Heinke, C. O. (2011). The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole. *The astronomer's telegram*. <https://www.astronomerstelegram.org/?read=3595>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

ATel On

Patreon
Mastodon
Twitter

The Astronomer's Telegram

Post | Search | Policies
Credential | Feeds | Email

9 Dec 2022; 10:18 UT

This space for free for your conference.

MIAPP workshop on
Interacting Supernovae
6 February - 3 March 2023
Garching, Germany

Ne/O Si

Thanks to Patrons, The Astronomer's Telegram is free to read, free to publish and always will be. Thank you.

[[Previous](#) | [Next](#) | [ADS](#)]

The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole

ATel #3595; *R. Wijnands, Y. J. Yang, N. Degenaar, D. Altamirano (University of Amsterdam), V. Tudose (ASTRON), C. O. Heinke (University of Alberta)*

on 26 Aug 2011; 10:49 UT

Credential Certification: Rudy Wijnands (rudy@space.mit.edu)

Subjects: Radio, X-ray, Request for Observations, Black Hole, Globular Cluster, Neutron Star, Transient

Referred to by ATel #: [3627](#)

[Tweet](#)

Gibaud et al. (2011, ATel #[3565](#)) reported the INTEGRAL discovery of a new transient, IGR J17361-4441, located in the Galactic globular cluster NGC 6388 (Ferrigno et al. 2011, ATel #[3566](#)). Swift/XRT has monitored this source from 16 to 25 August on a nearly daily basis (see ATel #[3566](#) for the first observation). The 0.5-10 keV unabsorbed flux in those observations was roughly constant, typically $(4-6)e^{-11}$ erg/s/cm², corresponding to a 0.5-10 keV X-ray luminosity of $(6-9)e^{35}$ erg/s (for a distance of 11.5 kpc), making this source a very-faint X-ray transient (VFXT; see Wijnands et al. 2006, A&A, 449, 1117, for a discussion about VFXTs). The spectrum is very hard, with a photon index of 0.6-1.0 (see also ATel #[3566](#)).

Considering its luminosity and location in a globular cluster, the system is most probably a neutron-star low-mass X-ray binary (NS-LMXB), although the hard X-ray spectrum is atypical for LMXB transients, which normally show indices of 1.6-2.2 in this luminosity range. However, there exist a small sub-group of NS-LMXBs with similarly hard spectra: accreting X-ray pulsars with high magnetic field ($>1e^{12}$ Gauss) neutron stars (e.g., Her X-1, GRO J1744-28; see Becker et al. 1977, ApJ, 214, 879; Giles et al. 1996, ApJ, 469, L25). Sensitive X-ray observations (e.g., Chandra, XMM-Newton) could detect pulsations with periods above ~ 0.1 second as observed for this class. This would be the first such transient found in a globular cluster.

Alternatively, the source could be an LMXB with an ordinary, low magnetic field ($<1e^{10}$ Gauss) neutron star. In that case, long observations might detect type-I X-ray bursts and high-sensitivity, high-time resolution observations (e.g., with XMM-Newton) might detect

Related

- [3958](#) IGR J17361-4441: ~ 200 days of monitoring with Swift/XRT
- [3627](#) Chandra localisation of IGR J17361-4441 in the globular cluster NGC 6388
- [3617](#) ATCA radio non-detection of IGR J17361-4441
- [3595](#) The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole
- [3566](#) Swift/XRT localization of IGR J17361-4441 in the globular cluster NGC 6388
- [3565](#) A new hard X-ray transient discovered by INTEGRAL: IGR J17361-4441

pulsations > 100 Hz. This source probably has a rather low time-averaged accretion rate and it such systems often harbor accreting millisecond X-ray pulsars (e.g. Wijnands 2008, AIPC, 1010, 382). Detecting either fast pulsations or type-I X-ray bursts from the system will prove it is an ordinary NS-LMXB, although its enigmatic X-ray spectrum still needs to be explained.

Thirdly, NGC 6388 may harbor an intermediate mass black hole (of $(6-20)E3$ solar masses, Lanzoni et al. 2007, ApJ, 670, 1065, Lutzgendorf et al. 2011, A&A, 533, 36). This outburst could be an accretion outburst from an intermediate mass black hole, as the X-ray position is currently consistent with the cluster center. A scheduled Chandra observation of this source will determine its position, and may rule out this hypothesis. Alternatively, sensitive radio observations might detect elevated radio emission at a level of a few tens of mJy (using the results of Merloni et al. 2003, MNRAS, 345, 1057; Maccarone 2004, MNRAS, 351, 1049, and the observed X-ray luminosity) which would also provide support for this hypothesis.

Since this VFXT in NGC 6388 is very unusual in its behavior, we strongly urge follow-up observations of this source at all wavelengths to gain more insight in its nature.

[[Telegram Index](#)]

R. E. Rutledge, Editor-in-Chief

rrutledge@astronomerstelegam.org

Derek Fox, Editor

dfox@astronomerstelegam.org