



## UvA-DARE (Digital Academic Repository)

### Understanding the complex dynamics of financial markets through microsimulation

Qiu, G.

**Publication date**  
2011

[Link to publication](#)

#### **Citation for published version (APA):**

Qiu, G. (2011). *Understanding the complex dynamics of financial markets through microsimulation*.

#### **General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

#### **Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

# Appendix B

## Analytical solution for a special case of the options market model

Here we show analytically that, if the mean of the speculators' expected prices  $M_{SP}^t$  coincides with the current price of the underlying  $S^t$ , the simulated option prices based on the S model converge to the corresponding BS prices.

The transaction quantity of a speculator is

$$Q_{SP}^{i,n,\phi,t} = \lambda_{SP}(\max(\phi(S_{SP}^{i,t} - K^n), 0) - V^{n,\phi,t}). \quad (\text{B.1})$$

In the S model, all the traders are speculators, so

$$\begin{aligned} Q_{SP}^{n,\phi,t} &= \sum_{i=1}^{N_{tr}} Q_{SP}^{i,n,\phi,t} \\ &= \lambda_{SP} \left[ \sum_{i=1}^{N_{tr}} \max(\phi(S_{SP}^{i,t} - K^n), 0) - N_{tr} V^{n,\phi,t} \right]. \end{aligned} \quad (\text{B.2})$$

The price updating rule is

$$V^{n,\phi,t+1} = V^{n,\phi,t} + \frac{\beta Q_{SP}^{n,\phi,t}}{N_{tr}}. \quad (\text{B.3})$$

The price of each option eventually converges, i.e.,  $V^{n,\phi,t+1} = V^{n,\phi,t}$ . Then,

$$Q^{n,\phi,t} = 0. \quad (\text{B.4})$$

---

Substituting Equation (B.4) in Equation (B.2), we have

$$\sum_{i=1}^{N_{tr}} \max(\phi(S_{SP}^{i,t} - K^n), 0) - N_{tr} V^{n,\phi,t} = 0, \quad (\text{B.5})$$

i.e. (law of large numbers),

$$\begin{aligned} V^{n,\phi,t} &= \frac{1}{N_{tr}} \left( \sum_{i=1}^{N_{tr}} \max(\phi(S_{SP}^{i,t} - K^n), 0) - V^{n,\phi,t} \right) \\ &\approx E[\max(\phi(S_{SP}^{i,t} - K^n), 0)]. \end{aligned} \quad (\text{B.6})$$

Since we assume that  $S_{SP}^{i,t} (i = 1, 2, \dots, N_{tr})$  follow a lognormal distribution, the right hand side of Equation (B.6) is equal to the corresponding Black-Scholes solution.