UvA-DARE (Digital Academic Repository)

Statins augment the chemosensitivity of colorectal cancer cells inducing epigenetic reprogramming and reducing colorectal cancer cell 'stemness' via the bone morphogenetic protein pathway

Kodach, L.L.; Jacobs, R.J.; Voorneveld, P.W.; Wildenberg, M.E.; Verspaget, H.W.; van Wezel, T.; Morreau, H.; Hommes, D.W.; Peppelenbosch, M.P.; van den Brink, G.R.; Hardwick, J.C.H.

DOI

[10.1136/gut.2011.237495](https://doi.org/10.1136/gut.2011.237495)

Publication date

2011

Document Version

Final published version

Published in

Gut

[Link to publication](#)

Citation for published version (APA):

Kodach, L. L., Jacobs, R. J., Voorneveld, P. W., Wildenberg, M. E., Verspaget, H. W., van Wezel, T., Morreau, H., Hommes, D. W., Peppelenbosch, M. P., van den Brink, G. R., & Hardwick, J. C. H. (2011). Statins augment the chemosensitivity of colorectal cancer cells inducing epigenetic reprogramming and reducing colorectal cancer cell 'stemness' via the bone morphogenetic protein pathway. *Gut*, 60(11), 1544-1553. <https://doi.org/10.1136/gut.2011.237495>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Statins augment the chemosensitivity of colorectal cancer cells inducing epigenetic reprogramming and reducing colorectal cancer cell 'stemness' via the bone morphogenetic protein pathway

Liudmila L Kodach,¹ Rutger J Jacobs,¹ Philip W Voorneveld,¹ Manon E Wildenberg,¹ Henricus W Verspaget,¹ Tom van Wezel,³ Hans Morreau,³ Daniel W Hommes,¹ Maikel P Peppelenbosch,⁴ Gijs R van den Brink,^{1,2,5} James C H Hardwick¹

► Additional figure and tables are published online only. To view these files please visit the journal online (<http://gut.bmj.com>).

¹Department of Gastroenterology and Hepatology, Leiden University Medical Center, Leiden, The Netherlands

²Tytgat Institute for Liver and Intestinal Research, Academic Medical Center, Amsterdam, The Netherlands

³Department of Pathology, Leiden University Medical Center, Leiden, The Netherlands

⁴Department of Gastroenterology and Hepatology, Erasmus MC-University Medical Center Rotterdam, Rotterdam, The Netherlands

⁵Department of Gastroenterology & Hepatology, Academic Medical Center, Amsterdam, The Netherlands

Correspondence to

James Hardwick, Leiden University Medical Center, Leiden University Medical Center, Albinusdreef 2, Leiden 2300RC, The Netherlands; j.c.h.hardwick@lumc.nl

LLK and RJJ contributed equally to this work.

Revised 2 March 2011
Accepted 19 March 2011
Published Online First
6 May 2011

ABSTRACT

Background Promoter hypermethylation is an important and potentially reversible mechanism of tumour suppressor gene silencing in cancer. Compounds that demethylate tumour suppressor genes and induce differentiation of cancer cells, but do not have toxic side effects, would represent an exciting option in cancer therapy. Statins are cholesterol-lowering drugs with an excellent safety profile and associated with a reduced incidence of various cancers including colorectal cancer (CRC). The authors have previously shown that statins act by activating tumour suppressive bone morphogenetic protein (BMP) signalling in CRC, increasing expression of *BMP2*. *BMP2* is silenced by hypermethylation in gastric cancer.

Aim To investigate whether *BMP2* is methylated in CRC, whether statins can reverse this, and what implications this has for the use of statins in CRC.

Methods Methylation-specific PCR, bisulphite sequencing, immunoblotting, reverse transcription PCR, quantitative PCR, fluorescence-activated cell sorting analysis, an in vitro DNA methyltransferase (DNMT) assay, and cell viability studies were performed on CRC cells. The effect of statins was confirmed in a xenograft mouse model.

Results *BMP2* is silenced by promoter hypermethylation in cell lines with the hypermethylator phenotype and in primary tumours. Treatment with lovastatin downregulates DNMT activity, leading to *BMP2* promoter demethylation and to upregulation of expression of *BMP2* as well as other genes methylated in CRC. Statins alter gene expression, indicating a shift from a stem-like state to a more differentiated state, thereby sensitising cells to the effects of 5-fluorouracil. In a xenograft mouse model, simvastatin treatment induces *BMP2* expression, leading to differentiation and reduced proliferation of CRC cells.

Conclusions Statins act as DNMT inhibitors, demethylating the *BMP2* promoter, activating BMP signalling, inducing differentiation of CRC cells, and reducing 'stemness'. This study indicates that statins may be able to be used as differentiating agents in combined or adjuvant therapy in CRC with the CpG island methylator phenotype.

INTRODUCTION

Aberrant DNA methylation of CpG islands in the promoter regions of many genes has been observed

Significance of this study

What is already known about this subject?

- The bone morphogenetic protein (BMP) pathway is an important tumour suppressor pathway in colorectal cancer (CRC).
- Statins are cholesterol-lowering drugs with an excellent safety profile and associated with a reduced incidence of various cancers including CRC.
- Statins induce apoptosis in CRC cells by activating the BMP pathway.
- The *BMP2* promoter is methylated in a large proportion of gastric cancers.

What are the new findings?

- *BMP2* is silenced by promoter hypermethylation in a subgroup of CRCs.
- Statin treatment inhibits DNA methyltransferase activity, demethylates the promoters of *BMP2* and other tumour suppressor genes methylated in CRC.
- Statins induce differentiation in CRC cells in vitro and in a xenograft model in vivo and reduce stemness.
- Statins increase the chemosensitivity of cancer cells to 5-fluorouracil.

How might it impact on clinical practice in the foreseeable future?

- Statins could be used as part of combined or adjuvant therapy in CRC with the CpG island methylator phenotype (CIMP).

in human colorectal cancer (CRC) and is associated with tumour suppressor gene silencing. Cancers that show extensive DNA methylation in the promoter regions of specific genes have been described as having the CpG island methylator phenotype (CIMP),¹ and these cancers are resistant to current chemotherapy.² Epigenetic alterations do not involve changes in the DNA sequence and are thus potentially reversible. This has already found clinical application in cancer therapy where demethylating agents have proven to be a valuable option in selected malignancies.

DNA methylation is regulated by DNA methyltransferases (DNMTs). DNMTs are enzymes that catalyse the addition of methyl groups to cytosine residues in DNA. The activity of DNMTs is elevated in CRC cells,³ and the inhibition of DNMT activity can strongly inhibit the formation of tumours in vivo.⁴ DNMT inhibitors have been intensively studied as promising new drugs for cancer therapy. 5-Azacytidine and decitabine have already entered clinical practice,⁵ but currently known DNMT inhibitors cause significant toxicity,^{6,7} as they become incorporated into RNA and interfere with protein translation.⁸ For this reason, compounds acting as DNMT inhibitors but not having toxic side effects would open up new opportunities in cancer therapy.

Epigenetic reprogramming involving changes in promoter methylation is also the mechanism underlying cell differentiation. Differentiating agents are already successfully used in combined chemotherapy where they are thought to force relatively poorly differentiated cancer stem cells to differentiate, making them more sensitive to the chemotherapeutic agents that are administered at the same time.⁹ Bone morphogenetic proteins (BMPs) have also been used for this purpose by intratumoural injection of recombinant protein in animal models.¹⁰ This methodology is not attractive for clinical application in humans, but small molecular compounds that activate the BMP pathway could be an attractive alternative strategy.

The BMP pathway plays an important role in intestinal epithelial homeostasis¹¹ and CRC.^{12,13} BMP signalling promotes intestinal differentiation and inhibits stem cell activation.^{14,15} Germline mutations in *BMPR1a* and *SMAD4* are the cause of familial juvenile polyposis syndrome,¹⁶ a syndrome with a high lifetime risk of developing CRC. Genome-wide association studies have identified gene alterations within multiple members of the BMP pathway as being associated with an increased risk of CRC, namely *BMP2*, *BMP4*, *Gremlin1* and *Smad7*.^{17,18} This makes the study of compounds that specifically modify the BMP pathway even more relevant.

3-Hydroxy-3-methylglutaryl-CoA reductase inhibitors, better known as statins, not only reduce serum cholesterol and decrease the incidence of cardiovascular and cerebrovascular events,^{19,20} but reduce the risk of developing CRC.^{21,22} In vitro studies show that statins inhibit cellular proliferation and induce apoptosis in CRC cells and in animal models.^{23,24} A screen of 30 000 compounds for their ability to enhance BMP2 expression for eventual use to enhance bone formation identified two statins as the two most active compounds.²⁵ We have subsequently shown that the actions of statins in CRC depend on their ability to upregulate BMP2 expression and activate the BMP pathway.²⁴ Interestingly, the *BMP2* gene is silenced by promoter hypermethylation in a large proportion of gastric cancers,²⁶ but whether this also occurs in CRCs is unknown.

In this study, we initially set out to investigate whether the *BMP2* promoter is hypermethylated in CRC. Having previously shown that statins upregulate BMP2 expression, we then investigated whether statin treatment influences the methylation status of the *BMP2* promoter and whether statins can also alter the methylation of promoters of other tumour suppressor genes methylated in CRC. We then assessed whether statins inhibit DNMT activity. To look at the overall effect of these epigenetic changes on CRC cell phenotype, we then investigated the effects of statins on differentiation markers and stem cell markers both in CRC cells in vitro and in a xenograft model in vivo. To establish whether the observed differentiation and 'stemness' changes are BMP pathway specific, we activated BMP signalling in CRC cells by transfection with *BMPR2* and observed

the same effects as seen in statin-treated cells. Finally, to investigate whether the differentiating effects of statins have potential clinical application, by modifying CRC chemosensitivity we assessed the effect of pretreatment with low doses of statins—too low to have cytotoxic effects on their own—on the sensitivity of CRC cells to 5-fluorouracil (5-FU). Our findings support the possible use of statins as agents to potentiate standard cytotoxic cancer therapy acting as demethylating agents that activate the BMP pathway.

MATERIALS AND METHODS

Cell culture, immunoblotting, DNA extraction and bisulphite modification, transfection and luciferase reporter assay, immunohistochemistry, RNA isolation and real-time reverse transcription (RT)-PCR are described in the online supplementary data section.

Methylation-specific PCR (MSP)

The primer sequences for MSP of *BMP2*, *HIC1* and *TIMP3* have been described previously.^{26,27} PCR was performed with 40 cycles of 94°C, 62°C (for *BMP2*), 65°C (for *HIC1* and *TIMP3*) and 72°C of 1 min each, preceded by a 5 min denaturing step at 94°C and followed by a 10 min extension step at 72°C. The products were electrophoresed on 5% agarose gel. Human genomic DNA from peripheral blood lymphocytes was used as an unmethylated control. Human genomic DNA treated in vitro with SssI methyltransferase (New England Biolabs, Beverly, Massachusetts, USA) was used as a positive control for the methylated reaction.

Bisulphite sequencing (BS)

For BS, cell line-derived DNA was treated with sodium bisulphite and amplified by PCR. The primers of *BMP2* for BS were 5'-GTATTTGGTTTATAGGGTTAGGAGAG-3' (forward) and 5'-CCAAATACTAACACACAACAACAAC-3' (reverse). PCR was performed with 35 cycles of 94°C, 62°C, and 72°C of 1 min each, preceded by a 5 min denaturing step at 94°C and followed by a 10 min extension step at 72°C. The PCR product was purified using the QIAquick Gel Extraction Kit (Qiagen, Venlo, The Netherlands). The purified PCR products were ligated into pCR2.1-TOPO using the TOPO-TA cloning system (Invitrogen, Breda, The Netherlands). Bacteria TOP10 were transformed with plasmids and cultured overnight, and the plasmid DNA was isolated using the Miniprep Kit (Qiagen). For each sample, five to ten separate clones were sequenced on an ABI 377 or 3100 automated sequencer (Applied Biosystems, Foster City, California, USA) using the ABI Big Dye Terminator Cycle Sequencing Kit (Applied Biosystems) and the original primers of *BMP2* for BS.

In vitro DNMT assay

HCT116 cells were treated with different concentrations of lovastatin or 5 µM 5-azacytidine (Sigma, St Louis, Missouri, USA) for 48 or 72 h. Cells were washed in ice-cold phosphate-buffered saline (PBS) and scraped into 200 µl ice-cold cell extract buffer (10 mM Hepes/KOH, pH 7.9, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM dithiothreitol (DTT) and 0.2 mM phenylmethanesulphonyl fluoride (PMSF)). The cells were kept on ice for 10 min, vortex-mixed for 10 s, and centrifuged at 4°C at 14000 rpm for 30 s. The supernatant was discarded, and the pellet was resuspended in 30 µl nuclear extraction buffer (20 mM Hepes/KOH, pH 7.9, 25% glycerol, 420 mM NaCl, 1.5 mM MgCl₂, 0.2 mM EDTA, 0.5 mM DTT and 0.2 mM PMSF), placed on ice for 20 min, and centrifuged at 4°C at 14000 rpm for 2 min. The supernatant was saved as the nuclear

extract and used to measure total DNMT activity using the EpiQuik DNA Methyltransferase Activity/Inhibition Assay Kit (ITK Diagnostics BV (Uithoorn, The Netherlands)) according to the manufacturer's instructions.

Xenograft mouse model

Eight female NMRI nu/nu mice were injected subcutaneously in the flank with 1×10^6 HCT116 cells in Matrigel (BD Bioscience, Breda, The Netherlands). Mice were fed ad libitum with food containing simvastatin (Arie Blok BV, Woerden, The Netherlands), thereby receiving 50 mg/kg/day for 3 weeks, initiated when the tumour volume reached 100–200 mm³. After the mice had been killed, the tumours were harvested and either frozen in liquid nitrogen and later homogenised in Trizol (Invitrogen) or embedded into paraffin blocks. We chose simvastatin because it was the second most potent statin in our in vitro experiments²⁴ and because it is licensed for use in humans in the Netherlands whereas lovastatin is not.

Chemosensitivity assay

Cells were plated in 96-well plates and treated with low-dose lovastatin (Sigma) (0.2 µM) or vehicle control in Dulbecco's modified Eagle's medium (DMEM) with 0.5% fetal calf serum (FCS). This low concentration of FCS allowed 5 days of exposure without the cells reaching confluence. Most studies use doses between 10 and 30 µM in vitro.²⁸ After 5 days of statin exposure, cells were incubated in fresh DMEM with 10% FCS for 24 h. Subsequently, cells were treated with 5-FU (Sigma) (1–50 µM) for 48 h in DMEM with 0.5% FCS. After treatment, MTT solution was added (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; final concentration 0.5 mg/ml, stock solution 5 mg/ml MTT in PBS), for 3 h. The medium was discarded and cells were lysed in acidified propan-2-ol. Absorbance was measured at 550–560 nm.

Fluorescence-activated cell sorting (FACS) analysis

Cells were treated with 0.2 µM lovastatin or vehicle control for 72 h, and then harvested, washed and stained in FACS buffer (PBS containing 1% bovine serum albumin) on ice with anti-CD166-PE (Becton Dickinson, Breda, The Netherlands). Cells were washed again, and CD166 cell surface levels were analysed by flow cytometry using a FACSCalibur (BD Bioscience) and FlowJo Software (Treestar, Ashland, Oregon, USA).

Statistical analysis

Statistical analysis was performed using the two-tailed Student *t* test, and $p < 0.05$ was considered significant. Data are shown as mean $\pm$ SEM.

RESULTS

BMP2 expression is often impaired due to promoter hypermethylation in CRC

We performed RT-PCR for *BMP2* in six CRC cell lines. HCT116 cells do not express *BMP2* at the mRNA level. RKO and SW48 (not shown) express very low levels of *BMP2* mRNA (figure 1A). These three cell lines are known to exhibit the CIMP phenotype, with hypermethylation of the promoters of several tumour suppressor genes. The main genetic characteristics of the cell lines used in this study are presented in online supplementary table 1.

We performed MSP for the *BMP2* promoter region starting 214 bp in front of exon 1 (figure 1B). This region contains a CpG island and has been shown to be methylated in gastric cancers.²⁶ The same cell lines that express reduced levels of *BMP2* mRNA

show CpG island methylation of the *BMP2* promoter region. MSP for HCT116 cells reveals only signals for methylated alleles of *BMP2*. RKO and SW48 seem to be partly methylated, as they exhibit both methylated and unmethylated signals, whereas only unmethylated alleles are found in SW480, DLD1, LOVO, HT29 and CACO2 cells (figure 1C). We verified the results of MSP by direct BS (figure 1D). The examined region of the promoter CpG island between positions –453 and –2 contains 50 CpG dinucleotides. HCT116 cells show dense methylation of the *BMP2* promoter with 96% of CpGs methylated. SW480 and CACO2 cells show only minimal methylation within the examined region, supporting the results of MSP.

It was not possible to perform direct BS on SW48 and RKO cell lines probably because of a mixture of methylated and unmethylated signals. Therefore we performed BS of multiple independent clones to determine the *BMP2* promoter methylation status in these cell lines. As seen in figure 1E, the level of *BMP2* promoter methylation in SW48 cells is 51%, and in RKO it is 72%.

To determine whether *BMP2* is methylated in a subgroup of patients with CRC, we performed MSP on bisulphate-modified DNA from 55 CRC tumours. Thirteen out of 55 tumours (23%) showed a methylated signal on MSP, confirming the relevance of our in vitro findings for patients (figure 1F). Further characterisation of these tumours by BRAF V600E mutation analysis was informative in 40 tumours and revealed nine BRAF V600E mutations. Three of the 13 cancers with *BMP2* methylation have BRAF mutations, one was uninformative for BRAF status and nine were wild-type (online supplementary table 2).

Lovastatin treatment leads to demethylation of the *BMP2* promoter region, upregulation of *BMP2* expression, and demethylation of the promoters of other genes methylated in CRC.

We evaluated the methylation status of the *BMP2* promoter after treatment of HCT116 cells with 2 µM lovastatin for 48 h and 72 h by MSP. As shown in figure 2A, lovastatin treatment leads to demethylation of the *BMP2* promoter, with the appearance of the unmethylated signal using MSP. To confirm this finding and to quantify the extent of demethylation, we performed BS of multiple independent clones from HCT116 cells and HCT116 cells treated for 72 h with lovastatin. The BS results show that *BMP2* promoter methylation decreases from 97% to 40% after treatment with 2 µM lovastatin for 72 h (figure 2B). We performed RT-PCR for *BMP2* to evaluate the differences in the expression of *BMP2* on mRNA level after treatment of HCT116 cells with 2 µM lovastatin for 3, 5 and 7 days. As shown in figure 2C, lovastatin treatment, as well as treatment with a strong demethylating agent, 5-deoxyazacytidine-C, upregulates the mRNA level of *BMP2* in a time-dependent manner.

We performed MSP analysis of three other genes methylated in HCT116 cells²⁷ before and after treatment with 2 µM lovastatin for 72 h. Lovastatin treatment leads to demethylation of the *hypermethylated in cancer 1 (HIC1)* and *tissue inhibitor of metalloproteinase 3 (TIMP3)* promoters, with reappearance of the unmethylated band after lovastatin treatment (figure 2D). We also performed MSP for *death-associated protein kinase (DAPK)* before and after lovastatin treatment, but did not see any difference in the methylation level of its promoter (data not shown). These three genes belong to a panel of genes hypermethylated in HCT116 cells and in CRCs.²⁷

Lovastatin inhibits DNMT activity in vitro

To further investigate the mechanism by which lovastatin leads to promoter demethylation, we performed an in vitro DNMT assay. The DNMT assay shows that treatment with different concentrations of lovastatin leads to a dose-dependent

Figure 1 (A) Reverse transcription (RT)-PCR analysis of *BMP2* expression in colon cancer cell lines (left) and corresponding glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) controls (right). HCT116 and RKO cell lines show absent *BMP2* expression. (B) Schematic representation of the large CpG island in the *BMP2* promoter region extending into exon 1. The double-headed black arrow indicates the region subjected to methylation-specific PCR analysis (MSP). (C) MSP analysis of the *BMP2* promoter region in colorectal cancer (CRC) cell lines. PCR products specific for unmethylated (u) and methylated (m) CpG sites were analysed in 2.5% agarose gels. In the positive controls, completely methylated and completely unmethylated DNA show only a methylated and unmethylated band, respectively. The same cell lines with absent *BMP2* expression by RT-PCR show full (HCT116) or partial (RKO) methylation of the *BMP2* promoter by MSP. Control is MSP performed without DNA. (D) Direct bisulphite sequencing (BS) of the *BMP2* promoter in SW480, CACO2 and HCT116 CRC cell lines. Solid and open circles represent methylated and unmethylated CpG sites, respectively. (E) BS of multiple independent clones from SW48 (B) and RKO (C) cell lines. Each horizontal row of circles represents the 52 CpG sites contained in the region. Solid and open circles represent methylated and unmethylated CpG sites, respectively. (F) MSP in human colorectal cancer tissue. The *BMP2* promoter is methylated in a subgroup of patients with CRC. T1–T6 show bands in the 'm' lanes representing *BMP2* promoter methylation. T7 and T8 show no *BMP2* promoter methylation. '-' is MSP without DNA.

downregulation of DNMT activity in HCT116 cells. Remarkably, even low concentrations of lovastatin (0.25 and 0.5 μ M), approximating those found in the serum of patients taking standard doses of statins, downregulate DNMT activity, implying a specific effect on DNMT and not a consequence of a general toxicity (figure 2E). The effect seems to be through inhibition of DNMT function, since protein and mRNA levels of DNMTs are not influenced by lovastatin treatment (online supplementary figure 1). These data further support the conclusion that lovastatin acts as a DNMT inhibitor and thus leads to promoter demethylation and re-expression of putative tumour suppressors such as *BMP2*.

Lovastatin induces differentiation and reduces 'stemness' in HCT116 cells

The BMP pathway is thought to induce differentiation of normal intestinal epithelial cells counteracting signals such as

WNT which impose a more stem-like phenotype.^{29 30} *BMP2* expression is upregulated by lovastatin, therefore we investigated whether lovastatin induces differentiation in CRC cells. We performed immunoblotting on HCT116 cells treated with lovastatin for markers of absorptive cell differentiation and quantitative RT-PCR for markers of goblet cell differentiation. We show that lovastatin induces dose-dependent upregulation of *BMP2* and villin, but not *CAII* expression, suggesting that the induced enterocyte differentiation is partial (figure 2F). Lovastatin also pushes CRC cells towards goblet cell differentiation, as judged by upregulation of the goblet cell markers, *Mucin2* and *Galectin4* (figure 3A). Goblet cell differentiation is controlled by Notch signalling.³¹ We see downregulation of the Notch pathway target, *HES1*, and upregulation of *KLF4*, a goblet cell-specific differentiation factor in the colon³² regulated by Notch signalling³³ in HCT116 cells treated with lovastatin. An important *Wnt* target and oncogene *c-Myc* and gene inducing

Figure 2 (A) Methylation-specific PCR analysis (MSP) of the *BMP2* promoter region for HCT116 cells treated with 2 μ M lovastatin for 48 and 72 h. PCR products specific for unmethylated (u) and methylated (m) CpG sites were analysed in 2.5% agarose gels. Unmeth DNA control, human genomic DNA from peripheral blood lymphocytes. Meth DNA control, human genomic DNA treated in vitro with SssI methyltransferase. (B) Demethylation of *BMP2* by lovastatin. HCT116 cells were treated with vehicle (HCT116) or 2 μ M lovastatin (HCT116+L) for 72 h, and bisulphite sequencing of multiple independent clones was performed. Each horizontal row of circles represents analysis of 52 CpG sites contained in the region. Solid and open circles represent methylated and unmethylated CpG sites, respectively. (C) Reverse transcription PCR analysis of *BMP2* expression in HCT116 cells treated at different time points with 2 μ M lovastatin or the demethylating agent, 5-deoxyazacytidine (AZA). Lovastatin and 5-deoxyazacytidine both led to re-expression of *BMP2* mRNA. (D) MSP of the *TIMP3* and *HIC1* promoter regions for HCT116 cells treated with different concentrations of lovastatin for 72 h. PCR products specific for unmethylated (u) and methylated (m) CpG sites were analysed in 2.5% agarose gels. (E) HCT116 cells were treated with 2 μ M lovastatin for 48 h, and then DNA methyltransferase (DNMT) Activity/Inhibition Assay was performed. Treatment with lovastatin inhibits DNMT activity in a time- and dose-dependent manner. DNMT positive control is provided by the manufacturer, and values obtained with DNMT positive control have been set at 100%. No nuclear extract was added in blank wells, and the measured absorbance was considered as a background level. (F) Immunoblots for *BMP2*, villin and *CaII* of HCT116 cells treated with different concentrations of lovastatin for 7 days. The protein expression was analysed using the corresponding specific antibody. Actin served as a loading control.

stemness³⁴ is also downregulated by lovastatin treatment (figure 3A), as well as the expression of two markers of cancer stem cells, *CD166* and *EpCAM*^{35–37} (figures 3B,C), further evidence that statins push CRC cells towards a more differentiated phenotype and away from a stem-like, crypt cell phenotype. The widely used marker of stem cells, *CD133*, is not an optimal marker in HCT116 cells as the *CD133* promoter is densely methylated in the HCT116 cell line and was therefore not used in this study.³⁸

Lovastatin induces differentiation in a xenograft model

Our previous work showed that simvastatin inhibited the growth of HCT116 xenografts in mice,²⁴ but the effect of

oral administration of statins on the differentiation of cells within the xenografts had not yet been studied. We performed immunohistochemical analysis on HCT116 xenografts in mice for the proliferation marker, Ki-67, and the differentiation markers, villin and *BMP2*. After 3 weeks of oral administration of simvastatin (50 mg/kg/day) HCT116 xenografts show significant downregulation of Ki-67 and upregulation of *BMP2* and villin protein expression (figure 4A,B). Simvastatin also induces differentiation of HCT xenografts towards the goblet cell lineage, as revealed by staining with periodic acid–Schiff (figure 4A). Simvastatin treatment also results in downregulation of *c-Myc* and *HES1* mRNA in xenografts (figure 4C).

Figure 3 (A) Quantitative reverse transcription (RT)-PCR analysis of *Villin*, *c-Myc*, *KLF4*, *HES1*, *Mucin2* and *Galectin4* expression in HCT116 cells treated with 0.2 or 2 μ M lovastatin for 72 h. (B) Quantitative RT-PCR analysis of stem cell marker expression in HCT116 cells treated with lovastatin. (C) Fluorescence-activated cell sorting analysis of CD166 expression in HCT116 cells treated with lovastatin. Data are from three experiments, mean $\pm$ SEM. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

Activation of BMP signalling leads to the differentiation of CRC cells

To test the hypothesis that statin-induced differentiation in CRC cells is due to the activation of BMP signalling, we transfected HCT116 cells with *BMP2* and activated BMP signalling sixfold (figure 5A). This results in upregulation of *Villin* and *Mucin2* expression, downregulation of Notch pathway activity, and downregulation of expression of cancer stem cells markers, *CD166* and *EpCAM*, as seen with statin treatment (figure 5B,C). These data suggest that activation of the BMP pathway induces differentiation and reduces stemness of CRC cells and implies that statins may induce the shift from a stem-like state to a more differentiated state of CRC cells by demethylation of the *BMP2* promoter and activation of BMP signalling.

Lovastatin enhances the cytotoxic effect of 5-FU

We tested the effects of lovastatin treatment on the chemosensitivity of CRC cells to the conventional chemotherapeutic drug, 5-FU (experimental setup is depicted in figure 6A). We treated HCT116 cells with a low concentration of lovastatin for 5 days in order to differentiate CRC cells. At this time point, no effects of this low concentration of lovastatin on cell viability was observed compared with untreated cells or ethanol-treated control cells (figure 6B). After the removal of lovastatin, cells were treated with different concentrations of 5-FU for 48 h. Lovastatin treatment significantly increases the sensitivity of cancer cells to 5-FU (figure 6C). Even at low concentrations of 5-FU that do not inhibit the viability of HCT116 cells when used alone, lovastatin pretreatment sensitises CRC cells and

Colon

Figure 4 (A) Immunohistochemistry for Ki-67, villin, BMP2 and periodic acid–Schiff (PAS) staining of goblet cells of HCT116 xenografts from mice treated with simvastatin and from controls. Original magnification: 20×. (B) The percentage of Ki-67-positive cells relative to all cells per 20× field (for every xenograft three images were scored). Error bars represent SEM (n=8). ***p<0.001. (C) Quantitative reverse transcription PCR analysis of *c-Myc* and *HES1* in HCT116 xenografts from mice treated with simvastatin and from controls. *p<0.05. Error bars represent SEM (n=8). GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

Figure 5 (A) HCT116 cells were transiently co-transfected with either pcDNA4/TO/BMPR2 plasmid or pcDNA4/TO control vector and BRE-Luc vectors, and the dual luciferase assay was performed 48 h after transfection. Data were normalised to *Renilla* luciferase activity. Data are from three experiments, mean $\pm$ SEM. (B) Immunoblots for villin of HCT116 cells transfected with BMPR2 or pcDNA. Protein expression was analysed using the corresponding specific antibody. Actin served as a loading control. (C) Quantitative reverse transcription PCR analysis of *Villin*, *HES1*, *Mucin2*, *CD166* and *EpCAM* of HCT116 cells transfected with pcDNA4/TO/BMPR2 plasmid or pcDNA4/TO control vector. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

leads to significant inhibition of cell viability. Importantly, concentrations of 5-FU of 1–10 μ M are clinically relevant, as they approximate serum (6–12 μ M)³⁹ and tissue (2–5 μ M)⁴⁰ levels in patients.

DISCUSSION

DNA hypermethylation of the promoter region of tumour suppressor genes occurs often in cancer and results in the transcriptional silencing and loss of function of critical tumour suppressor genes.⁴¹ In this study, we show that the *BMP2* promoter is methylated in a subgroup of CRC cell lines and primary cancer specimens. We use HCT116 cells for a large proportion of our studies, as *BMP2* is fully methylated in these cells. These cells have often been used for methylation studies^{42 43} and are a heavily methylated cell line as seen in CIMP.⁴⁴ However,

MLH1 is mutated in HCT116, rather than methylated, suggestive of Lynch syndrome origins.

We have analysed *BRAF* V600E mutation status in the primary cancer specimens in which we analysed *BMP2* methylation. *BRAF* V600E mutations are often used as a marker of CIMP status, but may underestimate the prevalence of CIMP by ~50%.⁴⁵ Of the cancers we tested, 25% show concomitant *BMP2* methylation and *BRAF* mutation. Although in cell lines *BMP2* promoter methylation is seen exclusively in those with the CIMP phenotype, analysis of *BRAF* only as a marker of CIMP is insufficient to confirm or refute this in tumour specimens.

We report that statins exhibit demethylating properties. Inhibition of DNMTs can be seen even at low statin concentrations (0.25 μ M), which are comparable to the serum levels of ~0.1 μ M measured in patients treated with standard doses for

Figure 6 (A) Schematic representation of the experimental setup. (B) HCT116 cells were plated in 96-well plates and either not treated or treated with ethanol (EtOH) or 0.2 μ M lovastatin (diluted in EtOH) for 5 days, and the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay was performed. Values are expressed as percentage of living cells relative to the control with control values set at 100%. Results represent the mean $\pm$ SEM (n=10). (C) After 5 days of treatment with lovastatin or EtOH and 1 day of no treatment, cells were treated with different concentrations of 5-fluorouracil (5-FU), and the MTT assay was performed. Values are expressed as a percentage of living cells relative to the control with control values set at 100%. Results represent the mean $\pm$ SEM (n=10). * p <0.05; *** p <0.001.

hypercholesterolaemia⁴⁶ and far lower than the maximum safely achievable levels in humans. These low concentrations of statins are safe and well tolerated by patients for years. Thus, in contrast with most known DNMT inhibitors, statins inhibit DNMTs and induce DNA demethylation at non-toxic doses. Our experiments in xenografts investigated whether the demethylating and differentiating properties of statins in vitro translate into the same differences in vivo. We chose oral administration of statins to more closely mimic their use in humans. We show that oral treatment with simvastatin induces BMP2 expression and leads to differentiation and to down-regulation of proliferation of colon cancer cells in a xenograft mouse model.

Lovastatin treatment induces DNA demethylation and the re-activation of *BMP2* gene expression, which is silenced by hypermethylation in CRC cells. Importantly, we find demethylation of two other methylated CRC genes, *TIMP3* and *HIC1*, after treatment with lovastatin, implying a more general effect on gene hypermethylation. We do not see demethylation of the *DAPK* promoter region. One possible explanation for this could be that other epigenetic factors that are not influenced by statin treatment—for example, histone modification—regulate methylation more prominently in this gene.⁴⁷

From these studies, it is unclear how statins inhibit DNMTs. We show that they have little or no influence on expression levels of DNMTs, and further studies are needed to determine the mechanism by which they inhibit DNMTs. It is also unclear from these studies what the influence of statins is on global methylation levels. Although CIMP CRCs exhibit widespread promoter hypermethylation, global levels of DNA methylation are often found to be lower in CRC. However, the importance and mechanism of action of global hypomethylation are less well established and the influence of statins, if any, on this phenomenon is a subject for future investigation.

Lovastatin treatment leads to increased expression of markers of intestinal epithelial cell differentiation and decreased expression of cancer stem cell markers. This could be due to demethylation of multiple genes; however, activation of the BMP pathway alone is sufficient to lead to the same effects on differentiation and stem cell markers as statin treatment, suggesting that the action of statins is primarily due to their effects on the BMP pathway. Low-dose statin treatment increases the sensitivity of CRC cells to the conventional chemotherapeutic drug, 5-FU. Increases in sensitivity to conventional chemotherapy have been seen with the use of other differentiation-inducing agents in CRC cells, such as the γ -secretase inhibitors⁴⁸ and PPAR- γ agonists,⁴⁹ and with multiple interventions aimed at differentiating CRC stem cells and thus improving their chemosensitivity, including the use of BMPs. Delivery of BMP to tumours is highly challenging and at present has only been successfully performed by intratumoural injection in mouse models of cancer. It is questionable whether delivery of sufficient amounts of BMPs at the required site can be achieved in human cancer therapy. Statins have been shown to increase levels of BMP2 in bone cells,²⁵ and we have shown that this also holds true for CRC cells. Although in bone cells the mechanism appears to be direct activation of the *BMP2* promoter, the promoter is often silenced in CRC, as we show here, but this can be reversed by statin therapy.

The CIMP is found not only in CRCs but also in colonic polyps^{50 51} and even in the normal colorectal mucosa in patients with hyperplastic polyposis.⁵² It has been observed that cigarette smoking is strongly associated with CIMP CRC, with a dose-response relationship with respect to the amount

smoked.⁵³ These groups of patients could potentially benefit from a chemopreventive agent exhibiting mild demethylating properties and an excellent safety profile. As statin treatment also reduces morbidity and mortality associated with cardiovascular disease, their demethylating effect could make them a particularly valuable chemopreventive agent in a well-defined group of patients.

In conclusion, the *BMP2* promoter is methylated in a subgroup of CRC cell lines and in patients with CRC. Lovastatin acts as a DNMT inhibitor and demethylates the *BMP2*, *TIMP3* and *HIC1* promoters. Statins decrease stemness and induce differentiation of CRC cells in vitro and in vivo, sensitising cells to 5-FU chemotherapy. Our study suggests a potential role for statins as chemopreventive or therapeutic agents in a subgroup of patients with CRC: those with, or prone to develop, CRC with the CIMP.

Funding JCHH is supported by The Netherlands Organization for Health Research and Development — (ZonMw VENI 916.76.087). LLK and RJJ are supported by the Dutch Cancer Society (KWF 2007-3725).

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Toyota M, Ahuja N, Ohe-Toyota M, *et al.* CpG island methylator phenotype in colorectal cancer. *Proc Natl Acad Sci U S A* 1999;**96**:8681–6.
2. Jover R, Nguyen TP, Perez-Carbonell L, *et al.* 5-Fluorouracil adjuvant chemotherapy does not increase survival in patients with CpG island methylator phenotype colorectal cancer. *Gastroenterology* 2011;**140**:1174–81.
3. De Marzo AM, Marchi VL, Yang ES, *et al.* Abnormal regulation of DNA methyltransferase expression during colorectal carcinogenesis. *Cancer Res* 1999;**59**:3855–60.
4. Laird PW, Jackson-Grusby L, Fazeli A, *et al.* Suppression of intestinal neoplasia by DNA hypomethylation. *Cell* 1995;**81**:197–205.
5. Silverman LR, Demakos EP, Peterson BL, *et al.* Randomized controlled trial of azacitidine in patients with the myelodysplastic syndrome: a study of the cancer and leukemia group B. *J Clin Oncol* 2002;**20**:2429–40.
6. Wijermans P, Lubbert M, Verhoef G, *et al.* Low-dose 5-aza-2'-deoxycytidine, a DNA hypomethylating agent, for the treatment of high-risk myelodysplastic syndrome: a multicenter phase II study in elderly patients. *J Clin Oncol* 2000;**18**:956–62.
7. Schrupp DS, Fischette MR, Nguyen DM, *et al.* Phase I study of decitabine-mediated gene expression in patients with cancers involving the lungs, esophagus, or pleura. *Clin Cancer Res* 2006;**12**:5777–85.
8. Cihak A. Biological effects of 5-azacytidine in eukaryotes. *Oncology* 1974;**30**:405–22.
9. Isik P, Cetin I, Tavit B, *et al.* All-transretinoic acid (ATRA) treatment-related pancarditis and severe pulmonary edema in a child with acute promyelocytic leukemia. *J Pediatr Hematol Oncol* 2010;**32**:e346–8.
10. Piccirillo SG, Reynolds BA, Zanetti N, *et al.* Bone morphogenetic proteins inhibit the tumorigenic potential of human brain tumour-initiating cells. *Nature* 2006;**444**:761–5.
11. Hardwick JC, van den Brink GR, Bleuming SA, *et al.* Bone morphogenetic protein 2 is expressed by, and acts upon, mature epithelial cells in the colon. *Gastroenterology* 2004;**126**:111–21.
12. Kodach LL, Wiercinska E, de Miranda NF, *et al.* The bone morphogenetic protein pathway is inactivated in the majority of sporadic colorectal cancers. *Gastroenterology* 2008;**134**:1332–41.
13. Kodach LL, Bleuming SA, Musler AR, *et al.* The bone morphogenetic protein pathway is active in human colon adenomas and inactivated in colorectal cancer. *Cancer* 2008;**112**:300–6.
14. Auclair BA, Benoit YD, Rivard N, *et al.* Bone morphogenetic protein signaling is essential for terminal differentiation of the intestinal secretory cell lineage. *Gastroenterology* 2007;**133**:887–96.
15. He XC, Zhang J, Li L. Cellular and molecular regulation of hematopoietic and intestinal stem cell behavior. *Ann N Y Acad Sci* 2005;**1049**:28–38.
16. Howe JR, Bair JL, Sayed MG, *et al.* Germline mutations of the gene encoding bone morphogenetic protein receptor 1A in juvenile polyposis. *Nat Genet* 2001;**28**:184–7.
17. Houlston RS, Webb E, Broderick P, *et al.* Meta-analysis of genome-wide association data identifies four new susceptibility loci for colorectal cancer. *Nat Genet* 2008;**40**:1426–35.
18. Broderick P, Carvajal-Carmona L, Pittman AM, *et al.* A genome-wide association study shows that common alleles of SMAD7 influence colorectal cancer risk. *Nat Genet* 2007;**39**:1315–17.
19. Downs JR, Clearfield M, Tyroler HA, *et al.* Air Force/Texas Coronary Atherosclerosis Prevention Study (AFCAPS/TEXCAPS): additional perspectives on tolerability of long-term treatment with lovastatin. *Am J Cardiol* 2001;**87**:1074–9.

20. **Furberg CD**, Adams HP Jr, Applegate WB, *et al.* Effect of lovastatin on early carotid atherosclerosis and cardiovascular events. Asymptomatic Carotid Artery Progression Study (ACAPS) Research Group. *Circulation* 1994;**90**:1679–87.
21. **Poynter JN**, Gruber SB, Higgins PD, *et al.* Statins and the risk of colorectal cancer. *N Engl J Med* 2005;**352**:2184–92.
22. **Bardou M**, Barkun A, Martel M. Effect of statin therapy on colorectal cancer. *Gut* 2010;**59**:1572–85.
23. **Agarwal B**, Rao CV, Bhendwal S, *et al.* Lovastatin augments sulindac-induced apoptosis in colon cancer cells and potentiates chemopreventive effects of sulindac. *Gastroenterology* 1999;**117**:838–47.
24. **Kodach LL**, Bleuming SA, Peppelenbosch MP, *et al.* The effect of statins in colorectal cancer is mediated through the bone morphogenetic protein pathway. *Gastroenterology* 2007;**133**:1272–81.
25. **Mundy G**, Garrett R, Harris S, *et al.* Stimulation of bone formation in vitro and in rodents by statins. *Science* 1999;**286**:1946–9.
26. **Wen XZ**, Akiyama Y, Baylin SB, *et al.* Frequent epigenetic silencing of the bone morphogenetic protein 2 gene through methylation in gastric carcinomas. *Oncogene* 2006;**25**:2666–73.
27. **Ohm JE**, McGarvey KM, Yu X, *et al.* A stem cell-like chromatin pattern may predispose tumor suppressor genes to DNA hypermethylation and heritable silencing. *Nat Genet* 2007;**39**:237–42.
28. **Agarwal B**, Bhendwal S, Halmos B, *et al.* Lovastatin augments apoptosis induced by chemotherapeutic agents in colon cancer cells. *Clin Cancer Res* 1999;**5**:2223–9.
29. **Vermeulen L**, De Sousa EMelo F, van der Heijden M, *et al.* Wnt activity defines colon cancer stem cells and is regulated by the microenvironment. *Nat Cell Biol* 2010;**12**:468–76.
30. **Sansom OJ**, Reed KR, Hayes AJ, *et al.* Loss of Apc in vivo immediately perturbs Wnt signaling, differentiation, and migration. *Genes Dev* 2004;**18**:1385–90.
31. **van Es JH**, van Gijn ME, Riccio O, *et al.* Notch/gamma-secretase inhibition turns proliferative cells in intestinal crypts and adenomas into goblet cells. *Nature* 2005;**435**:959–63.
32. **Katz JP**, Perreault N, Goldstein BG, *et al.* The zinc-finger transcription factor Klf4 is required for terminal differentiation of goblet cells in the colon. *Development* 2002;**129**:2619–28.
33. **Ghaleb AM**, Aggarwal G, Bialkowska AB, *et al.* Notch inhibits expression of the Kruppel-like factor 4 tumor suppressor in the intestinal epithelium. *Mol Cancer Res* 2008;**6**:1920–7.
34. **Takahashi K**, Tanabe K, Ohnuki M, *et al.* Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;**131**:861–72.
35. **Dalerba P**, Dylla SJ, Park IK, *et al.* Phenotypic characterization of human colorectal cancer stem cells. *Proc Natl Acad Sci U S A* 2007;**104**:10158–63.
36. **Ricci-Vitiani L**, Pagliuca A, Palio E, *et al.* Colon cancer stem cells. *Gut* 2008;**57**:538–48.
37. **Levin TG**, Powell AE, Davies PS, *et al.* Characterization of the intestinal cancer stem cell marker CD166 in the human and mouse gastrointestinal tract. *Gastroenterology* 2010;**139**:2072–82.
38. **Yi JM**, Tsai HC, Glöckner SC, *et al.* Abnormal DNA methylation of CD133 in colorectal and glioblastoma tumors. *Cancer Res* 2008;**68**:8094–103.
39. **Takimoto CH**, Yee LK, Venzon DJ, *et al.* High inter- and intrapatient variation in 5-fluorouracil plasma concentrations during a prolonged drug infusion. *Clin Cancer Res* 1999;**5**:1347–52.
40. **Peters GJ**, Lankelma J, Kok RM, *et al.* Prolonged retention of high concentrations of 5-fluorouracil in human and murine tumors as compared with plasma. *Cancer Chemother Pharmacol* 1993;**31**:269–76.
41. **Jones PA**, Baylin SB. The fundamental role of epigenetic events in cancer. *Nat Rev Genet* 2002;**3**:415–28.
42. **Rhee I**, Jair KW, Yen RW, *et al.* CpG methylation is maintained in human cancer cells lacking DNMT1. *Nature* 2000;**404**:1003–7.
43. **O’Gorman A**, Colleran A, Ryan A, *et al.* Regulation of NF-kappaB responses by epigenetic suppression of IkappaBalpha expression in HCT116 intestinal epithelial cells. *Am J Physiol Gastrointest Liver Physiol* 2010;**299**:G96–105.
44. **Paz MF**, Fraga MF, Avila S, *et al.* A systematic profile of DNA methylation in human cancer cell lines. *Cancer Res* 2003;**63**:1114–21.
45. **Goel A**, Nagasaka T, Arnold CN, *et al.* The CpG island methylator phenotype and chromosomal instability are inversely correlated in sporadic colorectal cancer. *Gastroenterology* 2007;**132**:127–38.
46. **Pan HY**, DeVault AR, Wang-Iverson D, *et al.* Comparative pharmacokinetics and pharmacodynamics of pravastatin and lovastatin. *J Clin Pharmacol* 1990;**30**:1128–35.
47. **Kondo Y**, Shen L, Cheng AS, *et al.* Gene silencing in cancer by histone H3 lysine 27 trimethylation independent of promoter DNA methylation. *Nat Genet* 2008;**40**:741–50.
48. **Akiyoshi T**, Nakamura M, Yanai K, *et al.* Gamma-secretase inhibitors enhance taxane-induced mitotic arrest and apoptosis in colon cancer cells. *Gastroenterology* 2008;**134**:131–44.
49. **Zhang YQ**, Tang XQ, Sun L, *et al.* Rosiglitazone enhances fluorouracil-induced apoptosis of HT-29 cells by activating peroxisome proliferator-activated receptor gamma. *World J Gastroenterol* 2007;**13**:1534–40.
50. **Chan AO**, Issa JP, Morris JS, *et al.* Concordant CpG island methylation in hyperplastic polyposis. *Am J Pathol* 2002;**160**:529–36.
51. **Wynter CV**, Walsh MD, Higuchi T, *et al.* Methylation patterns define two types of hyperplastic polyp associated with colorectal cancer. *Gut* 2004;**53**:573–80.
52. **Minoo P**, Baker K, Goswami R, *et al.* Extensive DNA methylation in normal colorectal mucosa in hyperplastic polyposis. *Gut* 2006;**55**:1467–74.
53. **Samowitz WS**, Albertsen H, Sweeney C, *et al.* Association of smoking, CpG island methylator phenotype, and V600E BRAF mutations in colon cancer. *J Natl Cancer Inst* 2006;**98**:1731–8.

Gut online

Visit **Gut online** and listen to the latest podcast, post comments and download any you might have missed. Keep informed and up to date by visiting gut.bmj.com.

Statins augment the chemosensitivity of colorectal cancer cells inducing epigenetic reprogramming and reducing colorectal cancer cell 'stemness' via the bone morphogenetic protein pathway

Liudmila L Kodach, Rutger J Jacobs, Philip W Voorneveld, et al.

Gut 2011 60: 1544-1553 originally published online May 6, 2011
doi: 10.1136/gut.2011.237495

Updated information and services can be found at:
<http://gut.bmj.com/content/60/11/1544.full.html>

These include:

Data Supplement

"online only data"

<http://gut.bmj.com/content/suppl/2011/10/12/gut.2011.237495.DC1.html>

References

This article cites 53 articles, 22 of which can be accessed free at:
<http://gut.bmj.com/content/60/11/1544.full.html#ref-list-1>

Article cited in:

<http://gut.bmj.com/content/60/11/1544.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Colon cancer](#) (1102 articles)

[Pancreatic cancer](#) (565 articles)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>