

The Cenozoic history of palms: global diversification, biogeography, and the decline of megathermal forests

Appendix 1

List of literature sources used to assemble the fossil pollen dataset

- Absy, M.L., Cozzuol, M.A., Senna, C.d.S.F.d. & Silva, S.A.F.d. (2003) Mudanças paleoambientais na Amazônia evidenciadas por registros de flora e fauna durante o Cenozóico. *I Congresso sobre Planejamento e Gestão das Zonas Costeiras dos Países de Expressão Portuguesa, IX Congresso da Associação Brasileira de Estudos do Quaternário, II Congresso do Quaternário dos Países de Língua Ibéricas*, Museu Paraense Emilio Goeldi, pp. 1–3.
- Absy, M.L. & Hammen, T. (1976) Some palaeoecological data from Rondonia, southern part of the Amazon Basin. *Acta Amazonica*, **6**, 293–299.
- Acharya, M. (2000) Early Eocene palynofossils from subsurface of mannargudi area, Tamil Nadu, India. *Geophytology*, **28**, 19–30.
- Adebayo, O.F., Orijemie, A.E. & Aturamu, A.O. (2012) Palynology of Bog-1 Well, Southeastern Niger Delta Basin, Nigeria. *International Journal of Science and Technology*, **2**, 214–222.
- Adegoke, O.S., Agumanu, A.E., Benkhelil, M.J. & Ajayi, P.O. (1986) New stratigraphic, sedimentologic and structural data on the Kerri-Kerri Formation, Bauchi and Borno States, Nigeria. *Journal of African Earth Sciences*, **5**, 249–277.
- Adegoke, O.S., Jan du Chêne, R.E., Agumanu, A.E. & Ajayi, P.O. (1978) Palynology and age of the Kerri-Kerri Formation, Nigeria. *Revista española de micropaleontología*, **10**, 267–282.
- Adeigbe, O.C., Ola-Buraimo, A.O. & Moronhunkola, A.O. (2013) Palynological characterization of the Tertiary offshore Emi-1 well, Dahomey Basin, southwestern Nigeria. *International Journal of Scientific and Technology Research*, **2**, 58–70.
- Adeniran, O.A. (2015) Upper cretaceous to Paleogene Palynosequence stratigraphy of H-1 well offshore eastern Dahomey Basin, southwestern Nigeria. *International Journal of Research and Innovations in Earth Science*, **2**, 81–88.
- Adeonipekun, P.A., Ehinola, O.A., Yussuph, I., Toluhi, A. & Oyelami, A. (2012) Bio-sequence stratigraphy of Shagamu Quarry outcrop, Benin basin, southwestern Nigeria. *World Applied Sciences Journal*, **18**, 91–106.
- Adeonipekun, P.A., Sowunmi, M.A. & Richards, K. (2017) A new Late Miocene to Pleistocene palynomorph zonation for the western offshore Niger Delta. *Palynology*, **41**, 2–16.
- Agrawal, S., Verma, P., Rao, M., Garg, R., Kapur, V.V. & Bajpai, S. (2017) Lignite deposits of the Kutch Basin, western India: Carbon isotopic and palynological signatures of the early Eocene hyperthermal event ETM2. *Journal of Asian Earth Sciences*, **146**, 296–303.
- Aguilera, O., Guimarães, J.T.F. & Moraes-Santos, H. (2013) Neogene Eastern Amazon carbonate platform and the palaeoenvironmental interpretation. *Swiss Journal of Palaeontology*, **132**, 99–118.

- Akgün, F., Akay, E. & Erdoğan, B. (2002) Tertiary terrestrial to shallow marine deposition in Central Anatolia: A palynological approach. *Turkish Journal of Earth Sciences*, **11**, 127–160.
- Akgün, F., Akkiraz, M.S., Üçbaş, S.D. & Bozcu, M. (2013) Oligocene vegetation and climate characteristics in north-west Turkey: data from the south-western part of the Thrace Basin. *Turkish Journal of Earth Sciences*, **22**, 277–303.
- Akgün, F., Kayseri, M.S. & Akkiraz, M.S. (2007) Palaeoclimatic evolution and vegetational changes during the Late Oligocene–Miocene period in Western and Central Anatolia (Turkey). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **253**, 56–90.
- Akkiraz, M.S. & Akgün, F. (2005) Palynology and age of the Early Oligocene units in Çardak–Tokça basin, southwest Anatolia: paleoecological implications. *Geobios*, **38**, 283–299.
- Akkiraz, M.S., Akgün, F. & Örcen, S. (2009) Palynomorph and foraminifer content of the Lower Miocene (Aquitanian) Kavak Formation outcropping in Burdur area. *Bulletin of the Turkish Association of Petroleum Geologists*, **21**, 31–53.
- Akkiraz, M.S., Akgün, F. & Örcen, S. (2010a) The Oligocene palynostratigraphy and paleoclimate of the İncesu Basin on the northern part of the western Taurids, Isparta Region, Turkey. *Abstract*, 1–2.
- Akkiraz, M.S., Akgün, F. & Örcen, S. (2010b) Palaeoecology of the Rupelian-early Chattian (Oligocene) aged Tokca Formation outcropping east of Civril (Denizli): Quantitative climatic correlations. *Turkiye Jeoloji Bulteni-Geological Bulletin of Turkey*, **53**, 63–95.
- Akkiraz, M.S., Akgün, F. & Örcen, S. (2011) Stratigraphy and palaeoenvironment of the Lower-“middle” Oligocene units in the northern part of the Western Taurides (İncesu area, Isparta, Turkey). *Journal of Asian Earth Sciences*, **40**, 452–474.
- Akkiraz, M.S., Akgün, F., Örcen, S., Bruch, A.A. & Mosbrugger, V. (2006) Stratigraphic and Palaeoenvironmental Significance of Bartonian–Priabonian (Middle–Late Eocene) Microfossils from the Başçeşme Formation, Denizli Province, Western Anatolia. *Turkish Journal of Earth Sciences*, **15**, 155–180.
- Akkiraz, M.S., Kayseri, M.S. & Akgün, F. (2008) Palaeoecology of coal-bearing Eocene sediments in Central Anatolia (Turkey) based on quantitative palynological data. *Turkish Journal of Earth Sciences*, **17**, 317–360.
- Akyol, E. (1971) Microflore de l’Oligocene Inférieur récoltée dans un sondage près d’Avcikoru, Sile-Istanbul. *Pollen et Spores*, **13**, 117–133.
- Alexová, V. (2011) *Palynological studies of some Middle Miocene localities of the Central Paratethys*. Master’s thesis, Masaryk University.
- Alizadeh, K. (2018) *Investigating Environmental (Climate and Vegetation) Change of Eastern Amazonia During Pleistocene and Holocene Using Multi-Proxy Analysis*. Ph.D. thesis, University of Göttingen.
- Alizadeh, K., Cohen, M. & Behling, H. (2015) Origin and dynamics of the northern South American coastal savanna belt during the Holocene—the role of climate, sea-level, fire and humans. *Quaternary Science Reviews*, **122**, 51–62.
- Alizadeh, K., Matthias, I., Rodríguez-Zorro, P.A., Hermanowski, B., da Costa, M.L. & Behling, H. (2017) Forest-savanna boundary shift on the plateau of Serra Sul dos Carajás (southeastern Amazonia) since the mid-Holocene; driving forces and limiting factors. *Quaternary International*, **449**, 12–21.
- Allen, L.O. & Meyer, K.J. (1988) Great Britain; the London Basin and East Anglia. *Geologisches Jahrbuch Reihe A*, **100**, 296–297.

- Alley, N.F. & Clarke, J.D.A. (1992) Stratigraphy and palynology of Mesozoic sediments from the Great Australian Bight area, southern Australia. *BMR journal of Australian geology and geophysics*, **13**, 113–129.
- Anderson, J.A.R. & Muller, J. (1975) Palynological study of a Holocene peat and a Miocene coal deposit from NW Borneo. *Review of Palaeobotany and Palynology*, **19**, 291–351.
- Annan, W. (2016) *Palynological Analysis Of The St-5 And St-6 Wells, Offshore Tano Basin, South Western Ghana: Implications For Stratigraphy And Paleoenvironment*. Ph.D. thesis, University of Ghana.
- Antoine, P.O., Abello, M.A., Adnet, S., Sierra, A.J.A., Baby, P., Billet, G., Boivin, M., Calderón, Y., Candela, A., Chabain, J. *et al.* (2016) A 60-million-year Cenozoic history of western Amazonian ecosystems in Contamana, eastern Peru. *Gondwana Research*, **31**, 30–59.
- Apaalse, L.A. & Atta-Peters, D. (2013) Cretaceous-Palaeogene palynology of the Keta-1 well offshore Keta Basin, southeastern Ghana. *World Applied Sciences Journal*, **23**, 1576–1583.
- Archangelsky, S. (1973a) Paleocene and Eocene pollen assemblages from Patagonia, Argentina. *The palynology of the cenophyte. Proc. III Intern. Palynol. Conf*, pp. 96–100.
- Archangelsky, S. (1973b) Palinología del Paleoceno de Chubut. I. Descripciones sistemáticas. *Ameghiniana*, **10**, 339–399.
- Archangelsky, S. & Zamalao, M.D.C. (1986) New palynological descriptions from the Salamanca and Bororo Formations, Paleocene Chubut Province, (Argentine Republic). *Ameghiniana*, **23**, 35–46.
- Asadu, A.N. & Omigie, J.I. (2015) Miospore Biostratigraphy of Oligocene–Lower Miocene Sediments in Well X, Deep Offshore Niger Delta. *International Journal of Science and Research*, **6**, ART20173249.
- Atta-Peters, D. & Salami, M.B. (2004) Late Cretaceous to early tertiary pollen grains from offshore Tano basin, Southwestern Ghana. *Revista española de micropaleontología*, **36**, 451–465.
- Auffret, J.P. & Gruas-Cavagnetto, C. (1975) Les formations paleogenes sous-marines de la Manche orientale; donnees palynologiques. *Bulletin de la Société géologique de France*, **7**, 641–655.
- Ayinla, H.A., Adeigbe, O.C., Ola-Buraimo, A.O., Bankole, S.A. & Adebowale, M. (2014) Palynostratigraphy and High Resolution Paleoenvironmental Reconstruction of Part of Kemar-1 Well, Bornu Basin, Northeastern Nigeria. *The Pacific Journal of Science and Technology*, **15**, 392–403.
- Ayinla, H.A., Ola-Buraimo, A.O., Adeigbe, O.C., Bankole, S.A. & Adebowale, M. (2013) Biostratigraphy and high resolution paleoenvironmental reconstruction of part of Kemar-1 well, Bornu Basin, Northeastern Nigeria. *Journal of Research in Environmental Science and Toxicology*, **2**, 53–63.
- Azema, C., Ollivier-Pierre, M.F. & Viaud, J.M. (1978) La palynologie Permet d'Attribuer a l'Ypresien la flore contenue dans les argiles noires du Loreau, en Soullans (Vendee). *Comptes Rendus de l'Academie des Sciences de Paris, Serie D*, **286**, 1141–1143.
- Baillie, P.W., Bacon, C.A. & Morgan, R. (1986) Geological observations on the Macquarie Harbour Beds at Coal Head, western Tasmania. *Unpublished Report Department of Mines Tasmania*, **73**.
- Baker, W.J., Coode, M.J.E., Dransfield, J., Dransfield, S., Harley, M.M., Hoffmann, P. & Johns, R.J. (1998) Patterns of distribution of Malesian vascular plants. R. Hall & J.D. Hallaway, eds., *Biogeography and geological evolution of SE Asia*, Backhuys Publishers Leiden, pp. 243–258.
- Baksi, S.K. & Deb, U. (1980a) Microfloristics and the Cretaceous-Tertiary boundary in the Bengal Basin, India. *International Palynological Conference, 5th. Cambridge, England, Abstracts*, p. 24.

- Baksi, S.K. & Deb, U. (1980b) Palynostratigraphic zonation of the Upper Cretaceous-Paleogene sequence of Bengal Basin. *Geophytology*, **10**, 199–224.
- Ballesteros, T., Montoya, E., Vegas-Vilarrúbia, T., Giralt, S., Abbott, M.B. & Rull, V. (2014) An 8700-year record of the interplay of environmental and human drivers in the development of the southern Gran Sabana landscape, SE Venezuela. *The Holocene*, **24**, 1757–1770.
- Banerjee, S. & Nandi, B. (1994) Palynology of the Middle Formation near Kolasib, northern Mizor India. *Geophytology*, **23**, 215–220.
- Bankole, S.I., Schrank, E. & Osterloff, P.L. (2014) Palynostratigraphy, palaeoclimates and palaeodepositional environments of the Miocene aged Agbada Formation in the Niger Delta, Nigeria. *Journal of African Earth Sciences*, **95**, 41–62.
- Baraniecka, M.D. (1981) Osady trzeciorzędowe SW części niecki warszawskiej na przykładzie profilu z Kaczorówka. *Geological Quarterly*, **25**, 365–386.
- Barberi, M. (2001) *Mudanças paleoambientais na região dos cerrados do Planalto Central durante o Quaternário tardio: o estudo da Lagoa Bonita, DF*. Ph.D. thesis, Universidade de São Paulo.
- Barberi, M., Salgado-Labouriau, M.L. & Suguio, K. (2000) Paleovegetation and paleoclimate of “Vereda de Águas Emendadas”, central Brazil. *Journal of South American Earth Sciences*, **13**, 241–254.
- Barbin, V. (1992) Paleoenvironment of the Brendola area (Late Priabonian stage stratotype, northern Italy) during the Eocene-Oligocene transition. *Neues Jahrbuch für Geologie und Paläontologie-Monatshefte*, 221–230.
- Barui, N.C. & Chanda, S. (1992) Late-Quaternary pollen analysis in relation to palaeoecology, biostratigraphy and dating of Calcutta peat. *Proceedings of the Indian National Science Academy. Part B Biological sciences*, **58**, 191–200.
- Basistová, P. & Doláková, N. (2016) Palynologické zhodnocení sediment z vrtu Iván IK-1 u Hustopecí. *Geologické výzkumy na Morave a ve Slezsku*, **18**.
- Bati, Z. & Sancay, R.H. (2007) Palynostratigraphy of Rupelian sediments in the Mus Basin, Eastern Anatolia, Turkey. *Micropaleontology*, **53**, 249–283.
- Bayona, G.A., Castillo, O.C.M., Molina, A.C., Jaramillo, C., Lamus, F., Moron, S.E., Quiroz, L., Ruiz, M.C., Valencia, V., Parra, M. *et al.* (2010) Estratigrafía, procedencia, subsidencia y exhumación de las unidades Paleógenas en el Sinclinal de Usme, sur de la zona axial de la Cordillera Oriental. *Geología Colombiana*, **35**, 5–35.
- Bechtel, A., Markic, M., Sachsenhofer, R.F., Jelen, B., Gratzer, R., Lücke, A. & Püttmann, W. (2004) Paleoenvironment of the upper Oligocene Trbovlje coal seam (Slovenia). *International Journal of Coal Geology*, **57**, 23–48.
- Behling, H. (2001) Late Quaternary environmental changes in the Lagoa da Curuçã region (eastern Amazonia, Brazil) and evidence of Podocarpus in the Amazon lowland. *Vegetation History and Archaeobotany*, **10**, 175–183.
- Behling, H. (2002a) Impact of the Holocene sea-level changes in coastal, eastern and central Amazonia. *Amazoniana: Limnologia et Oecologia Regionalis Systematis Fluminis Amazonas*, **17**, 41–52.
- Behling, H. (2002b) Late Quaternary vegetation and climate dynamics in southeastern Amazonia inferred from Lagoa da Confusao in Tocantins State, northern Brazil. *Amazoniana: Limnologia et Oecologia Regionalis Systematis Fluminis Amazonas*, **17**, 27–39.

- Behling, H., Berrio, J.C. & Hooghiemstra, H. (1999) Late Quaternary pollen records from the middle Caquetá river basin in central Colombian Amazon. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **145**, 193–213.
- Behling, H., Cohen, M.C.L. & Lara, R.J. (2004) Late Holocene mangrove dynamics of Marajo island in Amazonia, northern Brazil. *Vegetation History and Archaeobotany*, **13**, 73–80.
- Behling, H. & da Costa, M.L. (2000) Holocene environmental changes from the Rio Curuá record in the Caxiuanã region, eastern Amazon Basin. *Quaternary Research*, **53**, 369–377.
- Behling, H. & da Costa, M.L. (2001) Holocene vegetational and coastal environmental changes from the Lago Crispim record in northeastern Pará State, eastern Amazonia. *Review of Palaeobotany and Palynology*, **114**, 145–155.
- Behling, H. & Hooghiemstra, H. (1998) Late Quaternary palaeoecology and palaeoclimatology from pollen records of the savannas of the Llanos Orientales in Colombia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **139**, 251–267.
- Behling, H. & Hooghiemstra, H. (1999) Environmental history of the Colombian savannas of the Llanos Orientales since the Last Glacial Maximum from lake records El Pinal and Carimagua. *Journal of Paleolimnology*, **21**, 461–476.
- Behling, H. & Hooghiemstra, H. (2000) Holocene Amazon rainforest-savanna dynamics and climatic implications: high-resolution pollen record from Laguna Loma Linda in eastern Colombia. *Journal of Quaternary Science*, **15**, 687–695.
- Behling, H., Keim, G., Irion, G., Junk, W. & De Mello, J.N. (2001) Holocene environmental changes in the central Amazon Basin inferred from Lago Calado (Brazil). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **173**, 87–101.
- Bermúdez, M.A., Hoorn, C., Bernet, M., Carrillo, E., van Der Beek, P.A., Garver, J.I., Mora, J.L. & Mehrkian, K. (2017) The detrital record of late-Miocene to Pliocene surface uplift and exhumation of the Venezuelan Andes in the Maracaibo and Barinas foreland basins. *Basin Research*, **29**, 370–395.
- Berrio, J.C., Arbeláez, M.V., Duivenvoorden, J.F., Cleef, A.M. & Hooghiemstra, H. (2003) Pollen representation and successional vegetation change on the sandstone plateau of Araracuara, Colombian Amazonia. *Review of Palaeobotany and Palynology*, **126**, 163–181.
- Berrio, J.C., Hooghiemstra, H., Behling, H. & van der Borg, K. (2000) Late Holocene history of savanna gallery forest from Carimagua area, Colombia. *Review of Palaeobotany and Palynology*, **111**, 295–308.
- Berrio Mogollon, J.C. (2002) *Lateglacial and Holocene vegetation and climate change in lowland Colombia*. Ph.D. thesis, University of Amsterdam.
- Berry, C.N., Misra, C.M. & Chopra, A.S. (1996) Palynostratigraphy, paleoenvironment of Changartalai well in Punjab Basin. *15th Indian Colloquium Micropalaeontology and Stratigraphy, Dehradun*, pp. 643–650.
- Bertassoli Jr, D.J., Sawakuchi, A.O., Sawakuchi, H.O., Pupim, F.N., Hartmann, G.A., McGlue, M.M., Chiessi, C.M., Zabel, M., Schefuß, E., Pereira, T.S. *et al.* (2017) The fate of carbon in sediments of the Xingu and Tapajós clearwater rivers, Eastern Amazon. *Frontiers in Marine Science*, **4**, 44.
- Bhattacharya, K. & Chanda, S. (1992) Late Quaternary vegetational history of Upper Assam, India. *Review of palaeobotany and palynology*, **72**, 325–333.
- Biaggi, R.E. & Fisk, L.H. (1984) Palynology of the Miocene Latah Formation, Northeastern Washington. *Palynology*, **8**, 253–254.

- Biagolini, C.H., Bernardes-de Oliveira, M.E.C. & Caramês, A.G. (2013) Itaquaquetuba Formation, São Paulo basin, Brazil: new angiosperm components of Paleogene taphoflora. *Brazilian Journal of Geology*, **43**, 639–652.
- Bié, G.R., Digbehi, Z.B., Yao, K.R., Tea-Yassi, J., Kangah, K.D. & Tahi, I. (2012) Stratigraphie Palynologique du Maastrichtien Supérieur-Eocène Supérieur du Bassin Sédimentaire Offshore de Côte d’Ivoire, Afrique de l’Ouest. *International Journal of African Studies*, **6**, 40–57.
- Bigwood, A.J. & Hill, R.S. (1985) Tertiary araucarian macrofossils from Tasmania. *Australian journal of botany*, **33**, 645–656.
- Bio-Lokoto, A., Carbonnel, G., de Klasz, I., de Klasz, S., Lang, J. & Salard-Cheboldaeff, M. (1998) Données nouvelles biochronologiques et paléoécologiques sur le bassin sédimentaire côtier du Bénin entre le Crétacé supérieur et l’Éocène basal (Afrique de l’ouest). *Revue de micropaléontologie*, **41**, 29–57.
- Bogotá-Ángel, G., Huang, H., Jardine, P.E., Chazot, N., Salamanca, S., Banks, H., Pardo-Trujillo, A., Plata, A., Dueñas, H., Star, W. *et al.* (2021) Climate and geological change as drivers of Mauritiinae palm biogeography. *Journal of Biogeography*, **48**, 1001–1022.
- Bolaji, T.A., Ndukwe, O.S., Oyebamiji, A.R. & Ikegwuonu, O.N. (2020) palynological Age control and paleoenvironments of the paleogene Strata in eastern Dahomey Basin, Southwestern Nigeria. *Scientific Reports*, **10**, 1–11.
- Boonstra, M., Ramos, M.I.F., Lammertsma, E.I., Antoine, P.O. & Hoorn, C. (2015) Marine connections of Amazonia: Evidence from foraminifera and dinoflagellate cysts (early to middle Miocene, Colombia/Peru). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **417**, 176–194.
- Boreham, C.J., Blevin, J.E., Duddy, I., Newman, J., Liu, K., Middleton, H.M., Macphail, M.K. & Cook, A.C. (2002) Exploring the potential for oil generation, migration and accumulation in Cape Sorell–1, Sorell Basin, offshore West Tasmania. *The APPEA Journal*, **42**, 405–435.
- Borghain, B., Kalita, K.D., Bezbaruah, D. & Dutta, S. (2013a) Sequence-stratigraphy and palynostratigraphy of Eocene sediments in Bhogpara area, Dibrugarh district, upper Assam shelf basin, India. *South East Asian Journal of Sedimentary Basin Research*, 9–27.
- Borghain, B., Kalita, K.D., Bezbaruah, D. & Dutta, S. (2013b) Sequence-stratigraphy and palynostratigraphy of Eocene sediments in Bhogpara area, Dibrugarh district, upper Assam shelf basin, India. *South East Asian Journal of Sedimentary Basin Research*, **1**, 19–27.
- Borne, V., Chevalier, M. & Olivier-Pierre, M.F. (1989) Les premiers dépôts paléogènes (yprésiens) de la bordure méridionale du Massif armoricain; aspects sédimentologique, tectonique et paléogéographique. *Géologie de la France*, 1–2.
- Borne, V., Margerel, J.P. & Olivier-Pierre, M.F. (1991) L’évolution des paleoenvironnements au Paleogene dans l’Ouest de la France; le bassin de Saffre-Nord-sur-Erdre (Loire-Atlantique, France). *Bulletin de la Société géologique de France*, **162**, 739–751.
- Boudouresque, L. (1980) *Contribution de la paléopalynologie à la reconstitution floristique, stratigraphique et paléogéographique de la bordure occidentale du bassin des Iullemmeden au Crétacé supérieur et au Paléogène (Niger et Mali-Afrique de l’Ouest)*. Ph.D. thesis, University of Orléans et Niamey.
- Boulter, M.C. (1986) Pollen and spore events from the marine Tertiary of North Europe. *Journal of Micropalaeontology*, **5**, 75–84.
- Boulter, M.C. & Craig, D.L. (1979) A middle Oligocene pollen and spore assemblage from the Bristol Channel. *Review of Palaeobotany and Palynology*, **28**, 259–272.

- Boulter, M.C. & Hubbard, R.N.L.B. (1982) Objective paleoecological and biostratigraphic interpretation of Tertiary palynological data by multivariate statistical analysis. *Palynology*, **6**, 55–68.
- Boulter, M.C. & Wilkinson, G.C. (1977) A system of group names for some Tertiary pollen. *Palaeontology*, **20**, 559–579.
- Brelie, G.V.D., Doebel, F., Geissert, F., Weiler, H. & Weiler, W. (1973) Ein Aufschluß im unteren Bereich der Hydrobien-Schichten (Aquitain) beim Bau der Gruppen-Kläranlage “Queichtal”, Godramstein bei Landau (Pfalz). *Oberrheinische Geologische Abhandlungen*, **22**, 13–44.
- Brugger, S.O., Gobet, E., van Leeuwen, J.F., Ledru, M.P., Colombaroli, D., van der Knaap, W.O., Lombardo, U., Escobar-Torrez, K., Finsinger, W., Rodrigues, L. *et al.* (2016) Long-term man–environment interactions in the Bolivian Amazon: 8000 years of vegetation dynamics. *Quaternary Science Reviews*, **132**, 114–128.
- Bruno, D.Z., Elysée, G.K., Nrsquo, Y., Kouadio, A., Kizito, T.B.K.K., Raphael, Y.K., Ignace, T. *et al.* (2011) Palynostratigraphy and depositional palaeoenvironment of Cretaceous-Palaeogene (K-Pg) boundary deposits on Abidjan margin (Côte d’Ivoire). *Journal of Geography and Regional Planning*, **4**, 644–655.
- Burbridge, R.E., Mayle, F.E. & Killeen, T.J. (2004) Fifty-thousand-year vegetation and climate history of Noel Kempff Mercado National park, Bolivian Amazon. *Quaternary Research*, **61**, 215–230.
- Burn, M.J., Mayle, F.E. & Killeen, T.J. (2010) Pollen-based differentiation of Amazonian rainforest communities and implications for lowland palaeoecology in tropical South America. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **295**, 1–18.
- Bush, M.B. & Colinvaux, P.A. (1988) A 7000-year pollen record from the Amazon lowlands, Ecuador. *Vegetatio*, **76**, 141–154.
- Bush, M.B. & Colinvaux, P.A. (1990) A pollen record of a complete glacial cycle from lowland Panama. *Journal of Vegetation Science*, **1**, 105–118.
- Bush, M.B., Correa-Metrio, A., McMichael, C.H., Sully, S., Shadik, C.R., Valencia, B.G., Guilderson, T., Steinitz-Kannan, M. & Overpeck, J.T. (2016) A 6900-year history of landscape modification by humans in lowland Amazonia. *Quaternary Science Reviews*, **141**, 52–64.
- Bush, M.B., De Oliveira, P.E., Colinvaux, P.A., Miller, M.C. & Moreno, J.E. (2004) Amazonian paleoecological histories: one hill, three watersheds. *Palaeogeography, palaeoclimatology, palaeoecology*, **214**, 359–393.
- Bush, M.B., Miller, M.C., De Oliveira, P.E. & Colinvaux, P.A. (2000) Two histories of environmental change and human disturbance in eastern lowland Amazonia. *The Holocene*, **10**, 543–553.
- Bush, M.B., Silman, M.R. & Listopad, C.M.C.S. (2007) A regional study of Holocene climate change and human occupation in Peruvian Amazonia. *Journal of Biogeography*, **34**, 1342–1356.
- Bustin, R.M. & Chonchawalit, A. (1995) Formation and tectonic evolution of the Pattani Basin, Gulf of Thailand. *International Geology Review*, **37**, 866–892.
- Caputo, M.V. (2011) Discussão sobre a Formação Alter do Chão e o Alto de Monte Alegre. *Contribuição à Geologia da Amazônia, Manaus, SBG/Núcleo Norte*, **7**, 7–23.
- Caratini, C. (1975) Palynologie des deux formations detritiques eocenes dans le Sud du Bordelais (Gironde); mise en evidence d’une vegetation chaude et humide mais non typiquement tropicale. *Bulletin de la Société Géologique de France*, **7**, 797–802.
- Caratini, C., Tissot, C., Kar, R.K., Venkatachala, B.S. & Sarr, R. (1991) Paleocene palynoflora from Walalane borehole, Senegal. *Symposium on African palynology*, pp. 123–133.

- Carpenter, R.J., Jordan, G.J., Macphail, M.K. & Hill, R.S. (2012) Near-tropical Early Eocene terrestrial temperatures at the Australo-Antarctic margin, western Tasmania. *Geology*, **40**, 267–270.
- Cassino, R.F., Martinho, C.T. & da Silva Caminha, S.A.F. (2018) A Late Quaternary palynological record of a palm swamp in the Cerrado of central Brazil interpreted using modern analog data. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **490**, 1–16.
- Castro, D.F., Oliveira, P.E., Rossetti, D.F. & Pessenda, L.C.R. (2013) Late Quaternary landscape evolution of northeastern Amazonia from pollen and diatom records. *Anais da Academia Brasileira de Ciências*, **85**, 35–55.
- Cavagnetto, C. & Anadón, P. (1995) Une mangrove complexe dans le Bartonien du Bassin de l'Ebre (NE de l'Espagne). *Palaeontographica Abteilung B*, 147–165.
- Çelik, Y., Karayığit, A.İ., Querol, X., Oskay, R.G., Mastalerz, M. & Özer, M.S.K. (2017) Coal characteristics, palynology, and palaeoenvironmental interpretation of the Yeniköy coal of Late Oligocene age in the Thrace Basin (NW Turkey). *International Journal of Coal Geology*, **181**, 103–123.
- Chakraborty, M. (2004) Palynology of the Lakadong Sandstone (Late Palaeocene) exposed around Bhalukurung, North Cachar Hills, Assam. *Paleobotanist*, **53**, 1–4.
- Châteauneuf, J.J. (1980) Palynostratigraphie et paléoclimatologie de l'Eocène supérieur et de l'Oligocène du Bassin de Paris (France). *Mémoires du Bureau de recherches géologiques et minières*, **116**, 1–360.
- Chiaghanam, O.I., Chiadikobi, K.C., Ikegwuonu, O.N., Omoboriowo, A.O., Onyemesili, O.C. & Yikarebogha, Y. (2014) Sedimentology and Sequence Stratigraphy of the Eocene Nanka Formation (Ameki Group): An Evaluation of Ogbunike Reference Locality in Anambra Basin, South-Eastern Nigeria. *IOSR Journal of Applied Geology and Geophysics*, **2**, 01–10.
- Chiaghanam, O.I., Chiadikobi, K.C., Oguanya, C.E., Ikegwuonu, O.N. & Nwokeabia, C.N. (2017) Palynological and Paleoenvironmental Study of Paleogene in Bende–Umuahia, Niger Delta Basin, Nigeria. *Journal of Environment and Earth Science*, **7**, 1–11.
- Christopher, R.A., Goodman, D.K., Jansonius, J. & McGregor, D.C. (1996) Introduction to biostratigraphy and time scales. *Palynology: Principles and Applications. American Association of Stratigraphic Palynologists Foundation*, **2**, 463–492.
- Chukwura, E., Onuigbo, E.N., Okoro, A.U., Okafor, P.N. & Okolo, C.M. (2017) Palynological evaluation of the Imo Formation at Isiagu, Niger Delta Basin, Nigeria: age and paleoenvironmental significance. *Journal of Basic Physical Research*, **7**, 6–16.
- Ciuk, E. (1979) The Cainozoic Formations in the Wieruszow Area (Kalisz Voivodship). *Biuletyn - Instytutu Geologiczny*, **320**, 5–25.
- Cohen, M.C.L., Lara, R.J., Smith, C.B., Angélica, R.S., Dias, B.S. & Pequeno, T. (2008) Wetland dynamics of Marajó Island, northern Brazil, during the last 1000 years. *Catena*, **76**, 70–77.
- Colinvaux, P.A., de Oliveira, P.E., Moreno, J.E., Miller, M.C. & Bush, M.B. (1996) A long pollen record from lowland Amazonia: forest and cooling in glacial times. *Science*, **274**, 85–88.
- Colinvaux, P.A., Frost, M., Frost, I., Liu, K.B. & Steinitz-Kannan, M. (1988) Three pollen diagrams of forest disturbance in the western Amazon basin. *Review of Palaeobotany and Palynology*, **55**, 73–81.
- Collinson, M.E., Fowler, K. & Boulter, M.C. (1981) Floristic changes indicate a cooling climate in the Eocene of southern England. *Nature*, **291**, 315–317.

- Collinson, M.E., Knobloch, E. & Kvaček, Z. (1989) Vegetational change during the Palaeogene in the coastal wetlands of southern England. *Proceedings of the Symposium "Paleofloristic and Paleoclimatic Changes in the Cretaceous and Tertiary*, pp. 135–139.
- Colmenares, O.A. (1988) A palynological study of samples from three wells of the Boscan Field, Venezuela. *Revista Técnica Intevep*, **8**, 83–97.
- Colmenares, O.A. & Teran, L. (1993) A biostratigraphic study of Paleogene sequences in southwestern Venezuela. *Palynology*, **17**, 67–89.
- Contreras, L., Pross, J., Bijl, P., O'Hara, R.B., Raine, J.I., Sluijs, A. & Brinkhuis, H. (2014) Southern high-latitude terrestrial climate change during the Palaeocene–Eocene derived from a marine pollen record (ODP Site 1172, East Tasman Plateau). *Climate of the Past*, **10**, 1401–1420.
- Cornou, M.E., Quattrocchio, M.E. & Martínez, M.A. (2014) Palinoestratigrafía de la Formación Salto del Macho, Paleógeno de la Cuenca de Ñirihuau, Argentina. *Ameghiniana*, **51**, 556–571.
- Corredor, F. (2003) Eastward extent of the late Eocene-early Oligocene onset of deformation across the northern Andes: constraints from the northern portion of the Eastern Cordillera fold belt, Colombia, and implications for regional oil exploration. *Journal of South American Earth Sciences*, **16**, 445–457.
- Cripps, J.A., Widdowson, M., Spicer, R.A. & Jolley, D.W. (2005) Coastal ecosystem responses to late stage Deccan Trap volcanism: the post K–T boundary (Danian) palynofacies of Mumbai (Bombay), west India. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **216**, 303–332.
- Crouch, E.M. & Brinkhuis, H. (2005) Environmental change across the Paleocene–Eocene transition from eastern New Zealand: a marine palynological approach. *Marine Micropaleontology*, **56**, 138–160.
- Curry, D., Adams, C., Boulter, M.C., Dille, F.C., Eames, F.E., Funnell, B.M. & Wells, M.K. (1978) A correlation of Tertiary rocks in the British Isles. *Geological Society, Special Report*, **12**, 25–39.
- da Silva-Caminha, S.A.F., Jaramillo, C.A. & Absy, M.L. (2010) Neogene palynology of the Solimões basin, Brazilian Amazonia. *Palaeontographica Abteilung B*, **284**, 13–79.
- da Silva Meneses, M.E.N., da Costa, M.L. & Behling, H. (2013) Late Holocene vegetation and fire dynamics from a savanna-forest ecotone in Roraima state, northern Brazilian Amazon. *Journal of South American Earth Sciences*, **42**, 17–26.
- da Silveira, R.R. & Souza, P.A. (2017) Palinoestratigrafia da Formação Solimões na Região do Alto Solimões (Atalaia do Norte e Tabatinga), Amazonas, Brasil. *Geociências (São Paulo)*, **36**, 100–117.
- D'Apolito, C. (2016) *Landscape evolution in Western Amazonia: palynostratigraphy, palaeoenvironments and diversity of the miocene Solimões formation, Brazil*. Ph.D. thesis, University of Birmingham.
- D'Apolito, C., Absy, M.L. & Latrubesse, E.M. (2013) The Hill of Six Lakes revisited: new data and re-evaluation of a key Pleistocene Amazon site. *Quaternary Science Reviews*, **76**, 140–155.
- D'Apolito, C., da Silva-Caminha, S.A.F., Jaramillo, C., Dino, R. & Soares, E.A.A. (2019) The Pliocene–Pleistocene palynology of the Negro River, Brazil. *Palynology*, **43**, 223–243.
- De Coninck, J. (1972) Application stratigraphique des microfossiles organiques dans l'Yprésien du bassin belge. *Bulletin de la Société belge de géologie, de paléontologie et d'hydrologie*, **81**.
- De Coninck, J. (1975) Microfossiles à paroi organique de l'Yprésien du Bassin belge. *Professional Paper. Geological Survey of Belgium*.

- De Coninck, J. (1977) Organic walled microfossils from the Eocene of the Woensdrecht borehole, southern Netherlands. *Mededelingen Rijks Geologische Dienst, Nieuwe Serie*, **28**, 33–64.
- De Coninck, J. (1981) Espèces indicatrices de microfossiles à paroi organique des dépôts de l'Yprésien supérieur et du Lutétien dans le sondage de Kallo. *Bulletin de la Société belge de Géologie*, **89**, 309–317.
- De Coninck, J. (1985) Microfossiles à paroi organique dans les sables de Lede (Eocène Moyen) du sondage de Mol (Belgique). *Bulletin de la Société belge de Géologie*, **94**, 65–78.
- De Coninck, J. & Nolf, D. (1979) Note sur les couches de base de la Formation du Panisel entre Torhout et Tielt. *Bulletin de la Société belge de géologie*, **87**, 171–178.
- De Klasz, I. & Jan Du Chêne, R. (1978) Presence of Albian-Cenomanian in southwestern Nigeria and its paleogeographic implications. *Compte Rendu de la Société de Physique et d'Histoire Naturelle de Genève*, **13**, 10–15.
- de la Parra, F. (2009) *Palynological changes across the Cretaceous-Tertiary boundary in Colombia, South America*. Master's thesis, University of Florida.
- de Lima, M.R. & Salard-Chebouldaef, M. (1981) Palynologie des bassins de Gandarela et Fonseca (Eocène de l'état de Minas Gerais, Brésil). *Boletim IG*, **12**, 33–53.
- de Oliveira, P.E., Barreto, A.M.F. & Suguio, K. (1999) Late Pleistocene/Holocene climatic and vegetational history of the Brazilian caatinga: the fossil dunes of the middle São Francisco River. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **152**, 319–337.
- de Toledo, M.B. (2004) *Holocene vegetation and climate history of savanna-forest ecotones in northeastern Amazonia*. Ph.D. thesis, Florida Institute of Technology.
- de Toledo, M.B. & Bush, M.B. (2007) A mid-Holocene environmental change in Amazonian savannas. *Journal of Biogeography*, **34**, 1313–1326.
- de Toledo, M.B. & Bush, M.B. (2008a) A Holocene pollen record of savanna establishment in coastal Amapá. *Anais da Academia Brasileira de Ciências*, **80**, 341–351.
- de Toledo, M.B. & Bush, M.B. (2008b) Vegetation and hydrology changes in Eastern Amazonia inferred from a pollen record. *Anais da Academia Brasileira de Ciências*, **80**, 191–203.
- Demirtaş, F., Bozcu, M., Koşun, E. & Akkiraz, M.S. (2015) Petrography and palynology of Late Oligocene and Middle Miocene coals in the Gelibolu peninsula, NW Turkey. *Turkish Journal of Earth Sciences*, **24**, 383–397.
- Diederix, D.O.J., de Haan, R. & Tichler, W.O. (1972) Organic microfossils. *Deep Sea Drilling Project, Washington, Initial Reports*, **14**, 257–257.
- Dino, R., Soares, E.A.A., Antonioli, L., Riccomini, C. & Nogueira, A.C.R. (2012) Palynostratigraphy and sedimentary facies of Middle Miocene fluvial deposits of the Amazonas Basin, Brazil. *Journal of South American Earth Sciences*, **34**, 61–80.
- Do, B., Nguyen, Q.A., H., N.Q., Ngo, X.V., Nguyen, T.H., Do, V.H. & Nguyen, T.H. (2002) Stratigraphy of Tertiary sediments in the Southwest Vietnam Continental Shelf. *Journal of Geology (Vietnam)*, **273**, 1–12.
- Do Campo, M., Del Papa, C., Jiménez-Millán, J. & Nieto, F. (2007) Clay mineral assemblages and analcime formation in a Palaeogene fluvial-lacustrine sequence (Maíz Gordo Formation Palaeogen) from northwestern Argentina. *Sedimentary Geology*, **201**, 56–74.

- Do Carmo, F., Barberi, M. & Rubin, J.C.R.D. (2003) Análise palinológica de sedimentos do Quaternário Tardio, a partir de 44.000 anos AP, na região centro-sul do Estado de Goiás, Brasil. *IX Congresso da Associação Brasileira de Estudos do Quaternário*, **1**.
- Dogra, N.N., Singh, R.Y. & Misra, P.S. (1985) Palynology of Dharmasala beds, Himachal Pradesh. *Journal of Palaeontological Society of India*, **30**, 63–77.
- Doubinger, J. (1973) Pollen and spores from the Paleocene coal basin of Cerrejon (Guajira Province, Colombia). *Proceedings, 96 Congres Nat. l des Soc. Savantes (Toulouse, 1971)*, vol. 5, pp. 253–262.
- Doubinger, J. (1976) Spore–pollen assemblages from the Tertiary of Colombia. *Actas 1997 Congres de National des Societé Savantes, Nantes*, vol. 4, pp. 9–18.
- Duan, W. & Lei, Z. (1984) Tertiary Stratigraphy of Zhujiangkou Basin. *Oil & Gas Geology*, **2**.
- Dueñas, H. (1979) Estudio palinológico del pozo QE-22 Oligoceno superior a Mioceno inferior, Planeta Rica, Norte de Colombia. *Boletín Geológico*, **22**, 95–115.
- Dueñas, H. (1980) Palynology of oligocene-Miocene strata of borehole QE-22, Planeta Rica, Northern Colombia. *Review of Palaeobotany and Palynology*, **30**, 313–328.
- Dueñas, H. (1983) Fluctuaciones del nivel del mar durante el depósito de los sedimentos basales de la Formación Ciénaga de Oro. *Revista de la Academia Colombiana de Ciencias Exactas, Fisicas y Naturales*, **XV**, **58**, 67–76.
- Dueñas, H., Navarrete, R.E., Mojica, J., Pardo, M. & Camargo, R. (2000) Edad de la Formación Vijes en el Pozo V3A, Oligoceno del piedemonte oriental de la Cordillera Occidental, Departamento del Valle del Cauca, Colombia. *Geología Colombiana*, **25**, 25–43.
- Dueñas-Jimenez, H. (1986) *Geologia y palinologia de la Formacion Cienaga de Ororegion Caribe Colombiana*. Instituto Nacioal de Investigaciones Geológico-Mineras.
- Dupuis, C. & Gruas-Cavagnetto, C. (1985) The Woolwich Beds et le London Clay de Newhaven (East Sussex), données palynologiques et stratigraphiques nouvelles. *Bulletin d'information des géologues du bassin de Paris*, **22**, 19–33.
- Durand, S. (1969) Observations nouvelles sur la présence du pollen de palmier *Nypa* dans l'Eocène de l'ouest de la France et du sud de l'Angleterre. *Bulletin de la Societie Geologique et Mineralogique de Bretagne*, **100**, 49–57.
- Durand, S. & Ollivier-Pierre, M.F. (1972) Geologic study of the Cassel (Nord) Boring. Palynologic analysis. *Bulletin d'Information des Geologues du Bassin de Paris*, **32**, 41–49.
- Durugbo, E.U. (2016) Palynostratigraphy, palynofacies and thermal maturation of the Nsukka Formation from an excavation site in Okigwe, southeastern Nigeria. *Palaeontologia africana*, **50**, 76–92.
- Durugbo, E.U. & Aroyewun, R.F. (2012) Palynology and paleoenvironments of the upper araromi Formation, Dahomey Basin, Nigeria. *Asian Journal of Earth Sciences*, **5**, 50.
- Dutta, S., Mathews, R.P., Singh, B.D., Tripathi, S.M., Singh, A., Saraswati, P.K., Banerjee, S. & Mann, U. (2011a) Petrology, palynology and organic geochemistry of Eocene lignite of Matanomadh, Kutch Basin, western India: Implications to depositional environment and hydrocarbon source potential. *International Journal of Coal Geology*, **85**, 91–102.
- Dutta, S., Tripathi, S.M., Mallick, M., Mathews, R.P., Greenwood, P.F., Rao, M.R. & Summons, R.E. (2011b) Eocene out-of-India dispersal of Asian dipterocarps. *Review of Palaeobotany and Palynology*, **166**, 63–68.

- Ediger, V. (1982) Paleo-environmental analysis of Kuleli-Babaeski Ridge (NW Thrace) and a new approach to the evaluation of hydrocarbon potential of Northern Thrace Basin. *TPAO Rep*, **427**, 1–194.
- Eifert, T.L. (2009) *The Cretaceous-Paleogene transition in the northern Mississippi Embayment, SE Missouri: palynology, micropaleontology, and evidence of a mega-tsunami deposit*. Ph.D. thesis, Missouri University of Science and Technology.
- Eisawi, A. & Schrank, E. (2008) Upper Cretaceous to Neogene palynology of the Melut Basin, southeast Sudan. *Palynology*, **32**, 101–129.
- Eisawi, A.A.M., Ibrahim, A.B., Rahim, O.B.A. & Schrank, E. (2012) Palynozonation of the Cretaceous to Lower Paleogene strata of the Muglad Basin, Sudan. *Palynology*, **36**, 191–207.
- El Beialy, S.Y. (1998) Stratigraphic and palaeoenvironmental significance of Eocene palynomorphs from the Rusayl Shale Formation, Al Khawd, northern Oman. *Review of Palaeobotany and Palynology*, **102**, 249–258.
- Elsik, W.C. & Yancey, T.E. (2000) Palynomorph biozones in the context of changing paleoclimate, middle Eocene to lower Oligocene of the Northwest Gulf of Mexico. *Palynology*, **24**, 177–186.
- Engel, E. (1987) Zur Palynologie und Stratigraphie der Braunkohle von Borken (Hessen). *Geol. Jb. Hessen*, **115**, 171–203.
- Engelhardt, D.W. & Wrenn, J.H. (1994) Palynology and paleoenvironments of Eocene sediments, Saw area, Salin Basin, Myanmar (Burma). *Palynology*, **18**, 243–244.
- Erikson, J.P., Kelley, S.A., Osmolovsky, P. & Verosub, K.L. (2012) Linked basin sedimentation and orogenic uplift: the Neogene Barinas basin sediments derived from the Venezuelan Andes. *Journal of South American Earth Sciences*, **39**, 138–156.
- Escobar, L.E. (1982) Estudio palinológico de la Formación Amagá. *Boletín de Ciencias de la Tierra*, 117–129.
- Evans, D., Morton, A.C., Wilson, S., Jolley, D. & Barreiro, B.A. (1997) Palaeoenvironmental significance of marine and terrestrial Tertiary sediments on the NW Scottish shelf in BGS borehole 77/7. *Scottish Journal of Geology*, **33**, 31–42.
- Fechner, G.G. (1988) Selected palynomorphs from the Lower to Middle Eocene of the South Atlas border zone (Morocco) and their environmental significance. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **65**, 73–79.
- Fernández Marrón, M.T., Fonollá Ocete, J.F. & Jiménez Rodrigo, J.C. (2002) Primeros datos paleobotánicos y palinológicos de una flora miocénica procedente de la unidad intermedia en el sector central de la Cuenca de Madrid. *Geogaceta*, **31**, 177–180.
- Ferraz-Vicentini, K.R. & Salgado-Labouriau, M.L. (1996) Palynological analysis of a palm swamp in central Brazil. *Journal of South American Earth Sciences*, **9**, 207–219.
- Field, J.H., Kealhofer, L., Cosgrove, R. & Coster, A.C.F. (2016) Human-environment dynamics during the Holocene in the Australian Wet Tropics of NE Queensland: A starch and phytolith study. *Journal of Anthropological Archaeology*, **44**, 216–234.
- Fitzgerald, J.A. (1999) *Pollen and spore assemblages from the oligocene Lough Neagh Group*. Ph.D. thesis, University of Sheffield.
- Fontes, D., Cordeiro, R.C., Martins, G.S., Behling, H., Turcq, B., Sifeddine, A., Seoane, J.C.S., Moreira, L.S. & Rodrigues, R.A. (2017) Paleoenvironmental dynamics in South Amazonia, Brazil, during the last 35,000 years inferred from pollen and geochemical records of Lago do Saci. *Quaternary Science Reviews*, **173**, 161–180.

- Foster, C.B. (1979) Tertiary miospores from the Rundle oil shale deposit. *Queensland Government Mining Journal*, **80**, 442–443.
- Fournier, G.R. (1982) Palynostratigraphic analysis of cores from site 493, Deep Sea Drilling Project leg 66. *Deep Sea Drilling Project, Washington, Initial Reports*, **66**, 661–670.
- Fowler, M. (1988) Palynological study of the Defense borehole (Paris). *Revue de Micropaleontologie*, **19**, 27–46.
- França, M.C., Francisquini, M.I., Cohen, M.C.L., Pessenda, L.C.R., Rossetti, D.F., Guimarães, J.T.F. & Smith, C.B. (2012) The last mangroves of Marajó Island—Eastern Amazon: impact of climate and/or relative sea-level changes. *Review of Palaeobotany and Palynology*, **187**, 50–65.
- Frederiksen, N.O. (1969) *Stratigraphy and palynology of the Jackson Stage (upper Eocene) and adjacent strata of Mississippi and Western Alabama*. Ph.D. thesis, University of Wisconsin.
- Frederiksen, N.O. (1980) Sporomorphs from the Jackson Group (Upper Eocene) and adjacent strata of Mississippi and western Alabama. *United States Geological Survey, Professional Paper*, **1084**, 1–75.
- Frederiksen, N.O. (1983) Middle Eocene palynomorphs from San Diego, California. Part II. Angiosperm pollen and miscellanea. *American Association Stratigraphic Palynological Contributions Series*, **12**, 32–155.
- Frederiksen, N.O. (1984) *Stratigraphic, paleoclimatic, and paleobiogeographic significance of Tertiary sporomorphs from Massachusetts*. US Government Printing Office.
- Frederiksen, N.O. (1989) Eocene sporomorph biostratigraphy of Southern California. *Palaeontographica Abteilung B*, **211**, 135–179.
- Frederiksen, N.O. (1994) Middle and late Paleocene angiosperm pollen from Pakistan. *Palynology*, **18**, 91–137.
- Frederiksen, N.O. (1998) Upper Paleocene and lowermost Eocene angiosperm pollen biostratigraphy of the eastern Gulf Coast and Virginia. *Micropaleontology*, **44**, 45–68.
- Frederiksen, N.O. (2001) Pollen biostratigraphy of Lower Tertiary sediments from five cores from Screven and Burke Counties, Georgia. *United States Geological Survey, Professional Paper*, **1603**, H1–H2.
- Friaes, Y.S. (2013) *Dinâmica da vegetação da região de humaitá-am durante o Pleistoceno tardio e o Holoceno*. Master's thesis, Instituto de Geociências.
- Frieling, J., Huurdeman, E.P., Rem, C., Donders, T.H., Pross, J., Bohaty, S.M., Holdgate, G.R., Gallagher, S.J., McGowran, B. & Bijl, P.K. (2018) Identification of the Paleocene–Eocene boundary in coastal strata in the Otway Basin, Victoria, Australia. *Journal of Micropalaeontology*, **37**, 317–339.
- Frost, I. (1988) A Holocene sedimentary record from Anangucocha in the Ecuadorian Amazon. *Ecology*, **69**, 66–73.
- Fullagar, R., Field, J., Denham, T. & Lentfer, C. (2006) Early and mid Holocene tool-use and processing of taro (*Colocasia esculenta*), yam (*Dioscorea* sp.) and other plants at Kuk Swamp in the highlands of Papua New Guinea. *Journal of Archaeological Science*, **33**, 595–614.
- Gee, C. & Westgate, J.W. (1990) Paleoecological reconstruction of a middle Eocene mangrove community from southwest Texas using palynology. *Palynology*, **14**, 212.
- Germeraad, J.H. & de Haan, R. (1964) Palynological contributions to the stratigraphy, palaeogeography and climatology of the uppermost Cretaceous, Tertiary and Quaternary of Venezuela. Tech. rep., Shell Internal Report, The Hague.
- Germeraad, J.H., Hopping, C.A. & Muller, J. (1968) Palynology of Tertiary sediments from tropical areas. *Review of palaeobotany and palynology*, **6**, 189–348.

- Ghosh, A.K. (1977) Palynology of the Siwaliks. *Advances in Pollen-Spore Research*, **2**, 14–19.
- Gold, D.P., Fenton, J.P.G., Casas-Gallego, M., Novak, V., Pérez-Rodríguez, I., Cetean, C., Price, R., Nembhard, N. & Thompson, H. (2018) The biostratigraphic record of Cretaceous to Paleogene tectono-eustatic relative sea-level change in Jamaica. *Journal of South American Earth Sciences*, **86**, 140–161.
- González, C., Urrego, L.E. & Martínez, J.I. (2006) Late Quaternary vegetation and climate change in the Panama Basin: Palynological evidence from marine cores ODP 677B and TR 163-38. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **234**, 62–80.
- Govindan, A. & Mallikarjuna, U.B. (2019) An Integrated Stratigraphic and Paleoenvironmental Studies from Narmada-Tapti Block, South Cambay Basin, Western India. *Journal of the Geological Society of India*, **94**, 171–178.
- Grabowska, I. (1972) Palinologiczne opracowanie profili trzeciorzędowych z obszaru woj. olsztyńskiego. *Kwartalnik Geologiczny (Instytut Geologiczny)*, **16**, 1009–1010.
- Grabowska, I. & Piwocki, M. (1975) The age and origin of the Toruń clays from the vicinity of Toruń defined on the basis of palynological and lithological observations. *Biuletyn Instytutu Geologicznego*, **284**, 41–70.
- Graham, A. (1993) Contribution toward a Tertiary palynostratigraphy for Jamaica: the status of Tertiary paleobotanical studies in northern Latin America and preliminary analysis of the Guys Hill Member (Chapelton Formation, middle Eocene) of Jamaica. *Biostratigraphy of Jamaica*, **182**, 443–461.
- Grover, R., Das, K.K., Murthy, M.S. & Singh, J. (2010) Biostratigraphy and Paleoenvironmental Interpretation Along Eastern Margin in Gamij Area of Cambay Basin, Gujarat, India. *Search and Discovery*, 1–12.
- Grover, R. & Poovendan, A. (1996) A note on depositional history of deeper sands of Linch-Jotana-Mewad-Sobhasan Area, Mehsana Block, Cambay Basin. *Contributions to XV Indian Colloquium on Micropaleontology and Stratigraphy*, pp. 801–804.
- Gruas, C. & Bignot, G. (1985) La transgression cuisienne en Haute-Normandie à Sotteville-sur-Mer (76-France) et le diachronisme des faciès sparnaciens. *Revue de paléobiologie*, **4**, 117–132.
- Gruas-Cavagnetto, C. (1967) Quelques nouvelles espèces sparnaciennes de pollens et spores. *Bulletin de la Société Géologique de France*, **7**, 57–62.
- Gruas-Cavagnetto, C. (1976a) *Étude palynologique du Paléogène du Sud de l'Angleterre*, vol. 1. Centre national de la recherche scientifique.
- Gruas-Cavagnetto, C. (1976b) Great Britain; the Hampshire Basin. *Geologisches Jahrbuch Reihe A*, **100**, 294–296.
- Gruas-Cavagnetto, C. (1978) Étude palynologique de L'Éocène du Bassin Anglo-Parisien. *Mémoires de la Société Géologique de France (Nouvelle Série)*, **131**, 1–64.
- Gruas-Cavagnetto, C. (1991) Pollens et dinophycées de l'Ilerdien moyen (Eocène inférieur) de Fordones (Corbières, France). *Cahiers de micropaléontologie*, **6**, 51–85.
- Gruas-Cavagnetto, C., Châteauneuf, J.J., Meyer, K.J. & Sittler, C. (1988) France; the Paris Basin. *Geologisches Jahrbuch Reihe A*, **100**, 297–299.
- Guédé, K.É., Slimani, H., Chekar, M., M'Hamdi, A., Mouah, R., Digbehi, B.Z. *et al.* (2019) Late Cretaceous to Early Eocene dinoflagellate cysts from the “12 frères” borehole, Fresco, southwestern Côte d'Ivoire: biostratigraphy and paleobiogeographic implication. *Journal of African Earth Sciences*, **150**, 744–756.

- Guimarães, J.T., Cohen, M.C., França, M.C., Lara, R.J. & Behling, H. (2010) Model of wetland development of the Amapá coast during the late Holocene. *Anais da Academia Brasileira de Ciências*, **82**, 451–465.
- Guimarães, J.T.F., Cohen, M.C.L., França, M.C., Alves, I.C.C., Smith, C.B., Pessenda, L.C.R. & Behling, H. (2013a) An integrated approach to relate Holocene climatic, hydrological, morphological and vegetation changes in the southeastern Amazon region. *Vegetation history and archaeobotany*, **22**, 185–198.
- Guimarães, J.T.F., Cohen, M.C.L., França, M.C., Friaes, Y.S., Pessenda, L.C.R. & Behling, H. (2011) Intepoxy records of channel dynamics and Holocene vegetation changes in the southeastern Amazon region. *XIII ABEQUA Congress - The South American Quaternary: Challenges and Perspectives*, 1–5.
- Guimarães, J.T.F., Cohen, M.C.L., Pessenda, L.C.R., França, M.C., Smith, C.B. & Nogueira, A.C.R. (2012) Mid-and late-Holocene sedimentary process and palaeovegetation changes near the mouth of the Amazon River. *The Holocene*, **22**, 359–370.
- Guimarães, J.T.F. & Nogueira, A.C.R. (2015) Palynology of the Middle Miocene—Pliocene Novo Remanso Formation, Central Amazonia, Brazil. *Ameghiniana*, **52**, 107–134.
- Guimarães, J.T.F., Nogueira, A.C.R., Da Silva Jr, J.B.C., Soares, J.L. & Silveira, R. (2013b) Fossil fungi from Miocene sedimentary rocks of the central and coastal Amazon region, North Brazil. *Journal of Paleontology*, **87**, 484–492.
- Guimaraes, J.T.F., Sahoo, P.K., Souza-Filho, P.W.M., Figueiredo, M.M.J.C.D., Reis, L.S., Silva, M.S. & Rodrigues, T.M. (2019) Holocene history of a lake filling and vegetation dynamics of the Serra Sul dos Carajás, southeast Amazonia. *Anais da Academia Brasileira de Ciências*, **91**.
- Gupta, A., Kotnala, S.K. & Poovenan, A. (1996a) Palynostratigraphy and palaeoenvironmental analysis of subsurface Paleogene sequence of Jambusar–Dabka area, south Cambay Basin. *Contribution to XV Indian Colloquim on Micropalaeontology and Stratigraphy*, 615–633.
- Gupta, A., Kotnala, S.K. & Poovendan, A. (1996b) Palynostratigraphy and paleoenvironment of Basal Clastic (Panna Formation) sequence in Mahim area, Bombay Offshore. *Contributions to XV Indian Colloquium on Micropaleontology and Stratigraphy A*, **15**, 615–633.
- Gupta, A., Kotnala, S.K. & Poovendan, A. (1996c) Palynostratigraphy and paleoenvironmental analysis of subsurface Paleogene sequence of Jambusar–Dabka area, South Cambay Basin. *Contributions to XV Indian Colloquium on Micropaleontology and Stratigraphy*, pp. 615–633.
- Guzmán, A.E.G. (1967) *A palynological study on the upper Los Cuervos and Mirador formations*. Brill Archive.
- Haag, N.A. (2019) *Palinostratigrafia Do Meso-Cenozoico Da Bacia Do Acre (Amazônia Sul Ocidental, Brasil)*. Ph.D. thesis, Universidade de Coimbra.
- Haberle, S. (1997) Upper Quaternary vegetation and climate history of the Amazon Basin: correlating marine and terrestrial pollen records. R.D. Flood, D.J.W. Piper, A. Klaus & L.C. Peterson, eds., *Proceedings of the ocean drilling program, scientific results*, vol. 155, Ocean Drilling Program, pp. 381–396.
- Haberle, S.G. & Maslin, M.A. (1999) Late Quaternary vegetation and climate change in the Amazon Basin based on a 50,000 year pollen record from the Amazon Fan, ODP Site 932. *Quaternary Research*, **51**, 27–38.
- Hait, A.K. & Banerjee, M. (1994) Palynology of lignite sediments from Mizoram, eastern India with remarks on age and environment of deposition. *Journal of Palynology*, **30**, 113–135.
- Handley, L., Crouch, E.M. & Pancost, R.D. (2011) A New Zealand record of sea level rise and environmental change during the Paleocene–Eocene Thermal Maximum. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **305**, 185–200.

- Harley, M.M. (1991) Systematic implications of comparative morphology in selected Tertiary and extant pollen from the Palmae and the Sapotaceae. S. Blackmore & S. Barnes, eds., *Pollen and spores: patterns of diversification*, Clarendon Press, Oxford, pp. 225–238.
- Harley, M.M. & Morley, R.J. (1995) Ultrastructural studies of some fossil and extant palm pollen, and the reconstruction of the biogeographical history of subtribes Iguanurinae and Calaminae. *Review of Palaeobotany and Palynology*, **85**, 153–182.
- Harrington, G.J. (2003) Wasatchian (Early Eocene) pollen floras from the Red Hot Truck Stop, Mississippi, USA. *Palaeontology*, **46**, 725–738.
- Harrington, G.J. (2008) Comparisons between Palaeocene–Eocene paratropical swamp and marginal marine pollen floras from Alabama and Mississippi, USA. *Palaeontology*, **51**, 611–622.
- Haseldonckx, P. (1973) The palynology of some Palaeogene deposits between the Rio Esera and the Rio Segre, southern Pyrenees, Spain. *Leidse geologische mededelingen*, **49**, 145–163.
- Haseldonckx, P. (1977) The palynology of a Holocene marginal peat swamp environment in Johore, Malaysia. *Review of Palaeobotany and Palynology*, **24**, 227–238.
- Hayward, B.W. & Raine, J.I. (1978) Biostratigraphy of North Tasman-1 offshore Well. *New Zealand Geological Survey, Report Palynology*, 1–11.
- Hekel, H. (1972) Pollen and spore assemblages from Queensland Tertiary sediments. *Geological Survey of Queensland, Publication*, 1–34.
- Helenes, J. & Cabrera, D. (2003) Oligocene-miocene palynomorph assemblages from eastern Venezuela. *Palynology*, **27**, 5–25.
- Helenes, J., Somoza, D., Vasquez, J., De Guerra, C. & Corpoven, S.A. (1994) Chronostratigraphy and palynomorph assemblages of a Cretaceous/Oligocene unconformity in Eastern Venezuela. *Palynology*, **18**, 248.
- Hermanowski, B., da Costa, M.L. & Behling, H. (2012a) Environmental changes in southeastern Amazonia during the last 25,000 yr revealed from a paleoecological record. *Quaternary Research*, **77**, 138–148.
- Hermanowski, B., da Costa, M.L., Carvalho, A.T. & Behling, H. (2012b) Palaeoenvironmental dynamics and underlying climatic changes in southeast Amazonia (Serra Sul dos Carajás, Brazil) during the late Pleistocene and Holocene. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **365**, 227–246.
- Hernandez, P.E. (1998) *Sedimentology of a mixed carbonate-siliciclastic succession: The Guasare Formation, Maracaibo Basin, Venezuela*. Master's thesis, University of Calgary.
- Heryanto, R. (2006) Perbandingan karakteristik lingkungan pengendapan, batuan sumber, dan diagenesis Formasi Lakat di lereng timur laut dengan Formasi Talangakar di tenggara Pegunungan Tigapuluh, Jambi. *Indonesian Journal on Geoscience*, **1**, 173–184.
- Higgs, K.E., King, P.R., Raine, J.I., Sykes, R., Browne, G.H., Crouch, E.M. & Baur, J.R. (2012) Sequence stratigraphy and controls on reservoir sandstone distribution in an Eocene marginal marine-coastal plain fairway, Taranaki Basin, New Zealand. *Marine and Petroleum Geology*, **32**, 110–137.
- Hochuli, P. (1982) Organische Mikrofossilien aus Proben der Molasse von Vorarlberg und Teilen des Schweizerischen Mittellandes. *Travaux et Documents des Laboratoires de Géologie de Lyon*, **7**, 39–45.
- Hofmann, C.C., Egger, H. & King, C. (2015) LM and SEM investigations of pollen from PETM and EECO localities of Austria and Great Britain: new findings of Atherospermataceae, Annonaceae, Araceae and Areaceae from the Lower Eocene. *Plant systematics and evolution*, **301**, 773–793.

- Hofmann, C.C., Mohamed, O. & Egger, H. (2011) A new terrestrial palynoflora from the Palaeocene/Eocene boundary in the northwestern Tethyan realm (St. Pankraz, Austria). *Review of Palaeobotany and Palynology*, **166**, 295–310.
- Hofmann, C.C., Pancost, R., Ottner, F., Egger, H., Taylor, K., Mohamed, O. & Zetter, R. (2012) Palynology Biomarker and Clay mineralogy of the Early Eocene Climate Optimum (EECO) in the transgressive Krappfeld succession (Eastern Alps, Austria). *Austrian Journal of Earth Sciences*, **105**.
- Hoorn, C. (1988) Nota preliminar sobre la edad de los sedimentos Terciarios de la zona de Araracuara (Amazonas). *Boletín Geológico*, **29**, 85–95.
- Hoorn, C. (1993) Marine incursions and the influence of Andean tectonics on the Miocene depositional history of northwestern Amazonia: results of a palynostratigraphic study. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **105**, 267–309.
- Hoorn, C. (1994a) An environmental reconstruction of the palaeo-Amazon river system (Middle–Late Miocene, NW Amazonia). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **112**, 187–238.
- Hoorn, C. (1994b) Fluvial palaeoenvironments in the intracratonic Amazonas Basin (early Miocene–early middle Miocene, Colombia). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **109**, 1–54.
- Hoorn, C. (1994c) Miocene depositional history of northwestern Amazonia: results of a palynostratigraphic study. *Palynology*, **18**, 248–249.
- Hoorn, C. (2006) Mangrove forests and marine incursions in Neogene Amazonia (lower Apaporis River, Colombia). *Palaios*, **21**, 197–209.
- Hoorn, C., Bogotá-A, G.R., Romero-Baez, M., Lammertsma, E.I., Flantua, S.G.A., Dantas, E.L., Dino, R., do Carmo, D.A. & Chemale Jr, F. (2017) The Amazon at sea: Onset and stages of the Amazon River from a marine record, with special reference to Neogene plant turnover in the drainage basin. *Global and Planetary Change*, **153**, 51–65.
- Hoorn, C., Goodman, D.K. & Clarke, R.T. (2001) Pleistocene deposits of the Amazon Fan: a palynological analysis of holes 940a, 944a, and 946a (ODP leg 155). *Proceedings of the IX International Palynological Congress, Houston*, vol. 1996, p. 32.
- Hoorn, C., Straathof, J., Abels, H.A., Xu, Y., Utescher, T. & Dupont-Nivet, G. (2012) A late Eocene palynological record of climate change and Tibetan Plateau uplift (Xining Basin, China). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **344**, 16–38.
- Hopping, C. (1967) Palynology and the oil industry. *Review of Palaeobotany and Palynology*, **2**, 23–48.
- Huang, H., Morley, R., Licht, A., Dupont-Nivet, G., Grímsson, F., Zetter, R., Westerweel, J., Win, Z., Wa Aung, D. & Hoorn, C. (2020) Eocene palms from central Myanmar in a South-East Asian and global perspective: evidence from the palynological record. *Botanical Journal of the Linnean Society*, **194**, 177–206.
- Huang, L. & Zhong, B. (1996) Paleogene stratigraphy and Paleocyanography in Pearl River Mouth Basin. Y. Hao *et al.*, eds., *Research on Micropalaeontology and Paleocyanography in Pearl River Mouth Basin, South China Sea*, China University of Geosciences Press, pp. 99–106.
- Hunt, C.O. & Premathilake, R. (2012) Early Holocene vegetation, human activity and climate from Sarawak, Malaysian Borneo. *Quaternary International*, **249**, 105–119.
- Igbinigie, N.S. & Akenzua-Adamczyk, A.H. (2018) Palynological studies of Maastrichtian to paleocene sediments exposed at Okpekpe, western flank of Anambra Basin, Edo State, Nigeria. *Journal of Applied Sciences and Environmental Management*, **22**, 1563–1566.

- Igbini, N.S., Ogbamikhumi, A. & Komlan-Dodoh, N.O. (2017) Palynological Evaluation of Potential Source Rocks at Imiegba, Benin Flank of the Anambra Basin, Nigeria. *Indo-Iranian Journal of Scientific Research*, **1**, 88–94.
- Ikegwonu, O.N. & Umeji, O.P. (2016) Palynological age and palaeoenvironment of deposition of Mid-Cenozoic sediments around Umuahia, Niger delta basin, southeastern Nigeria. *Journal of African Earth Sciences*, **117**, 160–170.
- Ikegwonu, O.N., Umeji, O.P., Chiaghanam, O.I., Nwozor, K.K., Ndukwe, O.S. & Chiadikobi, K.C. (2020) Palynomorph assemblage biozonation of Paleogene strata in Bende–Umuahia Area, Niger Delta Basin, southeastern Nigeria. *Journal of Palaeogeography*, **9**, 1–13.
- Irion, G., Bush, M.B., De Mello, J.N., Stüben, D., Neumann, T., Müller, G., Junk, J.W. *et al.* (2006) A multiproxy palaeoecological record of Holocene lake sediments from the Rio Tapajós, eastern Amazonia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **240**, 523–535.
- Jaillard, E. (1995) Síntesis estratigráfica del Cretaceo y Paleógeno de la cuenca oriental del Perú. *Convenio ORSTOM-Petroproduccion, Quito*.
- Jain, K.P., Kar, R.K. & Sah, S.C.D. (1973) A palynological assemblage from Barmer, Rajasthan. *Geophytology*, **3**, 150–165.
- Jais, J.H.M. (1999) *Oligocene to Pliocene quantitative stratigraphic palynology of the southern Malay Basin, offshore Malaysia*. Ph.D. thesis, University of Sheffield.
- Jan du Chêne, R.E., Klasz, D.E. & Archibong, E.E. (1978a) Biostratigraphic study of the borehole OJO-1, SW Nigeria, with special emphasis on the Cretaceous microflora. *Revue de Micropaléontologie*, **21**, 123–139.
- Jan du Chêne, R.E., Onyke, M.S. & Sowunmi, M.A. (1978b) Some new Eocene pollen of the Ogwashi-Asaba Formation, south-eastern Nigeria. *Revista Española de Micropaleontología*, **10**, 285–322.
- Jan du Chêne, R.E., Perch-Nielsen, K. & Petters, S.W. (1978c) New stratigraphic data on the Paleogene Ewekoro and Akinbo Formations (southwestern Nigeria). *Archives des sciences*, **31**, 205–212.
- Jan du Chêne, R.E. & Salami, M.B. (1978) Palynology and micropaleontology of the Upper Eocene of the well nsukwa-1 (Niger Delta, Nigeria). *Archives des Sciences*, **13**, 5–9.
- Jaramillo, C., Hoorn, C., Silva, S.A.F., Leite, F., Herrera, F., Quiroz, L., Dino, R. & Antonioli, L. (2010) The origin of the modern Amazon rainforest: implications of the palynological and palaeobotanical record. C. Hoorn & F. Wesselingh, eds., *Amazonia, landscape and species evolution*, Wiley-Blackwell, Chichester, U.K., pp. 317–334.
- Jaramillo, C., Romero, I., D’Apolito, C., Bayona, G., Duarte, E., Louwye, S., Escobar, J., Luque, J., Carrillo-Briceño, J.D., Zapata, V. *et al.* (2017) Miocene flooding events of western Amazonia. *Science Advances*, **3**, e1601693.
- Jaramillo, C., Sepulchre, P., Cardenas, D., Correa-Metrio, A., Moreno, J.E., Trejos, R., Vallejos, D., Hoyos, N., Martínez, C., Carvalho, D. *et al.* (2020) Drastic Vegetation Change in the Guajira Peninsula (Colombia) During the Neogene. *Paleoceanography and Paleoclimatology*, **35**, e2020PA003933.
- Jaramillo, C.A., Bayona, G., Pardo-Trujillo, A., Rueda, M., Torres, V., Harrington, G.J. & Mora, G. (2007) The palynology of the Cerrejón Formation (upper Paleocene) of northern Colombia. *Palynology*, **31**, 153–189.
- Jaramillo, C.A., Munoz, F., Cogollo, M. & Parra, F. (2005) Quantitative biostratigraphy for the Paleocene of the Llanos foothills, Colombia: improving palynological resolution for oil exploration. A.J. Powell & J.B. Riding, eds., *Recent Developments in Applied Biostratigraphy*, The Micropalaeontological Society, pp. 145–159.

- Jaramillo, C.A., Rueda, M. & Torres, V. (2011) A palynological zonation for the Cenozoic of the Llanos and Llanos Foothills of Colombia. *Palynology*, **35**, 46–84.
- Jardine, P.E. (2011) *Spatial and temporal diversity trends in an extra-tropical, megathermal vegetation type: the early Palaeogene pollen and spore record from the US Gulf Coast*. Ph.D. thesis, University of Birmingham.
- Jarzen, D.M. & Dilcher, D.L. (2006) Middle Eocene terrestrial palynomorphs from the Dolime Minerals and Gulf Hammock quarries, Florida, USA. *Palynology*, **30**, 89–110.
- Jelen, B., Kedves, M., Skaberne, D., Brezigar, A., Buser, S., Cimerman, F., Drobne, K., Monostori, M., Pavlovec, R. & Pavšič, J. (1994) Dorog type (Middle Eocene) Spore-pollen assemblage in the Socka Beds of Slovenia I. *Plant Cell Biology and Development*, **5**, 20–28.
- Jiménez, H.D. & van der Hammen, T. (2007) Significado geológico y asociaciones palinológicas de las Formaciones Diablo inferior (Mioceno tardío) y San Fernando superior (Mioceno medio), Piedemonte Cuenca de Los Llanos Orientales, Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, **31**, 481–499.
- Jolley, D.W. (1998) Early Eocene palynofloras from Holes 915A, 916A, 917A and 918D, East Greenland. *Proceedings of the Ocean Drilling Program: Scientific results*, vol. 152, Ocean Drilling Program, pp. 221–231.
- Jones, E.L. (1961) *Plant microfossils of the laminated sediments of the Lower Eocene Wilcox Group in south-central Arkansas*. Ph.D. thesis, University of Oklahoma.
- Jones, J.G. & Gennett, J.A. (1991) Pollen and Spores from the Type Section of the Middle Eocene Stone City Formation, Burlison Co., Texas. *Transactions of the Gulf Coast Association of Geological Societies*, **41**, 348–352.
- Jones, S.E., Hunt, C. & Reimer, P.J. (2013) A 2300 yr record of sago and rice use from the southern Kelabit Highlands of Sarawak, Malaysian Borneo. *The Holocene*, **23**, 708–720.
- Kachniasz, K.E. & da Silva-Caminha, S.A.F. (2017) Palinoestratigrafia da Formação Solimões: Comparação entre bioestratigrafia tradicional e o método de associações unitárias. *Revista Brasileira de Paleontologia*, **19**, 481–490.
- Kapoor, P.N. (1994) Fluorescence palynology: a tool for quantitative estimation of organic matter maturation. *Geoscience Journal*, **15**, 1–10.
- Kapoor, P.N., Prasad, B., Swamy, S.N. & Shukla, M. (1999) Palynostratigraphy and hydrocarbon source potential of Cretaceous sediments in Krishna Sub-Basin, India. *Geoscience Journal*, **20**, 11–24.
- Kapoor, R., Dogra, N.N. & Singh, R.Y. (1988) Palynology of the Kasauli Formation in the type area in Solan District, Himachal Pradesh. *Journal of the Palaeontological Society of India*, **33**, 105–115.
- Kar, R.K. (1985) The fossil floras of Kachchh-IV. Tertiary palynostratigraphy. *Palaeobotanist*, **34**, 1–279.
- Kar, R.K. (1989) Palynological studies of the Barail Group (Oligocene) in the type area, Assam. *Palaeobotanist*, **38**, 229–242.
- Kar, R.K. (1991) Occurrence of *Acrostichum* spores from the Langpar Formation, Early Palaeocene of Meghalaya, India. *Geophytology*, **21**, 33–36.
- Kar, R.K. (1993) Occurrence of *Normapolles* pollen from the Miocene sediments of Kerala, India. *Journal of Palynology*, **29**, 29–39.
- Kar, R.K. & Bhattacharya, M. (1990) Palynology of Rajpardi lignite, Cambay Basin and Gujra Dam and Akri lignite, Kutch Basin. *Palaeobotanist*, **39**, 250–263.

- Kar, R.K., Handique, G.K., Kalita, C.K., Mandal, J., Sarkar, S., Kumar, M. & Gupta, A. (1993) Palynostratigraphical studies on subsurface Tertiary sediments in Upper Assam Basin, India. *Palaeobotanist*, **42**, 183–198.
- Kar, R.K. & Jain, K.P. (1981) Palynology of Neogene sediments around Quilon and Varkala, Kerala coast, South India-2. Spores and pollen grains. *Palaeobotanist*, **27**, 113–131.
- Karaman, M.E., Meriç, E. & Tansel, İ. (1989) Gönen–Atabey arasındaki bölgenin jeolojisi. *Cum. Üniv. Müh. Fak. Derg., Sivas*.
- Kasiński, J.R. & Słodkowska, B. (2016) Factors controlling Cenozoic anthracogenesis in the Polish Lowlands. *Geological Quarterly*, **60**, 959–974.
- Kaska, H.V. (1989) A spore and pollen zonation of Early Cretaceous to Tertiary nonmarine sediments of central Sudan. *Palynology*, **13**, 79–90.
- Kayseri, M.S. & Akgün, F. (2008) Palynostratigraphic, palaeovegetational and palaeoclimatic investigations on the Miocene deposits in Central Anatolia (Çorum Region and Sivas Basin). *Turkish Journal of Earth Sciences*, **17**, 361–403.
- Kayseri, M.S. & Akgun, F. (2010) The late Burdigalian-Langhian time interval in Turkey and the palaeoenvironment and palaeoclimatic implications and correlation of Europe and Turkey: late Burdigalian-Langhian palynofloras and palaeoclimatic properties of the Mugla-Milas (Kultak). *Turkiye Jeoloji Bulteni-Geological Bulletin of Turkey*, **53**, 1–44.
- Kayseri-Özer, M.S. (2013) Spatial distribution of climatic conditions from the Middle Eocene to Late Miocene based on palynoflora in Central, Eastern and Western Anatolia. *Geodinamica Acta*, **26**, 122–157.
- Kedves, M. (1968) Etudes palynologiques des couches du Tertiaire Inférieur de la région parisienne. *Pollen et Spores*, **10**, 315–334.
- Kedves, M. (1974) Paleogene fossil sporomorphs of the Bakony Mountains: Part II. *Studia Biologica Hungarica*, **13**, 5–124.
- Kedves, M. (1986a) Introduction to the Palynology of Pre-quaternary Deposits. *Studia Biologica Academiae Scientiarum Hungaricae*, 144.
- Kedves, M. (1986b) Paleogene fossil sporomorphs of the Bakony Mountains: Part IV. *Studia Biologica Hungarica*, **21**, 1–121.
- Kelly, T.J., Lawson, I.T., Roucoux, K.H., Baker, T.R., Honorio-Coronado, E.N., Jones, T.D. & Rivas Panduro, S. (2018) Continuous human presence without extensive reductions in forest cover over the past 2500 years in an aseasonal Amazonian rainforest. *Journal of Quaternary Science*, **33**, 369–379.
- Kelly, T.J., Lawson, I.T., Roucoux, K.H., Baker, T.R., Jones, T.D. & Sanderson, N.K. (2017) The vegetation history of an Amazonian domed peatland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **468**, 129–141.
- Kemp, E.M. & Harris, W.K. (1975) The vegetation of Tertiary islands on the Ninetyeast Ridge. *Nature*, **258**, 303–307.
- Kershaw, A.P. (1983) A Holocene pollen diagram from Lynch's Crater, north-eastern Queensland, Australia. *New Phytologist*, **94**, 669–682.
- Kershaw, A.P. & Sluiter, I.R. (1982) Late Cenozoic pollen spectra from the Atherton Tableland, north-eastern Australia. *Australian Journal of Botany*, **30**, 279–295.

- Khan, A.M. (1976) Palynology of Tertiary sediments from Papua New Guinea. I. New form genera and species from Upper Tertiary sediments. *Australian journal of botany*, **24**, 753–781.
- Khan, S. (1995) Results of palynological studies on rock samples from the Surghar range. Palynostratigraphy of U.S. Geological Survey-Geological Survey of Pakistan samples from the Makarwal area. *United States Geological Survey, Bulletin*, **2096**, 33.
- Khanolkar, S. & Sharma, J. (2019) Record of early to middle Eocene paleoenvironmental changes from lignite mines, western India. *Journal of Micropalaeontology*, **38**, 1–24.
- Kirchner, M. (1984) The Upper Oligocene microflora of the Southern Bavarian Pitch Coal Mine District. *Palaeontographica Abteilung B*, **192**, 85–162.
- Knobloch, E. & Konzalová, M. (1998) Comparison of the Eocene plant assemblages of Bohemia (Czech Republic) and Saxony (Germany). *Review of Palaeobotany and Palynology*, **101**, 29–41.
- Kogbe, C.A. (1981) Attempt to correlate the stratigraphic sequence in the middle Benue basin with those of the Anambra and upper Benue basins. *Earth Evolution Science*, **1**, 139–143.
- Kogbe, C.A. & Sowunmi, M.A. (1975) The age of the Gwandu formation (Continental Terminal) in Northwestern Nigeria as suggested by spore-pollinitic analysis. *Savanna*, **4**, 47–51.
- Konzalova, M. (1971) Arecales (Palmae) in the North-Bohemian Tertiary. *Sbor. geol. VČd, paleont*, **13**, 143–158.
- Konzalová, M. (1972) Paleotropic and stratigraphically important plant remains (spores and pollen), from the Early Tertiary of the northwestern Bohemia. *Review of Palaeobotany and Palynology*, **47**, 239–242.
- Konzalová, M. (1990) Palynological investigation of the basal Tertiary sediments (Paleogene) in western Bohemia and their correlation with the Carpathian region. *Západné Karpaty Sér. Paleontol.*, **14**, 73–102.
- Konzalová, M. & Ziembińska-TworzydŁo, M. (2008) Some monocot pollen taxa from the Lower Miocene basal coaly deposits of the Czech and Polish parts of the Žytawa (Zittau) Basin. *Acta Musei Nationalis Pragae, Series B – Historia Naturalis*, **64**, 149–162.
- Koshal, V.N. & Uniyal, S.N. (1984) Paleocene-Early Eocene palynofossils in the subsurface of North Cambay Basin, Gujarat (western India). *Indian Colloquium on Micropalaeontology and Stratigraphy, Proceedings, 10th*, **10**, 233–244.
- Kovar-Eder, J., Kvaček, Z., Martinetto, E. & Roiron, P. (2006) Late Miocene to Early Pliocene vegetation of southern Europe (7–4 Ma) as reflected in the megafossil plant record. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **238**, 321–339.
- Krutzsch, W. (1961) Beitrag zur Sporenpaläontologie der präoberoligozänen kontinentalen und marinen Tertiärlagerungen Brandenburgs. *Berichte Geologische Gesellschaft*, **5**, 290–343.
- Krutzsch, W. (1976) *The microflora of the Brown Coal of the Geisel Valley; Part IV: The stratigraphic position of the Geisel Valley profile in the Eocene, and the spore-stratigraphic subdivision of the Middle Eocene*. Abhandlungen des Zentralen Geologischen Instituts.
- Krutzsch, W. & Vanhoome, R. (1977) Die Pollenflora von Epinois und Loksbergen in Belgien. *Palaeontographica Abteilung B*, **163**, 1–110.
- Kulkarni, A.R. & Phadtare, N.R. (1981) Pollen of *Nypa* from lignitic beds of Ratnagiri district, Maharashtra. *Phytomorphology*, **31**, 48–51.
- Kumar, A. (1993) Biozonation and depositional environments of the Tertiary outcrop section along the Silchar-Haflong Road. Cachar District, Southern Assam. *Proceedings 2nd Seminar on Petroliferous Basins of India*, vol. 1, pp. 459–476.

- Kumar, A. (2016) New palynological evidence for the age of the Beda Formation, Sirte Basin, Libya. *Palaeontologia Electronica*, **19**, 1–14.
- Kumar, A. & Takahashi, K. (1991) Palynology of the Tertiary sediments of southern Assam, India. *Bulletin of the Faculty of Liberal Arts, Nagasaki University, (Natural Science)*, **31**, 515–659.
- Kumar, M. (1994) Palynological and palaeoecological studies of Palaeocene coal seams in Jarain and Laitrymbai areas, Jaintia Hills, Meghalaya, India. *Geophytology*, **23**, 203–207.
- Kumar, M., Mandal, J., Dutta, S.K., Bhuyan, D., Das, B. & Saikia, B. (2001) Palynostratigraphy of the subsurface sediments of Upper Assam basin, India. *Geobios*, **34**, 241–251.
- Kumar, M., Spicer, R.A., Spicer, T.E.V., Shukla, A., Mehrotra, R.C. & Monga, P. (2016) Palynostratigraphy and palynofacies of the early Eocene Gurha lignite mine, Rajasthan, India. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **461**, 98–108.
- Kumar, M., Srivastava, G., Spicer, R.A., Spicer, T.E.V., Mehrotra, R.C. & Mehrotra, N.C. (2012) Sedimentology, palynostratigraphy and palynofacies of the late Oligocene Makum Coalfield, Assam, India: A window on lowland tropical vegetation during the most recent episode of significant global warmth. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **342**, 143–162.
- Kumar, P. (2012) *Palynological investigation of coal-bearing deposits of the Thar Coal Field Sindh, Pakistan*. Master's thesis, Lund University.
- Kumaran, K.P.N., Soman, K., Kamble, C.V. & Joseph, A. (1995) Palynofloral analysis of sections from Bharathi and Kundara clay mines of Kerala basin: palaeological and tectonic perspective. *Current science*, **69**, 1023–1027.
- Kwetche, P., Ntamak-Nida, M.J., Nitchou, A.L.D., Etame, J., Owono, F.M., Mbesse, C.O., Kissaaka, J.B.I., Ngon, G.N., Bourquin, S. & Bilong, P. (2018) Facies analysis and sequence stratigraphy of Missole Outcrops: N'Kapa Formation of the south-eastern edge of Douala Sub-Basin (Cameroon). *Earth Sciences Research*, **7**, 35–54.
- Lang, J., Kogbe, C., Alidou, S., Alzouma, K.A., Bellion, G., Dubois, D., Durand, A., Guiraud, R., Houessou, A., De Klasz, I. *et al.* (1990) The continental terminal in West Africa. *Journal of African Earth Sciences (and the Middle East)*, **10**, 79–99.
- Lara, M., Cardona, A., Monsalve, G., Yarce, J., Montes, C., Valencia, V., Weber, M., de La Parra, F., Espitia, D. & López-Martínez, M. (2013) Middle Miocene near trench volcanism in northern Colombia: A record of slab tearing due to the simultaneous subduction of the Caribbean Plate under South and Central America? *Journal of South American Earth Sciences*, **45**, 24–41.
- Larny, A. (1986) Plio-Pleistocene Palynology and Visual Kerogen Studies, Trinidad, WI, With Emphasis on the Columbus Basin. *Transactions of the First Geological Conference of the Geological Society of Trinidad & Tobago, Port-of-Spain, Trinidad & Tobago*, 115–125.
- Latrubesse, E.M., Cozzuol, M., da Silva-Caminha, S.A.F., Rigsby, C.A., Absy, M.L. & Jaramillo, C. (2010) The Late Miocene paleogeography of the Amazon Basin and the evolution of the Amazon River system. *Earth-Science Reviews*, **99**, 99–124.
- Latrubesse, E.M., da Silva, S.A.F., Cozzuol, M. & Absy, M.L. (2007) Late Miocene continental sedimentation in southwestern Amazonia and its regional significance: Biotic and geological evidence. *Journal of South American Earth Sciences*, **23**, 61–80.

- Leal, A., Bilbao, B. & Berrío, J.C. (2013) A contribution to pollen rain characterization in forest-savanna mosaics of the Venezuelan Guayana and its use in vegetation reconstructions from sedimentary records. *American Journal of Plant Sciences*, **4**, 33–52.
- Leal, A., Perez, T. & Bilbao, B. (2011) Contribution to Early Holocene vegetation and climate history of Eastern Orinoco Llanos, Venezuela, from a paleoecological record of a *Mauritia* Lf swamp. *Acta Amazonica*, **41**, 513–520.
- Leal, A.V. & Bilbao, B.A. (2011) Vegetation changes during the late Holocene in morichal of the Orinoco Llanos, Venezuela. *Acta Botanica Venezuelica*, **34**, 237–256.
- Leandro, L.M., Vieira, C.E.L., Santos, A. & Fauth, G. (2019) Palynostratigraphy of two Neogene boreholes from the northwestern portion of the Solimões Basin, Brazil. *Journal of South American Earth Sciences*, **89**, 211–218.
- Ledru, M.P., Ceccantini, G., Gouveia, S.E.M., López-Sáez, J.A., Pessenda, L.C.R. & Ribeiro, A.S. (2006) Millennial-scale climatic and vegetation changes in a northern Cerrado (Northeast, Brazil) since the Last Glacial Maximum. *Quaternary Science Reviews*, **25**, 1110–1126.
- Ledru, M.P., Salgado-Labouriau, M.L. & Lorscheitter, M.L. (1998) Vegetation dynamics in southern and central Brazil during the last 10,000 yr BP. *Review of Palaeobotany and Palynology*, **99**, 131–142.
- Lee, C.Y., Liew, P.M. & Lee, T.Q. (2010) Pollen records from southern Taiwan: implications for East Asian summer monsoon variation during the Holocene. *The Holocene*, **20**, 81–89.
- Leidelmeyer, P. (1966) The Paleocene and lower Eocene pollen flora of Guyana. *Leidse geologische mededelingen*, **38**, 49–57.
- Leite, F.P.R. (1997) *Palinofloras neógenas da Formação Pirabas e Grupo Barreiras, área litorânea nordeste do Estado do Pará, Brasil*. Master's thesis, Universidade de São Paulo.
- Leite, F.P.R. (2006) *Palinogia da formação Solimões, neógeno da Bacia do Solimões, Estado do Amazonas, Brasil: implicações paleoambientais e bioestratigráficas*. Ph.D. thesis, Universidade de Brasília.
- Leite, F.P.R., Paz, J., do Carmo, D.A. & Silva-Caminha, S.A. (2017) The effects of the inception of Amazonian transcontinental drainage during the Neogene on the landscape and vegetation of the Solimões Basin, Brazil. *Palynology*, **41**, 412–422.
- Lelono, E.B. (2000) *Palynological Study of the Eocene Nanggulan Formation, Central Java, Indonesia*. Ph.D. thesis, Royal Holloway, University of London.
- Lelono, E.B. (2007a) Gondwanan palynomorphs from the Paleogene sediment of East Java: ?The evidence of earlier arrival. *Lemigas Scientific Contributions*, **30**, 1–12.
- Lelono, E.B. (2007b) Palinomorf Eosen dari Selat Makasar. *Lembaran Publikasi Minyak dan Gas Bumi*, **41**, 1–10.
- Lelono, E.B. (2007c) Palynological Investigation Of The Oligocene Sediment In East Java Sea. *Lemigas Scientific Contributions*, **30**, 7–17.
- Lelono, E.B. & Morley, R.J. (2011) Oligocene palynological succession from the East Java Sea. *Geological Society, London, Special Publications*, **355**, 333–345.
- Lenz, O.K. (2000) *Paläoökologie eines Küstenmoores aus dem Eozän Mitteleuropas am Beispiel der Wulfersdorfer Flöze und deren Begleitschichten (Helmstedter Oberflözgruppe, Tagebau Helmstedt)*. Ph.D. thesis, University of Göttingen.

- Lenz, O.K. & Riegel, W. (2001) Isopollen maps as a tool for the reconstruction of a coastal swamp from the middle Eocene at Helmstedt (northern Germany). *Facies*, **45**, 177–194.
- Lenz, O.K., Wilde, V. & Riegel, W. (2007) Recolonization of a Middle Eocene volcanic site: quantitative palynology of the initial phase of the maar lake of Messel (Germany). *Review of Palaeobotany and Palynology*, **145**, 217–242.
- Leško, B., Ďurkovič, T., Gašparíková, V., Samuel, O. & Snopková, P. (1984) A geological evaluation of Hanušovce-1 borehole. *Mineralia Slovaca*, **16**, 217–255.
- Li, J., Batten, D.J. & Zhang, Y. (2008) Palynological indications of environmental changes during the Late Cretaceous–Eocene on the southern continental margin of Laurasia, Xizang (Tibet). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **265**, 78–86.
- Li, Z. (1996) Paleogene stratigraphy and Paleooceanography in Pearl River Mouth Basin. Y. Hao, ed., *Research on Micropalaeontology and Paleooceanography in Pearl River Mouth Basin, South China Sea*, China University of Geosciences Press, pp. 43–51.
- Li, Z. (1998) Eocene palynology of Well LF13-2-1 in Pearl River Mouth Basin. *China Offshore Oil and Gas (Geology)*, **12**, 168–173.
- Liew, P.M., Wu, M.H., Lee, C.Y., Chang, C.L. & Lee, T.Q. (2014) Recent 4000 years of climatic trends based on pollen records from lakes and a bog in Taiwan. *Quaternary International*, **349**, 105–112.
- Lima Jr, W.J.S., Cohen, M.C.L., Rossetti, D.F. & França, M.C. (2018) Late Pleistocene glacial forest elements of Brazilian Amazonia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **490**, 617–628.
- Linhares, A.P., de Souza Gaia, V.d.C. & Ramos, M.I.F. (2017) The significance of marine microfossils for paleoenvironmental reconstruction of the Solimões Formation (Miocene), western Amazonia, Brazil. *Journal of South American Earth Sciences*, **79**, 57–66.
- Liu, K.B. & Colinvaux, P.A. (1988) A 5200-year history of Amazon rain forest. *Journal of Biogeography*, 231–248.
- Lorente, M.A. (1986) Palynology and palynofacies of the Upper Tertiary in Venezuela. *Dissertationes. Botanicæ*, **9**, 1–222.
- Lucas, F.A. (2017) Miospores and Geological Boundaries in Maastrichtian to Lutetian Succession of Ajire-1 well, Anambra Basin, Nigeria. *International Journal of Science and Advanced Innovative Research*, **2**, 74–84.
- Lucas, F.A. & Omodolor, H.E. (2018) Palynofacies Analysis, Organic Thermal Maturation and Source Rock Evaluation of Sedimentary Succession from Oligocene to Early Miocene Age in X2 Well, Greater Ughelli Depobelt, Niger Delta Basin, Nigeria. *Journal of Geosciences and Geomatics*, **6**, 85–93.
- Macphail, M. (1999) Palynostratigraphy of the Murray Basin, inland southeastern Australia. *Palynology*, **23**, 197–240.
- Macphail, M. & Gibson, D. (2014) Testing the Gippsland Basin zonation in northern Australia: palynostratigraphical analysis of a 23 Ma $^{40}\text{Ar}/^{39}\text{Ar}$ dated claystone from Toowoomba, southeast Queensland. *Palynology*, **38**, 117–128.
- Macphail, M. *et al.* (2007) *Australian palaeoclimates: Cretaceous to Tertiary a review of palaeobotanical and related evidence to the year 2000*. CRC LEME.
- Macphail, M.K. (2013) *Preliminary Palynostratigraphic Age Determination on Core from Strahan BH-1, Macquarie Harbour, Western Tasmania*. Tasmanian Geological Survey.

- Macphail, M.K. & Hill, R.S. (2019) What was the vegetation in northwest Australia during the Paleogene, 66–23 million years ago? *Australian Journal of Botany*, **66**, 556–574.
- Macphail, M.K. & Jordan, G.J. (2015) Tropical palms and arums at near-polar latitudes: fossil pollen evidence from the Tamar and Macquarie grabens, northern Tasmania. *Papers and Proceedings of the Royal Society of Tasmania*, vol. 149, pp. 23–28.
- Maezumi, S.Y., Alves, D., Robinson, M., de Souza, J.G., Levis, C., Barnett, R.L., de Oliveira, E.A., Urrego, D., Schaan, D. & Iriarte, J. (2018) The legacy of 4,500 years of polyculture agroforestry in the eastern Amazon. *Nature Plants*, **4**, 540–547.
- Maloney, B. (1999) A 10,600 year pollen record from Nong Thale Song Hong, Trang province, south Thailand. *Bulletin of the Indo-Pacific Prehistory Association*, **18**, 129–138.
- Mandal, J. (1996) Palynofossils from the Tertiary (Barail Group) of Nagaland: Palaeoecological interpretation and age. *Palaeobotanist*, **45**, 98–108.
- Mandal, J., Chandra, A. & Bhattacharyya, A.P. (2003) Palynology of the Baratang Formation, Andaman-Nicobar Islands and the significance of reworked palynomorphs. *Palaeobotanist*, **52**, 97–112.
- Mandal, J., Chandra, A. & Kar, R.K. (1996) Palynological findings from the mud volcanoes of Baratang Island (Andaman and Nicobar Islands), India. *Geophytology*, **25**, 77–81.
- Mandal, J. & Guleria, J.S. (2006) Palynology of Vastan lignite (Surat District), Gujarat: its age, palaeoecology and depositional environment. *Palaeobotanist*, **55**, 1–66.
- Mandal, J.P. (1986) Palynological study of Sutunga coal seam, Jaintia Hills, Meghalaya. *Palaeobotanist*, **35**, 196–199.
- Mandal, J.P. (1990) Palynological investigation of Palaeocene sediments from Thanjinath, Meghalaya. *Palaeobotanist*, **37**, 324–330.
- Mandaokar, B. (1993) A palynological investigation of the Tirak Parbat Formation (Oligocene) of Dangri Kumari Colliery, Dibrugarh district, Assam, India. *Tertiary Research*, **14**, 127–139.
- Mandaokar, B. (2000) Palynology of coal bearing sediments of the Tikak Parbat Formation (Oligocene) from Namchik River section, Changlang District, Arunachal Pradesh, India. *Tertiary Research*, **20**, 37–46.
- Mandaokar, B.D. (1990) Palynology of Miocene rocks around Maibong, Assam. *Geophytology*, **20**, 24–29.
- Mandaokar, B.D. (2002) Palynoflora from the Keifang Formation (early Miocene) Mizoram, India and its environmental significance. *Journal of the Palaeontological Society of India*, **47**, 77–83.
- Mandaokar, B.D. (2004) Age and depositional environment of the Upper Bhuban Formation of Champhai area (Eastern Mizo hills) India-A palynological approach. *Palaeobotanist*, **53**, 143–153.
- Mandaokar, B.D. & Mukherjee, D. (2014) Palynostratigraphy of the Cuddalore Formation (Early Miocene) of Panruti, Tamil Nadu, India. *Journal of the Palaeontological Society of India*, **59**, 69–80.
- Marquillas, R.A., Del Papa, C. & Sabino, I.F. (2005) Sedimentary aspects and paleoenvironmental evolution of a rift basin: Salta Group (Cretaceous–Paleogene), northwestern Argentina. *International Journal of Earth Sciences*, **94**, 94–113.
- Mathews, R.P., Singh, H., Singh, V.P., Singh, B.D. & Singh, A. (2015) Organic composition and palaeoenvironment of Valia Lignite Deposit (Cambay Basin), Gujarat, western India: inferences from palynology and petrography. *67th Annual Meeting of the International Committee for Coal and Organic Petrology*, 117.

- Mathews, R.P., Tripathi, S.M., Banerjee, S. & Dutta, S. (2013) Palynology, palaeoecology and palaeodepositional environment of Eocene lignites and associated sediments from Matanomadh mine, Kutch Basin, western India. *Journal of the Geological Society of India*, **82**, 236–248.
- Mathur, Y.K. & Chopra, A.S. (1987) Palynofossils from the Cenozoic subsurface sediments of the Bengal Basin, India. *Geoscience journal*, **8**, 109–151.
- Mathur, Y.K. & Chowdhary, L.R. (1977) Palaeoecology of the Kalol Formation, Cambay Basin, India. *Proceedings of the IVth. Colloquium of Indian Micropalaentology and Stratigraphy, Dehradun*, 164–178.
- Mathur, Y.K. & Jain, A.K. (1980) Palynology and age of the Dras Volcanics near Shergol, Ladakh, Jammu and Kashmir, India. *Geoscience journal*, **1**, 55–74.
- Maxwell, A.L. (2001) Holocene monsoon changes inferred from lake sediment pollen and carbonate records, northeastern Cambodia. *Quaternary Research*, **56**, 390–400.
- Mayle, F.E., Burbridge, R. & Killeen, T.J. (2000) Millennial-scale dynamics of southern Amazonian rain forests. *Science*, **290**, 2291–2294.
- Mbesse, C.O. (2013) *La limite Paléocène-Eocène dans le Bassin de Douala Biostratigraphie et essai de reconstitution des paléoenvironnements*. Ph.D. thesis, University of Liège.
- McIntyre, D.J. (1965) Some new pollen species from New Zealand Tertiary deposits. *New Zealand journal of botany*, **3**, 204–214.
- Mebradu, S., Akpedeye, I. & Cnojakpor, O.K. (1985) The stratigraphy and palynology of Owan/Okpanam/Oba tertiary sediments, southern Nigeria. *Geophytology*, **15**, 146–150.
- Medeiros, V.B., Oliveira, P.E., Santos, R.A., Barreto, A.M.F., Oliveira, M.A.T.D. & Pinaya, J.L.D. (2018) New Holocene pollen records from the Brazilian Caatinga. *Anais da Academia Brasileira de Ciências*, **90**, 2011–2023.
- Medus, J. (1975a) Palynology of Tertiary sediments of Southern Senegal. *Pollen et Spores*, **17**, 545–608.
- Medus, J. (1975b) Quelques palynoflores du Tertiaire inférieur du sud de la France et du nord de l'Espagne. *Revista española de micropaleontología*, **7**, 113–126.
- Medus, J. (1982) Restionales pollen in Paleocene sediments of Senegal. *Cahiers de Micropaléontologie*, **2**, 75–80.
- Medus, J. & Paradis, G. (1987) Des palynoflores du Paléocène inférieur dans le Tertiaire des environs d'Abidjan, Côte d'Ivoire. *Géologie Méditerranéenne*, **14**, 73–79.
- Mehrotra, N.C. (2002a) Bombay Basin. *Memoir of the Geological Society of India*, **48**, 27–42.
- Mehrotra, N.C. (2002b) Cambay Basin. *Memoir of the Geological Society of India*, **48**, 13–26.
- Mehrotra, N.C., Berry, C.M., Nautiyal, D.D. & Rawat, R.S. (1995) Palynostratigraphic and source rock evaluation studies on Oligocene-Miocene subsurface sediments of Surat Depression. *Proceedings of Petrotech-95, New Delhi, Technology Trends in Petroleum Industry*, **2**, 43–52.
- Mello, C.L., Sant'Anna, L.G. & Bergqvist, L.P. (2000) The palaeontological site of Fonseca, Minas Gerais state, Brazil (fossil plants of the Tertiary of Brazil). C. Schobbenhaus, D.A. Campos, E.T. Queiroz, M. Winge & M. Berbert-Born, eds., *Sítios Geológicos e Paleontológicos do Brasil*, see <http://www.unb.br/ig/sigep/sitio086/sitio086.htm>, online.
- Meneses, M.E.N.S., Costa, M.L., Enters, D. & Behling, H. (2015) Environmental changes during the last millennium based on multi-proxy palaeoecological records in a savanna-forest mosaic from the northernmost Brazilian Amazon region. *Anais da Academia Brasileira de Ciências*, **87**, 1623–1651.

- Méon, H. (1991) Études sporopolliniques à la limite Crétacé-Tertiaire.: la coupe du Kef (Tunisie nord-occidentale); étude systématique, stratigraphie, paléogéographie et évolution climatique. *Palaeontographica Abteilung B Paläophytologie*, **223**, 107–168.
- Michoux, D. (1985) Palynostratigraphy of the Eocene of Montfort-en-Chalosse (Landes, France). *Revue de Micropaléontologie*, **28**, 138–153.
- Mildenhall, D.C. (1989) Summary of the age and paleoecology of the miocene Manuherikia Group, Central Otago, New Zealand. *Journal of the Royal Society of New Zealand*, **19**, 19–29.
- Mildenhall, D.C. & Pocknall, D.T. (1989) Miocene-Pleistocene spores and pollen from Central Otago, South Island, New Zealand. *New Zealand Geological Survey Paleontological Bulletin*.
- Milne, L.A. (1988) Palynology of a Late Eocene Lignitic sequence from the western margin of the Eucla Basin, Western Australia. *Association of Australasian Palaeontologists, Memoir*, **5**, 285–310.
- Misra, B.K. & Navale, G.K.B. (1990) Panandhro lignite from Kutch (Gujarat), India: Petrological nature, genesis, rank and sedimentation. *Palaeobotanist*, **39**, 236–249.
- Misra, C.M., Kumar, A., Prasad, B., Pundir, B.S. & Saxena, R.K. (1994) A note on the bioevents across the K/T transition in the well Modi-A, Krishna-Godavari basin, India. *Geoscience Journal*, **15**, 33–36.
- Mogollón Daza, A.A. (2015) *Revisión geológica de las arenas pertenecientes a la Formación Burguita del campo Bejucal del distrito Barinas, división Boyacá*. Master's thesis, Tesis en opción al título de Máster en Geología. Mención Prospección y
- Mohr, B. & Fechner, G. (1986) Eine eozäne Mikroflora (Sporomorphae und Dinoflagellaten-Zysten) aus der Südatlas-Randzone westlich Boumalne du Dadès (Marokko). *Berliner Geowissenschaftliche Abhandlungen, Reihe A, Geologie und Paleontologie*, **66**, 381–414.
- Moltzer, J.G. & Binda, P.L. (1981) Micropaleontology and palynology of the middle and upper members of the Shumaysi Formation, Saudi Arabia. *Bulletin Faculty of Earth Science (Jeddah)*, **4**, 57–76.
- Monga, P., Kumar, M. & Joshi, Y. (2015a) Morphological variations and depositional processes of microforaminiferal linings in the early Tertiary sediments of northeastern and northwestern India. *Palaeobotanist*, **64**, 129–138.
- Monga, P., Kumar, M., Prasad, V. & Joshi, Y. (2015b) Palynostratigraphy, palynofacies and depositional environment of a lignite-bearing succession at Surkha Mine, Cambay Basin, north-western India. *Acta Palaeobotanica*, **55**, 183–207.
- Montes, C., Guzman, G., Bayona, G., Cardona, A., Valencia, V. & Jaramillo, C. (2010) Clockwise rotation of the Santa Marta massif and simultaneous Paleogene to Neogene deformation of the Plato-San Jorge and Cesar-Ranchería basins. *Journal of South American Earth Sciences*, **29**, 832–848.
- Montoya, E., Pedra-Mendez, J., Garcia-Falco, E., Gomez-Paccard, M., Giralt, S., Vegas-Vilarrubia, T., Stauffer, F.W. & Rull, V. (2019) Long-term vegetation dynamics of a tropical megadelta: Mid-Holocene palaeoecology of the Orinoco Delta (NE Venezuela). *Quaternary Science Reviews*, **221**, 105874.
- Montoya, E., Rull, V., Nogué, S. & Díaz, W.A. (2009) Paleoecología del Holoceno en la Gran Sabana, SE Venezuela: Análisis preliminar de polen y microcarbones en la Laguna Encantada. *Collectanea Botanica*, **28**, 65–79.
- Montoya, E., Rull, V., Stansell, N.D., Abbott, M.B., Nogué, S., Bird, B.W. & Díaz, W.A. (2011) Forest-savanna-morichal dynamics in relation to fire and human occupation in the southern Gran Sabana (SE Venezuela) during the last millennia. *Quaternary Research*, **76**, 335–344.

- Morley, R.J. (1977) Palynology of Tertiary and Quaternary sediments in southeast Asia. *Proceedings of the Indonesian Petrological Association 6th Annual Convention*, pp. 255–275.
- Morley, R.J. (1998) Palynological evidence for Tertiary plant dispersals in the SE Asian region in relation to plate tectonics and climate. R. Hall & J.D. Halloway, eds., *Biogeography and geological evolution of SE Asia*, Backhuys Publishers Leiden, pp. 211–234.
- Morley, R.J., Bui, V.D., Nguyen, T.T., Kullman, A., Bird, R.T., Nguyen, H.C. *et al.* (2019) High-resolution Palaeogene sequence stratigraphic framework for the Cuu Long Basin, offshore Vietnam, driven by climate change and tectonics, established from sequence biostratigraphy. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **530**, 113–135.
- Morley, R.J. & Morley, H.P. (2010) Neogene Climate History of the Makassar Straits, With Emphasis on the Attaka Region, East Kalimantan, Indonesia. *Neogene climate history of the Makassar Straits with emphasis on the Attaka Field. Proceedings of the 34th Indonesian Petroleum Association Convention & Exhibition, Jakarta, IPA10-G-208*, pp. 1–17.
- Morley, R.J., Morley, H.P. & Restrepo-Pace, P. (2013) Unravelling the tectonically controlled stratigraphy of the West Natuna Basin by means of palaeo-derived mid Tertiary climate changes. *Proceedings of the Indonesian Petroleum Association 29th Annual Convention & Exhibition. Jakarta: Indonesian Petroleum Association*, pp. 264–288.
- Morley, R.J., Swiecicki, T. & Pham, D.T.T. (2011) A sequence stratigraphic framework for the Sunda region, based on integration of biostratigraphic, lithological and seismic data from Nam Con Son Basin, Vietnam. *Proceedings, Indonesian Petroleum Association: Thirty-Fifth Annual Convention & Exhibition*, 1–22.
- Movshovich, E.B. (1975) Fundamentals of elaboration of a general classification for gas and oil traps. *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, **1975**, 89–98.
- Muller, J. (1968) Palynology of the Pedawan and plateau sandstone formations (Cretaceous-Eocene) in Sarawak, Malaysia. *Micropaleontology*, 1–37.
- Muller, J., de Di Giacomo, E. & Van Erve, A.W. (1987) *A palynological zonation for the Cretaceous, Tertiary, and Quaternary of northern South America*. AASP.
- Mürriger, F. & Pflanzl, G. (1955) *Pollenanalytische Datierungen einiger hessischer Braunkohlen*. Hessisches Landesamt für Bodenforschung.
- Nagy, E. (1963) Spores et pollens nouveaux d'une coupe de la Briqueterie d'Eger (Hongrie). *Pollen et Spores*, **5**, 397–412.
- Nagy, E. (1973) Palynological data for the Neogene of Cserehát. *Acta botanica*, **19**, 453–460.
- Nagy, E. (1979a) Changes in paleoenvironment and paleoclimate during the Miocene in Hungary. *Annales Géologiques des Pays Helléniques*, vol. 1979, pp. 879–888.
- Nagy, E. (1979b) Palynological evaluation of the holostratotype of the Egerian. *Acta Biol. Szeged*, **25**, 45–52.
- Nagy, L. & Palfalvy, I. (1963) Az egri téglagyári szelvény ősnóvénytani vizsgálata. *Magy. All. Földt. Int. évi Jel.(1960)*, 223–263.
- Nandi, B. (1972) Some observations on the microflora of middle Siwalik sediments of Mohand (East) Field, Himachal Pradesh. *Seminar Palaeo-palynol. Indian Stratigr Calcutta*, pp. 375–385.
- Nandi, B. (1980) Further contribution on the palynostratigraphy of the Siwalik Group. *Proc 4th Int Palynol Conf Lucknow*, vol. 2, pp. 727–734.

- Nandi, B. (1990) Palynostratigraphy of upper cretaceous sediments, Meghalaya, northeastern India. *Review of palaeobotany and palynology*, **65**, 119–129.
- Nandi, B. & Sharma, R. (1984) Palynology and biostratigraphy of the Boldamgiri Formation, Garo Hills, Meghalaya. *Proceedings of the Symposium on Evolutionary Botany and Biostratigraphy/edited by AK Sharma, GC Mitra, Manju Banerjee; sponsored by University Grants Commission [and] Department of Botany, University of Calcutta, 1979*, Today and Tomorrow's Printers & Publishers, pp. 565–580.
- Ngon, G.F.N., Etame, J., Ntamak-Nida, M.J., Mbesse, C.O., Mbai, J.S., Bayiga, É.C. & Gerard, M. (2016) Geochemical and palaeoenvironmental characteristics of Missole I iron duricrusts of the Douala sub-basin (Western Cameroon). *Comptes Rendus Geoscience*, **348**, 127–137.
- Nickel, B. (1996) Palynofazies und Palynostratigraphie der Pechelbronn Schichten im nördlichen Oberrheintalgraben. *Palaeontographica Abteilung B*, 1–151.
- Nogueira, A.C.R., Silveira, R. & Guimarães, J.T.F. (2013) Neogene–Quaternary sedimentary and paleovegetation history of the eastern Solimões Basin, central Amazon region. *Journal of South American Earth Sciences*, **46**, 89–99.
- Nott, J.F. & Owen, J.A.K. (1992) An Oligocene palynoflora from the middle Shoalhaven catchment NSW and the Tertiary evolution of flora and climate in the southeast Australian highlands. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **95**, 135–151.
- Oboh, F.E. & Morris, L.M.R. (1995) Correlation between Sequoia type pollen and lower Oligocene transgressive deposits in the eastern Gulf Coast. *Palaios*, 371–382.
- Oboh, F.E. & Reeves Morris, L.M. (1994) Early Oligocene palynosequences in the eastern Gulf Coast, USA. *Palynology*, **18**, 213–235.
- Oboh-Ikuenobe, F.E., Obi, C.G. & Jaramillo, C.A. (2005) Lithofacies, palynofacies, and sequence stratigraphy of Palaeogene strata in Southeastern Nigeria. *Journal of African Earth Sciences*, **41**, 79–101.
- Oboh-Ikuenobe, F.E., Spencer, M.K., Campbell, C.E. & Haselwander, R.D. (2012) A portrait of Late Maastriichtian and Paleocene palynoflora and paleoenvironment in the northern Mississippi Embayment, southeastern Missouri. *Palynology*, **36**, 63–79.
- Oboh-Ikuenobe, F.E., Yepes, O. & Gregg, J.M. (1998) Palynostratigraphy, palynofacies, and thermal maturation of Cretaceous–Paleogene sediments from the Côte D'Ivoire–Ghana transform margin. *Proceedings of the Ocean Drilling Program: Scientific Results*, vol. 159, pp. 277–318.
- Ochoa, D., Hoorn, C., Jaramillo, C., Bayona, G., Parra, M. & De la Parra, F. (2012) The final phase of tropical lowland conditions in the axial zone of the Eastern Cordillera of Colombia: Evidence from three palynological records. *Journal of South American Earth Sciences*, **39**, 157–169.
- O'Keefe, J.M.K. (2008) *Paleogene Mirelands of the Upper Mississippi Embayment, Western Kentucky*. Ph.D. thesis, University of Kentucky.
- Okeke, K.K. & Umeji, O.P. (2016) Palynostratigraphy, palynofacies and palaeoenvironment of deposition of Selandian to Aquitanian sediments, southeastern Nigeria. *Journal of African Earth Sciences*, **120**, 102–124.
- Ola, P.S. & Adeoti, B. (2017) Palynomorphs of the First Down-Hole Shale Occurrence in Three Contiguous Basins in Nigeria: Implication on the K-Pg Boundary. *Geosciences Research*, **2**.
- Ola-Buraimo, A.O. (2012) Lithostratigraphy and palynostratigraphy of Tuma-1 well, Bornu Basin, northeastern Nigeria. *Journal of Biological and Chemical Research*, **29**, 206–223.

- Ola-Buraimo, A.O. (2013) Biostratigraphy of section of Murshe-1 well, Bornu Basin, northeastern Nigeria. *British Journal of Applied Science and Technology*, 361–375.
- Oliveira, P.E.D. (1992) *A palynological record of late Quaternary vegetational and climatic change in southeastern Brazil*. Ph.D. thesis, The Ohio State University.
- Ollivier-Pierre, M.F. (1970) *Contribution à l'étude palynologique du niveau sapropélien de la Sennetière en la Bernerie (Loire atlantique)*. Ph.D. thesis, Université de Rennes.
- Ollivier-Pierre, M.F. (1978) Spores and pollen. *Bulletin de la Société Géologique et Minéralogique de Bretagne, Nouvelle série*, **8**, 34–56.
- Ollivier-Pierre, M.F. (1980) Etude palynologique (spores et pollens) de gisements paléogènes du Massif armoricain. Stratigraphie et paléogéographie. *Mémoires de la Société géologique et minéralogique de Bretagne Rennes*, 1–239.
- Ollivier-Pierre, M.F. (1988) France; the Armorician Massif. *Geologisches Jahrbuch Reihe A*, **100**, 299–300.
- Ollivier-Pierre, M.F. (1989) The study of pollen. *Geologie de la France*, **1**, 93–94.
- Ollivier-Pierre, M.F., Chateaufort, J.J., Farjanel, G. & Estéoule-Choux, J. (1985) From marine environment to continental environment, an example: the ypresian laminated clays of la baie de Bourgneuf and of le lac de Grand-Lieu (Vendée). *Sciences Géologiques, Bulletins et Mémoires*, **38**, 45–59.
- Ollivier-Pierre, M.F., Maupin, C., Esteoule-Choux, J. & Sittler, C. (1993) Transgression et paléoenvironnement à l'Oligocène en Bretagne (France). Sédimentologie, micropaléontologie, palynologie et palynofaciès du Rupélien du Bassin de Rennes. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **103**, 223–250.
- Oloto, I.N. (1990) Palynological assemblage from the Danian of South-West Nigeria. *Acta Palaeobotanica*, **30**, 23–30.
- Oloto, I.N., Omoboriwo, A.O. & Soronnadi-Ononiwu, E.C. (2014) Palynology and Paleoenvironmental Studies of Bende-1 Well. Anambra and Niger-Delta Basins, Nigeria. *International Journal of Science Inventions Today*, **3**, 259–270.
- Oloto, I.N. & Promise, W. (2014) Biostratigraphic study and paleo environmental reconstruction of cores from offshore (South Western) Niger Delta, Nigeria. *International Journal of Scientific and Technology Research*, **3**, 279–286.
- Onyekuru, S.O., Iwueke, E.L., Ikoru, D.O., Opara, K.D. & Fagorite, V.I. (2019) The Stratigraphic Significances of Lignite Deposits in Parts of Orlu, Southeastern Nigeria. *International Journal of Innovative Environmental Studies Research*, **7**, 1–11.
- Osokpor, J., Ekwere, U.J. & Otele, A. (2019) Paleoclimatic cycles, sea level history and sequence stratigraphic elements in Eocene–Oligocene sediments of BIMOL-1 well northern Niger Delta Basin, Nigeria. *Journal of Applied Sciences and Environmental Management*, **23**, 241–248.
- Osorio-Granada, E., Pardo-Trujillo, A., Restrepo-Moreno, S.A., Gallego, F., Muñoz, J., Plata, A., Trejos-Tamayo, R., Vallejo, F., Barbosa-Espitia, A., Cardona-Sánchez, F.J. *et al.* (2020) Provenance of Eocene–Oligocene sediments in the San Jacinto Fold Belt: Paleogeographic and geodynamic implications for the northern Andes and the southern Caribbean. *Geosphere*, **16**, 210–228.
- Oyedé, M., De Klasz, S., Land, J., Carbonnel, G. & Grosdidier, E. (1992) Datation de la discordance dite Oligocène sur la partie terrestre du bassin sédimentaire côtier béninois (Afrique de l'Ouest). *Comptes rendus de l'Académie des sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre*, **315**, 971–977.

- Özer, M.S.K., Akiün, F., Mayda, S. & Kaya, T. (2014a) Palynofloras and vertebrates from Muş gla-Ören region (SW Turkey) and palaeoclimate of the Middle Burdigalian Langhian period in Turkey. *Bulletin of Geosciences*, **89**, 1.
- Özer, M.S.K., Sancay, R.H., Şen, Ş., Sözeri, K., Metais, G., Ayyildiz, T. & Varol, B.E. (2017) Paleoenvironment of the Late Oligocene from the Kağızman-Tuzluca Basin (northeastern Anatolia) based on the micro-and macrofloras. *Turkish Journal of Earth Sciences*, **26**, 227–248.
- Özer, M.S.K., Sozbilir, H. & Akgün, F. (2014b) Miocene palynoflora of the Kocaçay and Cumaovasi basins: A contribution to the synthesis of Miocene palynology, palaeoclimate, and palaeovegetation in western Turkey. *Turkish Journal of Earth Sciences*, **23**.
- Palamarev, E., Ivanov, D. & Kithanov, G. (1998) New data about the fossil flora from Bodovdol Basin and its biostratigraphic significance. *Review of the Bulgarian Geological Society*, **59**, 13–21.
- Palma-Heldt, S. (1980) Contribution to the palynological knowledge of the Tertiary Coal Beds of Arauco-Concepcion, Chile. *Actas del Segundo Congreso Argentino de Paleontología y Bioestratigrafía y Primer Congreso Latinoamericano de Paleontología*, **2**, 175–192.
- Pardo-Trujillo, A., Jaramillo, C.A. & Oboh-Ikuenobe, F.E. (2003) Paleogene palynostratigraphy of the eastern middle Magdalena Valley, Colombia. *Palynology*, **27**, 155–178.
- Pardo-Trujillo, A., Jaramillo, C.A. & Rangel, J.O. (2014) Palynology and Paleoenvironments of Eastern Cordillera Paleogene deposits of Colombia: 35 million years of Neotropical vegetation history. *Colombia Diversidad Biológica XIV: La región de la Orinoquia de Colombia*, 1–30.
- Pardo-Trujillo, A. & Roche, E. (2009) *Paleocene-Eocene palynology and palynofacies from northeastern Colombia and western Venezuela*. Cuadernos de Investigación 41, Universidad de Caldas.
- Pardo-Trujillo, A. & Sánchez, C. (2009) Cyclospora en el Cenozoico de Colombia y su supuesto origen marino. *Caldasia*, **31**, 105–121.
- Parra, F.J., Navarrete, R.E., di Pasquo, M.M., Roddaz, M., Calderón, Y. & Baby, P. (2020) Neogene palynostratigraphic zonation of the Marañon Basin, Western Amazonia, Peru. *Palynology*, **44**, 675–695.
- Parra, M., Mora, A., Jaramillo, C., Strecker, M.R., Sobel, E.R., Quiroz, L., Rueda, M. & Torres, V. (2009) Orogenic wedge advance in the northern Andes: Evidence from the Oligocene-Miocene sedimentary record of the Medina Basin, Eastern Cordillera, Colombia. *Geological Society of America Bulletin*, **121**, 780–800.
- Parra, M., Mora, A., Jaramillo, C., Torres, V., Zeilinger, G. & Strecker, M.R. (2010) Tectonic controls on Cenozoic foreland basin development in the north-eastern Andes, Colombia. *Basin Research*, **22**, 874–903.
- Paul, S., Sharma, J., Singh, B.D., Saraswati, P.K. & Dutta, S. (2015) Early Eocene equatorial vegetation and depositional environment: Biomarker and palynological evidences from a lignite-bearing sequence of Cambay Basin, western India. *International Journal of Coal Geology*, **149**, 77–92.
- Pérez, G.S. (1992) Palinología de la formación Guaduas. Estratigráfica y sistemática. *Boletín Geológico*, **32**, 45–126.
- Petrescu, I. & Codrea, V. (2004) The signification of the Paleocene microflora from Jibou (Sălaj District) for the Paleogene palinological heritage of Romania. *Acta Palaeontologica Romaniaae*, **4**, 351–360.
- Petrescu, L. & Givulescu, R. (1986) Flore et végétation de la Vallée du Jiu (Bassin de Petrosani)-Roumanie. *Revue de paléobiologie*, **5**, 109–116.

- Petrosyants, M.A. & Trofimov, D.M. (1971) Spore pollen characteristics of Upper Cretaceous deposits of the Mali-Niger Syncline, (South Sahara). *Byulleten Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Geologicheskii*, **46**, 75–85.
- Petrosyants, M.A. & Trofimov, D.M. (1975) Palynological characteristics of Danian-Paleocene strata in the South-West Sahara (Taudeni and Mali-Nigerian Synclises). *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, 108–115.
- Pflug, H. (1986) Palyno-stratigraphy of the Eocene/Oligocene in the area of Helmstedt. *North Hessa and in the adjacent Southern Range. In: Tobien, A.(ed.) Nordwestdeutschland im Tertiär. Beiträge zur regionalen Geologie der Erde*, **38**, 567–583.
- Phadtare, N.R. & Kulkarni, A.R. (1980) Palynological investigation of Ratnagiri lignite, Maharashtra. *Geophytology*, **10**, 158–170.
- Phadtare, N.R. & Kulkarni, A.R. (1984) Affinity of the genus Quilonipollenites with the Malaysian palm *Eugeissona* Griffith. *Pollen et Spores*, **26**, 217–225.
- Pickett, J.W., Macphail, M.K., Partridge, A.D. & Pole, M.S. (1997) Middle Miocene palaeotopography at Little Bay, near Maroubra, New South Wales. *Australian Journal of Earth Sciences*, **44**, 509–518.
- Piraquive Bermúdez, D. (2016) *Reconstrucción paleoecológica del holoceno medio (5500 AÑOS AP) en ambientes de cananguchales (Mauritia flexuosa L. f.) del municipio de Leticia, Amazonas (Colombia)*. Master's thesis, Pontificia Universidad Javeriana.
- Planderová, E. (1990) Microfloristic changes during the Miocene in Central Paratethys region. *Proceedings of the Symposium on Paleofloristic and Paleoclimatic changes in the Cretaceous and Tertiary. Praha*, pp. 215–222.
- Playford, G. (1982) Neogene palynomorphs from the Huon Peninsula, Papua New Guinea. *Palynology*, **6**, 29–54.
- Pocknall, D.T. (1982) Palynology of late Oligocene Pomahaka Estuarine Bed sediments, Waikoikoi, Southland, New Zealand. *New Zealand journal of botany*, **20**, 263–287.
- Pocknall, D.T., Erlich, R.N., Stein, J.A., Bergen, J.A., Lorente, M.A., Goodman, D.K. & Clarke, R.T. (1996a) The palynofloral succession across the Cretaceous to Paleocene transition zone, Mérida Andes, western Venezuela. *Proceedings of the IX International Palynological Congress, Houston, Texas, USA*, pp. 171–179.
- Pocknall, D.T., Wood, L.J., Geen, A.F., Harry, B.E., Hedlund, R.W., Goodman, D.K. & Clarke, R.T. (1996b) Integrated paleontological studies of Pliocene to Pleistocene deposits of the Orinoco delta, Eastern Venezuela and Trinidad. *Proceedings of the IX International Palynological Congress, Houston, Texas, USA*, pp. 319–326.
- Pole, M.S. & Macphail, M.K. (1996) Eocene *Nypa* from Regatta Point, Tasmania. *Review of Palaeobotany and Palynology*, **92**, 55–67.
- Poliakova, A. & Behling, H. (2016) Pollen and fern spores recorded in recent and late Holocene marine sediments from the Indian Ocean and Java Sea in Indonesia. *Quaternary International*, **392**, 251–314.
- Potonié, R. (1960) Sporologie der eozänen Kohle von Kalewa in Burma. *Senckenbergiana Lethaea*, **41**, 451–481.
- Potonié, R. (1966) Synopsis of the genera of the sporae dispersae. *Part*, **4**, 1–244.
- Potter, J.F. (1977) Eocene, lower Bracklesham Beds, iron workings in Surrey. *Proceedings of the Geologists' Association*, **88**, 229–241.
- Potter, J.F. & Fowler, K. (1986) A Bracklesham-Bagshot Beds unconformity in the London Basin. *Proceedings of the Geologists' Association*, **97**, 87–90.

- Povilauskas, L. (2019) Palynostratigraphy of the Cretaceous–Paleogene in the Austral basin, SW Santa Cruz province, Argentina. *Revista brasileira de paleontologia*, **20**, 299–320.
- Pradidtan, S. (1989) Characteristics and controls of lacustrine deposits of some Tertiary basins in Thailand. *International symposium on intermontane basins: geology and resources*, pp. 133–145.
- Prasad, B. & Dey, A.K. (1986) The occurrence of Eocene sediments in Arunachal Pradesh: a palynological evidence. *Bulletin of the Oil and Natural Gas Corporation (India)*, **23**, 67–74.
- Prasad, B., Jain, A.K. & Mathur, Y.K. (1995) A standard palynozonation scheme for the Cretaceous and pre-Cretaceous subsurface sediments of Krishna-Godavari Basin, India. *Geoscience Journal*, **10**, 155–220.
- Prasad, V., Farooqui, A., Tripathi, S.K.M., Garg, R. & Thakur, B. (2009) Evidence of late Palaeocene-early Eocene equatorial rain forest refugia in southern Western Ghats, India. *Journal of Biosciences*, **34**, 777–797.
- Prasad, V., Garg, R., Khowaja-Ateequzzaman, S.I.B. & Joachimski, M. (2006) Apectodinium acme and the palynofacies characteristics in the latest Palaeocene-earliest Eocene of northeastern India: Biotic response to Palaeocene-Eocene Thermal maxima (PETM) in low latitude. *Journal of the Palaeontological Society of India*, **51**, 75–91.
- Prasad, V., Singh, I.B., Bajpai, S., Garg, R., Thakur, B., Singh, A., Saravanan, N. & Kapur, V.V. (2013) Palynofacies and sedimentology-based high-resolution sequence stratigraphy of the lignite-bearing muddy coastal deposits (early Eocene) in the Vastan Lignite Mine, Gulf of Cambay, India. *Facies*, **59**, 737–761.
- Prasad, V., Utescher, T., Sharma, A., Singh, I.B., Garg, R., Gogoi, B., Srivastava, J., Uddandam, P.R. & Joachimski, M.M. (2018) Low-latitude vegetation and climate dynamics at the Paleocene-Eocene transition—A study based on multiple proxies from the Jathang section in northeastern India. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **497**, 139–156.
- Punyasena, S.W., Mayle, F.E. & McElwain, J.C. (2008) Quantitative estimates of glacial and Holocene temperature and precipitation change in lowland Amazonian Bolivia. *Geology*, **36**, 667–670.
- Quattrocchio, M., Volkheimer, W. & Del Papa, C. (1997) Palynology and paleoenvironment of the “Faja Gris”; Mealla Formation (Salta Group) at Garabatal Creek (NW Argentina). *Palynology*, **21**, 231–247.
- Quattrocchio, M.E., Martínez, M.A., Hinojosa, L.F. & Jaramillo, C. (2013) Quantitative analysis of Cenozoic palynofloras from Patagonia, southern South America. *Palynology*, **37**, 246–258.
- Raine, J.I. & Wilson, G.J. (1988) Palynology of the Mt Somers (South Island, New Zealand) early Cenozoic sequence. *New Zealand journal of geology and geophysics*, **31**, 385–390.
- Raiverman, V. & Misra, V.N. (1975) Suru tectonic axis, Kargil area, Ladakh. *Bull. Geol. Min. Met. Soc. India*, **48**, 1–16.
- Rákosi, L. (1966) Pollen analysis of the sedimentary record of boring szentendre 2. *Magyar Allami Földt. Intéz. Évi Jel*, **1964**, 377–387.
- Rákosi, L. (1973) A Dorogi-medence paleogén képződményeinek palinológiája. *Földt. Int. Évkönyve*, **55**, 497–575.
- Rákosi, L. (1980) The question of the Eocene/Oligocene boundary on the basis of palynological studies. *Discussiones Palaeontologicae*, **25**, 117–125.
- Rákosi, L. & Toth, K. (1980) Contribution à la litho- et biostratigraphie de l'Éocène au Sud de la Montagne Bakony. *Annual Report of the Hungarian Geological Institute*, 239–249.

- Ramadhan, R. (2013) *Studi Sikuenstratigrafi Berdasarkan Hasil Analisis Data Palinologi Pada Sumur Y, Cekungan Sumatera Selatan*. Ph.D. thesis, Diponegoro University.
- Ramanujam, C.G.K. (1966) Palynology of the Miocene lignite from south Arcot district, Madras, India. *Pollen et Spores*, **8**, 149–203.
- Ramanujam, C.G.K. (1982) Tertiary palynology and palynostratigraphy of southern India. *Cenozoic Stratigraphy and Palynology in India. Palaeontol Soc India Spec Publ*, **1**, 57–64.
- Ramanujam, C.G.K. (1987) Palynology of the Neogene Warkalli beds of Kerala state in south India. *Journal of the Palaeontological Society of India*, **32**, 26–46.
- Ramanujam, C.G.K. & Rao, K.P. (1977) Palynological approach to the study of Warkalli deposits of Kerala in south India. *Geophytology*, **7**, 160–164.
- Ramanujam, C.G.K. & Reddy, P.R. (1984) Palynoflora of Neyveli lignite. Floristic and palaeoenvironmental analysis. *Journal of palynology*, **20**, 58–74.
- Ramanujam, C.G.K., Reddy, P.R. & Ramakrishna, H. (1997) Dicolpate palm pollen from the Neogene deposits of Godavari-Krishna Basin, AP. *Journal of Palynology*, **33**, 129–136.
- Ramanujam, C.G.K., Reddy, P.R. & Rao, G.M. (1991) Palynoassemblages of the subsurface Tertiary strata at Pattanakad, Alleppey District, Kerala State. *Journal of the Palaeontological Society of India*, **36**, 51–58.
- Ramanujam, C.G.K., Siva, S, P. & Reddy, P.R. (1984) Quantification of the palynoassemblages of the first and second mine areas of Neyvelli lignite. *Indian colloquium on micropaleontology and stratigraphy*, pp. 269–276.
- Rao, K.P. (1974) *Palynology of the Tertiary deposits of Kerala State, South India*. Ph.D. thesis, Osmania University.
- Rao, K.P. & Ramanujam, C.G.K. (1975) A palynological approach to the study of Quilon Beds of Kerala State in South India. *Current Science*, 730–732.
- Rao, K.P. & Ramanujam, C.G.K. (1978) Palynology of the Neogene Quilon Beds of Kerala State in South India. I. Spores of pteridophytes and pollen of Monocotyledons. *Palaeobotanist*, **25**, 397–427.
- Rao, M.R. (1989) Palynological investigation of Arthungal bore-hole, Alleppey district, Kerala. *Palaeobotanist*, **38**, 243–255.
- Rao, M.R. (1995) Palynostratigraphic zonation and correlation of the Eocene-Early Miocene sequence in Alleppey district, Kerala, India. *Review of Palaeobotany and Palynology*, **86**, 325–348.
- Rao, M.R. (2000) Palynological investigation of the Kherapara Formation (Oligocene) exposed along Tura-Dalu road near Kherapara, west Garo Hills District, Meghalaya, India. *Palaeobotanist*, **49**, 293–309.
- Rao, M.R. & Nair, K.K. (1998) Palynological investigation of Miocene sediments exposed in Kannanellur-Kundara area, Quilon District, Kerala. *Geophytology*, **27**, 49–59.
- Rao, M.R. & Rajendran, C.P. (1996) Palynological investigations of Tertiary Lignite and associated sediments from Cannanore, Kerala Basin, India. *Palaeobotanist*, **43**, 63–82.
- Rao, M.R., Sahni, A., Rana, R.S. & Verma, P. (2013) Palynostratigraphy and depositional environment of Vastan lignite mine (Early Eocene), Gujarat, western India. *Journal of Earth System Science*, **122**, 289–307.
- Rao, M.R. & Verma, P. (2014) Palynological investigation of Neogene (Early Miocene) sediments of Mangalore basin, India: Palaeoenvironmental and palaeoclimatic implications. *Journal of the Geological Society of India*, **84**, 55–67.

- Räsänen, M.E., Gingras, M.K., Vieira Jr, V., Barberi, M., Irion, G. *et al.* (2006) Sedimentology and ichnology of tide-influenced Late Miocene successions in western Amazonia: The gradational transition between the Pebas and Nauta formations. *Journal of South American Earth Sciences*, **21**, 96–119.
- Ratanasthien, B. (1989) Depositional environment of Mae Lamao Basin as indicated by palynology and coal petrography. *International symposium on intermontane basins: geology and resources*, pp. 205–215.
- Rawat, M.S., Mukherjee, J. & Venkatachala, B.S. (1977) Palynology of the Kadi formation, Cambay basin, India. *Proceedings of the IV Colloquium on Indian Micropaleontology and Stratigraphy. Dehra Dun, India*, pp. 179–192.
- Rawat, R.S., Misra, C.M., Sharma, J. & Roy, H. (1996) Palynostratigraphy and paleoenvironmental interpretation of Basal Clastic (Panna Formation) Sequence of Mahim area, Bombay Offshore. J. Pandey, R.J. Azmi, A. Bhandari & A. Dave, eds., *Contributions to XV Indian Colloquium on Micropaleontology and Stratigraphy, XV Indian Colloquium on Micropaleontology and Stratigraphy*, pp. 657–662.
- Raymer, J.D. (2010) *Cretaceous/Paleogene boundary biostratigraphy and palynofacies of the Alo-1 well, South-eastern Nigeria*. Master's thesis, Missouri University of Science and Technology.
- Raymer, J.D., Francisca, E. & Jaramillo, C. (2008) Palynology of Cretaceous/Tertiary Boundary Sediments in West Africa. *2008 Joint Meeting of The Geological Society of America, Soil Science Society of America, American Society of Agronomy, Crop Science Society of America, Gulf Coast Association of Geological Societies with the Gulf Coast Section of SEPM*.
- Raymond, A., Phillips, M.K. & Gennett, J.A. (1995) Palynology and paleoecology of lignites from the Manning Formation (Jackson Group) outcrop in the Lake Somerville Spillway. P.D. Warwick & S.S. Crowley, eds., *Coal Geology of the Paleocene–Eocene Calvert Bluff Formation (Wilcox Group) and the Eocene Manning Formation (Jackson Group) in East-Central Texas, U.S. Geological Survey Open-File Report (1995)*, U.S. Geological Survey, pp. 95–595.
- Rebata-H, L.A., Gingras, M.K., Raesaenen, M.E. & Barberi, M. (2006) Tidal-channel deposits on a delta plain from the Upper Miocene Nauta Formation, Marañón Foreland Sub-basin, Peru. *Sedimentology*, **53**, 971–1013.
- Regali, M.D.S.P. (1974) Palinologia dos sedimentos Meso-Cenozoicos do Brasil. *Boletim Técnico Petrobrás*, **17**, 263–301.
- Reimann, K.U. & Thaug, A. (1981) Results of palynostratigraphical investigations of the Tertiary sequence in the Chindwin Basin/Northwestern Burma. *Proceedings of IV International Palynological Conference. Lucknow: Birbal Sahni Institute of Palaeobotany*, pp. 380–395.
- Reis, L.S., Guimarães, J.T.F., Souza-Filho, P.W.M., Sahoo, P.K., de Figueiredo, M.M.J.C., de Souza, E.B. & Giannini, T.C. (2017) Environmental and vegetation changes in southeastern Amazonia during the late Pleistocene and Holocene. *Quaternary International*, **449**, 83–105.
- René, B.G., Bruno, D.Z., Raphael, Y.K., Juliette, T.Y., David, K.K. & Ignacec, T. (2012) Stratigraphie Palynologique du Maastrichtien Supérieur-Eocène Supérieur du Bassin Sédimentaire Offshore de Côte d'Ivoire, Afrique de l'Ouest. *International Journal of African Studies*, **6**, 40–57.
- Richards, K. (2001) Palynology and sequence stratigraphy of early Miocene-late Oligocene deltaics in the Sihapas group, central Sumatra, Indonesia. *Proceedings of the 11th International Palynological Congress, American Association of Stratigraphic Palynologist Foundation, Houston, Texas, USA*, pp. 253–262.
- Riegel, W., Lenz, O.K. & Wilde, V. (2015) From open estuary to meandering river in a greenhouse world: an ecological case study from the middle Eocene of Helmstedt, northern Germany. *Palaios*, **30**, 304–326.

- Riegel, W., Wilde, V. & Lenz, O.K. (2012) The early Eocene of Schöningen (N-Germany)—an interim report. *Austrian Journal of Earth Sciences*, **105**, 88–109.
- Riveline, J. (1985) Age and characteristics of the Eocene transgression at Gánt (Vértés Mts, Transdanubia, Hungary). *Acta Geologica Hungarica*, **28**, 29–48.
- Roche, E. (1973) Stratigraphic markers (pollen and spores) of the Paleocene and Lower Eocene of Belgium. *Bulletin de la classe des sciences*, **59**, 956–969.
- Roche, E. (1974) Paléobotanique, paléoclimatologie et dérive des continents. *Sciences Géologiques, bulletins et mémoires*, **27**, 9–24.
- Roche, E. (1982) Palynological study (pollen and spores) of the Eocene of Belgium. *Service Géologique de Belgique, Professional Paper*, 1–60.
- Roche, E. (1988) Belgium. *Geologisches Jahrbuch Reihe A*, **100**, 302–304.
- Roche, E. & Schuler, M. (1976) Palynological analysis (pollen and spores) of different levels of the Belgian Tertiary. Stratigraphic and paleoecological interpretation. *Service Géologique de Belgique, Professional Paper*, 1–57.
- Roche, E. & Schuler, M. (1979) Palynological analysis of the Boom Clay (Belgium). *Service Géologique de Belgique, Professional Paper*, **7**, 1–20.
- Roche, E. & Schuler, M. (1980) Palynological study of "Kallo Complex". *Service Géologique de Belgique, Professional Paper*, **8**, 1–13.
- Roche, E. *et al.* (1973) Etude palynologique des couches yprésiennes du sondage de Kallo. *Bulletin de la Société belge de géologie, de paléontologie et d'hydrologie*, **82**, 487–495.
- Roddaz, M., Hermoza, W., Mora, A., Baby, P., Parra, M., Christophoul, F., Brusset, S. & Espurt, N. (2010) Cenozoic sedimentary evolution of the Amazonian foreland basin system. C. Hoorn & F. Wesselingh, eds., *Amazonia, Landscape and Species Evolution: A Look into the Past, 1st edition.*, Wiley-Blackwell, Chichester, U.K., pp. 61–88.
- Rodríguez-Forero, G., Oboh-Ikuenobe, F.E., Jaramillo-Munoz, C., Rueda-Serrano, M.J. & Cadena-Rueda, E. (2012) Palynology of the Eocene Esmeraldas Formation, Middle Magdalena Valley Basin, Colombia. *Palynology*, **36**, 96–111.
- Rodríguez-Zorro, P.A., da Costa, M.L. & Behling, H. (2017) Mid-Holocene vegetation dynamics with an early expansion of *Mauritia flexuosa* palm trees inferred from the Serra do Tepequém in the savannas of Roraima State in Amazonia, northwestern Brazil. *Vegetation History and Archaeobotany*, **26**, 455–468.
- Rodríguez-Zorro, P.A., Turcq, B., Cordeiro, R.C., Moreira, L.S., Costa, R.L., McMichael, C.H. & Behling, H. (2018) Forest stability during the early and late Holocene in the igapó floodplains of the Rio Negro, northwestern Brazil. *Quaternary Research*, **89**, 75–89.
- Roman, S. & Ticleanu, N. (1983) Considérations sur la flore du Badénien de la Roumanie. *Dări de seamă ale sedintelor. Institutul de geologie si geofizică. 3. Paleontologie*, **70**, 207–217.
- Roslim, A., Hofmann, C.C., Kocsis, L., Briguglio, A. & Goeting, S. (2019) Preliminary palynological results from onshore sediments of Brunei Darussalam. *Geophysical Research Abstracts*, vol. 21, p. 1.
- Rossetti, D.F., Gribel, R., Rennó, C.D., Cohen, M.C., Moulatlet, G.M., de Oliveira Cordeiro, C.L. & Rodrigues, E.d.S.F. (2017) Late Holocene tectonic influence on hydrology and vegetation patterns in a northern Amazonian megafan. *Catena*, **158**, 121–130.

- Roucoux, K.H., Lawson, I.T., Jones, T.D., Baker, T.R., Coronado, E.N.H., Gosling, W.D. & Lahteenoja, O. (2013) Vegetation development in an Amazonian peatland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **374**, 242–255.
- Rugmai, W., Grote, P.J., Chonglakmani, C., Zetter, R. & Ferguson, D.K. (2008) A Late Pleistocene palynoflora from the coastal area of Songkhla Lake, southern Thailand. *Science Asia*, **34**, 137–145.
- Rull, V. (1992) Successional patterns of the Gran Sabana (southeastern Venezuela) vegetation during the last 5000 years, and its responses to climatic fluctuations and fire. *Journal of Biogeography*, **19**, 329–338.
- Rull, V. (1997a) Oligo-Miocene palynology of the Rio Chama sequence (Western Venezuela), with comments on fossil algae as paleoenvironmental indicators. *Palynology*, **21**, 213–229.
- Rull, V. (1997b) Sequence analysis of western Venezuelan Cretaceous to Eocene sediments using palynology: Chronopaleoenvironmental and paleovegetational approaches. *Palynology*, **21**, 79–90.
- Rull, V. (1998a) Biogeographical and evolutionary considerations of *Mauritia* (Arecaceae), based on palynological evidence. *Review of Palaeobotany and Palynology*, **100**, 109–122.
- Rull, V. (1998b) Middle Eocene mangroves and vegetation changes in the Maracaibo Basin, Venezuela. *Palaios*, **13**, 287–296.
- Rull, V. (1999a) Palaeofloristic and palaeovegetational changes across the Paleocene/Eocene boundary in northern South America. *Review of Palaeobotany and Palynology*, **107**, 83–95.
- Rull, V. (1999b) A palynological record of a secondary succession after fire in the Gran Sabana, Venezuela. *Journal of Quaternary Science*, **14**, 137–152.
- Rull, V. (2000) Ecostratigraphic study of Paleocene and Early Eocene palynological cyclicity in northern South America. *Palaios*, **15**, 14–24.
- Rull, V. (2001a) A morphometric study of early Miocene *Mauritiidites* from Northern South America. *Grana*, **40**, 163–167.
- Rull, V. (2001b) A quantitative palynological record from the early Miocene of western Venezuela, with emphasis on mangroves. *Palynology*, **25**, 109–126.
- Rull, V. (2003) Contribution of quantitative ecological methods to the interpretation of stratigraphically homogeneous pre-quaternary sediments: A palynological example from the oligocene of Venezuela. *Palynology*, **27**, 75–98.
- Rull, V. (2007) Holocene global warming and the origin of the Neotropical Gran Sabana in the Venezuelan Guayana. *Journal of Biogeography*, **34**, 279–288.
- Russo Ermolli, E. (1991) Datation palynologique de gisements tertiaires de l'entre Sambre-et-Meuse. Essai de reconstitution des paleoenvironnements et des paleoclimats. *Professional paper-Service geologique de Belgique*.
- Sa, N.d.P. & Carvalho, M.d.A. (2017) Miocene fern spores and pollen grains from the Solimoes Basin, Amazon Region, Brazil. *Acta Botanica Brasiliica*, **31**, 720–735.
- Sabiham, S. & Furukawa, H. (1986) A study of floral composition of peat soil in the lower batang hari river basin of Jambi, Sumatra. *Japanese Journal of Southeast Asian Studies*, **24**, 113–132.
- Sahay, V.K. (2011) The hydrocarbon potential, thermal maturity, sequence stratigraphic setting and depositional palaeoenvironment of carbonaceous shale and lignite successions of Panandhro, northwestern Kutch Basin, Gujarat, Western India. *Central European Journal of Geosciences*, 1–17.

- Sahay, V.K., Samant, B. & Mude, S.N. (2016) Paleodepth and paleodepositional environment of green shale of Gypseous Shale Member (Naredi Formation), Kutch, western India. *Indian Journal of Geosciences*, **70**, 153–160.
- Salamanca Villegas, S., van Soelen, E.E., Teunissen van Manen, M.L., Flantua, S.G.A., Santos, R.V., Roddaz, M., Dantas, E.L., van Loon, E., Sinninghe Damsté, J.S., Kim, J.H. *et al.* (2016) Amazon forest dynamics under changing abiotic conditions in the early Miocene (Colombian Amazonia). *Journal of Biogeography*, **43**, 2424–2437.
- Salami, M.B. (1982) Palynological evidence of paleoenvironment of the Araromi Shale of Southwestern Nigeria. *Acta Delivsim Simposio Palinologia*, pp. 363–385.
- Salard-Cheboldaeff, M. (1979a) Maastrichtian and Tertiary Palynology of Cameroon—Qualitative study and range of the main species. *Review of Palaeobotany and Palynology*, **28**, 365–387.
- Salard-Cheboldaeff, M. (1979b) Palynologie Maestrichtienne et Tertiaire du Cameroun. Etude qualitative et repartition verticale des principales especes. *Review of Palaeobotany and Palynology*, **28**, 365–388.
- Salard-Cheboldaeff, M. (1981) Maastrichtian and Tertiary palynology of Cameroon. Botanical results. *Review of palaeobotany and palynology*, **32**, 401–439.
- Salard-Cheboldaeff, M. (1990) Intertropical African palynostratigraphy from Cretaceous to late Quaternary times. *Journal of African Earth Sciences (and the Middle East)*, **11**, 1–IN6.
- Salard-Cheboldaeff, M., Miko-Angoula, I. & Lang, J. (1995) Approche palynologique des paléoenvironnements du continental terminal dans le Bassin des Iullemmeden (Niger). *Revue de Micropaléontologie*, **38**, 329–338.
- Salazar, M., Moscardelli, L., Fisher, W. & Lorente, M.A. (2011) Tectonostratigraphic evolution of the Morichito piggyback basin, Eastern Venezuelan Basin. *Marine and Petroleum Geology*, **28**, 109–125.
- Salgado-Labouriau, M.L., Barberi, M., Ferraz-Vicentini, K.R. & Parizzi, M.G. (1998) A dry climatic event during the late Quaternary of tropical Brazil. *Review of Palaeobotany and Palynology*, **99**, 115–129.
- Salgado-Labouriau, M.L., Casseti, V., Ferraz-Vicentini, K.R., Martin, L., Soubiès, F., Suguio, K. & Turcq, B. (1997) Late Quaternary vegetational and climatic changes in cerrado and palm swamp from Central Brazil. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **128**, 215–226.
- Salujha, S.K., Basavaraju, M.H. & Kingra, G.S. (1990) Palynological study of Tertiary sediments met within the Well Adamtila-A (Cachar) with remarks on age and palaeoenvironment. *Bulletin of the Oil and Natural Gas Commission, Dehradun, India*, **27**, 63–78.
- Salujha, S.K. & Kindra, G.S. (1984) Palynostratigraphy of the Atharmura anticline, Tripura. *Indian colloquium on micropaleontology and stratigraphy*, pp. 391–410.
- Salujha, S.K. & Kindra, G.S. (1990) Palynostratigraphy of the well Baramura-2, Tripura (India). *Geoscience journal*, **11**, 19–30.
- Salujha, S.K., Kindra, G.S. & Rehman, K. (1972) Palynology of the South Shillong Front, Part 1. Seminar on Paleopalynology and Indian Stratigraphy. *Proceedings, University of Calcutta, Calcutta, India*, 265–291.
- Salujha, S.K., Kindra, G.S. *et al.* (1978) Palynostratigraphy of Tertiary sediments of the Gojalia Anticline, Tripura. Part 1: Discussion. *Indian Colloquium on Micropalaeontology and Stratigraphy, Proceedings, 6th*, **6**, 121–126.
- Salujha, S.K., Rehman, K. & Kindra, D.S. (1973) Distinction between the Bhuban and Bokabil sediments on the southern edge of shilling Plateau based on palynoassemblage. *Bulletin of the ONGC*, 109–117.

- Samant, B. (1994) Age of the Bhaunagar lignite deposit of Gujarat. *Journal of Palynology*, **30**, 49.
- Samant, B. & Phadtare, N.R. (1997) Stratigraphic palynoflora of the Early Eocene Rajpardi lignite, Gujarat and the lower age limit of the Tarkeswar Formation of South Cambay Basin, India. *Palaeontographica Abteilung B*, 1–108.
- Sanabria, L.L.O., Pereira, E. & Antonioli, L. (2015) Revisão estratigráfica dos depósitos miocênicos aflorantes na região do alto de Olivença, litoral Sul do Estado da Bahia, Brasil. *Revista Brasileira de Paleontologia*, **18**, 475–488.
- Sancay, R.H., Zühtü, B., Uğraş, I., Kirici, S. & Nihal, A. (2006) Palynomorph, foraminifera, and calcareous nannoplankton biostratigraphy of Oligo–Miocene sediments in the Muş Basin, Eastern Anatolia, Turkey. *Turkish Journal of Earth Sciences*, **15**, 259–319.
- Sánchez, F., Fernández, J., Gassón, R., Bezada, M. & Leal, A. (2017) Paleoecología y ocupación humana durante el Holoceno en los Llanos del Orinoco: una revisión y nuevos datos. *BioLlania (Special Edition)*, **15**, 297–333.
- Sarmiento, G. (1991) Palinología de la Formación Guaduas– estratigrafía y sistemática. *Boletín Geológico Ingeominas*, **32**, 45–126.
- Saxena, R.K. (1986) Significance of spores and pollen grains in Palaeocene biostratigraphy. *Proceedings of the Symposium on Palaeocene of India: limits and subdivisions*, pp. 68–82.
- Saxena, R.K. (1991) Neyveli lignites and associated sediments—their palynology, palaeoecology, correlation and age. *Palaeobotanist*, **40**, 345–353.
- Saxena, R.K., Khare, S. & Misra, N.K. (1990) Echimoporoopollis, a new pollen genus from the neyveli formation of Jayamkonda Cholapuram well-12 Tiruchirapalli District, Tamil Nadu. *Palaeobotanist*, **30**, 46–49.
- Saxena, R.K. & Misra, N.K. (1989) Palynological investigation of the Ratnagiri beds of Sindhu Durg district, Maharashtra. *Palaeobotanist*, **38**, 263–276.
- Saxena, R.K. & Rao, M.R. (1996) Palynological investigation of the Boldamgiri Formation (Early Miocene) in type area, Garo Hills, Meghalaya. *Geophytology*, **26**, 43–56.
- Saxena, R.K., Tripathi, S.K.M. & Prasad, V. (1996) Palynofloral investigation of the Tura Formation (Palaeocene) in Nongwal Bibra area, East Garo Hills, Meghalaya. *Geophytology*, **26**, 19–31.
- Saxena, R.K. & Trivedi, G.K. (2009) Palynological investigation of the Kopili Formation (Late Eocene) in North Cachar Hills, Assam, India. *Acta Palaeobotanica*, **49**, 253–277.
- Scafati, L., Melendi, D.L. & Volkheimer, W. (2009) A Danian subtropical lacustrine palynobiota from South America (Bororó Formation, San Jorge Basin, Patagonia-Argentina). *Geologica Acta: an international earth science journal*, **7**, 35–62.
- Schiferl, J.D., Bush, M.B., Silman, M.R. & Urrego, D.H. (2018) Vegetation responses to late Holocene climate changes in an Andean forest. *Quaternary Research*, **89**, 60–74.
- Schuler, M. (1983) Pollens et spores des séries salifères (Eocène/Oligocène) du bassin potassique de Mulhouse (France). Relations entre la microflore et les dépôts évaporitiques. *Sciences Géologiques, bulletins et mémoires*, **36**, 255–265.
- Schuler, M. (1990) Environnements et paléoclimats paléogènes. Palynologie et biostratigraphie de l'Eocène et de l'Oligocène inférieur dans les fossés rhénan, rhodanien et de Hesse. *Documents-BRGM*.

- Schuler, M. & Doubinger, J. (1970) Palynological studies in the Amaga Basin (Colombia). *Pollen et spores*, **12**, 429–450.
- Shanmukhappa, M. & Uniyal, S.N. (2008) Biostratigraphy and palaeoenvironmental analysis of subsurface Palaeogene sediments in western part of Ahmedabad Block, North Cambay Basin. *Palaeobotanist*, **57**, 465–477.
- Sharma, J. & Saraswati, P.K. (2015) Lignites of Kutch, western India: Dinoflagellate biostratigraphy and palaeoclimate. *Revue de micropaléontologie*, **58**, 107–119.
- Sharma, P. (2000) On the presence of Late Palaeocene in the subsurface of Bikaner District, Rajasthan. *Geophytology*, **28**, 51–55.
- Sharma, R. & Nandi, B. (1982) A preliminary note on the floral evaluation in the Palaeogene of Garo Hills, Meghalaya. *Science & Culture*, **48**, 254–255.
- Shaw, C.L. (2000) Eocene angiospermous palynomorphs of Taiwan. *Taiwania*, **45**, 167–180.
- Shivanna, M. & Singh, H. (2016) Depositional environment and hydrocarbon potential of marginal marine sediments of Eocene from western India: A palynofacies perspective. *Marine and Petroleum Geology*, **73**, 311–321.
- Sia, S.G., Abdullah, W.H., Konjing, Z. & John, J. (2019) Floristic and climatic changes at the Balingian Province of the Sarawak Basin, Malaysia, in response to Neogene global cooling, aridification and grassland expansion. *Catena*, **173**, 445–455.
- Sia, S.G., Abdullah, W.H., Konjing, Z. & Koraini, A.M. (2014) The age, palaeoclimate, palaeovegetation, coal seam architecture/mire types, paleodepositional environments and thermal maturity of syn-collision paralic coal from Mukah, Sarawak, Malaysia. *Journal of Asian Earth Sciences*, **81**, 1–19.
- Silva, M.N.A., Cohen, M.C.L., Rossetti, D.F. & Pessenda, L.C.R. (2018) Did sea-level changes affect the Brazilian Amazon forest during the Holocene? *Radiocarbon*, **60**, 91–112.
- Silveira, R.R. & Souza, P.A. (2015) Palinologia (grãos de pólen de angiospermas) das formações Solimões e Içá (bacia do Solimões), nas regiões de Coari e Alto Solimões, Amazonas. *Revista Brasileira de Paleontologia*, **18**, 455–474.
- Silveira, R.R.d. (2005) *Cronoestratigrafia e interpretação paleoambiental de depósitos Miocenos da Formação Solimões, região de Coari, AM*. Master's thesis, Universidade Federal do Amazonas.
- Simon, P., Caratini, C., Charpy, N. & Tissot, C. (1984) Sédimentologie et palynologie du Crétacé terminal et du Tertiaire de la région de Bonoua (Côte d'Ivoire). *Géologie Méditerranéenne*, **11**, 77–85.
- Singh, A., Mahesh, S., Singh, H., Tripathi, S.K.M. & Singh, B.D. (2013) Characterization of Mangrol lignite (Gujarat), India: petrography, palynology, and palynofacies. *International journal of coal geology*, **120**, 82–94.
- Singh, A., Shivanna, M., Mathews, R.P., Singh, B.D., Singh, H., Singh, V.P. & Dutta, S. (2017) Paleoenvironment of Eocene lignite bearing succession from Bikaner-Nagaur Basin, western India: organic petrography, palynology, palynofacies and geochemistry. *International Journal of Coal Geology*, **181**, 87–102.
- Singh, H. (2015) Palynofloral investigation of the Akli Formation (Palaeocene) of Giral lignite mine, Barmer district, Rajasthan. *Geophytology*, **45**, 209–214.
- Singh, H. & Mahesh, S. (2015) Palynofacies characterization for evaluation of hydrocarbon source rock potential of Lower Paleogene (Thanetian-Ypresian) sub-surface sediments of Barmer Basin, western Rajasthan, India. *Marine and Petroleum Geology*, **59**, 442–450.

- Singh, R.S. (1989) Palynology of Langrin Coalfield, South Shillong Plateau, Meghalaya. *Palaeobotanist*, **38**, 217–228.
- Singh, R.S. & Kar, R. (2003) Palynological assemblage from Deccan intertrappean Bed, Lalitpur, Uttar Pradesh, India. *Gondwana Geological Magazine*, **6**, 217–223.
- Singh, R.S. & Kar, R.K. (2002) Palaeocene palynofossils from the Lalitpur intertrappean beds, Uttar Pradesh, India. *Journal of the Geological Society of India*, **60**, 213–216.
- Singh, R.Y., Dogra, N.N. & Modak, S.K. (1986) Palynostratigraphy of the Oligocene sediments of Namdang River, Assam. *Journal of palynology*, **22**, 105–124.
- Singh, R.Y., Dogra, N.N. & Vimal, K.P. (1985) Palynology of the Barail sediments in the states of Assam and Meghalaya, India. *Journal of palynology*, **21**, 28–55.
- Singh, T. & Tripathi, S. (1989) Siwalik sediments of Arunachal Himalaya: Palynology, palaeoecology and palaeogeography. *Palaeobotanist*, **38**, 325–332.
- Sinha, R.N. (1974) Stratigraphy of the Tertiary sediments exposed in the Surma Valley, Assam-Tripura State, India. *Proc. 4th Colloq. Micropal. & Strat.*, pp. 237–249.
- Sittler, C. (1984) An attempt at a palynological zonation of Paleogene deposits of Basins of Tributaries of the Rhone Valley and South Mediterranean. *Geologie de la France*, 85–90.
- Slansky, M., Bassompierre, P., Roux, J.C., Sangnier, P., Cavelier, C., Faure, A., Sommer, F., Toureno, J., Trauth, N., Monciardini, C., Chateauneuf, J.J. & Pomerol, C. (1971) Étude géologique détaillée du Crétacé supérieur et du Paléogène dans le sondage des Hogues (Eure). *Bulletin du B.R.G.M. (deuxieme serie) Section I*, **1**, 1–24.
- Smith, C.B., Cohen, M.C.L., Pessenda, L.C.R., França, M.C., Guimarães, J.T.F., de Fátima Rossetti, D. & Lara, R.J. (2011) Holocene coastal vegetation changes at the mouth of the Amazon River. *Review of palaeobotany and palynology*, **168**, 21–30.
- Snopková, P. (1984) Paleogene sporomorphs from West Carpathians. *Zapadne Karpaty, Serie Paleontologia*, **5**, 7–74.
- Soares, E.A.A., D'Apolito, C., Jaramillo, C., Harrington, G., Caputo, M.V., Barbosa, R.O., dos Santos, E.B., Dino, R. & Gonçalves, A.D. (2017) Sedimentology and Palynostratigraphy of a Pliocene-Pleistocene (Piacenzian to Gelasian) deposit in the lower Negro River: Implications for the establishment of large rivers in Central Amazonia. *Journal of South American Earth Sciences*, **79**, 215–229.
- Soares, E.A.A., Dino, R., Soares, D.P., Antonioli, L. & Silva, M.A.L.d. (2015) New sedimentological and palynological data from surface Miocene strata in the central Amazonas Basin area. *Brazilian Journal of Geology*, **45**, 337–357.
- Song, Z., Wang, W. & Huang, F. (2004) Fossil pollen records of extant angiosperms in China. *The Botanical Review*, **70**, 425–458.
- Sowunmi, M.A. (1981) Aspects of late Quaternary vegetational changes in West Africa. *Journal of Biogeography*, 457–474.
- Srivastava, J. & Prasad, V. (2015) Effect of global warming on diversity pattern in *Nypa* mangroves across Paleocene–Eocene transition in the paleo-equatorial region of the Indian sub-continent. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **429**, 1–12.
- Srivastava, S.K. & Binda, P.L. (1991) Depositional history of the early Eocene Shumaysi Formation, Saudi Arabia. *Palynology*, **15**, 47–61.

- Stead, D.T. & Awad, M.Z. (2005) Palynological zonation of Cenozoic non-marine sediments, Muglad Basin, Sudan. *Recent Developments in Applied Biostratigraphy. The Micropalaeontological Society, Special Publications. Bath (UK): Geological Society Publishing House*, 161–178.
- Stoian, L.M. (2002) Palynological analyses and dating of eight core samples from Caroline 1 well, Gambier–Otway Basins, South Australia. *Department for Primary Industries and Resources, Report Book*, **2002**, 023.
- Stover, L.E. & Evans, P.R. (1973) Upper Cretaceous-Eocene spore-pollen zonation, offshore Gippsland Basin. *Geological Society of Australia Special Publication*, **4**, 55–72.
- Stover, L.E. & Partridge, A.D. (1973) Tertiary and Late Cretaceous spores and pollen from the Gippsland Basin, southeastern Australia. *Proceedings of the Royal Society of Victoria*, **85**, 237–286.
- Suedy, S.W.A., Muhadiono, M., Sabiham, S. & Qoyim, I. (2012) Fosil Polen Mangrove Berumur Pliosen Dari Formasi Tapak Daerah Kedung Randu, Banyumas. *Bioma: Berkala Ilmiah Biologi*, **14**, 17–24.
- Sun, X., Li, M., Zhang, Y., Lei, Z., Kong, Z., Li, P., Ou, C. & Liu, Q. (1981) Pollen and Spores. Y. Hou, ed., *Tertiary Paleontology of North Continental Shelf of South China Sea*, Guangdong Science and Technology Press, pp. 1–58.
- Sun, X., Luo, Y., Huang, F., Tian, J. & Wang, P. (2003) Deep-sea pollen from the South China Sea: Pleistocene indicators of East Asian monsoon. *Marine Geology*, **201**, 97–118.
- Sunardi, E. (2014) Studi Sedimentasi akhir Kuartar Sub-Cekungan Leles, Garut. *Jatinangor: Seminar Nasional Ke-II FTG Universitas Padjadjaran*, pp. 95–98.
- Sunardi, E. (2015) The Lithofacies Association of Brown Shales In Kiliran Jao Subbasin, West Sumatra Indonesia. *Indonesian Journal on Geoscience*, **2**, 77–90.
- Suyono, S. (2013) Stratigraphy and Tectonics of the East Ketungau Basin, West Kalimantan during Palaeogene. *Indonesian Journal on Geoscience*, **8**, 205–214.
- Swindles, G.T., Kelly, T.J., Roucoux, K.H. & Lawson, I.T. (2018a) Response of testate amoebae to a late Holocene ecosystem shift in an Amazonian peatland. *European Journal of Protistology*, **64**, 13–19.
- Swindles, G.T., Morris, P.J., Whitney, B., Galloway, J.M., Gałka, M., Gallego-Sala, A., Macumber, A.L., Mullan, D., Smith, M.W., Amesbury, M.J. *et al.* (2018b) Ecosystem state shifts during long-term development of an Amazonian peatland. *Global Change Biology*, **24**, 738–757.
- Tabara, D. (2006) Contributions to the palynological study of Bârnova-Muntele Formation (Moldavian Platform). *Analele Stiintifice de Universitatii AI Cuza din Iasi. Sect. 2, Geologie*, **52**, 85.
- Takahashi, K. (1982) Miospores from the Eocene Nanggulan Formation in the Yogyakarta Region, Central Java. *Transactions and proceedings of the Paleontological Society of Japan. New series*, **1982**, 303–326.
- Tambareau, Y., Gruas-Cavagnetto, C., Feist, M. & Villatte, J. (1991) Flores et faunes continentales ilardiennes du versant sud de la Montagne Noire et de la Montagne d'Alaric. *Revue de Micropaléontologie*, **34**, 69–89.
- Tantawy, A.A., Keller, G., Adatte, T., Stinnesbeck, W., Kassab, A. & Schulte, P. (2001) Maastrichtian to Paleocene depositional environment of the Dakhla Formation, Western Desert, Egypt: sedimentology, mineralogy, and integrated micro- and macrofossil biostratigraphies. *Cretaceous Research*, **22**, 795–827.
- Thanikaimoni, G., Caratini, C., Sivak, J. & Tissot, C. (1987) Striatoechiperiporites, a new pollen genus from the Eocene of Niger. *Pollen et Spores*, **29**, 411–419.
- Thiele-Pfeiffer, H. (1980) Die miozäne Mikroflora aus dem Braunkohlentagebau oder bei Wackersdorf/Oberpfalz. *Palaeontographica Abteilung B Paläophytologie*, **174**, 95–224.

- Torricelli, S., Knezaurek, G. & Biffi, U. (2006) Sequence biostratigraphy and paleoenvironmental reconstruction in the Early Eocene Figols Group of the Tremp–Graus Basin (south-central Pyrenees, Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, **232**, 1–35.
- Tripathi, S.K.M. (1994) Palynology of subsurface palaeocene-eocene sediments near Kapurdi, Barmer District, Rajasthan, India. *Palaeobotanist*, **43**, 45–53.
- Tripathi, S.K.M. (1997) Palynological changes across subsurface Palaeocene-Eocene sediments at Barmer, Rajasthan, India. *Palaeobotanist*, **46**, 168–171.
- Tripathi, S.K.M., Kumar, M. & Srivastava, D. (2009) Palynology of lower palaeogene (Thanetian–Ypresian) coastal deposits from the Barmer Basin (Akli Formation, Western Rajasthan, India): palaeoenvironmental and palaeoclimatic implications. *Geologica Acta*, 147–160.
- Tripathi, S.K.M., Singh, U.K. & Sisodia, M.S. (2003) Palynological investigation and environmental interpretation on Akli Formation (late Palaeocene) from Barmer Basin, western Rajasthan, India. *Palaeobotanist*, **52**, 87–95.
- Trivedi, G.K. (2009) Palynofloral diversity in the Kopili Formation (Late Eocene) from north-east India. *Acta Musei Nationalis Pragae Series B-Historia Naturalis*, **65**, 9–24.
- Trivedi, G.K. & Ranhotra, P.S. (2015) Palynofloral evidence for palaeoecology and depositional environment of the Kopili Formation (Late Eocene), Jaintia Hills, Meghalaya. *Journal of the Geological Society of India*, **86**, 33–40.
- Trivedi, G.K. & Saxena, R.K. (2000) Palynofloral investigation of the Kopili Formation (late Eocene) exposed near Umrongso in North Cachar Hills District, Assam, India. *Palaeobotanist*, **49**, 269–280.
- Truswell, E.M. (1997) Palynomorph assemblages from marine Eocene sediments on the west Tasmanian continental margin and the South Tasman Rise. *Australian Journal of Earth Sciences*, **44**, 633–654.
- Truswell, E.M. & Owen, J.A. (1988) Eocene pollen from Bungonia, New South Wales. *Association of Australasian Palaeontologists, Memoir*, **5**, 259–284.
- Truswell, E.M., Sluiter, I.R. & Harris, W.K. (1984) Palynology of the Oligocene-Miocene sequence in the Cakvale-1 corehole, western Murray Basin, south-Australia. *BMR Journal of Australian Geology & Geophysics*, **9**, 267–295.
- Tschudy, R.H. (1973) Stratigraphic distribution of significant Eocene palynomorphs of the Mississippi embayment. *US Geological Survey Journal of Research*, **743**, B1–B24.
- Tschudy, R.H. & Patterson, S.H. (1975) Palynological evidence for Late Cretaceous, Paleocene, and early and middle Eocene ages for strata in the kaolin belt, central Georgia. *US Geological Survey Journal of Research*, **3**, 437–445.
- Umeji, O.P. & Nwajide, C.S. (2007) Age control and designation of the standard stratotype of Nsukka Formation of Anambra Basin, southeastern Nigeria. *Journal of Mining and Geology*, **43**, 147–166.
- Upadhyay, R., Kar, R. & Sinha, A.K. (2004) Palynological evidence for the Palaeocene evolution of the forearc basin, Indus Suture Zone, Ladakh, India. *Terra Nova*, **16**, 216–225.
- Urrego, D.H., Bush, M.B. & Silman, M.R. (2010) A long history of cloud and forest migration from Lake Consuelo, Peru. *Quaternary Research*, **73**, 364–373.
- Urrego, L.E., Molina, L.A., Urrego, D.H. & Ramírez, L.F. (2006) Holocene space–time succession of the Middle Atrato wetlands, Chocó biogeographic region, Colombia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **234**, 45–61.

- Vajda-Santivanez, V. (1999) Miospores from upper cretaceous-Paleocene strata in northwestern Bolivia. *Palynology*, **23**, 181–196.
- Vallé, F. (2016) *A combination of palynology and geochemical proxies to investigate Pliocene environmental and climate changes in tropical West Africa and east equatorial Atlantic*. Ph.D. thesis, Universität Bremen.
- van der Hammen, T. & Garcia de Mutis, C. (1965) The Paleocene pollen flora of Colombia. *Leidse Geologische Mededelingen*, **35**, 105–114.
- van der Hammen, T., Urrego, L.E., Espejo, N., Duivenvoorden, J.F. & Lips, J.M. (1992) Late-glacial and Holocene sedimentation and fluctuations of river water level in the Caquetá River area (Colombian Amazonia). *Journal of Quaternary Science*, **7**, 57–67.
- van der Hammen, T. & Wymstra, T.A. (1964) A palynological study on the Tertiary and Upper Cretaceous of British Guiana. *Leidse Geologische Mededelingen*, **30**, 183–241.
- Van der Kaars, W.A. & Dam, M.A.C. (1995) A 135,000-year record of vegetational and climatic change from the Bandung area, West-Java, Indonesia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **117**, 55–72.
- Varma, Y.N.R., Ramanujam, C.G.K. & Patil, R.S. (1986) Palynoflora of Tertiary sediments of Tonakkal area, Kerala. *Journal of Palynology*, **22**, 39–53.
- Vélez, M.I., Hooghiemstra, H., Metcalfe, S., Wille, M. & Berrío, J.C. (2006) Late Glacial and Holocene environmental and climatic changes from a limnological transect through Colombia, northern South America. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **234**, 81–96.
- Vélez, M.I., Wille, M., Hooghiemstra, H. & Metcalfe, S. (2005) Integrated diatom-pollen based Holocene environmental reconstruction of lake Las Margaritas, eastern savannas of Colombia. *The Holocene*, **15**, 1184–1198.
- Venkatachala, B.S. (1973) Palynological evidence on the age of Cuddalore Sandstone. *Geophytology*, **3**, 145–149.
- Venkatachala, B.S. & Chowdhary, L.R. (1977) Palaeoecology of the Kadi Formation, Cambay Basin, India. *Proceedings of the IVth Colloquium of Indian Micropalaentology and Stratigraphy*, pp. 259–277.
- Venkatachala, B.S., Kar, R.K., Rajanikanth, A. & Ly, A. (1998) Palynology of the Cretaceous-Tertiary transition in an Atlantic Bore Hole off Senegal, West Africa. *Géologie Méditerranéenne*, **25**, 33–41.
- Venkatachala, B.S. & Sharma, K.D. (1984) Palynological zonation in subsurface sediments in Narsapur well No. 1, Godavari-Krishna Basin, India. *Indian colloquium on micropaleontology and stratigraphy*, pp. 445–466.
- Vergara, L.E. & Ch, G.R. (1997) The Upper Cretaceous and lower Paleocene of the Eastern Bogota Plateau and Llanos Thrustbelt, Colombia: alternative appraisal to the nomenclature and sequence stratigraphy. *Geología Colombiana*, **22**, 51–79.
- Verma, P., Garg, R., Rao, M.R. & Bajpai, S. (2019) Palynofloral diversity and palaeoenvironments of early Eocene Akri lignite succession, Kutch Basin, western India. *Palaeobiodiversity and Palaeoenvironments*, 1–23.
- Verma, P., Garg, R., Rao, M.R. & Bajpai, S. (2020) Palynofloral diversity and palaeoenvironments of early Eocene Akri lignite succession, Kutch Basin, western India. *Palaeobiodiversity and Palaeoenvironments*, **100**, 605–627.
- Verma, P., Mandal, N., Rao, M.R., Thakkar, M.G. & Bajpai, S. (2013) Palynological record from Chhasra Formation (Early Miocene), Eastern Kutch, Gujarat. *Palaeobotanist*, **62**, 149–155.
- Villamizar Meza, F.J. (2014) *Sismoestratigrafía del oligoceno para el sector centro-oeste de la cuenca de los llanos orientales (Municipio de Puerto Gaitán-Meta) Colombia*. Master's thesis, Universidad Nacional de Colombia.

- Volkheimer, W., Novara, M.G., Narváez, P.L. & Marquillas, R.A. (2006) Palynology and paleoenvironmental significance of the Tunal Formation (Danian) at its type locality, El Chorro creek (Salta, Argentina). *Ameghiniana*, **43**, 567–584.
- Volkheimer, W., Scafati, L. & Melendi, D.L. (2007) Palynology of a Danian warm climatic wetland in central northern Patagonia, Argentina. *Revista Española de Micropaleontología*, **39**, 117–134.
- Von der Brelie, G., Hager, H. & Weiler, H. (1981) Pollenflora und Phytoplankton in den Kölner Schichten sowie deren Lithostratigraphie im Siegburger Graben. *Fortschr. Geol. Rheinld. Westf.*, **29**, 21–58.
- Wakefield, M. & Monteil, E. (2002) Biosequence stratigraphical and palaeoenvironmental findings from the Cretaceous through Tertiary succession, Central Indus Basin, Pakistan. *Journal of Micropalaeontology*, **21**, 115–130.
- Wang, W.M. (2006) Correlation of pollen sequences in the Neogene palynofloristic regions of China. *Palaeoworld*, **15**, 77–99.
- Warwick, P.D., Javed, S., Mashhadi, S.T.A., Shakoor, T., Khan, A.M. & Khan, A.L. (1995) *Lithofacies and palynostratigraphy of some Cretaceous and Paleocene rocks, Surghar and Salt Range coal fields, northern Pakistan*. 2006, US Government Printing Office.
- Watanasak, M. (1988) *Mid-Tertiary palynology of onshore and offshore Thailand*. Ph.D. thesis, University of Adelaide.
- Weng, C., Bush, M.B. & Athens, J.S. (2002) Holocene climate change and hydrarch succession in lowland Amazonian Ecuador. *Review of Palaeobotany and Palynology*, **120**, 73–90.
- Westgate, J.W. & Gee, C.T. (1990) Paleoecology of a middle Eocene mangrove biota (vertebrates, plants, and invertebrates) from southwest Texas. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **78**, 163–177.
- Wijmstra, T.A. (1969) Palynology of the Alliance well. *Geologie en Mijnbouw*, **48**, 125–133.
- Wijmstra, T.A. (1977) *The palynology of the Guiana coastal basin*. Ph.D. thesis, University of Amsterdam.
- Wijninga, V.M. (1996a) Neogene ecology of the Salto de Tequendama site (2475 m altitude, Cordillera Oriental, Colombia): the paleobotanical record of montane and lowland forests. *Review of Palaeobotany and Palynology*, **92**, 97–156.
- Wijninga, V.M. (1996b) Palynology and paleobotany of the Early Pliocene section Río Frío 17 (Cordillera Oriental, Colombia): biostratigraphical and chronostratigraphical implications. *Review of Palaeobotany and Palynology*, **92**, 329–350.
- Wilkinson, G.C. & Boulter, M.C. (1980) Oligocene pollen and spores from the western part of the British Isles. *Palaeontographica Abteilung B Paläophytologie*, **175**, 27–83.
- Wille, M., Hooghiemstra, H., van Geel, B., Behling, H., de Jong, A. & van der Borg, K. (2003) Submillennium-scale migrations of the rainforest–savanna boundary in Colombia: 14C wiggle-matching and pollen analysis of core Las Margaritas. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **193**, 201–223.
- Williams, R.H. (2016) *Paleoenvironment reconstruction in extreme climates using organic geochemical investigations*. Ph.D. thesis, Massachusetts Institute of Technology.
- Willwms, W.A.E. (1982) Microfossil assemblages, zonations and planktonic datum levels in the Ieper Formation (Ypresian ss, Early Eocene) in Belgium. *Professional paper-Service géologique de Belgique*, 1–16.
- Winantris, Syafri, I. & Rinawan, R. (2006) Kandungan Mikrofosil Dalam Formasi Pembawa Batubara Dari Daerah Perian Kecamatan Muara Muntai, Kabupaten Kutai Kertanegara, Kalimantan Timur. *Bulletin of Scientific Contribution*, **4**, 8–18.

- Winantris, W., Hamdani, H., Harlia, E. *et al.* (2017) Paleoenvironment of Tanjung Formation Barito Basin-Central Kalimantan Based on palynological data. *Journal of Geoscience, Engineering, Environment, and Technology*, **2**, 110–116.
- Wirmann, D., Eagar, S.H., Harper, M.A., Leroy, É. & Sémah, A.M. (2011) First insights into mid-Holocene environmental change in central Vanuatu inferred from a terrestrial record from Emaotfer Swamp, Efaté Island. *Quaternary Science Reviews*, **30**, 3908–3924.
- Wu, G., Qin, J. & Mao, S. (2003) Deep-water Oligocene pollen record from South China Sea. *Chinese Science Bulletin*, **48**, 2511–2515.
- Wu, P., Hou, D., Gan, J., Li, X., Ding, W., Liang, G. & Wu, B. (2018) Paleoenvironment and controlling factors of Oligocene source rock in the eastern deep-water area of the Qiongdongnan basin: evidences from organic geochemistry and palynology. *Energy & Fuels*, **32**, 7423–7437.
- Xie, J., Qin, J., Li, J., Cai, K. & Liu, X. (2014) Palynological assemblages of the Oligocene and Pliocene in the Yinggehai–Qiongdongnan Basin of the northern South China Sea and their paleoenvironmental implications. *Acta Micropalaeontologica Sinica*, **31**, 85–97.
- Yakzan, A.M., Harun, A., Nasib, B.M. & Morley, R.J. (1996) Integrated biostratigraphic zonation for the Malay Basin. *The Bulletin of the Geological Society of Malaysia*, **39**, 157–184.
- Yang, S. (1993) Paleocene Palynology of the Shangyang Formation, Hepu Basin, Guangxi, China. *Acta Micropalaeontologica Sinica*, **10**, 213–222.
- Yang, Y., Zhang, H., Chang, F., Meng, H., Pan, A., Zheng, Z. & Xiang, R. (2016) Vegetation and climate history inferred from a Qinghai Crater Lake pollen record from Tengchong, southwestern China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **461**, 1–11.
- Yans, J., Marandat, B., Masure, E., Serra-Kiel, J., Schnyder, J., Storme, J.Y., Marivaux, L., Adnet, S., Vianey-Liaud, M. & Tabuce, R. (2014) Refined bio-(benthic foraminifera, dinoflagellate cysts) and chemostratigraphy ($\delta^{13}\text{C}_{\text{org}}$) of the earliest Eocene at Albas-Le Clot (Corbières, France): implications for mammalian biochronology in Southern Europe. *Newsletters on Stratigraphy*, **47**, 331–353.
- Yulianto, E., Rahardjo, A.T., Noeradi, D., Siregar, D.A. & Hirakawa, K. (2005a) A Holocene pollen record of vegetation and coastal environmental changes in the coastal swamp forest at Batulicin, South Kalimantan, Indonesia. *Journal of Asian Earth Sciences*, **25**, 1–8.
- Yulianto, E., Tsuji, H., Sukpti, W.S. & Tanaka, N. (2005b) A holocene pollen and charcoal record from a tropical lowland swamp in Rawa Danau, West Java, Indonesia. *Tropics*, **14**, 271–281.
- Zamaloa, M.d.C. & Romero, E.J. (1990) Some spores and pollen from the Cullen Formation (upper Eocene to middle Oligocene), Tierra del Fuego, Argentina. *Palynology*, **14**, 123–133.
- Zeng, H., Ambrose, W.A. & Villalta, E. (2001) Seismic sedimentology and regional depositional systems in Mioceno Norte, Lake Maracaibo, Venezuela. *The Leading Edge*, **20**, 1260–1269.
- Zetter, R., Hofmann, C.C., Piller, W.E. & Rasser, M.W. (2001) New aspects of the palynoflora of the lowermost Eocene (Krappfeld Area, Carinthia). W.E. Piller & M.W. Rasser, eds., *Paleogene of the Eastern Alps*, Österreichische Akademie der Wissenschaften, pp. 473–507.
- Zhang, Y. (1981) Tertiary spores and pollen grains from the Leizhou Peninsula. *Acta palaeontologica sinica*, **20**, 449–457.
- Zhong, X. (1985) Pearl River Mouth Basin of South China Sea. *ESCAP Atlas of Stratigraphy IV. Peoples Republic of China*, United Nations. Economic and Social Commission for Asia and the Pacific, pp. 45–49.