



UvA-DARE (Digital Academic Repository)

Prevention and therapy of periodontal diseases and oral malodour

van der Sluijs, E.

[Link to publication](#)

Citation for published version (APA):

van der Sluijs, E. (2017). Prevention and therapy of periodontal diseases and oral malodour: Brush, rinse and cool DIDES

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



3

Prebrushing rinse with water on plaque removal: a split-mouth design

E. van der Sluijs

D.E. Slot

N.L. Hennequin-Hoenderdos

M.P.C. van Leeuwen

G.A. van der Weijden

Abstract

Aim

The aim was to evaluate if there is an additional beneficial effect on dental plaque removal of rinsing the oral cavity with water before toothbrushing.

Methods

In total 48 non-dental, systemically healthy participants ≥ 18 years were included in this randomized controlled clinical trial using a split-mouth design. The participants were requested to refrain from any form of oral hygiene for 48 hours. First dental plaque scores (PI) were assessed full-mouth. Two randomly assigned contra-lateral quadrants were brushed. Next the participant rinsed for 1 minute with 15 ml water. Subsequently, the opposite two contra-lateral quadrants were brushed. Brushing was performed without toothpaste. Subsequently the second full-mouth PI assessment was performed. The brushing and rinsing procedure was done under supervision and brushing time was tracked by a timer, each quadrant was brushed for 30 s. For the buccal, lingual, and approximal surfaces and tooth type a sub-analysis was performed.

Results

At baseline there was no statistically significant difference between the two sets of contra-lateral quadrants. When a water rinse was used before toothbrushing the PI-score was reduced by 58%. If water rinse was used post-brushing the PI-score reduced by 57%. The difference of 0.04 in mean plaque index score reduction between the two brushing regimens was not significant ($P = 0.162$).

Conclusion

When a 2 minutes brushing exercise was performed, on average more than 55% dental plaque was removed. Pre-rinsing with water did not contribute significantly to toothbrush efficacy.

Introduction

In a healthy situation, dental plaque and adjacent tissues maintain a delicate balance. When healthy dental plaque changes into a pathogenic bio-film, this can initiate the periodontal disease process. Toothbrushing is today the most widespread mechanical means of controlling plaque at home (1) and the cornerstone to prevent and control periodontal diseases (2,3). Supragingival plaque is exposed to saliva and to the natural self-cleansing mechanisms existing in the oral cavity. However, although such mechanisms may eliminate food debris, they do not adequately remove dental plaque (2,4). With respect to daily home care the American Dental Association recommends to brush at least twice a day, for 2 min. After tooth brushing, it is common practice to rinse the mouth with tap water with the intention to remove dentifrice remnants and loosened plaque (5). Other researchers have proposed to rinse before brushing in order to hydrate the biofilm, reduce adherence and render plaque more readily detached by mechanical cleaning methods. Changing oral care habits by introducing a prebrushing rinse would mean a change in the world of oral home care (6). A systematic review evaluated the effectiveness of a marketed prebrushing rinse in plaque removal (7). The review authors concluded that the prebrushing rinse appears to offer some benefit in plaque reductions. This specific mouthwash was compared to control groups which were rinsing, with either a placebo or water. A stepwise toothbrush exercise showed that prerinsing with water prior to brushing provided additional plaque removal (8). However, no comparison to regular toothbrushing was performed. The only study that did compare a prebrushing rinse with water to regular toothbrushing was the study by O'Mullane et al. (9). They did, however, not evaluate the effect of a brushing exercise per sé but assessed plaque scores over time with different interventions. Although not significant, the numerical data may be considered indicative of a difference between no rinse and prerinsing with tap water. The purpose of the present study was to compare more specifically Prerinsing with water prior to toothbrushing with common daily toothbrushing that is followed by rinsing with water afterwards.

Materials and methods

Ethical procedures

The study followed instructions based on the Helsinki principles (2013), approximating Good Clinical Practice guidelines and held the approval of the medical ethical committee of the Amsterdam Medical Centre #2014_118. The study was also registered at the Dutch Trial Register (#NTR4604). The study took place in the time period of June until September 2014 at the Academic Centre for Dentistry of Amsterdam (ACTA), the Netherlands, Department of Periodontology. Before enrolment, all volunteers were given verbal and written instructions as well as a description of the aim, rationale and duration of study participation. The investigator explained the details of the trial and potential risk involved. All eligible subjects that agreed to participate signed an informed consent form prior to the study procedures, in addition they were informed that they were free to withdraw from the trial at any time.

Sample size

Sample size calculations were performed using the PS Power and Sample Size Program (10). Earlier data indicated that the difference in the response of the primary outcome in matched pairs is normally distributed with standard deviation 0.30 [pooled estimate out of Van der Weijden et al. (11–13)]. If the true difference in the mean response of matched pairs is 0.14, 38 participants were needed to be able to reject the null hypothesis that the response difference is zero with probability α (power) 0.8. The Type I error probability associated with this test of the null hypothesis was 0.05. Additional participants were entered to compensate for potential dropout.

Screening and inclusion

The participants were non-dental students from different universities and colleges in and around Amsterdam who were invited by e-mail and flyer advertisement. The first selection was performed by a general telephone screening. Additionally, in the clinic the participants were screened by a dental hygienist (MPCL). Inclusion criteria were: ≥ 18 years old, classified as systemically healthy as assessed by the medical questionnaire, periodontally healthy as assessed by scoring the DPSI ≤ 3 min (14, 15), ≥ 5 teeth per quadrant and right-handed brushers. Excluded were those that presented with an orthodontic appliance or a removable (partial) denture, overt caries, any pathological alterations of the oral mucosa, pregnancy and use of medications within 2 weeks before the appointment including antibiotics or chronic use of non-steroidal anti-inflammatory drugs (excluding birth control pills).

Design

This study had a single-(examiner) blind design using a split mouth model where contralateral quadrants were randomly assigned to a treatment (16). The study coordinator (EVDS) supervised the assigned brushing procedures.

Randomization and blinding

Every participant received a unique subject identification number. No stratification was applied. Randomization was performed using true random numbers, which were generated by sampling and processing a source of atmospheric noise (17). The randomization code was kept in a sealed envelope in the investigator site file and was only accessible by the coordinator who was therefore responsible for allocation concealment. Participants were instructed not to reveal their intervention in any way to the clinical examiner. Records of earlier examinations were not available to the examiner at the time of re-examination.

Primary outcome

The main study parameter was the level of full-mouth dental plaque as measured by the Silness & L oe Plaque index (PI) (18) and modified as described by Van der Weijden et al. (11). Where 0 = no plaque, 1 = a film of plaque adhering to the free gingival margin and adjacent area of the tooth 2 = moderate accumulation of soft deposits on the tooth and gingival margin that can be seen with the naked eye, 3 = abundance of soft matter on the tooth and gingi-

val margin (19). Throughout the study examinations were carried out by one and the same experienced examiner (NLHH) under the same conditions. At the time of examination, the examiner was unaware of the randomization scheme.

Study procedure

A 48-h non-brushing period was used in order to accumulate an adequate amount of plaque to assess the incremental effect of a single brushing exercise (20). An SMS (Short Message Service) was sent to remind each participant before the visit. First, a full-mouth assessment of dental plaque score was performed. Then the participants went to a separate room, away from the examiner. The intervention was performed under the supervision of the study coordinator. Before the brushing exercise a short verbal instruction was given and no mirror was available. Two randomly assigned contra-lateral quadrants were brushed without dentifrice using a new and prewetted toothbrush (Oral-B indicator P35 soft). Each side of the quadrants was brushed for 15 s, in total 30 s per quadrant. If necessary the participants were able to expectorate saliva after each period of 15 s. Next, the participants rinsed with 15 ml mineral water (Christaline SAINTE-SOPHIE bottled Spring water) for 1 min. The instruction was to take the total amount of water that was inside the cup into their mouth and rinse laboriously without swallowing. Additional subsequent rinsing was not allowed. Next the opposing two quadrants were brushed using a new and prewetted toothbrush following the same procedure as previously described. The supervised brushing and rinsing procedure was tracked by a timer. Following the brushing–rinsing–brushing exercise, a full-mouth plaque assessment was performed to obtain the post-brushing plaque scores. Additionally, the participant completed a questionnaire designed to evaluate their opinion. Their perception was questioned concerning the 48-h non-brushing period, the toothbrush and the procedure used. The participants placed a vertical mark on a 10-cm-long uncalibrated line. The VAS-score line had two extremes at either end of the line with the negative extreme on the left side. The investigator then measured the distance along the line from the left extreme to the marking made by the participant (21). At the end of the study, participants received monetary compensation for their contribution.

Statistical analysis

The statistical analysis using SPSS software (Statistical Package for Social Sciences, version 20.0 for Windows) was performed before the randomization code was broken. The primary analysis was the overall dental plaque measurements. The mean and standard deviations were calculated for all sites for each participant, and in addition, the mean incremental difference and proportional change were computed. For a sub-analysis, all vestibular sites, all lingual sites and all approximal sites were calculated. The central incisors were excluded from the measurement to avoid crossover effects of brushing the adjacent teeth. The total mean and variation of the dental plaque scores were analysed for each participant and subsequently by group. Data were tested for normal distribution by the Kolmogorov–Smirnov test and found to correspond with the normal distribution assumption. Parametric statistical technique by the use of a paired T-test was applied. Values of $P < 0.05$ were accepted as statistically significant. The mean scores and standard deviation

on the VAS questions were calculated. As the VAS questions were descriptive and did not differentiate between groups, no statistical analyses were performed concerning these data.

Results

In total, 50 participants were assessed for eligibility (Fig. 1) (22). Two participants were excluded due to the presence of overt caries and periodontitis. In total, 48 participants were included. The mean age of the 48 participants was 22.5 (2.81) with a range of 20–34. In total, there were more female participants in this clinical trial (N = 28, 58%) than men (N = 20, 42%). No adverse events were reported.

Primary outcome

Table 1 shows the mean and standard deviation of the total dental plaque scores. The split-mouth design implicates that each participant contributed with 2 baseline and 2 post-brushing values. The rinsing and brushing procedure of this study was carefully standardized and controlled. For the post-brushing rinse group, the baseline plaque score was 1.69 (0.32) and for the prebrushing rinse group 1.66 (0.33) which was not significantly different. The post-brushing rinse assessment revealed a reduction in plaque scores of 0.95 (0.20) in the prerinse group compared to a reduction of 0.91 (0.21) in the post-rinse group. Both groups showed a significant reduction in dental plaque. However, there was no statistically significant difference between the groups on the end scores ($P = 0.935$). The difference of 0.04 in mean plaque index score reduction between the two brushing regimens was not significant ($P = 0.162$). When the mouth was rinsed with water before brushing, plaque scores reduced by 58% as compared to a 57% plaque score reduction with rinsing after brushing ($P = 0.722$).

Subanalysis

Subanalysis (Table 2) was performed for the various surfaces, analysing buccal sites, lingual sites or approximal sites separately. The baseline approximal scores between interventions were not balanced well ($P = 0.027$). Regarding the lingual sites, 68% and 64% plaque score reduction was found for prebrushing or post-brushing water rinsing, respectively, which was close to being significant ($P = 0.049$). No further statistical differences were found for pre- and post-brushing rinse data.

Secondary outcome

In total, six questions were asked after the assessment (Table 3). The participants felt rather uncomfortable to refrain 48 h of any oral hygiene before the appointment (VAS is 2.57). The participants felt that the toothbrush bristles are neither too hard or too soft (4.87) nor felt pleased. They scored the use of this manual toothbrush with a 6.04, which is close to the mean value of 5. A brushing exercise without the use of dentifrice was judged as unpleasant (3.07). Prerinsing with water was felt moderately pleasant (6.66), and the 1 min rinsing duration seems appropriate (6.18).

Figure 1 Flow chart depicting subject enrollment and measurement (22).

* these quadrants are used as post brushing rinse

‡ these quadrants are pre brushed rinse

Q = Quadrant




Table 1 Overall dental plaque scores on baseline and end presented by mean and SD (standard deviation) when post- and pre- brushing rinse with water was performed.

N=48	Post-brushing rinse	Pre-brushing rinse	Statistical analysis*
Baseline	1.69 (0.32)	1.66 (0.33)	0.095
End	0.75 (0.34)	0.76 (0.33)	0.935
Incremental difference	0.95 (0.20)	0.91 (0.21)	0.162
% Reduction	58% (15)	57% (15)	0.722

* Paired sample T-test

Discussion

The aim of the present study was to compare rinsing with water prior to toothbrushing with toothbrushing followed by rinsing on the removal of dental plaque. The plaque scores were reduced by more than 55%, whereas no difference was obtained between the two interventions. Similar results were also found for the subanalysis on various surfaces.

Study design used

Bentley and Disney (16) evaluated the comparison of partial and full-mouth scores of plaque and gingivitis without compromising study quality. Their analyses showed that all-teeth, the Ramfjord teeth, half mouth and molars/anteriors revealed a similar pattern as whole-mouth and proximal areas. The present study used an examiner blind split-mouth model with randomly assigned contra-lateral quadrants for each allocation. The advantage of this design is that it reduces interindividual variability from the estimates of the treatment effect, because participants acted as their own control (23). Also the impact of the ‘Hawthorne effect’ of present study would have impacted both interventions similarly because the same operator performs both interventions in the same oral cavity. The split-mouth randomization was applied in the current design in such a manner that all participants contributed with two quadrants to the prebrushing rinse and with two to the post-brushing rinse data. The rationale for using contra-lateral quadrants is that the skills of the individual using the brush may affect plaque removal. To rule out the influence of dexterity, contra-lateral quadrants were randomly assigned. This has been used in previous brushing evaluations also using the same plaque index (24,25). Additionally, only right-handed brushers were included, because a previous investigation showed a significant difference between right- and left-handed participants (13).

Rinsing procedure used

The results of a meta-analysis have demonstrated that a commercially available prebrushing rinse has a positive effect on reducing plaque score levels (7). The present study evaluated solely the additional effect of water as a prebrushing rinse. Based on the data of a previous study (26), a rinse volume of 15 ml was used for this experiment. On a 0–10 VAS scale (extremely unpleasant to very pleasant), participants felt that rising with water was moderately pleasant with a mean VAS score of 6.66.

Table 2 Sub-analysis evaluating dental plaque scores for the buccal, lingual and approximal sites separately. Data are presented by mean and SD (standard deviation) in relation to a specific sequence in brushing order.

Sites	N=48	Post-brushing rinse	Pre-brushing rinse	Statistical analysis*
Buccal	Baseline	1.49 [0.58]	1.48 [0.55]	0.957
	End	0.30 [0.27]	0.38 [0.29]	0.121
	Difference	1.18 [0.51]	1.11 [0.42]	0.294
Lingual	Baseline	1.53 [0.45]	1.45 [0.46]	0.125
	End	0.49 [0.33]	0.52 [0.36]	0.531
	Difference	1.04 [0.36]	0.93 [0.36]	0.049**
Approximal	Baseline	1.79 [0.29]	1.74 [0.31]	0.027**
	End	0.91 [0.43]	0.89 [0.40]	0.506
	Difference	0.88 [0.24]	0.85 [0.23]	0.218

* Paired sample T-test

** Significant difference between the groups (P<0.05)

Rinsing duration has also been evaluated in previous work. Paraskevas et al. (27) showed that the intra-oral spread of the mouthrinses in relation to the rinsing periods of 15, 30 and 60 s. Rinsing with 30 s was sufficient to allow all plaque-covered surfaces of the dentition to come into contact with the solution. To be on the safe side regarding spread and optimal wetting of the dental plaque for the present study, 1 min prerinsing duration was used. The participants evaluated this duration on a scale from too short up to too long as 6.18. This is just above the neutral score of 5.00 of being neither too short nor too long.

Brushing procedure used

Recently, in a metareview concerning the evidence as presented in systematic reviews, the efficacy of available home-care toothbrush regimens for mechanical plaque removal on plaque and gingivitis in adults was evaluated. The conclusion was that tooth brushing is effective in reducing levels of dental plaque (28). The efficacy in plaque removal following a brushing exercise is a reduction from baseline plaque scores of 42% between pre- and post-brushing on average, with a variation of 30–53% dependent on the plaque index used. This variation based on plaque indices is also supported by the results of a study by Binney et al. (8) showing that the overall effect of brushing upon plaque reduction with the Quigley & Hein Plaque Index (29) was 27%, whereas in the same participant, a 64% reduction in ‘plaque area’ scores was observed. The results of the present study have shown that using a manual toothbrush and assessing plaque according to Silness & Loe plaque index 57% and 58% reductions are observed for the pre- and post-brushing water rinse, respectively. An explanation for more plaque removal in the present study as compared to Slot & Van der Weijden (28) may in part be explained by the Silness & Loe plaque index used (18). Even a difference in outcomes between the Turesky (30) and Lobene (31) modification of the Quigley & Hein Plaque Index (29) has been shown (20). Also, the supervised brushing may have improved efficacy. This supervision is often performed in research to make sure that the

Table 3 Questionnaire related to the self-perception of the participants using a Visual Analogue Scale (VAS) presented as mean and standard deviation (SD) with negative extremes on the left and positive extremes on the right (from 0 to 10).

N=48		Extreme		Mean (SD)
		From	To	
1	What do you think of refraining 48 hours before the appointment?	Uncomfortable	Comfortable	2.57 [1.57]
2	I think the toothbrush is cleaning my teeth well enough?	Not agree	Totally agree	6.04 [2.05]
3	I think the bristles are...	Very hard	Very soft	4.87 [1.65]
4	I think brushing without a dentifrice is...	Unpleasant	Pleasant	3.07 [2.45]
5	I think rinsing with water is...	Unpleasant	Pleasant	6.66 [1.68]
6	I think the time of rinsing (1 minute) is...	Too short	Too long	6.18 [1.35]

study procedures including brushing duration are performed according to the protocol. Finally, also the non-oral hygiene period prior to the clinical assessment may have affected the plaque score reduction. The longer the plaque could accumulate freely on the dental surfaces, the higher the plaque score reduction. An explanation could be that if there is more plaque present on the tooth surface, there is simply more plaque to remove (20).

In the present study, no dentifrice was used during the brushing procedure. Based on the VAS questionnaire regarding this aspect, the participants' perception was that this was not considered to be pleasant. Already, half a century ago, this was also evaluated. The general reactions of participants who used no dentifrice were most unfavourable (32). The use of a dentifrice is in general stimulated by many factors such as pleasantness of flavour and aftertaste, foaming quality, mouth feel during application, removal of tooth stain, feeling of freshness and cleanliness (33). This also contributes to confidence of not offending others by 'bad breath' (34). However, the effect of dentifrice on plaque removal compared to brushing with only water showed that the reduction in plaque index by brushing with a dentifrice was less marked than brushing with water (28). Already, in 1967, Bergenholtz et al. (35) reported that brushing without dentifrice was more efficient than similar brushing with dentifrice. The findings in Binney et al. (8) demonstrated that none of the commercial dentifrices tested possessed plaque removal properties of clinical significance and certainly little greater than what might be achieved with water or a slurry of toothpaste.

Post-brushing rinsing

Rinsing with water after brushing with fluoride toothpaste can reduce the benefit of fluoride toothpaste. Davies et al. (36) recommend discouraging rinsing with water after brushing. As an alternative, a fluoride mouthrinse can be used after brushing. Because most people are used to rinsing after brushing, this will certainly help compliance. This recommendation could have a beneficial effect for caries control.

Limitation and directions for further research

- The current study presents the results of young healthy adults. Older adults with wider interdental spaces might show different results.
- The effect of post-brushing rinsing with water may have an additional effect on plaque removal. This aspect has not been studied before so that the potential impact on outcomes is unclear.
- In the present study, we used a new- and prewetted toothbrush for each intervention. O’Hehir and Suvan (37) suggested that lingual surfaces should be dry-brushed first. In a practice-based evaluation, they noted significant improvements. Both these aspects could be the topic of future evaluations.

Conclusion

When a 2 min brushing exercise was performed, plaque score reduced by more than 55%. The study does not support that the use of a prebrushing rinse provides a greater effect than mechanical plaque removal by a manual toothbrush.

Acknowledgements

The study was sponsored by an unrestricted educational grant by Procter & Gamble Company Worldwide Clinical Investigations – Oral Care. The manual toothbrushes were also provided free of charge by Procter & Gamble.

Conflict of interest and source of funding statement

The authors declare that they have no conflict of interest. Authors and their research group at ACTA have formerly received external advisor fees, lecturer fees and/or research grants from toothbrush manufacturers. Among these were Braun AG, Colgate, Dentaïd, GABA, Lactona, Oral-B, Philips, Procter & Gamble, Sara Lee, Sunstar and Unilever.

Authors contributions

Conception or design of the study: EVDS, DES, NLHH, GAW

Analysis and/or interpretation of the data: EVDS, DES, MPCVL, GAW,

Drafted the manuscript: EVDS

Critically revised the manuscript: DES, NLHH, MPCVL, GAW

All authors gave their final approval and agreed to be accountable for all aspects of the work

References

1. Lang NP, Karring T. Proceeding of the 1st European Workshop on Periodontology. Axelsson P, Session III Mechanical plaque control. London: Quintessence Publishing Co; 1994, 219–243.
2. Van der Weijden GA, Hioe KP. A systematic review of the effectiveness of self-performed mechanical plaque removal in adults with gingivitis using a manual toothbrush. *J Clin Periodontol* 2005; 32: (Suppl 6): 214–228.
3. Hayasaki H, Saitoh I, Nakakura-Ohshima K et al. Tooth brushing for oral prophylaxis. *Jpn Dent Sci Rev* 2014; 50: 69–77.
4. Dawes C, Jenkins GN, Tonge CH. The nomenclature of the integuments of the enamel surface of teeth. *Br Dent J* 1963; 115: 65–68.
5. Gerardu VA, van Loveren C, Heijnsbroek M, Buijs MJ, van der Weijden GA, ten Cate JM. Effects of various rinsing protocols after the use of amine fluoride/stannous fluoride toothpaste on the acid production of dental plaque and tongue flora. *Caries Res* 2006; 40: 245–250.
6. Lippert F. An introduction to toothpaste – its purpose, history and ingredients. *Monogr Oral Sci* 2013; 23: 1–14.
7. Angelillo IF, Nobile CGA, Pavia M. Evaluation of the effectiveness of a pre-brushing rinse in plaque removal: a meta-analysis. *J Clin Periodontol* 2002; 29: 301–309.
8. Binney A, Addy M, Newcombe RG. The plaque removal effects of single rinsings and brushings. *J Periodontol* 1993; 64: 181–185.
9. O'Mullane DM, Kavanagh D, Ellwood RP. A three year clinical trial of a combination of trimetaphosphate and sodium fluoride in silica toothpastes. *J Dent Res* 1997; 76: 1776–1778.
10. Dupont WD, Plummer WD. PS power and sample size program available for free on the internet. *Control Clin Trials* 1997; 18: 274.
11. Van der Weijden GA, Danser MM, Nijboer A, Timmerman MF, Van der Velden U. The plaque-removing efficacy of an oscillating/ rotating toothbrush. A short term study. *J Clin Periodontol* 1993; 20: 273–278.
12. Van der Weijden GA, Timmerman MF, Nijboer A, Lie MA, Van der Velden U. A comparative study of electric toothbrushes for the effectiveness of plaque removal in relation to toothbrushing duration. Timerstudy. *J Clin Periodontol* 1993; 20: 476–481.
13. Van der Weijden GA, Timmerman MF, Reijerse E et al. The long-term effect of an oscillating/rotating electric toothbrush on gingivitis. An 8-month clinical study. *J Clin Periodontol* 1994; 21: 139–145.
14. Gómez SM, Danser MM, Sipos PM, Rowshani B, van der Weijden GA. Tongue coating and salivary bacterial counts in healthy/gingivitis subjects and periodontitis patients. *J Clin Periodontol* 2001; 28: 970–978.
15. Van der Velden U. The Dutch periodontal screening index validation and its application in the Netherlands. *J Clin Periodontol* 2009; 36: 1018–1024.
16. Bentley CD, Disney JA. A comparison of partial and full mouth scoring of plaque and gingivitis in oral hygiene studies. *J Clin Periodontol* 1995; 22: 131–153.
17. Randomness and Integrity Services Ltd. Dublin, Ireland. Available at: www.random.org (accessed 1 December 2015).
18. Silness J, Løe H. Periodontal disease in pregnancy (II). Correlation between oral hygiene and periodontal condition. *Acta Odontol Scand* 1964; 22: 121–135.
19. Danser MM, Driesen GM, Timmerman MF, van der Velden U, van der Weijden GA. A laboratory and clinical investigation comparing 2 oscillating/ rotating electric toothbrushes. *J Clin Periodontol* 2000; 27: 277–283.
20. Rosema N, Slot DE, van Palenstein Helderman WH, Wiggelinkhuizen L, Van der Weijden GA. The efficacy of powered toothbrushes following a brushing exercise: a systematic review. *Int J Dent Hyg* 2016; 14: 29–41.
21. Mottola CA. Measurement strategies: the visual analogue scale. *Decubitus* 1993; 6: 56–58.
22. CONSORT Group. The CONSORT statement 2010. Available at: <http://www.consort-statement.org/>

- (accessed 1 December 2015).
23. Lesaffre E, Philstrom B, Needleman I, Worthington H. The design and analysis of split-mouth studies: what statisticians and clinicians should know. *Stat Med* 2009; 28: 3470–3482.
 24. Van der Weijden GA, Timmerman MF, Novotny AG, Rosema NA, Verkerk AA. Three different rinsing times and inhibition of plaque accumulation with chlorhexidine. *J Clin Periodontol* 2005; 32: 89–92.
 25. Graveland MP, Rosema NA, Timmerman MF, Van der Weijden GA. The plaque-removing efficacy of a finger brush (I-Brush). *J Clin Periodontol* 2004; 31: 1084–1087.
 26. Keukenmeester RS, Slot DE, Rosema NA, Van der Weijden GA. Determination of a comfortable volume of mouthwash for rinsing. *Int J Dent Hyg* 2012; 10: 169–174.
 27. Paraskevas S, Danser MM, Timmerman MF, Van der Velden U, van der Weijden GA. Optimal rinsing time for intra-oral distribution (spread) of mouthwashes. *J Clin Periodontol* 2005; 32: 665–669.
 28. Slot DE, Van der Weijden FA. Group A. Initiator paper. Plaque control: home remedies practiced in developing countries. *J Int Acad Periodontol* 2015; 17: 4–15.
 29. Quigley GA, Hein JW. Comparative cleansing efficiency of manual and power brushing. *J Am Dent Assoc* 1962; 65: 26–29.
 30. Turesky S, Gilmore ND, Glickman L. Reduced plaque formation by the chloromethyl analogue of vitamin C. *J Periodontol* 1970; 41: 41–43.
 31. Lobene RR, Soparkar PM, Newman BS. Use of dental floss. Effect on plaque and gingivitis. *Clin Prev Dent* 1982; 4: 5–8.
 32. Dudding NJ, Dahl LO, Muhler JC. Patient reactions to brushing teeth with water, dentifrice, or salt and soda. *J Periodontol* 1960; 31: 386–392.
 33. Sälzer S, Rosema NA, Martin EC et al. The effectiveness of dentifrices without and with sodium lauryl sulfate on plaque, gingivitis and gingival abrasion—a randomized clinical trial. *Clin Oral Investig* 2015; 22 [Epub ahead of print].
 34. Pader M. *Oral Hygiene Products and Practice*. New York: Marcel Dekker, Inc; 1988.
 35. Bergenholtz A, Hugoson A, Sohlberg F. An evaluation of the plaque-removing ability of some aids to oral hygiene. *Sven Tandlak Tidsskr* 1967; 60: 447–454.
 36. Davies RM, Davies GM, Ellwood RP. Prevention. Part 4: toothbrushing: what advice should be given to patients? *Br Dent J* 2003; 9: 135–141.
 37. O’Hehir TE, Suvan JE. Dry brushing lingual surfaces first. *J Am Dent Assoc* 1998; 129: 614.