



UvA-DARE (Digital Academic Repository)

Division by zero in common meadows

Bergstra, J.A.; Ponse, A.

DOI

[10.48550/arXiv.1406.6878](https://doi.org/10.48550/arXiv.1406.6878)

Publication date

2021

Document Version

Submitted manuscript

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Bergstra, J. A., & Ponse, A. (2021). *Division by zero in common meadows*. (v4 ed.) ArXiv. <https://doi.org/10.48550/arXiv.1406.6878>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Division by Zero in Common Meadows*

Jan A. Bergstra and Alban Ponse

Section Theory of Computer Science
 Informatics Institute, Faculty of Science
 University of Amsterdam, The Netherlands
<https://staff.fnwi.uva.nl/{j.a.bergstra/,a.ponse/}>

Abstract

Common meadows are fields expanded with a total inverse function. Division by zero produces an additional value denoted with \mathbf{a} that propagates through all operations of the meadow signature (this additional value can be interpreted as an error element). We provide a basis theorem for so-called common cancellation meadows of characteristic zero, that is, common meadows of characteristic zero that admit a certain cancellation law.

Keywords and phrases: Meadow, common meadow, division by zero, additional value, abstract datatype.

Contents

1	Introduction	2
1.1	Common Meadows versus Involutive Meadows	2
1.2	Motivating a Preference for Common Meadows	3
2	Common Meadows	4
2.1	Meadow Signatures	4
2.2	Axioms for Common Meadows	4
2.3	Conditional Equations	8
3	Models and Model Classes	9
3.1	Common Cancellation Meadows	10
3.2	A Basis Theorem for Common Cancellation Meadows of Characteristic Zero	11
4	Concluding Remarks	14
	References	15

*The preceding version v2 (22 December 2014) appeared in: Rocco De Nicola and Rolf Hennicker (eds.), *Software, Services and Systems: Essays Dedicated to Martin Wirsing*, LNCS 8950, pp. 46-61, Springer, 2015. Main changes: axiom (12) in Table 2 is new and the proof of Thm.3.2.1 has been corrected.

1 Introduction

Elementary mathematics is uniformly taught around the world with a focus on natural numbers, integers, fractions, and fraction calculation. The mathematical basis of that part of mathematics seems to reside in the field of rational numbers. In elementary teaching material the incorporation of rational numbers in a field is usually not made explicit. This leaves open the possibility that some other abstract datatype or some alternative abstract datatype specification improves upon fields in providing a setting in which such parts of elementary mathematics can be formalized.

In this paper we will propose the signature for — and model class of — *common meadows* and we will provide a *loose* algebraic specification of common meadows by way of a set of equations. In the terminology of Broy and Wirsing [10, 17], the semantics of a loose algebraic specification S is given by the class of all models of S , that is, the semantic approach is not restricted to the isomorphism class of initial algebras. For a loose specification it is expected that its initial algebra is an important member of its model class, worth of independent investigation. In the case of common meadows this aspect is discussed in the last remark of Section 4 (Concluding remarks).

A common meadow (using inversive notation) is an extension of a field equipped with a multiplicative inverse function $(\dots)^{-1}$ and an additional element \mathbf{a} that serves as the inverse of zero and propagates through all operations. It should be noticed that the use of the constant \mathbf{a} is a matter of convenience only because it merely constitutes a derived constant with defining equation $\mathbf{a} = 0^{-1}$. This implies that all uses of \mathbf{a} can be removed from the story of common meadows (a further comment on this can be found in Section 4).

The inverse function of a common meadow is not an involution because $(0^{-1})^{-1} = \mathbf{a}$. We will refer to meadows with zero-totalized inverse, that is, $0^{-1} = 0$, as *involution meadows* because inverse becomes an involution. By default a “meadow” is assumed to be an involutive meadow.

The key distinction between meadows and fields, which we consider to be so important that it justifies a different name, is the presence of an operator symbol for inverse in the signature (inversive notation, see [4]) or for division (divisive notation, see [4]), where divisive notation x/y is defined as $x \cdot y^{-1}$. A major consequence is that fractions can be viewed as terms over the signature of (common) meadows. Another distinction between meadows and fields is that we do not require a meadow to satisfy the separation axiom $0 \neq 1$.

The paper is structured as follows: below we conclude this section with a brief introduction to some aspects of involutive meadows that will play a role later on, and a discussion on why common meadows can be preferred over involutive meadows. In Section 2 we formally define common meadows and present some elementary results. In Section 3 we define “common cancellation meadows” and provide a basis theorem for common cancellation meadows of characteristic zero, which we consider our main result. Section 4 contains some concluding remarks.

1.1 Common Meadows versus Involution Meadows

Involution meadows, where instead of choosing $1/0 = \mathbf{a}$, one calculates with $1/0 = 0$, constitute a different solution to the question how to deal with the value of $1/0$ once the design decision has been made to work with the signature of meadows, that is to include a function name for inverse or for division (or both) in an extension of the syntax of fields. Involution meadows

$(x + y) + z = x + (y + z)$	$x \cdot y = y \cdot x$
$x + y = y + x$	$1 \cdot x = x$
$x + 0 = x$	$x \cdot (y + z) = x \cdot y + x \cdot z$
$x + (-x) = 0$	$(x^{-1})^{-1} = x$
$(x \cdot y) \cdot z = x \cdot (y \cdot z)$	$x \cdot (x \cdot x^{-1}) = x$

Table 1: The set Md of axioms for (involutive) meadows

feature a definite advantage over common meadows in that, by avoiding an extension of the domain with an additional value, theoretical work is very close to classical algebra of fields. This conservation property, conserving the domain, of involutive meadows has proven helpful for the development of theory about involutive meadows in [2, 1, 6, 4, 9, 8]. Earlier and comparable work on the equational theory of fields was done by Komori [13] and Ono [16]: in 1975, Komori introduced the name *desirable pseudo-field* for what was introduced as a “meadow” in [8].¹

An equational axiomatization Md of involutive meadows is given in Table 1, where $^{-1}$ binds stronger than \cdot , which in turn binds stronger than $+$. From the axioms in Md the following equations are derivable:

$$\begin{array}{ll}
0 \cdot x = 0, & 0^{-1} = 0, \\
x \cdot (-y) = -(x \cdot y), & (-x)^{-1} = -(x^{-1}), \\
-(-x) = x, & (x \cdot y)^{-1} = x^{-1} \cdot y^{-1}.
\end{array}$$

Involutive *cancellation meadows* are involutive meadows in which the following cancellation law holds:

$$(x \neq 0 \wedge x \cdot y = x \cdot z) \rightarrow y = z. \quad (\text{CL})$$

Involutive cancellation meadows form an important subclass of involutive meadows: in [1, Thm.3.1] it is shown that the axioms in Table 1 constitute a complete axiomatization of the equational theory of involutive cancellation meadows. We will use a consequence of this result in Section 3.

A definite disadvantage of involutive meadows against common meadows is that $1/0 = 0$ is quite remote from common intuitions regarding the partiality of division.

1.2 Motivating a Preference for Common Meadows

Whether common meadows are to be preferred over involutive meadows depends on the applications one may have in mind. We envisage as an application area the development of alternative foundations of elementary mathematics from a perspective of abstract datatypes, term rewriting, and mathematical logic. For that objective we consider common meadows to be the preferred option over involutive meadows. At the same time it can be acknowledged that a systematic investigation of involutive meadows constitutes a necessary stage in

¹[8] was published in 2007; the finding of [13, 16] is mentioned in [4] (2011) and was found via Ono’s 1983-paper [16].

the development of a theory of common meadows by facilitating in a simplified setting the determination of results which might be obtained about common meadows. Indeed each result about involutive meadows seems to suggest a (properly adapted) counterpart in the setting of common meadows, while proving or disproving such counterparts is not an obvious matter.

2 Common Meadows

In this section we formally define “common meadows” by fixing their signature and providing an equational axiomatization. Then, we consider some conditional equations that follow from this axiomatization. Finally, we discuss some conditional laws that can be used to define an important subclass of common meadows.

2.1 Meadow Signatures

The signature Σ_f^S of fields (and rings) contains a sort (domain) S , two constants 0 , and 1 , two two-place functions $+$ (addition) and \cdot (multiplication) and the one-place function $-$ (minus) for the inverse of addition.

We write Σ_{md}^S for the signature of meadows in inversive notation:

$$\Sigma_{md}^S = \Sigma_f^S \cup \{-^{-1}: S \rightarrow S\},$$

and we write $\Sigma_{md,\mathbf{a}}^S$ for the signature of meadows in inversive notation with an \mathbf{a} -totalized inverse operator:

$$\Sigma_{md,\mathbf{a}}^S = \Sigma_{md}^S \cup \{\mathbf{a}: S\}.$$

The interpretation of \mathbf{a} is called the additional value and we write $\hat{\mathbf{a}}$ for this value. Application of any function to the additional value returns that same value.

When the name of the carrier is fixed it need not be mentioned explicitly in a signature. Thus, with this convention in mind, Σ_{md} represents Σ_{md}^S and so on. If we want to make explicit that we consider terms over some signature Σ with variables in set X , we write $\Sigma(X)$.

Given a field several meadow signatures and meadows can be connected with it. This will now be exemplified with the field \mathbb{Q} of rational numbers. The following meadows are distinguished in this case:

\mathbb{Q}_0 , the meadow of rational numbers with zero-totalized inverse: $\Sigma(\mathbb{Q}_0) = \Sigma_{md}^{\mathbb{Q}}$.

$\mathbb{Q}_{\mathbf{a}}$, the meadow of rational numbers with \mathbf{a} -totalized inverse: $\Sigma(\mathbb{Q}_{\mathbf{a}}) = \Sigma_{md,\mathbf{a}}^{\mathbb{Q}}$. The additional value $\hat{\mathbf{a}}$ interpreting \mathbf{a} has been taken outside $|\mathbb{Q}|$ so that $|\mathbb{Q}_{\hat{\mathbf{a}}}| = |\mathbb{Q}| \cup \{\hat{\mathbf{a}}\}$.

2.2 Axioms for Common Meadows

The axioms in Table 2 define the class (variety) of *common meadows*, where we adopt the convention that $-^{-1}$ binds stronger than \cdot , which in turn binds stronger than $+$. Some comments: Axiom (14) implies \mathbf{a} 's propagation through all operations, and for the same reason, axiom (10) has its particular form. Axiom (4) is a variant of the common axiom on additional inverse, which also serves \mathbf{a} 's propagation. Axioms (11) and (12) are further equations needed for manipulation of $(\dots)^{-1}$ -expressions.

$(x + y) + z = x + (y + z)$	(1)
$x + y = y + x$	(2)
$x + 0 = x$	(3)
$x + (-x) = 0 \cdot x$	(4)
$(x \cdot y) \cdot z = x \cdot (y \cdot z)$	(5)
$x \cdot y = y \cdot x$	(6)
$1 \cdot x = x$	(7)
$x \cdot (y + z) = x \cdot y + x \cdot z$	(8)
$-(-x) = x$	(9)
$x \cdot x^{-1} = 1 + 0 \cdot x^{-1}$	(10)
$(x \cdot y)^{-1} = x^{-1} \cdot y^{-1}$	(11)
$(1 + 0 \cdot x)^{-1} = 1 + 0 \cdot x$	(12)
$0^{-1} = \mathbf{a}$	(13)
$x + \mathbf{a} = \mathbf{a}$	(14)

Table 2: $\text{Md}_{\mathbf{a}}$, an independent set of axioms for common meadows

We note that the axiom set $\text{Md}_{\mathbf{a}}$ is independent, which can be easily demonstrated with the tool *Mace4*, and that the typical identities for common meadows established by the following Propositions 2.2.1 and 2.2.2 were checked with the theorem prover *Prover9*, see [15] for both these tools.

Proposition 2.2.1. *Equations that follow from $\text{Md}_{\mathbf{a}}$ (see Table 2):*

$0 \cdot 0 = 0$	(e1)
$0 \cdot (x \cdot x) = 0 \cdot x$	(e2)
$-0 = 0,$	(e3)
$0 \cdot x = 0 \cdot (-x)$	(e4)
$0 \cdot (x \cdot y) = 0 \cdot (x + y)$	(e5)
$-(x \cdot y) = x \cdot (-y)$	(e6)
$(-x) \cdot (-y) = x \cdot y$	(e7)
$(-1) \cdot x = -x$	(e8)
$1^{-1} = 1$	(e9)
$(x \cdot x^{-1}) \cdot x^{-1} = x^{-1}$	(e10)
$(-x)^{-1} = -(x^{-1})$	(e11)
$(x \cdot x^{-1})^{-1} = x \cdot x^{-1}$	(e12)
$(x^{-1})^{-1} = x + 0 \cdot x^{-1}$	(e13)

and

$$x \cdot \mathbf{a} = \mathbf{a} \tag{e14}$$

$$-\mathbf{a} = \mathbf{a} \tag{e15}$$

$$\mathbf{a}^{-1} = \mathbf{a} \tag{e16}$$

Proof. Most derivations are trivial.

(e1). By axioms (3), (7), (8), (2) we find $x = (1+0) \cdot x = x+0 \cdot x = 0 \cdot x+x$, hence $0 = 0 \cdot 0+0$, so by axiom (3), $0 = 0 \cdot 0$.

(e2). First derive $0 \cdot (x \cdot x) = (0 \cdot x) \cdot (0 \cdot x)$ and $0 \cdot x = 0 \cdot 1 + 0 \cdot 0 \cdot x = 0 \cdot (1 + 0 \cdot x)$. Hence,
 $0 \cdot (x \cdot x) = 0 \cdot (1+0 \cdot x) \cdot (1+0 \cdot x) \stackrel{(12)}{=} 0 \cdot (1+0 \cdot x) \cdot (1+0 \cdot x)^{-1} \stackrel{(10)}{=} 0 \cdot (1+0 \cdot (1+0 \cdot x)^{-1}) \stackrel{(12)}{=} 0 \cdot (1 + 0 \cdot (1 + 0 \cdot x)) = 0 \cdot (1 + 0 \cdot x) = 0 \cdot x$.

(e3). By axioms (3), (2), (4) and (e1) we find $-0 = (-0) + 0 = 0 + (-0) = 0 \cdot 0 = 0$.

(e4). By axioms (2), (4), (9) we find $0 \cdot x = x + (-x) = (-x) + -(-x) = 0 \cdot (-x)$.

(e5). First note $0 \cdot x + 0 \cdot x = (0 + 0) \cdot x = 0 \cdot x$. By axioms (2) – (4), (6), (8), and (e2) we find $0 \cdot (x + y) = 0 \cdot ((x + y) \cdot (x + y)) = (0 \cdot x + 0 \cdot (x \cdot y)) + (0 \cdot y + 0 \cdot (x \cdot y)) = (0 + 0 \cdot y) \cdot x + (0 + 0 \cdot x) \cdot y = 0 \cdot (x \cdot y) + 0 \cdot (x \cdot y) = 0 \cdot (x \cdot y)$.

(e6). We give a detailed derivation:

$$\begin{aligned} -(x \cdot y) &= -(x \cdot y) + 0 \cdot -(x \cdot y) && \text{by } x = x + 0 \cdot x \\ &= -(x \cdot y) + 0 \cdot (x \cdot y) && \text{by (e4)} \\ &= -(x \cdot y) + x \cdot (0 \cdot y) && \text{by axioms (5) and (6)} \\ &= -(x \cdot y) + x \cdot (y + (-y)) && \text{by axiom (4)} \\ &= -(x \cdot y) + (x \cdot y + x \cdot (-y)) && \text{by axiom (8)} \\ &= -(x \cdot y) + x \cdot y + x \cdot (-y) && \text{by axiom (1)} \\ &= 0 \cdot (x \cdot y) + x \cdot (-y) && \text{by axioms (2) and (4)} \\ &= 0 \cdot (x \cdot -y) + x \cdot (-y) && \text{by axioms (6) and (5), and (e4)} \\ &= x \cdot (-y). && \text{by } x = 0 \cdot x + x \end{aligned}$$

(e7). By (e6), $(-x) \cdot (-y) = -((-x) \cdot y) = -(y \cdot (-x)) = -(-(y \cdot x)) = x \cdot y$.

(e8). From (e6) with $y = 1$ we find $-x = -(x \cdot 1) = x \cdot (-1) = (-1) \cdot x$.

(e9). By (e1) and axioms (3) and (12), $1^{-1} = (1 + 0 \cdot 0)^{-1} = 1 + 0 \cdot 0 = 1$.

(e10). By axioms (10) and (e2), $(x \cdot x^{-1}) \cdot x^{-1} = (1 + 0 \cdot x^{-1}) \cdot x^{-1} = x^{-1} + 0 \cdot x^{-1} = x^{-1}$.

(e11). By (e10) and (e7), $(-1)^{-1} = -1 \cdot (-1)^{-1} \cdot (-1)^{-1} = -1 \cdot ((-1) \cdot (-1))^{-1} = -1 \cdot 1^{-1} = -1$.
Hence, $(-x)^{-1} = (-1 \cdot x)^{-1} = (-1)^{-1} \cdot x^{-1} = -1 \cdot x^{-1} = -(x^{-1})$.

(e12). By axioms (10) and (12), $x \cdot x^{-1} = 1 + 0 \cdot x^{-1} = (1 + 0 \cdot x^{-1})^{-1} = (x \cdot x^{-1})^{-1}$.

(e13). By (e10), $(x^{-1})^{-1} = x^{-1} \cdot (x^{-1})^{-1} \cdot (x^{-1})^{-1} \stackrel{(11)}{=} (x \cdot x^{-1})^{-1} \cdot (x^{-1})^{-1} \stackrel{(e12)}{=} (x \cdot x^{-1}) \cdot (x^{-1})^{-1} = x \cdot (x \cdot x^{-1})^{-1} \stackrel{(e12)}{=} x \cdot (x \cdot x^{-1}) = x \cdot (1 + 0 \cdot x^{-1}) = x + 0 \cdot x \cdot x^{-1} = x + 0 \cdot (1 + 0 \cdot x^{-1}) = x + 0 \cdot x^{-1}$.

(e14). By axioms (8) and (14), $\mathbf{a} \cdot (1 + x) = \mathbf{a} + \mathbf{a} \cdot x = \mathbf{a}$, hence $\mathbf{a} \cdot x = \mathbf{a} \cdot ((1 + (-1)) + x) = \mathbf{a} \cdot (1 + (-1 + x)) = \mathbf{a}$, and thus $x \cdot \mathbf{a} = \mathbf{a}$ by axiom (6).

(e15). By axioms (6) and (5), and (e6) and (e14), $-\mathbf{a} = -(\mathbf{a} \cdot 1) = \mathbf{a} \cdot (-1) = \mathbf{a}$.

(e16). By axioms (13) and (11), and (e14), $\mathbf{a}^{-1} = (0 \cdot \mathbf{a})^{-1} = 0^{-1} \cdot \mathbf{a}^{-1} = \mathbf{a} \cdot \mathbf{a}^{-1} = \mathbf{a}$.

□

The next proposition establishes a generalization of a familiar identity concerning the addition of fractions.

Proposition 2.2.2. $\text{Md}_{\mathbf{a}} \vdash x \cdot y^{-1} + u \cdot v^{-1} = (x \cdot v + u \cdot y) \cdot (y \cdot v)^{-1}$.

Proof. We first derive

$$\begin{aligned}
x \cdot y \cdot y^{-1} &= x \cdot (1 + 0 \cdot y^{-1}) && \text{by axiom (10)} \\
&= x + 0 \cdot x \cdot y^{-1} \\
&= x + 0 \cdot x + 0 \cdot y^{-1} && \text{by (e5)} \\
&= x + 0 \cdot y^{-1}.
\end{aligned} \tag{15}$$

Hence,

$$\begin{aligned}
(x \cdot v + u \cdot y) \cdot (y \cdot v)^{-1} &= x \cdot y^{-1} \cdot v \cdot v^{-1} + u \cdot v^{-1} \cdot y \cdot y^{-1} \\
&= (x \cdot y^{-1} + 0 \cdot v^{-1}) + (u \cdot v^{-1} + 0 \cdot y^{-1}) && \text{by (15)} \\
&= (x \cdot y^{-1} + 0 \cdot y^{-1}) + (u \cdot v^{-1} + 0 \cdot v^{-1}) \\
&= x \cdot y^{-1} + u \cdot v^{-1}.
\end{aligned}$$

□

We end this section with two more propositions that characterize typical properties of common meadows and that are used in the proof of Theorem 3.2.1. The first of these establishes that each (possibly open) term over $\Sigma_{\text{md}, \mathbf{a}}$ has a simple representation in the syntax of meadows.

Proposition 2.2.3. *For each term t over $\Sigma_{\text{md}, \mathbf{a}}(X)$ with variables in X there exist terms r_1, r_2 over $\Sigma_f(X)$ such that $\text{Md}_{\mathbf{a}} \vdash t = r_1 \cdot r_2^{-1}$ and $\text{VAR}(t) = \text{VAR}(r_1) \cup \text{VAR}(r_2)$.*

Proof. By induction on the structure of t , where the $\text{VAR}(t)$ -property follows easily in each case.

If $t \in \{0, 1, x, \mathbf{a}\}$, this follows trivially (for the first three cases use $1^{-1} = 1$).

Case $t \equiv t_1 + t_2$. By Proposition 2.2.2.

Case $t \equiv t_1 \cdot t_2$. Trivial.

Case $t \equiv -t_1$. By Proposition 2.2.1 (e6).

Case $t \equiv t_1^{-1}$. By induction there exist $r_i \in \Sigma_f(X)$ such that $\text{Md}_{\mathbf{a}} \vdash t_1 = r_1 \cdot r_2^{-1}$. Now derive $t_1^{-1} = r_1^{-1} \cdot (r_2^{-1})^{-1} = r_1^{-1} \cdot (r_2 + 0 \cdot r_2^{-1}) = r_2 \cdot r_1^{-1} + 0 \cdot r_1^{-1} + 0 \cdot r_2^{-1} = r_2 \cdot r_1^{-1} + 0 \cdot r_2^{-1}$ and apply Proposition 2.2.2.

□

The next proposition shows how a term of the form $0 \cdot t$ with t a (possibly open) term over $\Sigma_f(X)$ can be simplified (note that $0 \cdot x = 0$ is not valid, since $0 \cdot \mathbf{a} = \mathbf{a}$).

Proposition 2.2.4. *For each term t over $\Sigma_f(X)$, $\text{Md}_{\mathbf{a}} \vdash 0 \cdot t = 0 \cdot \sum_{x \in \text{VAR}(t)} x$, where $\sum_{x \in \emptyset} x = 0$.*

Proof. By induction on the structure of t , where equation (e5) (Proposition 2.2.1) covers the multiplicative case. \square

2.3 Conditional Equations

We discuss a number of conditional equations that will turn out useful, and we start off with a few that follow directly from $\text{Md}_{\mathbf{a}}$.

Proposition 2.3.1. *Conditional equations that follow from $\text{Md}_{\mathbf{a}}$ (see Table 2):*

$$x \cdot y = 1 \rightarrow 0 \cdot y = 0, \tag{ce1}$$

$$x \cdot y = 1 \rightarrow x^{-1} = y, \tag{ce2}$$

$$0 \cdot x = 0 \cdot y \rightarrow 0 \cdot (x \cdot y) = 0 \cdot x, \tag{ce3}$$

$$0 \cdot x \cdot y = 0 \rightarrow 0 \cdot x = 0, \tag{ce4}$$

$$0 \cdot (x + y) = 0 \rightarrow 0 \cdot x = 0, \tag{ce5}$$

$$0 \cdot x^{-1} = 0 \rightarrow 0 \cdot x = 0, \tag{ce6}$$

$$0 \cdot x = \mathbf{a} \rightarrow x = \mathbf{a}. \tag{ce7}$$

Proof. Most derivations are trivial.

(ce1). By equations (e2) and (e5), $0 \cdot x \cdot y = 0 \cdot x \cdot y \cdot y = 0 \cdot x \cdot y + 0 \cdot y \cdot y = (0 \cdot x + 0 \cdot y) \cdot y$, and hence by assumption, $0 = 0 \cdot 1 = 0 \cdot x \cdot y = (0 \cdot x + 0 \cdot y) \cdot y = 0 \cdot x \cdot y + 0 \cdot y \cdot y = 0 + 0 \cdot y = 0 \cdot y$.

(ce2). By assumption and axioms (11) and (12), $x^{-1} \cdot y^{-1} = 1$, and thus by (ce1), $0 \cdot x^{-1} = 0$, so by axiom (10), $y = (1 + 0 \cdot x^{-1}) \cdot y = (x \cdot x^{-1}) \cdot y = (x \cdot y) \cdot x^{-1} = x^{-1}$.

(ce3). By assumption, equation (e5), and axiom (8), $0 \cdot (x \cdot y) = 0 \cdot x + 0 \cdot y = 0 \cdot x + 0 \cdot x = 0 \cdot x$.

(ce4). By assumption, $0 \cdot x = 0 \cdot x + 0 \cdot x \cdot y = x \cdot (0 + 0 \cdot y) = 0 \cdot (x \cdot y) = 0$.

(ce5). Apply equation (e5) to (ce4).

(ce6). By axiom (10) and assumption, $x \cdot x^{-1} = 1 + 0 \cdot x^{-1} = 1$, so by (ce1), $0 \cdot x = 0$.

(ce7). By $x = x + 0 \cdot x$ and assumption, $x = x + \mathbf{a} = \mathbf{a}$.

\square

Note that (ce1) and (ce2) immediately imply

$$x \cdot y = 1 \rightarrow 0 \cdot x^{-1} = 0.$$

In Table 3 we define various conditional laws that we will use to single out certain classes of common meadows in Section 3: the Normal Value Law (NVL), the Additional Value Law (AVL), and the Common Inverse Law (CIL). Here we use the adjective “normal” to express that values different from \mathbf{a} (more precisely, the interpretation of \mathbf{a}) are at stake. We conclude this section by interrelating these laws.

$x \neq \mathbf{a} \rightarrow 0 \cdot x = 0$	Normal Value Law	(NVL)
$x^{-1} = \mathbf{a} \rightarrow 0 \cdot x = x$	Additional Value Law	(AVL)
$x \neq 0 \wedge x \neq \mathbf{a} \rightarrow x \cdot x^{-1} = 1$	Common Inverse Law	(CIL)

Table 3: Some conditional laws for common meadows

Proposition 2.3.2.

1. $\text{Md}_{\mathbf{a}} + \text{NVL} \vdash (x \cdot y = \mathbf{a} \wedge x \neq \mathbf{a}) \rightarrow y = \mathbf{a}$,
2. $\text{Md}_{\mathbf{a}} + \text{NVL} \vdash x^{-1} \neq \mathbf{a} \rightarrow 0 \cdot x = 0$,
3. $\text{Md}_{\mathbf{a}} + \text{NVL} + \text{AVL} \vdash \text{CIL}$,
4. $\text{Md}_{\mathbf{a}} + \text{CIL} \vdash \text{NVL}$,
5. $\text{Md}_{\mathbf{a}} + \text{CIL} \vdash \text{AVL}$.

Proof.

1. By NVL, $x \neq \mathbf{a} \rightarrow 0 \cdot x = 0$, so $0 \cdot y = (0 \cdot x) \cdot y = 0 \cdot (x \cdot y) = 0 \cdot \mathbf{a} = \mathbf{a}$ and hence $y = (1 + 0) \cdot y = y + 0 \cdot y = y + \mathbf{a} = \mathbf{a}$.
2. By NVL, $0 \cdot x^{-1} = 0$ and hence by axiom (10), $x \cdot x^{-1} = 1$ and by (ce1), $0 \cdot x = 0$.
3. From $x \neq \mathbf{a}$ we find $0 \cdot x = 0$. There are two cases: $x^{-1} = \mathbf{a}$ which implies by AVL that $x = 0$ contradicting the assumptions of CIL, and $x^{-1} \neq \mathbf{a}$ which implies by NVL that $0 \cdot x^{-1} = 0$, and this implies $x \cdot x^{-1} = 1$ by axiom (10).
4. Assume that $x \neq \mathbf{a}$. If $x = 0$ then also $0 \cdot x = 0$. If $x \neq 0$ then by CIL, $x \cdot x^{-1} = 1$, so $0 \cdot x = 0$ by (ce1).
5. We distinguish three cases: $x = 0$, $x = \mathbf{a}$, and $x \neq 0 \wedge x \neq \mathbf{a}$. In the first two cases it immediately follows that $0 \cdot x = x$. In the last case it follows by CIL that $x \cdot x \cdot x^{-1} = x$, so $x^{-1} = \mathbf{a}$ implies $x = \mathbf{a}$, and thus $x = 0 \cdot x$.

□

3 Models and Model Classes

In this section we define “common cancellation meadows” as common meadows that satisfy the so-called “inverse cancellation law”, a law that is equivalent with the Common Inverse Law CIL. Then, we provide a basis theorem for common cancellation meadows of characteristic zero.

3.1 Common Cancellation Meadows

In [1, Thm.3.1] we prove a generic basis theorem that implies that the axioms in Table 1 constitute a complete axiomatization of the equational theory of the involutive cancellation meadows (over signature Σ_{md}). The cancellation law used in that result (that is, CL in Section 1.1) has various equivalent versions, and a particular one is $x \neq 0 \rightarrow x \cdot x^{-1} = 1$, a version that is close to CIL.

Below we define common cancellation meadows, using a cancellation law that is equivalent with CIL, but first we establish a correspondence between models of $\text{Md}_{\mathbf{a}} + \text{NVL} + \text{AVL}$ and involutive cancellation meadows.

Proposition 3.1.1.

1. *Every field can be extended with an additional value $\hat{\mathbf{a}}$ and subsequently it can be expanded with a constant \mathbf{a} and an inverse function in such a way that the equations of common meadows as well as NVL and AVL are satisfied, where the interpretation of \mathbf{a} is $\hat{\mathbf{a}}$.*
2. *A model of $\text{Md}_{\mathbf{a}} + \text{NVL} + \text{AVL}$ extends a field with an additional value $\hat{\mathbf{a}}$ (the interpretation of \mathbf{a}) and expands it with the \mathbf{a} -totalized inverse.*

Proof. Statement 1 follows immediately. To prove 2, consider the substructure of elements b of the domain that satisfy $0 \cdot b = 0$. Only $\hat{\mathbf{a}}$ is outside this subset. For b with $0 \cdot b = 0$ we must check that $0 \cdot b^{-1} = 0$ unless $b = 0$. To see this distinguish two cases: $b^{-1} = \mathbf{a}$ (which implies $b = 0$ with help of AVL), and $b^{-1} \neq \mathbf{a}$ which implies $0 \cdot b^{-1} = 0$ by NVL. \square

As a consequence, we find the following result.

Theorem 3.1.2. *The models of $\text{Md}_{\mathbf{a}} + \text{NVL} + \text{AVL}$ that satisfy $0 \neq 1$ are in one-to-one correspondence with the involutive cancellation meadows satisfying Md (see Table 1).*

Proof. An involutive cancellation meadow can be expanded to a model of $\text{Md}_{\mathbf{a}} + \text{NVL} + \text{AVL}$ by extending its domain with a constant $\hat{\mathbf{a}}$ in such a way that the equations of common meadows as well as NVL and AVL are satisfied, where the interpretation of \mathbf{a} is $\hat{\mathbf{a}}$ (cf. Proposition 3.1.1.1).

Conversely, given a model \mathbb{M} of $\text{Md}_{\mathbf{a}} + \text{NVL} + \text{AVL}$, we construct a cancellation meadow \mathbb{M}' as follows: $|\mathbb{M}'| = |\mathbb{M}| \setminus \{\hat{\mathbf{a}}\}$ with $\hat{\mathbf{a}}$ the interpretation of \mathbf{a} , and $0^{-1} = 0$ (by $0 \neq 1$, $|\mathbb{M}'|$ is non-empty). We find by NVL that $0 \cdot x = 0$ and by CIL (thus by NVL + AVL, cf. Proposition 2.3.2.3) that $x \neq 0 \rightarrow x \cdot x^{-1} = 1$, which shows that \mathbb{M}' is a cancellation meadow. \square

We define a *common cancellation meadow* as a common meadow that satisfies the following *inverse cancellation law* (ICL):

$$(x \neq 0 \wedge x \neq \mathbf{a} \wedge x^{-1} \cdot y = x^{-1} \cdot z) \rightarrow y = z. \quad (\text{ICL})$$

The class CCM of common cancellation meadows is axiomatized by $\text{Md}_{\mathbf{a}} + \text{CIL}$ in Table 2 and Table 3, respectively. In combination with $\text{Md}_{\mathbf{a}}$, the laws ICL and CIL are equivalent: first, $\text{Md}_{\mathbf{a}} + \text{ICL} \vdash \text{CIL}$ because

$$(x \neq 0 \wedge x \neq \mathbf{a}) \xrightarrow{(\text{e10})} (x \neq 0 \wedge x \neq \mathbf{a} \wedge x^{-1} \cdot x \cdot x^{-1} = x^{-1} \cdot 1) \xrightarrow{\text{ICL}} x \cdot x^{-1} = 1.$$

Conversely, $\text{Md}_{\mathbf{a}} + \text{CIL} \vdash \text{ICL}$:

$$(x \neq 0 \wedge x \neq \mathbf{a} \wedge x^{-1} \cdot y = x^{-1} \cdot z) \rightarrow x \cdot x^{-1} \cdot y = x \cdot x^{-1} \cdot z \xrightarrow{\text{CIL}} y = z.$$

$\underline{n+1} \cdot (\underline{n+1})^{-1} = 1$	$(n \in \mathbb{N})$	(C_0)
$\underline{0} = 0$	(axioms for numerals, $n \in \mathbb{N}$ and $n \geq 1$)	
$\underline{1} = 1$		
$\underline{n+1} = \underline{n} + 1$		

Table 4: C_0 , the set of axioms for meadows of characteristic zero and numerals

3.2 A Basis Theorem for Common Cancellation Meadows of Characteristic Zero

As in our paper [2], we use numerals \underline{n} and the axiom scheme C_0 defined in Table 4 to single out common cancellation meadows of characteristic zero. In this section we prove that $\text{Md}_{\mathbf{a}} + C_0$ constitutes an axiomatization for common cancellation meadows of characteristic zero. In [2, Cor.2.7] we prove that $\text{Md} + C_0$ (for Md see Table 1) constitutes an axiomatization for involutive cancellation meadows of characteristic zero. We define CCM_0 as the class of common cancellation meadows of characteristic zero.

We further write $\frac{t}{r}$ (and sometimes t/r in plain text) for $t \cdot r^{-1}$.

Theorem 3.2.1. $\text{Md}_{\mathbf{a}} + C_0$ is a basis for the equational theory of CCM_0 .

Proof. Soundness holds by definition of CCM_0 .

In order to prove completeness, we consider two cases.

Case 1. Assume $\text{CCM}_0 \models t = \mathbf{a}$. By Proposition 2.2.3 we can bring t in the form t_1/t_2 with t_1, t_2 polynomials over $\Sigma_f(X)$, thus

$$\text{CCM}_0 \models t = \frac{t_1}{t_2}. \quad (16)$$

Then in each model $\mathbb{M} \in \text{CCM}_0$, $t_1 = \mathbf{a}$ or $t_2 = \mathbf{a}$ or $t_2 = 0$. By definition of CCM_0 , the first two cases are excluded because in \mathbb{M} each variable of both t_1 and t_2 can be interpreted as a non- $\hat{\mathbf{a}}$ value (e.g. 0), and then t_1 (t_2) evaluates to a value different from $\hat{\mathbf{a}}$ in \mathbb{M} . Thus (16) implies that $t_2 = 0$ holds in each field. Apparently it holds in each field that

$$t_2 = 0 \cdot \sum_{x \in \text{VAR}(t_2)} x$$

and it must be the case that in the polynomial t_2 , all coefficients are 0. Furthermore, if for a closed term s over Σ_f it holds that $s = 0$ in each field, then there exists an equational proof of $s = 0$ in which substitutions happen first and only closed instances of the axioms for commutative rings are used.² This implies that $\text{Md}_{\mathbf{a}} \vdash s = 0$ because by Proposition 2.2.4, $\text{Md}_{\mathbf{a}} \vdash 0 \cdot s = 0$ and thus each application of $\text{Md}_{\mathbf{a}}$ -axiom (4) (that is $x + (-x) = 0 \cdot x$), reduces to $u + (-u) = 0$ with u a closed term over Σ_f .

It follows that $\text{Md}_{\mathbf{a}} \vdash t_2 = 0$, so by axiom (13) and equation (e14), $\text{Md}_{\mathbf{a}} \vdash t = t_1/t_2 = \mathbf{a}$.

²Without loss of generality it can be assumed that in equational proofs, substitutions happen first, see, e.g. [12].

Case 2. Assume $\text{CCM}_0 \models t = r$ and $\text{CCM}_0 \not\models t = \mathbf{a}$ (and thus $\text{CCM}_0 \not\models r = \mathbf{a}$). By Proposition 2.2.3 we can bring t in the form t_1/t_2 and r in the form r_1/r_2 with t_i, r_i polynomials over $\Sigma_f(X)$, thus

$$\text{CCM}_0 \models \frac{t_1}{t_2} = \frac{r_1}{r_2}. \quad (17)$$

Using axiom (12) it can be guaranteed that in t_2 and r_2 no summands with coefficient 0 occur.³ We will first argue that (17) implies that the following three equations are valid in CCM_0 :

$$0 \cdot t_2^{-1} = 0 \cdot r_2^{-1}, \quad (18)$$

$$0 \cdot t_1 + 0 \cdot t_2 = 0 \cdot r_1 + 0 \cdot r_2, \quad (19)$$

$$t_2 \cdot r_2 \cdot (t_1 \cdot r_2 + (-r_1) \cdot t_2) + 0 \cdot t_2^{-1} + 0 \cdot r_2^{-1} = 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1}. \quad (20)$$

Ad (18). Assume this is not the case, then there exists a common cancellation meadow $\overline{\mathbb{M}} \in \text{CCM}_0$ and an interpretation of the variables in t_2 and r_2 such that one of t_2^{-1} and r_2^{-1} is interpreted as $\hat{\mathbf{a}}$ (the interpretation of \mathbf{a}), and the other is not. This contradicts (17).

Ad (19). This equation characterizes that t_1/t_2 and r_1/r_2 contain the same variables, and is related to Proposition 2.2.4. Assume this is not the case, say t_1 and/or t_2 contains a variable x that does not occur in r_1 and r_2 . Since $\text{CCM}_0 \not\models r_1/r_2 = \mathbf{a}$, there is an instance of r_i 's variables, say \overline{r}_i such that $\text{CCM}_0 \models \overline{r}_1/\overline{r}_2 \neq \mathbf{a}$. But then x can be instantiated with \mathbf{a} , which contradicts (17).

Ad (20). It follows from (17) that in (20) both the lefthand-side and the righthand-side equal zero in all involutive cancellation meadows. By Theorem 3.1.2 we find $\text{CCM} \models (20)$, and hence $\text{CCM}_0 \models (20)$.

We now argue that (18) – (20) are derivable from $\text{Md}_{\mathbf{a}} + \text{C}_0$, and that from those (17) is derivable from $\text{Md}_{\mathbf{a}} + \text{C}_0$.

Ad (18). The statement $\text{CCM}_0 \models 0 \cdot t_2^{-1} = 0 \cdot r_2^{-1}$ implies that t_2 and r_2 have the same zeros in the algebraic closure $\overline{\mathbb{Q}}$ of \mathbb{Q} (if this were not the case, then $\overline{\mathbb{Q}}_{\mathbf{a}} \not\models 0 \cdot t_2^{-1} = 0 \cdot r_2^{-1}$, but $\overline{\mathbb{Q}}_{\mathbf{a}} \in \text{CCM}_0$). By the Nullstellensatz (see, e.g. [14, Thm.1.5 (Ch.IX)]), there exists $m \geq 1$ such that $(t_2)^m \in I$, the ideal generated by r_2 , so $(t_2)^m$ is of the form $r_2 \cdot s$ for some polynomial s . Thus each factor of r_2 is a factor of $r_2 \cdot s$, and hence a factor of $(t_2)^m$. For the same reason, each factor of r_2 is one of t_2 . We may assume that the gcd of t_2 's coefficients is 1, and similar for r_2 : if not, then $t_2 = k \cdot t'$ with t' a polynomial with that property, and since k is a fixed numeral, we find $0 \cdot k = 0$ (also in fields with a characteristic that is a factor of k), and hence $0 \cdot t_2 = 0 \cdot t'$. By unique factorisation ([14, Cor.2.4 (Ch.IV)]), we find that t_2 and r_2 have equal primitive polynomials. Application of C_0 (for the case $t_2 = k \cdot t'$) and equation (e2) (that is, $0 \cdot (x \cdot x) = 0 \cdot x$) then yields

$$\text{Md}_{\mathbf{a}} + \text{C}_0 \vdash 0 \cdot t_2^{-1} = 0 \cdot r_2^{-1}. \quad (21)$$

Ad (19). From Proposition 2.2.4 and validity of (19) it follows that

$$\text{Md}_{\mathbf{a}} \vdash 0 \cdot t_1 + 0 \cdot t_2 = 0 \cdot \sum_{x \in \text{VAR}(t_1/t_2)} x = 0 \cdot \sum_{x \in \text{VAR}(r_1/r_2)} x = 0 \cdot r_1 + 0 \cdot r_2. \quad (22)$$

³ This was the reason to revise the preceding version of this paper (v3), in which $\text{Md}_{\mathbf{a}}$ is defined differently: axiom (12) was replaced by $1^{-1} = 1$ and equations (e2) and (e13) (Prop.2.2.1) were also axioms; this resulted in a weaker system from which axiom (12) cannot be derived, whereas the absence of summands with coefficient 0 in t_2 and r_2 can be proven with the current modification of $\text{Md}_{\mathbf{a}}$. For example, using $x + 0 \cdot y = x \cdot (1 + 0 \cdot y)$ (which follows easily), $(3x + 2) \cdot (6x^5 + 0 \cdot xy + 2yz^3 + 4)^{-1} = (3x + 2) \cdot ((6x^5 + 2yz^3 + 4) \cdot (1 + 0 \cdot xy))^{-1} = ((3x + 2) \cdot (1 + 0 \cdot xy)) \cdot (6x^5 + 2yz^3 + 4)^{-1}$.

Ad (20). We first derive

$$\begin{aligned}
\mathbf{Md}_a \vdash 0 \cdot t_1 + 0 \cdot t_2^{-1} &= 0 \cdot t_1 + 0 \cdot (1 + 0 \cdot t_2^{-1}) \\
&= 0 \cdot t_1 + 0 \cdot t_2 \cdot t_2^{-1} && \text{by axiom (10)} \\
&= 0 \cdot t_1 + 0 \cdot t_2 + 0 \cdot t_2^{-1},
\end{aligned}$$

and in a similar way one derives $\mathbf{Md}_a \vdash 0 \cdot r_1 + 0 \cdot r_2^{-1} = 0 \cdot r_1 + 0 \cdot r_2 + 0 \cdot r_2^{-1}$. Hence, we find with (21) and (22) that

$$\mathbf{Md}_a + \mathbf{C}_0 \vdash 0 \cdot t_1 + 0 \cdot t_2^{-1} = (0 \cdot t_1 + 0 \cdot t_2^{-1}) + (0 \cdot r_1 + 0 \cdot r_2^{-1}) \quad (23)$$

$$= 0 \cdot r_1 + 0 \cdot r_2^{-1}. \quad (24)$$

From $\text{CCM}_0 \models (20)$ it follows from the completeness result on the class of involutive meadows of characteristic zero (see [2, Cor.2.7]) that $\mathbf{Md} + \mathbf{C}_0 \vdash (20)$, and also that $\mathbf{Md} + \mathbf{C}_0 \vdash t_2 \cdot r_2 \cdot (t_1 \cdot r_2 + (-r_1) \cdot t_2) = 0$. Because all coefficients in this identity are integer expressions, \mathbf{C}_0 plays no role in these proofs, so that $\mathbf{Md} \vdash t_2 \cdot r_2 \cdot (t_1 \cdot r_2 + (-r_1) \cdot t_2) = 0$. Moreover, $0 \cdot s = 0$ can be applied to each coefficient s , so that $\mathbf{Md}_a \vdash t_2 \cdot r_2 \cdot (t_1 \cdot r_2 + (-r_1) \cdot t_2) = 0 \cdot (t_1 \cdot t_2 \cdot r_1 \cdot r_2)$, from which one easily finds $\mathbf{Md}_a \vdash (20)$.

Finally, we show the derivability of $t_1/t_2 = r_1/r_2$ in $\mathbf{Md}_a + \mathbf{C}_0$. Multiplying both sides of (20) with $(t_2 \cdot r_2)^{-1} \cdot (t_2 \cdot r_2)^{-1}$ implies by (e10), $0 \cdot x + 0 \cdot x = 0 \cdot x$, and equation (e2) that

$$\mathbf{Md}_a \vdash (t_2 \cdot r_2)^{-1} \cdot (t_1 \cdot r_2 + (-r_1) \cdot t_2) + 0 \cdot t_2^{-1} + 0 \cdot r_2^{-1} = 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1},$$

which implies by Proposition 2.2.2 that

$$\mathbf{Md}_a \vdash \frac{t_1}{t_2} + \frac{-r_1}{r_2} + 0 \cdot t_2^{-1} + 0 \cdot r_2^{-1} = 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1},$$

and thus

$$\mathbf{Md}_a \vdash \frac{t_1}{t_2} + \frac{-r_1}{r_2} + 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1} = 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1}. \quad (25)$$

Hence,

$$\begin{aligned}
\mathbf{Md}_a + \mathbf{C}_0 \vdash \frac{t_1}{t_2} &= \frac{t_1}{t_2} + 0 \cdot t_1 + 0 \cdot t_2^{-1} \\
&= \frac{t_1}{t_2} + 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1} && \text{by (23)} \\
&= \frac{t_1}{t_2} + \left(\frac{r_1}{r_2} + \frac{-r_1}{r_2}\right) + 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1} \\
&= \left(\frac{t_1}{t_2} + \frac{-r_1}{r_2}\right) + \frac{r_1}{r_2} + 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1} \\
&= \frac{r_1}{r_2} + 0 \cdot t_1 + 0 \cdot t_2^{-1} + 0 \cdot r_1 + 0 \cdot r_2^{-1} && \text{by (25)} \\
&= \frac{r_1}{r_2} + 0 \cdot r_1 + 0 \cdot r_2^{-1} && \text{by (24)} \\
&= \frac{r_1}{r_2}.
\end{aligned}$$

□

4 Concluding Remarks

Open Question. It is an open question whether there exists a basis result for the equational theory of CCM. We notice that in [5] a basis result for one-totalized non-involutive cancellation meadows is provided, where the multiplicative inverse of 0 is 1 and cancellation is defined as usual (that is, by the cancellation law CL in Section 1.1).

Common Intuitions and Related Work. Common meadows are motivated as being the most intuitive modelling of a totalized inverse function to the best of our knowledge. As stated in Section 1 (Introduction), the use of the constant \mathbf{a} is a matter of convenience only because it merely constitutes a derived constant with defining equation $\mathbf{a} = 0^{-1}$, which implies that all uses of \mathbf{a} can be removed.⁴ We notice that considering $\mathbf{a} = 0^{-1}$ as an error-value supports the intuition for the equations of $\text{Md}_{\mathbf{a}}$.

As a variant of involutive and common meadows, *partial meadows* are defined in [4]. The specification method used in this paper is based on meadows and therefore it is more simple, but less general than the construction of Broy and Wirsing [10] for the specification of partial datatypes.

The construction of common meadows is related to the construction of *wheels* by Carlström [11]. However, we have not yet found a structural connection between both constructions which differ in quite important details. For instance, wheels are involutive whereas common meadows are non-involutive.

Quasi-Cancellation Meadows of Characteristic Zero. Following Theorem 3.2.1, a common meadow of characteristic zero can alternatively be defined as a structure that satisfies all equations true of all common cancellation meadows of characteristic zero. We write CM_0 for the class of all common meadows of characteristic zero.

With this alternative definition in mind, we define a *common quasi-cancellation meadow of characteristic zero* as a structure that satisfies all conditional equations which are true of all common cancellation meadows of characteristic zero. We write CQCM_0 for the class of all common quasi-cancellation meadows of characteristic zero.

It is easy to show that CQCM_0 is strictly larger than CCM_0 . To see this one extends the signature of common meadows with a new constant c . Let $L_{ccm,0}$ be the set of conditional equations true of all structures in CCM_0 . We consider the initial algebra of $L_{ccm,0}$ in the signature extended with c . Now neither $L_{ccm,0} \vdash c = \mathbf{a}$ can hold (because c might be interpreted as say 1), nor $L_{ccm,0} \vdash 0 \cdot c = 0$ can hold (otherwise $L_{ccm,0} \vdash 0 = 0 \cdot \mathbf{a} = \mathbf{a}$ would hold). For that reason in the initial algebra of $L_{ccm,0}$ in the extended signature interprets c as an entity e in such a way that neither $c = \mathbf{a}$ nor $0 \cdot c = 0$ is satisfied. For that reason c will be interpreted by a new entity that refutes CIL.

CM_0 is strictly larger than CQCM_0 . To see this let $E_{ccm,0}$ denote the set of equations valid in all common cancellation meadows of characteristic zero. Again we add an extra constant b to the signature of common meadows. Consider the initial algebra I of $E_{ccm,0} + (b^{-1} = \mathbf{a})$ in the extended signature. In I the interpretation of b is a new object because it cannot be proven equal to 0 and not to \mathbf{a} and not to any other closed term over the signature of common meadows. Now we transform $E_{ccm,0} + (b^{-1} = \mathbf{a})$ into its set of closed consequences $E_{ccm,0}^{cl,b}$ over

⁴We notice that $0 = 1 + (-1)$, from which it follows that 0 can also be considered a derived constant over a reduced signature. Nevertheless, the removal of 0 from the signature of fields is usually not considered helpful.

the extended signature. We claim that $b = 0 \cdot b$ cannot be proven from $E_{ccm,0} + (b^{-1} = \mathbf{a})$. If that were the case at some stage in the derivation an \mathbf{a} must appear from which it follows that $b = \mathbf{a}$ is provable as well, because \mathbf{a} is propagated by all operations. But that cannot be the case as we have already concluded that b differs from \mathbf{a} in the initial algebra I_0 of $E_{ccm,0}^{cl,b}$. Thus, $b \neq \mathbf{a} \rightarrow 0 \cdot b = 0$ (an instance of NVL) is not valid in I_0 .

However, at this stage we do not know the answers to the following two questions:

- Is there a finite equational basis for the class CM_0 of common meadows of characteristic zero?
- Is there a finite conditional equational basis for the class CQCM_0 of common quasi-cancellation meadows of characteristic zero?

The Initial Common Meadow. In [7] we introduce *fracpairs* with a definition that is very close to that of the field of fractions of an integral domain. Fracpairs are defined over a commutative ring R that is *reduced*, i.e., R has no nonzero nilpotent elements. A fracpair over R is an expression $\frac{p}{q}$ with $p, q \in R$ (so $q = 0$ is allowed) modulo the equivalence generated by

$$\frac{x \cdot z}{y \cdot (z \cdot z)} = \frac{x}{y \cdot z}.$$

This rather simple equivalence appears to be a congruence with respect to the common meadow signature $\Sigma_{md,\mathbf{a}}$ when adopting natural definitions:

$$\begin{aligned} 0 &= \frac{0}{1}, & 1 &= \frac{1}{1}, & \mathbf{a} &= \frac{1}{0}, & \left(\frac{p}{q}\right) + \left(\frac{r}{s}\right) &= \frac{p \cdot s + r \cdot q}{q \cdot s}, \\ \left(\frac{p}{q}\right) \cdot \left(\frac{r}{s}\right) &= \frac{p \cdot r}{q \cdot s}, & -\left(\frac{p}{q}\right) &= \frac{-p}{q}, & \text{and} & \left(\frac{p}{q}\right)^{-1} &= \frac{q \cdot q}{p \cdot q}. \end{aligned}$$

In [7] we prove that the initial common meadow is isomorphic to the initial algebra of fracpairs over the integers \mathbb{Z} .⁵ Moreover, we prove that the initial algebra of fracpairs over \mathbb{Z} constitutes a homomorphic pre-image of the common meadow $\mathbb{Q}_{\mathbf{a}}$, and we define “rational fracpairs” over \mathbb{Z} that constitute an initial algebra that is isomorphic to $\mathbb{Q}_{\mathbf{a}}$. Finally, we consider some term rewriting issues for meadows.

These results reinforce our idea that common meadows can be used in the development of alternative foundations of elementary (educational) mathematics from a perspective of abstract datatypes, term rewriting and mathematical logic.

Acknowledgement. We thank Bas Edixhoven (Leiden University) for helpful comments concerning the proof of Theorem 3.2.1, including his suggestion to use [14] as a reference for this proof. Furthermore, we thank two anonymous reviewers for valuable comments. Finally, we thank John Tucker (Swansea) for valuable discussions that gave rise to re(formulating) axiom (12) of Table 2 in this version (v4).

⁵It should be mentioned that $\text{Md}_{\mathbf{a}}$ as presented in [7] differs from the current version, as explained in Footnote 3 (page 12). However, this makes no difference for the isomorphism result mentioned: with both versions of $\text{Md}_{\mathbf{a}}$ it easily follows that for every closed term p over Σ_f , $0.p = 0$, which is the essential property.

References

- [1] Bergstra, J.A., Bethke, I., and Ponse, A. (2013). Cancellation meadows: a generic basis theorem and some applications. *The Computer Journal*, 56(1):3-14.
DOI: <https://doi.org/10.1093/comjnl/bxs028>.
- [2] Bergstra, J.A., Bethke, I., and Ponse, A. (2015). Equations for formally real meadows. *Journal of Applied Logic*, 13(2):1-23.
DOI: <https://doi.org/10.1016/j.jal.2015.01.004>.
Also available at [arXiv:1310.5011v4](https://arxiv.org/abs/1310.5011v4) [math.RA, cs.LO], 13 January 2015.
- [3] Bergstra, J.A., Broy, M., Tucker, J.V., and Wirsing, M. (1981). On the power of algebraic specifications. In J. Gruska and M. Chytil (eds.), *Proc. MFCS'81*, Štrbské Pleso, Czechoslovakia. LNCS 118, pp. 193-204. Springer.
- [4] Bergstra, J.A. and Middelburg, C.A. (2011). Inversive meadows and divisive meadows. *Journal of Applied Logic*, 9(3):203-220.
DOI: <https://doi.org/10.1016/j.jal.2011.03.001>.
- [5] Bergstra, J.A. and Middelburg, C.A. (2015). Division by zero in non-involutive meadows. *Journal of Applied Logic*, 13(1):1-12.
DOI: <https://doi.org/10.1016/j.jal.2014.10.001>.
Also available at [arXiv:1406.2092v1](https://arxiv.org/abs/1406.2092v1) [math.RA], 9 June 2014.
- [6] Bergstra, J.A. and Ponse, A. (2017). Probability Functions in the context of signed involutive meadows. In P. James and M. Roggenbach (eds.), *WADT 2016*, LNCS 10644, pp. 73-87, Springer. DOI: https://doi.org/10.1007/978-3-319-72044-9_6. An extended version is available at [arXiv:1307.5173v4](https://arxiv.org/abs/1307.5173v4) [math.LO], 22 December 2016.
- [7] Bergstra, J.A. and Ponse, A. (2016). Fracpairs and fractions over a reduced commutative ring. *Indagationes Mathematicae*, 27(3):727-748.
DOI: <https://doi.org/10.1016/j.indag.2016.01.007>.
Also available at [arXiv:1411.4410v2](https://arxiv.org/abs/1411.4410v2) [math.RA], 22 January 2016.
- [8] Bergstra, J.A. and Tucker, J.V. (2007). The rational numbers as an abstract data type. *Journal of the ACM*, 54(2), Article 7 (25 pages).
- [9] Bethke, I. and Rodenburg, P.H. (2010). The initial meadows. *Journal of Symbolic Logic*, 75(3):888-895. DOI: <https://doi.org/10.2178/jsl/1278682205>.
- [10] Broy, M. and Wirsing, M. (1981). On the algebraic specification of nondeterministic programming languages. In E. Astesiano and C. Böhm (eds.), *Proc. CAAP 1981*, Genova, Italy. LNCS 112, pp. 162-179. Springer.
- [11] Carlström, J. (2004). Wheels - on division by zero. *Mathematical Structures in Computer Science*, 14(1):143-184. DOI: <https://doi.org/10.1017/S0960129503004110>.
- [12] Groote, J.F. (1990). A new strategy for proving omega-completeness applied to process algebra. In J.C.M. Baeten and J.W. Klop (eds.), *Theories of Concurrency: Unification and Extension* (CONCUR 1990, Amsterdam), LNCS 458, pp. 314-331. Springer.
- [13] Komori, Y. (1975). Free algebras over all fields and pseudo-fields. Report 10, pp. 9-15, Faculty of Science, Shizuoka University, Japan.

- [14] Lang, S. (2002). *Algebra*. Graduate Texts in Mathematics, Vol. 211, third revised edition. Springer.
- [15] McCune, W. (2008). The GUI: Prover9 and Mace4 with a Graphical User Interface. Prover9-Mace4-v05B.zip (March 14, 2008). Available at <https://www.cs.unm.edu/~mccune/prover9/gui/v05.html>.
- [16] Ono, H. (1983). Equational theories and universal theories of fields. *Journal of the Mathematical Society of Japan*, 35(2):289-306.
- [17] Wirsing, M. (1990). Algebraic specifications. In J. van Leeuwen (ed.), *Handbook of Theoretical Computer Science. Volume B: Formal models and semantics*, North-Holland, pp. 675-788.