UvA-DARE (Digital Academic Repository)

Discovery of a bow shock around Vela X-1

Kaper, L.; van Loon, J.T.; Augusteijn, T.; Goudfrooij, P.; Patat, F.; Waters, L.B.F.M.; Zijlstra, A.A.

DOI

[10.1086/310454](https://doi.org/10.1086/310454)

Publication date

1997

Published in

Astrophysical Journal

[Link to publication](#)

Citation for published version (APA):

Kaper, L., van Loon, J. T., Augusteijn, T., Goudfrooij, P., Patat, F., Waters, L. B. F. M., & Zijlstra, A. A. (1997). Discovery of a bow shock around Vela X-1. *Astrophysical Journal*, 475, L37-L40. <https://doi.org/10.1086/310454>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

DISCOVERY OF A BOW SHOCK AROUND VELA X-1

L. KAPER,¹ J. TH. VAN LOON,¹ T. AUGUSTEIJN,² P. GOUDFROOIJ,¹ F. PATAT,¹ L. B. F. M. WATERS,³ AND A. A. ZIJLSTRA¹

Received 1996 July 2; accepted 1996 October 31

ABSTRACT

We report the discovery of a symmetric bow shock around the well-known high-mass X-ray binary (HMXB) Vela X-1. Wind bow shocks are a ubiquitous phenomenon around OB-runaway stars, but now such a structure is found around an HMXB. The presence of a bow shock indicates that the system has a high (supersonic) velocity with respect to the interstellar medium. From the symmetry of the bow shock, the direction of motion and, moreover, the origin and age of the system can be derived. Our observation supports Blaauw's scenario for the formation of an OB-runaway star by the supernova explosion of the binary companion.

Subject headings: stars: binaries: close — stars: individual (HD 77581) — stars: mass loss — pulsars: individual (Vela X-1) — supernovae: general — open clusters and associations: individual (Vel OB1)

1. INTRODUCTION

The high-mass X-ray binary (HMXB) HD 77581 (Vela X-1) consists of a B0.5 Ib supergiant (Vidal, Wickramasingh, & Peterson 1973; Jones & Liller 1973) and an X-ray pulsar companion ($P_{\text{pulse}} = 283$ s; McClintock et al. 1976) in an 8.9 day orbit. Pulse-timing analysis of the X-ray pulsar has resulted in the accurate determination of its orbital parameters (Nagase 1989). The length of the X-ray eclipses (Watson & Griffiths 1977) indicates that the inclination of the orbital plane is close to 90° . Also, the radial-velocity curve of the supergiant companion is measured (van Paradijs et al. 1977; van Kerkwijk et al. 1995a), so that the masses of both binary components ($23.5^{+2.2}_{-1.5}$ and $1.88^{+0.69}_{-0.47} M_\odot$, 3σ limits) can be derived. It turns out that Vela X-1 is the most massive neutron star known (van Kerkwijk, van Paradijs, & Zuiderwijk 1995b), although the 3σ error bar still includes the canonical neutron star mass of $1.4 M_\odot$.

Evolutionary scenarios for massive binaries (see, e.g., van den Heuvel 1993 for a review) predict that the HMXB Vela X-1 originates from a $25 + 22.5 M_\odot$ binary (cf. Van Rensbergen, Vanbeveren, & De Loore 1996). When the initially most massive star (the primary) becomes a supergiant and starts filling its Roche lobe, a phase of mass transfer occurs, rejuvenating the secondary that eventually becomes the most massive star in the system. The primary evolves into a helium star that leaves a neutron star (or a black hole) after a supernova explosion. When less than half of the total mass of the system is lost, the system (now containing an OB star and a compact companion) remains bound (Blaauw 1961) and gets a kick velocity. The binary system then leaves the location where it was born. As soon as the secondary OB star evolves into a supergiant, accretion of its dense stellar wind onto the compact star will power a strong X-ray source. Thus, according to this model of massive binary evolution, all HMXBs should be runaways.

1.1. Blaauw's Scenario

Zwicky (1957) and Blaauw (1961) were the first to realize that a supernova explosion of one of the binary components can result in a high space velocity of the companion star. This mechanism for the formation of OB-runaway stars is presently known as Blaauw's scenario. In the original scenario a phase of mass transfer was not included, so that the massive binary had a higher probability to disrupt after the supernova explosion of the primary. But the modern version of this scenario predicts that the resulting runaway star has a high probability to remain bound to its compact companion (van den Heuvel 1993; Hills 1983). Although many people expect that a significant fraction of OB runaways have a compact companion, searches for compact stars around OB runaways have up to now not been successful (e.g., Philp et al. 1996).

As a result of a phase of mass transfer in a compact binary, one might expect that the abundances of nuclear processed elements are enhanced in the atmosphere of an OB-runaway star. Also, the angular momentum associated with the accreted material will increase the rotation rate of the OB runaway. For a small sample of bright OB runaways, Blaauw (1993) shows that these stars indeed exhibit a tendency toward higher helium abundance and higher projected rotational velocity. He further argues that some OB runaways seem to be younger than the age of the OB association they originate from (these stars are called blue stragglers). These observations provide, however, only circumstantial evidence for the supernova origin of (some) OB runaways.

1.2. Cluster Ejection Mechanism

An alternative explanation for the existence of OB-runaway stars is the cluster ejection model (Poveda, Ruiz, & Allen 1967): dynamical interaction in a compact cluster of stars results in the ejection of one or more of the members. From their extensive radial velocity survey of bright OB-runaway stars, Gies & Bolton (1986) concluded that the cluster ejection model should be favored. Their conclusion is mainly based on arguments refuting a supernova origin for the observed OB-runaway stars. Apart from the lack of observational evidence for the presence of compact companions around OB runaways, the existence of two runaway double-lined spectroscopic binaries cannot be explained with the supernova model.

¹ European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching, Germany.

² European Southern Observatory, Casilla 19001, Santiago 19, Chile.

³ Astronomical Institute "Anton Pannekoek," University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, Netherlands.

TABLE 1
KINEMATIC PROPERTIES OF HD 77581

	$\mu_{\alpha} \cos \delta$ (" yr ⁻¹)	μ_{δ} (" yr ⁻¹)	v_r (km s ⁻¹)	v_* (km s ⁻¹)
a) Turon et al. 1992	+0.004(3)	+0.009(3)	-1	89(40)
b) Sahu 1992	-0.0014(19)	+0.0107(22)	-4 ^a	98(26)

NOTE.—Proper motion, radial velocity, and space velocity of HD 77581 at the distance (1820 pc) of Vel OB1; for the space velocity, the average radial velocity of the OB stars in Vel OB1 (26.7 km s⁻¹; Humphreys 1978) is taken into account.

^a Radial velocity taken from Gies & Bolton 1986.

Also, the kinematical age of OB-runaway stars (i.e., the time needed to reach their present position with respect to the “parent” OB association) is often close to the age of the OB association itself, which is in support of the cluster ejection model. On the basis of the observed radial velocities they argued that the populations of HMXBs and OB runaways appear to be different. Van Oijen (1989), however, found strong indications that HMXBs are high-velocity objects.

In this Letter, we will provide compelling observational evidence that at least one HMXB is an OB-runaway system that obtained its high space velocity through the supernova explosion of the compact star’s progenitor.

2. WIND BOW SHOCK AROUND VELA X-1

Kinematic studies of OB stars are hampered by the large distances at which these stars are usually found, making it very difficult to measure proper motions accurately (although this situation will be significantly improved after release of the *HIPPARCOS* data). But it turns out that many OB stars with high space velocity create an unmistakable sign in the surrounding space. When an OB star moves supersonically through the interstellar medium (ISM), the interaction of its stellar wind with the ISM gives rise to a bow shock. Van Buren & McCray (1988) examined the *Infrared Astronomical Satellite* (*IRAS*) all-sky survey at the location of several OB-runaway stars and found extended arclike structures associated with many of them. The infrared emission results from interstellar dust that is swept up by the bow shock and heated by the radiation field of the OB star. In a subsequent study (Van Buren, Noriega-Crespo, & Dgani 1995), wind bow shocks were detected around one-third of a sample of 188 candidate OB-runaway stars. Thus, the detection of a wind bow shock can be considered as an observational confirmation of the runaway status of an OB star.

To this aim, we searched for the presence of a wind bow shock around the HMXB Vela X-1. This is a good candidate for an OB-runaway system. Its runaway nature was proposed in a recent paper by Van Rensbergen et al. (1996), based on the observed annual proper motion of HD 77581 listed in the *HIPPARCOS* Input Catalogue (Turon et al. 1992; see Table 1). The direction of the annual proper motion suggests that HD 77581 might originate from the OB association Vel OB1, which is located at a distance of 1820 pc (Humphreys 1978) from the Sun. At that distance, HD 77581 (Vela X-1) would have a space velocity of about 90 km s⁻¹ and would have left Vel OB1 2 ± 1 Myr ago. They further showed that this kinematical age is consistent with the predicted time of supernova explosion of Vela X-1’s progenitor.

Figure 1 (Plate L7) shows a narrowband H α image of a $10' \times 10'$ field centered on Vela X-1, obtained with the 1.54 m

FIG. 2.—Flux-calibrated long-slit spectrum of the bow shock, obtained with the Boller & Chivens spectrograph mounted on the 1.52 m ESO telescope. The spectrum was collapsed in the spatial direction along two regions in the slit (length 4:2) which are shown in the subpanel. The spectral resolution is 5.5 Å (FWHM). Region A corresponds to the secondary filament, which appears to produce the maximum H α intensity. Region B is the extension of the wind bow shock to the west. The forbidden [O III] lines are indicative of shock excitation.

Danish telescope at the European Southern Observatory (ESO) on 1996 February 14. The bright star HD 77581 ($V = 6.9$) has been removed (including its reflections) from the H α frame (exposure time 20 minutes) by subtracting a properly scaled *R*-band image of the same field. The uncalibrated H α image clearly reveals the presence of a bow shock, the apex being at $0'.9 \pm 0'.1$ to the north of HD 77581. At the distance of HD 77581 (see below), this corresponds to a projected distance of 0.48 ± 0.05 pc. With the help of the flux-calibrated long-slit spectrum (see Fig. 2) and knowledge of the position and dimensions of the slit, we estimate (within a factor of 2) the image peak H α intensity to be $\sim 10^{-15}$ ergs s⁻¹ cm⁻² arcsec⁻². The western (right) part of the bow shock consists of two separate filaments. Obviously, one filament is the continuation of the bow shock to the west. Filamentary structure has been observed in other wind bow shocks as well and might be related to their instability (Dgani, Van Buren, & Noriega-Crespo 1996). The *IRAS* Small Scale Structure Catalog (*IRAS* Small Scale Structure Catalog 1988) lists an extended infrared source at the position of Vela X-1. The extended source can be identified as an H II region with a Strömgren radius of $\sim 0'.2$ (~ 7 pc). We produced a high-resolution *IRAS* map (not shown here) of this area on the sky and found that inside the H II region the bow shock is well resolved at infrared wavelengths too, most pronounced in the 60 μ m wavelength band.

3. BOW SHOCK STRUCTURE

The structure of a wind bow shock is determined by the balance between the ram pressures of the wind and the

ambient medium (Baranov, Krasnobaev, & Kulikovskii 1971). One can show that

$$4\pi\rho_a v_*^2 R_0^2 = \dot{M}_w v_w.$$

Using the observed value for the distance R_0 between the OB star and the stagnation point (apex), the mass-loss rate, $\dot{M}_w = 10^{-6} M_\odot \text{ yr}^{-1}$, and terminal velocity, $v_w = 1105 \text{ km s}^{-1}$, of the stellar wind of HD 77581 (Kaper, Hammerschlag-Hensberge, & Van Loon 1993), and 90 km s^{-1} for the space velocity v_* of the system, we derive for the density of the ambient medium $\rho_a \approx 2.6 \times 10^{-24} \text{ g cm}^{-3}$ (number density $\sim 1 \text{ cm}^{-3}$).

A rough estimate of the local ISM density can be obtained from the size of the H II region and also from its observed infrared flux. Following Osterbrock (1989) and assuming that the H II region is a pure hydrogen nebula, optically thick in the Lyman series, a Strömgren radius of 7 pc and a blackbody input spectrum ($T_{\text{eff}} = 20,000 \text{ K}$) results in a neutral hydrogen number density n_H of 6 cm^{-3} . The total IR flux produced by dust inside the H II region is a measure for the number of Ly α photons produced by recombinations. The flux observed by *IRAS* (integrated over the H II region and corrected for the *IRAS* bandwidths) is $3 \times 10^{-8} \text{ ergs cm}^{-2} \text{ s}^{-1}$. A blackbody fit to the flux values shows that the dust temperature is 78 K and that the total dust emission is $5 \times 10^{-8} \text{ ergs cm}^{-2} \text{ s}^{-1}$, about 1.7 times the flux detected by *IRAS*. The energy per second per unit volume produced by recombination to the ground level is $3.3 \times 10^{-24} \times n_H^2 \text{ ergs cm}^{-3} \text{ s}^{-1}$. Comparison with the measured IR flux gives $n_H = 16 \text{ cm}^{-2}$. If the emission from the bow shock is excluded, this yields $n_H = 11 \text{ cm}^{-2}$. Given the uncertainties, the estimate of the local ISM density with help of the bow shock and stellar wind parameters is probably the most accurate.

We obtained a long-slit spectrum of the bow shock with the ESO 1.52 m telescope on the night of 1995 May 28 (Fig. 2). We show a flux-calibrated spectrum for two regions covered by the slit; the orientation and position of the slit are indicated in the subpanel. The spectra include the H α and H β lines and forbidden emission lines like [O III] $\lambda\lambda 4959, 5007$ and [N II] $\lambda 6583$. Region A corresponds to the secondary filament, which appears to produce the maximum H α intensity. The H α /H β ratio in region A is 6.5; if this large ratio is due to extinction, $E(B - V)$ would be equal to 0.72, which is consistent with the reddening of HD 77581 (Sadakane et al. 1985). In region B, which covers a part of the wind bow shock, the Balmer lines are much weaker with respect to the forbidden [O III] lines ([O III] $\lambda 5007$ /H $\beta \geq 4.1$) than in region A (≈ 1.5). Compared to other usual shock-excited emission lines, the [O III] lines originate close to the shock front where the ionization conditions depend critically on shock conditions (Dopita 1977; Hollis et al. 1992). Therefore, the spectra are in accordance with both regions being part of the wind bow shock. The filament covered by region A might be experiencing a somewhat weaker shock, yielding a much larger emission measure in H α .

4. DISCUSSION AND CONCLUSIONS

The present positions of HD 77581 and the OB association Vel OB1 are shown in Figure 3. The separation between the two is about 7° . The direction of the proper motion of HD 77581, (a) as listed in the HIC and (b) as measured by Sahu (1992), is indicated. The dotted line represents the proposed path of HD 77581 based on the symmetry axis of the bow

FIG. 3.—Position of HD 77581 (Vela X-1) on the sky with respect to the OB association Vel OB1. Right ascension and declination are given in degrees; north is up, and east is to the left. The direction of proper motion, as listed in the HIC (a) and as obtained by Sahu 1992 (b), is shown. The proposed path of the system, based on the symmetry axis of the bow shock, is given by a dotted line. The numbers along this line indicate the position where the system would have been (1, 2, and 3 Myr ago, respectively), given a velocity of 90 km s^{-1} .

shock, which is in a direction in between the proper-motion vectors **a** and **b**. The bow shock reflects the relative motion between the star and the surrounding medium, while the proper motion measures the star's motion with respect to other stars. Since the gas has a 10 km s^{-1} dispersion, 5° – 10° deviations can be expected, which would favor **b**. If the proposed path is correct, the origin of the binary system would be on the outskirts of Vel OB1. The usually adopted distance toward Vela X-1 of $1.9 \pm 0.1 \text{ kpc}$ (Sadakane et al. 1985) is in accordance with the stellar parameters of HD 77581. The accretion-induced X-ray luminosity of Vela X-1 (Kaper et al. 1993) also agrees very well with the distance toward Vel OB1. Thus, our observations confirm the proposed runaway status of HD 77581 and support its suggested origin in Vel OB1.

In conclusion, the discovered wind bow shock around Vela X-1 provides compelling observational evidence for a runaway HMXB. The direction of motion of the system is derived from the symmetry of the bow shock, and it is very likely that the OB association Vel OB1 is the origin of the system. From the observed proper motion one can compute that the system left the outskirts of Vel OB1 about 2.5 Myr ago, which corresponds to the expected time of the supernova explosion of Vela X-1's progenitor. Therefore, our observations support Blaauw's scenario for the formation of the runaway HMXB HD 77581 (Vela X-1). It remains to be shown whether all HMXBs are runaway systems. We predict that the answer to this question is positive and expect that in at least a third of the cases a wind bow shock is associated with them.

This paper is based on observations obtained at the European Southern Observatory in La Silla, Chile. We thank J. Bergeron, L. Lucy, and J. van Paradijs for carefully reading

the manuscript. G. Van de Steene produced the blackbody fit to the observed *IRAS* fluxes. The referee is acknowledged for

his/her helpful remarks and a critical reading of the manuscript.

REFERENCES

- Baranov, V. B., Krasnobaev, K. V., & Kulikovskii, A. G. 1971, *Soviet Phys. Dokl.*, 15, 791
- Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265
- Blaauw, A. 1993, in *ASP Conf. Ser. 35, Massive Stars: Their Lives in the Interstellar Medium*, ed. J. P. Cassinelli & E. B. Churchward (San Francisco: ASP) 207
- Dgani, R., Van Buren, D., & Noriega-Crespo, A. 1996, *ApJ*, 461, 372
- Dopita, M. 1977, *ApJS*, 33, 437
- Gies, D. R., & Bolton, C. T. 1986, *ApJS*, 61, 419
- Hills, J. G. 1983, *ApJ*, 267, 322
- Hollis, J. M., Oliverson, R. J., Wagner, R. M., & Feibelman, W. A. 1992, *ApJ*, 93, 217
- Humphreys, R. M. 1978, *ApJS*, 38, 309
- IRAS* Small Scale Structure Catalog. 1988, prepared by G. Helou & D. Walker (Washington: GPO), Vol. 7
- Jones, C., & Liller, W. 1973, *ApJ*, 184, 121
- Kaper, L., Hammerschlag-Hensberge, G., & van Loon, J. Th. 1993, *A&A*, 279, 485
- McClintock, J. E., et al. 1976, *ApJ*, 206, 99
- Nagase, F. 1989, *PASJ*, 41, 1
- Osterbrock, D. E. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Mill Valley: Univ. Science)
- Philp, C. J., Evans, C. R., Leonard, P. J. T., & Frail, D. A. 1996, *AJ*, 111, 1220
- Poveda, A., Ruiz, J., & Allen, C. 1967, *Bol. Obs. Tonantzintla Tacubaya*, 4, 860
- Sadakane, K., et al. 1985, *ApJ*, 288, 284
- Sahu, M. S. 1992, Ph.D. thesis, Univ. Groningen
- Turon, C., et al. 1992, *The HIPPARCOS Input Catalogue* (ESA SP-1136; Noordwijk: ESA), Vol. 3
- Van Buren, D., & McCray, R. 1988, *ApJ*, 329, 93
- Van Buren, D., Noriega-Crespo, A., & Dgani, R. 1995, *AJ*, 110, 2914
- van den Heuvel, E. P. J. 1993, in *Saas-Fée Advanced Course on Interacting Binaries*, ed. S. N. Shore, (Berlin: Springer), 263
- van Kerkwijk, M. H., et al. 1995a, *A&A*, 303, 483
- van Kerkwijk, M. H., van Paradijs, J., & Zuiderwijk, E. J. 1995b, *A&A*, 303, 497
- Van Oijen, J. G. J. 1989, *A&A*, 217, 115
- van Paradijs, J., et al. 1977, *A&A*, 30, 195
- Van Rensbergen, W., Vanbeveren, D., & De Loore, C. 1996, *A&A*, 305, 825
- Vidal, N. V., Wickramasinghe, D. T., & Peterson, B. A. 1973, *ApJ*, 265, 1036
- Watson, M. G., & Griffiths, R. E. 1977, *MNRAS*, 178, 513
- Zwicky, F. 1957, *Morphological Astronomy* (Berlin: Springer), 258

FIG. 1.— R -band corrected $H\alpha$ image of the field around Vela X-1. North is up, and east is to the left. The images were obtained with the DFOSC instrument and a CCD detector in the Cassegrain focus of the 1.54 m Danish telescope at the European Southern Observatory. A wind bow shock is detected about $0^{\circ}.9$ north of the (saturated) 6th magnitude star HD 77581. The western (right) arm of the bow shock is split into two filaments, a feature more often encountered in wind bow shocks.

KAPER et al. (see 475, L38)