



UvA-DARE (Digital Academic Repository)

Antimicrobial drug resistance at the human-animal interface in Vietnam

Nguyen, V.T.

[Link to publication](#)

Citation for published version (APA):

Nguyen, V. T. (2017). Antimicrobial drug resistance at the human-animal interface in Vietnam

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

CHAPTER 3
**NON-TYPHOIDAL *SALMONELLA* COLONIZATION IN CHICKENS AND
HUMANS IN THE MEKONG DELTA OF VIETNAM**

Chapter 3: Non-typhoidal *Salmonella* colonization in chickens and humans in the Mekong delta of Vietnam

N. V. Trung^{1,2,3}, J. J. Carrique-Mas^{3,4}, N. H. Nghia³, L. T. P. Tu³, H. H. Mai⁵, H. T. Tuyen³, J. Campbell^{3,4}, N. T. Nhung³, H. N. Nhung³, P. V. Minh³, T. T. B. Chieu³, T. Q. Hieu⁵, N. T. N. Mai⁶, S. Baker^{3,4}, J. A. Wagenaar^{7,8}, N. T. Hoa^{3,4} and C. Schultsz^{1,2,3}

¹ Department of Medical Microbiology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

² Department of Global Health-Amsterdam Institute for Global Health and Development, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

³ Oxford University Clinical Research Unit, Centre for Tropical Medicine, Ho Chi Minh City, Vietnam

⁴ Centre for Tropical Medicine, Nuffield Department of Medicine, University of Oxford, Oxford, UK

⁵ Sub-Department of Animal Health, My Tho, Tien Giang, Vietnam

⁶ Preventive Medicine Center, Tien Giang, Vietnam

⁷ Department of Infectious Diseases and Immunology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands

⁸ Central Veterinary Institute of Wageningen UR, Lelystad, The Netherlands

Abstract

Salmonellosis is a public health concern both in the developed and developing countries. Although the majority of human non-typhoidal *Salmonella enterica* (NTS) cases are the result of foodborne infections or person-to-person transmission, NTS infections may also be acquired by environmental and occupational exposure to animals. While a considerable number of studies have investigated the presence of NTS in farm animals and meat/carcasses, very few studies have investigated the risk of NTS colonization in humans as a result of direct animal exposure. We investigated asymptomatic NTS colonization in 204 backyard chicken farms, 204 farmers and 306 matched individuals not exposed to chicken farming, in southern Vietnam. Pooled chicken faeces, collected using boot- or handheld-swabs on backyard chicken farms, and rectal swabs from human participants were tested. NTS colonization prevalence was 45.6%, 4.4% and 2.6% for chicken farms, farmers and unexposed individuals, respectively. Our study observed a higher prevalence of NTS colonization among chicken farmers (4.4%) compared with age-, sex- and location- matched rural and urban individuals not exposed to chickens (2.9% and 2.0%). A total of 164 chicken NTS strains and 17 human NTS strains were isolated and 28 serovars were identified. *Salmonella* Weltevreden was the predominant serovar in both chickens and humans. NTS isolates showed resistance (20-40%) against tetracycline, chloramphenicol, sulfamethoxazole-trimethoprim and ampicillin. Our study reflects the epidemiology of NTS colonization in chickens and humans in the Mekong delta of Vietnam and emphasizes the need of larger, preferably longitudinal studies to study the transmission dynamics of NTS between and within animal and human host populations.

Keywords: Non-typhoidal *Salmonella*, colonization, antimicrobial resistance, chickens, humans, Vietnam

Introduction

Salmonellosis caused by non-typhoidal *Salmonella enterica* (NTS) is a potentially zoonotic infection commonly associated with gastroenteritis and represents a significant public health problem both in the developing and developed countries [1].

Although the majority of human NTS cases is the result of foodborne infection [2] or person-to-person transmission [3], humans can also become infected with NTS as a result of environmental and occupational exposure to animals, including farm animals [4-6]. This type of exposure is particular prevalent in developing countries, where a large fraction of the population is involved in raising livestock and/or poultry. Farms in these countries are typically backyard or small-scale and farming procedures mostly involve low levels of biosecurity and personal protection. This results in very close contact between animals and humans. To date, very few studies have investigated the role of exposure to farm animals in asymptomatic NTS infection.

We hypothesized that exposure to chickens through farming results in increased risk of asymptomatic colonisation with NTS in Vietnam, a country with a majority of rural population. In this country, small and backyard chicken farming is very common, with most farms typically having less than 50 chickens [7]. To investigate this hypothesis, we investigated chicken flocks and farmers to study the prevalence and serovar distribution of NTS organisms. The prevalence in farmers was compared with that among age- and gender-matched individuals living in the same areas but not engaged in poultry farming. In addition, given the extensive antimicrobial drug usage in these chicken farms [8], we analysed and compared the frequencies of resistance against key antimicrobial drugs of the most relevant classes, used both in veterinary and human medicine.

CHAPTER 3

Materials and Methods

Study population

The target population consisted of chicken farms, including chicken flocks and their farmers, as well as individuals not involved in poultry farming from 2 rural districts and the capital of the Mekong Delta province of Tien Giang (Vietnam).

Sampling of farms was conducted at random within two size strata: 10–200 chickens ('household farms', N=102), and 201–2,000 chickens ('small farms', N=102). In selected farms, both flocks and the person responsible for raising the chickens ('farmer', N=204) were recruited. In addition to the 204 farmers, 306 participants not involved in chicken or livestock farming were randomly selected ('unexposed individuals'). A subgroup of unexposed individuals ('rural subjects') were selected from the same commune as the farmer, and were matched to the farmer by age and gender (N=204, one per farmer). Another subgroup of unexposed individuals ('urban subjects') were selected from the provincial capital, and were matched to the farmers by age and gender (N=102, one every two farmers).

Farm selection and chicken flock sampling was carried out by staff at the Sub-Department of Animal Health (SDAH) in Tien Giang. Selection and sampling of human subjects was performed from the population census provided by the Preventive Medicine Centre (PMC) of Tien Giang [9].

Written informed consents were obtained from all participants prior to participation in the study. Participants that refused to participate were replaced by the next available eligible participant. The study was approved by the Peoples' Committee of Tien Giang Province, the Department of Health in Tien Giang and the Oxford University Tropical Research Ethics Committee (OxTREC, No. 48/11).

Sample collection

Farm and human household visits were evenly distributed over a year period from March 2012 to April 2013 in order to avoid seasonal effects. Pooled chicken faeces samples were collected from chicken houses using boot swabs (flocks reared on barn systems) or handheld gauze swabs

CHAPTER 3

(caged layer flocks and flocks on stilted mesh houses) as described previously [10]. The sample collection was conducted by a trained sampling team from Tien Giang SDAH.

Rectal swab samples were obtained from all human participants by trained staff from Tien Giang PMC, using Fecalswab (Copan, Italy).

All samples were stored at 4°C, transported to the laboratory at the Oxford University Clinical Research Unit in Ho Chi Minh City and cultured within 24 hours after sample collection.

Laboratory methods

Salmonella was isolated using the modification of the ISO 6579:2002 (Annex D) method for chicken faecal samples, involving: (a) pre-enrichment in 225 ml of buffered peptone water (BPW) (37°C, 18 hours); (b) plating 100 µl of the culture onto modified semi-solid Rappaport–Vassiliadis medium (Oxoid; UK) (41°C, 24 hours) and (c) plating onto Rambach agar (37°C, 24 hours) [11]. For human samples, rectal swabs were cultured on MacConkey agar, xylose-lysinedeoxycholate agar and selenite broth according to guidelines of World Health Organization [12].

Suspected NTS colonies were selected for each sample and confirmed by slide agglutination with relevant poly O antiserum [13]. All isolates confirmed as NTS were further tested for their antimicrobial susceptibility.

Antimicrobial susceptibility testing was performed by disc diffusion method and breakpoints were interpreted using the Clinical and Laboratory Standards Institute guidelines for Enterobacteriaceae [14]. Eleven antimicrobials were tested including chloramphenicol (30 mg), ceftazidime (30 mg), ceftriaxone (30 mg), amoxicillin/clavulanic acid (30 mg), meropenem (10 mg), ciprofloxacin (5 mg), tetracycline (30 mg), trimethoprim/sulfamethoxazole (1.25/23.75 mg), amikacin (30 mg), gentamicin (10 mg) and ampicillin (10 mg). Quality controls for susceptibility testing and identification were performed every week according to the CLSI guidelines. An MDR strain was defined as a strain resistant to at least three different antimicrobial classes. Chicken farms and farmers were the study unit of analysis. A chicken farm was defined as ‘positive’ for NTS if NTS was isolated from at least one of the boot- or handheld gauze swabs.

CHAPTER 3

Confirmed NTS isolates were genotyped using multilocus sequence typing (MLST) as described previously [15]. Briefly, pure colonies of overnight culture on nutrition agar were subjected to DNA extraction using Wizard Genomic DNA extraction kit (Promega, USA). Seven MLST loci (*aroC*, *dnaN*, *hemD*, *hisD*, *purE*, *sucA* and *thrA*) were amplified and sequenced in forward and reverse directions using the Big Dye Cycle Sequencing kit (Applied Biosystems, USA) on an ABI 3770 automatic sequencer according to the manufacturer's instructions. Sequence data of seven loci was trimmed and blasted to determine sequence type as well as serotype based on data available on the MLST database (<http://mlst.warwick.ac.uk/mlst/dbs/Senterica/>)

Differences in proportions were compared using the Chi-square test and the Fisher's Exact Test, when the Chi-square test was not relevant. A p-value <0.05 was considered statistically significant.

Results


Of 204 chicken farms, 93 (45.6%, 95% CI= 38.8–52.4%) tested positive for NTS. There was no statistically significant difference in the NTS farm-level prevalence between household farms (46/102, 45.1%) and small farms (47/102, 46.1%). NTS was recovered from 9 chicken farmers (4.4%, 95% CI= 1.6–7.2%), 6 rural subjects (2.9%, 95% CI= 0.6–5.3%) and 2 urban subjects (2.0%, 95% CI= 0.0–4.7%). The prevalence of NTS did not statistically differ between chicken farmers and unexposed individuals. The prevalence of NTS among farmers of NTS-positive chicken flocks (5/93, 5.4%; 95% CI= 0.8–10.0%) was similar to the prevalence of NTS among farmers of NTS-negative chicken flocks (4/111, 3.6%; 95% CI= 0.1–7.1%).

Among 164 chicken NTS isolates, the highest observed levels of resistance were against tetracycline (39.6%), chloramphenicol (28.0%), sulfamethoxazole-trimethoprim (26.8%), ampicillin (26.2%) and amoxicillin plus clavulanic acid (12.8%). The proportion of strains resistant against ciprofloxacin and gentamicin was 1.8%. No resistance against meropenem, ceftazidime and ceftriaxone was observed. A total of 9 NTS isolates from farmers and 8 NTS isolates from individuals unexposed to chicken farming were tested for antimicrobial resistance, which indicated similar levels of resistance against all tested antimicrobials. 27.4% (45/164), 22.2% (2/9) and 12.5% (1/8) of NTS isolates from chickens, farmers and unexposed individuals were multidrug resistant, respectively (Figure 1). Interestingly, one of the two multidrug resistant

CHAPTER 3

isolates from farmers was identical to the most common resistance pattern found in chicken flocks (chloramphenicol – ampicillin – tetracycline – trimethoprim/sulfamethoxazole) (Data not shown).

Figure 1: Percentage of NTS isolates resistant to a panel of 11 antimicrobials



C: chloramphenicol (30µg), CAZ: ceftazidime (30µg), CRO: ceftriaxone (30µg), AMC: amoxicillin/clavulanic acid (30 µg), MEM: meropenem (10µg), CIP: ciprofloxacin (5µg), TE: tetracycline (30µg), SXT: trimethoprim-sulphamethoxazole (10µg), AK: amikacin (30µg), CN: gentamicin (10µg), AMP: ampicillin (10µg), MDR: Multidrug resistance (resistant against at least three classes of antimicrobial).

MLST was performed on 163/164 chicken isolates from 93 chicken farms (one isolate could not be recovered after storage) and on all 17 isolates from humans. *Salmonella* Weltevreden was the most common serovar detected in chicken farms (10.3% of farms), farmers (2.0%), rural subjects (1.0%) and urban subjects (2.0%). Besides *Salmonella* Weltevreden, the predominant serovars in chickens were *Salmonella* Enteritidis, *Salmonella* Paratyphi B var Java monophasic, *Salmonella* Albany, *Salmonella* Derby, *Salmonella* Give, *Salmonella* Newport and *Salmonella* Typhimurium. *Salmonella* Enteritidis and *Salmonella* Typhimurium were only found in chickens and were detected in 4.4% and 1.5% of the chicken farms, respectively (Table 1).

NTS were detected both in chickens and farmers on 5/204 farms (2.5%, 95% CI= 0.3-4.6%). MLST revealed that the serovar of NTS isolates obtained from the farmer and their chicken on the same farm were identical in one farm (*Salmonella* Weltevreden), but differed between farmer and their chickens for the other 4 farms (Table 2).

Table 1. Distribution of different serovars of NTS isolated from chickens and humans in southern Vietnam.

NTS serovar	No. culture positive (%)			
	Chicken farms ^a (n=204)	Chicken farmers (n=204)	Rural subjects (n=204)	Urban subjects (n=102)
Weltevreden	21 (10.3)	4 (2.0)	2 (1.0)	2 (2.0)
Enteritidis	9 (4.4)	0	0	0
Paratyphi B var Java monophasic	9 (4.4)	0	0	0
Albany	7 (3.4)	0	0	0
Derby	6 (2.9)	0	1 (0.5)	0
Give	6 (2.9)	2 (1.0)	0	0
Newport	4 (2.0)	0	1 (0.5)	0
Typhimurium	3 (1.5)	0	0	0
Braenderup	1 (0.5)	1 (0.5)	0	0
Orientalis	0	1 (0.5)	0	0
Rubislaw	0	1 (0.5)	0	0
Ohio	0	0	1 (0.5)	0
Other serovars ^b	28 (13.7)	0	0	0
Untypeable	19 (9.3)	0	0	0
Any serovar	93 (45.6)	9 (4.4)	6 (2.9)	2 (2.0)

^aOn 19 farms, multiple serovars were present.

^bOther serovars: Anatum, Senftenberg, Stanley, Virchow (each serovar was present on 3 farms); Kentucky, London, Montevideo, Typhimurium monophasic (each serovar was present on 2 farms); Cerro, Indian, Litchfield, Mbandaka, Meleagridis, Oslo, Poona, Tennessee (each serovar was present on one farm).

Table 2. Serovar and antimicrobial resistance pattern of NTS isolated from chicken flocks and farmers from the same farm.

Farm ID	Source	Isolate number	<i>Salmonella</i> serovar	Antimicrobial resistance pattern ^a
CG 37	Farmer	1	Weltevreden	Fully susceptible
	Chicken	1	Untypeable	Fully susceptible
CT 67	Farmer	1	Rubislaw	C-AMC-TE-AMP
	Chicken	1	Albany	Fully susceptible
MT 26	Farmer	1	Weltevreden	Fully susceptible
	Chicken	1	Weltevreden	Fully susceptible
	Chicken	2	Weltevreden	Fully susceptible
MT 28	Farmer	1	Give	C-TE-SXT-AMP
	Chicken	2	Enteritidis	TE
MT 53	Farmer	1	Weltevreden	Fully susceptible
	Chicken	1	Senftenberg	CIP-TE-SXT
	Chicken	2	Senftenberg	Fully susceptible
	Chicken	3	Cerro	Fully susceptible

^a Isolates were tested for susceptibility to 11 antimicrobials using disk diffusion method and interpreted according to breakpoints as defined by Clinical and Laboratory Standard Institute (11). C: chloramphenicol (30µg), AMC: amoxicillin/clavulanic acid (30 µg), CIP: ciprofloxacin (5µg), TE: tetracycline (30µg), SXT: trimethoprim-sulphamethoxazole (10µg), AMP: ampicillin (10µg).

Discussion

To our knowledge, this is the first field survey reporting on prevalence of asymptomatic NTS colonization in humans occupationally exposed and unexposed to chickens in Vietnam. The observed prevalence of asymptomatic NTS colonization in humans was 3.3%, a figure considerably higher than the reported prevalence of asymptomatic NTS in developed countries (0.3% – 0.4%) [16-18]. However, our results were similar to the results from a study performed in 2004 in Hanoi, Vietnam (3.1%) [19] and from Thailand in 2003 (4.7%) [20].

We found a higher prevalence of NTS colonization among chicken farmers (4.4%) compared with unexposed individuals (2.6%). The prevalence of NTS colonization in the rural subjects was also higher (2.9%) than in the urban subjects (2.0%). Similarly, farmers of NTS-positive chicken flocks had a higher prevalence of infection compared with farmers of NTS-negative chicken flocks (5.4% versus 3.6%). However, none of these differences were statistically significant.

Our study demonstrated a high prevalence (45.6%) and high diversity of NTS serovars in both household-size and small-size chicken farms in the Mekong Delta of Vietnam, similar to the results from a recent survey (64.7%) carried out in Dong Thap, another Mekong Delta province [21]. In unconfined flocks, swabs may potentially gathered faecal material from other animals in the farm. However, in those cases, sampling was carried out near the perching and eating areas where chicken droppings were visible, so it is expected that the overwhelming majority of faecal material and *Salmonella* strains were of chicken origin. In terms of antimicrobial sensitivity, multidrug resistance was commonly observed in the chicken NTS isolates. Of antimicrobials of critical importance, the prevalence of resistance against ampicillin was particularly high among tested isolates (26.2%). However, levels of resistance against aminoglycosides (gentamicin) and fluoroquinolones (ciprofloxacin) were low (<2%).

We acknowledge several limitations in our study. Firstly, the number of NTS isolates from humans was small, limiting the power to demonstrate any statistical difference between study cohorts. Secondly, the cross-sectional study design may preclude the demonstration of transmission of any particular serovar, which is highly depending on the dynamics of NTS transmission between chickens and humans. It is therefore possible that the presence of NTS in the farmer results from earlier infection and further clearance in the current chicken flock, or

CHAPTER 3

from transmission from a previous flock. In addition, differences in NTS isolation methods for chicken and human samples may have had an impact on the sensitivity of detection. A higher sensitivity of detection of the ISO 6572: 2002 (Annex D) method compared with the WHO method for human samples is to be expected, since the former include a pre-enrichment, selective enrichment phase, which allows the detection of low numbers of *Salmonella* such as those likely to be found in asymptomatic chicken samples [22].

In spite of these limitations, we found that 1/21 (4.7%) of the farmers with *Salmonella* Weltevreden infected chicken flocks was also *Salmonella* Weltevreden positive compared with 3/183 (1.6%) of the farmers without *Salmonella* Weltevreden infected chicken flocks. Interestingly, the *Salmonella* Weltevreden isolated from the farmer had an identical antimicrobial resistance pattern to the isolate from his/her chickens (fully susceptible). It has been suggested that *Salmonella* Weltevreden may have acquired properties which facilitate adaptation to a broader range of hosts [23], and *Salmonella* Weltevreden has been shown to be able to persist in manure and soil for prolonged periods of time [24].

We believe that our study reflects the epidemiological situation of NTS in the Mekong delta of Vietnam, characterized by a high prevalence of infection in chicken flocks and a relatively high prevalence of colonization of human adults. Our study also underscores the need for additional larger and preferably longitudinal studies to investigate transmission dynamics of NTS between and within animal and human host populations.

References

1. Majowicz, S.E., et al., *The global burden of nontyphoidal Salmonella gastroenteritis*. Clin Infect Dis, 2010. **50**(6): p. 882-9.
2. Cao, V., et al., *Antimicrobial susceptibility of Neisseria gonorrhoeae strains isolated in 2004-2006 in Bangui, Central African Republic; Yaounde, Cameroon; Antananarivo, Madagascar; and Ho Chi Minh Ville and Nha Trang, Vietnam*. Sex Transm Dis, 2008. **35**(11): p. 941-5.
3. Thompson, C.N., et al., *Epidemiological features and risk factors of Salmonella gastroenteritis in children resident in Ho Chi Minh City, Vietnam*. Epidemiol Infect, 2013. **141**(8): p. 1604-13.
4. Baker, M.G., et al., *A recurring salmonellosis epidemic in New Zealand linked to contact with sheep*. Epidemiol Infect, 2007. **135**(1): p. 76-83.
5. Fone, D.L. and R.M. Barker, *Associations between human and farm animal infections with Salmonella typhimurium DT104 in Herefordshire*. Commun Dis Rep CDR Rev, 1994. **4**(11): p. R136-40.
6. Hoelzer, K., A.I. Moreno Switt, and M. Wiedmann, *Animal contact as a source of human non-typhoidal salmonellosis*. Vet Res, 2011. **42**: p. 34.
7. Burgos, S., et al., *Characterization of poultry production systems in Vietnam*. Int J of Poult Sci, 2007. **6**(10): p. 709-712.

CHAPTER 3

8. Carrique-Mas, J.J., et al., *Antimicrobial usage in chicken production in the Mekong Delta of Vietnam*. Zoonoses Public Health, 2015. **62 Suppl 1**: p. 70-8.
9. Anon., *General Statistics Office of Vietnam. Results of the 2006 Rural, Agricultural and Fishery Census*. Statistical Publishing House, Hanoi, Vietnam. 2007.
10. Nguyen, V.T., et al., *Prevalence and risk factors for carriage of antimicrobial-resistant Escherichia coli on household and small-scale chicken farms in the Mekong Delta of Vietnam*. J Antimicrob Chemother, 2015. **70(7)**: p. 2144-52.
11. Carrique-Mas, J.J., et al., *Comparison of three plating media for the isolation of Salmonella from poultry environmental samples in Great Britain using ISO 6579:2002 (Annex D)*. J Appl Microbiol, 2009. **107(6)**: p. 1976-83.
12. WHO, *World Health Organization Global Foodborne Infections Network. Isolation of Salmonella and Shigella from Faecal Specimens*. 2010: Atlanta, GA, USA.
13. Anon., *Antigenic Formulae of the Salmonella Serovars*. 9th edn. World Health Organization Collaborating Center for Reference and Research on Salmonella. 2007.
14. CLSI, *Performance standards for antimicrobial susceptibility testing; twenty-first international supplement*, in *CLSI document M100-S21*. 2011: Clinical and Laboratory Standards Institute.
15. Achtman, M., et al., *Multilocus sequence typing as a replacement for serotyping in Salmonella enterica*. PLoS Pathog, 2012. **8(6)**: p. e1002776.
16. de Wit, M.A., et al., *Sensor, a population-based cohort study on gastroenteritis in the Netherlands: incidence and etiology*. Am J Epidemiol, 2001. **154(7)**: p. 666-74.
17. Hellard, M.E., et al., *Prevalence of enteric pathogens among community based asymptomatic individuals*. J Gastroenterol Hepatol, 2000. **15(3)**: p. 290-3.
18. Nataro, J.P., et al., *Diarrheagenic Escherichia coli infection in Baltimore, Maryland, and New Haven, Connecticut*. Clin Infect Dis, 2006. **43(4)**: p. 402-7.
19. Do, T.T., et al., *Epidemiology and aetiology of diarrhoeal diseases in adults engaged in wastewater-fed agriculture and aquaculture in Hanoi, Vietnam*. Trop Med Int Health, 2007. **12 Suppl 2**: p. 23-33.
20. Sirinavin, S., L. Pokawattana, and A. Bangtrakulnondh, *Duration of nontyphoidal Salmonella carriage in asymptomatic adults*. Clin Infect Dis, 2004. **38(11)**: p. 1644-5.
21. Tu, L.T., et al., *High levels of contamination and antimicrobial-resistant non-typhoidal Salmonella serovars on pig and poultry farms in the Mekong Delta of Vietnam*. Epidemiol Infect, 2015: p. 1-13.
22. Carrique-Mas, J.J. and R.H. Davies, *Sampling and bacteriological detection of Salmonella in poultry and poultry premises: a review*. Rev Sci Tech, 2008. **27(3)**: p. 665-77.
23. Brankatschk, K., et al., *Comparative genomic analysis of Salmonella enterica subsp. enterica serovar Weltevreden foodborne strains with other serovars*. Int J Food Microbiol, 2012. **155(3)**: p. 247-56.
24. Arthurson, V., A. Sessitsch, and L. Jaderlund, *Persistence and spread of Salmonella enterica serovar Weltevreden in soil and on spinach plants*. FEMS Microbiol Lett, 2011. **314(1)**: p. 67-74.