



UvA-DARE (Digital Academic Repository)

Attrition analysed in five waves of a longitudinal yearly survey of smokers: Findings from the ITC Netherlands Survey

Zethof, D.; Nagelhout, G.E.; de Rooij, M.; Driezen, P.; Fong, G.T.; van den Putte, B.; Hummel, K.; de Vries, H.; Thompson, M.E.; Willemsen, M.C.

DOI

[10.1093/eurpub/ckw037](https://doi.org/10.1093/eurpub/ckw037)

Publication date

2016

Document Version

Final published version

Published in

European Journal of Public Health

License

Article 25fa Dutch Copyright Act (<https://www.openaccess.nl/en/in-the-netherlands/you-share-we-take-care>)

[Link to publication](#)

Citation for published version (APA):

Zethof, D., Nagelhout, G. E., de Rooij, M., Driezen, P., Fong, G. T., van den Putte, B., Hummel, K., de Vries, H., Thompson, M. E., & Willemsen, M. C. (2016). Attrition analysed in five waves of a longitudinal yearly survey of smokers: Findings from the ITC Netherlands Survey. *European Journal of Public Health*, 26(4), 693-699. <https://doi.org/10.1093/eurpub/ckw037>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Attrition analysed in five waves of a longitudinal yearly survey of smokers: findings from the ITC Netherlands survey

Dennis Zethof¹, Gera E. Nagelhout^{2,3}, Mark de Rooij⁴, Pete Driezen⁵, Geoffrey T. Fong^{5,6}, Bas van den Putte^{7,8}, Karin Hummel², Hein de Vries², Mary E. Thompson⁹, Marc C. Willemsen^{2,10}

1 Department of Psychology, Alumni, Leiden University, Leiden, The Netherlands

2 Department of Health Promotion, Maastricht University (CAPHRI), Maastricht, The Netherlands

3 Department of Family Medicine, Maastricht University (CAPHRI), Maastricht, The Netherlands

4 Department of Psychology, Leiden University, Leiden, The Netherlands

5 Department of Psychology and School of Public Health and Health Systems, University of Waterloo, Waterloo, Ontario, Canada

6 Ontario Institute for Cancer Research, Toronto, Ontario, Canada

7 Department of Communication, University of Amsterdam (ASCoR), Amsterdam, The Netherlands

8 Trimbos Institute, Netherlands Institute for Mental Health and Addiction, Utrecht, The Netherlands

9 Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, Ontario, Canada

10 Dutch Alliance for a Smokefree Society, The Hague, The Netherlands

Correspondence: Gera Nagelhout, Department of Health Promotion, Maastricht University (CAPHRI), Maastricht, The Netherlands, Tel: +316 244 170 80, Fax: +31 43 367 10 32, e-mail: gera.nagelhout@maastrichtuniversity.nl

Background: Attrition bias can affect the external validity of findings. This article analyses attrition bias and assesses the effectiveness of replenishment samples on demographic and smoking-related characteristics for the International Tobacco Control Netherlands Survey, a longitudinal survey among smokers. **Methods:** Attrition analyses were conducted for the first five survey waves (2008–12). We assessed, including and excluding replenishment samples, whether the demographic composition of the samples changed between the first and fifth waves. Replenishment samples were tailored to ensure the sample remained representative of the smoking population. We also constructed a multivariable survival model of attrition that included all five waves with replenishment samples. **Results:** Of the original 1820 respondents recruited in 2008, 46% participated again in 2012. Demographic differences between waves due to attrition were generally small and replenishment samples tended to minimize them further. The multivariable survival analysis revealed that only two of the 10 variables analysed were significant predictors of attrition: a weak effect for gender (men dropped out more often) and weak to moderate effects for age (respondents aged 15–24 years dropped out more than aged 25–39 years, who dropped out more than those aged 40+ years). **Conclusions:** Weak to moderate attrition effects were found for men and younger age groups. This information could be used to minimize respondent attrition. Our findings suggest that sampling weights and tailored replenishment samples can effectively compensate for attrition effects. This is already being done for the International Tobacco Control Netherlands Survey, including the categories that significantly predicted attrition in this study.

Introduction

Respondent attrition is one of the potential problems in longitudinal survey research. When attrition is random, the smaller sample size will reduce the precision of estimators. But when it is nonrandom, attrition can change the sample composition and result in biased estimators. This undermines the generalizability (i.e. external validity) of research findings.^{1–4}

The International Tobacco Control Policy Evaluation Project (ITC Project)⁵ was developed to meet the demands for a rigorous evaluation of national tobacco control policies. To date, 22 countries have participated in the ITC Project. This project employs a quasi-experimental pre–post design in which countries that have implemented a particular policy or policies can be compared to other countries that did not implement policies during the same time period. Such countries serve as control groups which allow the ITC Project to assess the psychosocial and behavioural effects of national level tobacco control policies⁵ in ways that increase internal validity relative to other evaluation designs (e.g. a single country pre–post design).⁶

The ITC Netherlands Survey uses replenishment samples to maintain a relatively constant sample size over time. These

replenishment samples address the problem of reduced precision due to attrition.⁷ In spite of this, the original sample may become increasingly unrepresentative of the target population due to long-term attrition.⁸ The ITC Netherlands Survey examines discrepancies in four demographics variables (gender, age, geographic region and household size) by comparing the sample composition against the composition of the smoking population. This makes it possible to customize replenishment samples and account for attrition in these demographic groups. This also compensates for disproportionate losses between these categories. However, this may not fully address the problem of nonrandom attrition because other factors may be affected. Thus, nonrandom attrition could still bias the sample and affect the generalizability of findings.

Previous studies of the effects of attrition on generalizability have found mixed results. Most studies examining attrition bias in longitudinal research only found small effects of attrition on generalizability. For example, regardless of large attrition rates (22–35%), Alderman *et al.*⁹ found that attrition was not a general or pervasive problem for attaining consistent estimates across data from several different surveys. In spite of a large attrition rate of 74%, Lohse *et al.*¹⁰ only found small, but significant, attrition effects; no effects were found on the primary outcome measures.

Other studies reported only minor problems arising from nonrandom attrition. Dennis and Li¹¹ found that dropouts and nondropouts had similar attitudes, values and behaviours with respect to areas like politics, health status and lifestyle. The differences they found became smaller after controlling for demographic variables. Another study found that nonrandom attrition had only a minor effect on key study indicators.⁷

Attrition had more substantial effects on sample composition in other studies. For instance, Mirowsky and Reynolds¹² found that suffering from depression predicted attrition. Although Bellón *et al.*¹³ did not support this finding, they identified a number of other psychosocial and socio-demographic predictors of attrition. Burkam and Lee¹⁴ reported that attrition resulted in an overestimation of the negative effects of race and ethnicity on academic achievement. We found no clear distinctions between the different analyses methods used in the studies described above and the attrition effects that they found.

In this study, we used data from the longitudinal ITC Netherlands Survey to examine attrition bias and whether replenishment samples maintained the representativeness of the cohort. The ITC Netherlands Survey is a web-based survey that has been conducted annually since 2008. Respondents were drawn from a large probability-based database.¹⁵ This study has three main objectives, to assess: (i) whether respondent attrition in the ITC Netherlands Survey biases the composition of the sample, (ii) whether attrition undermines the generalizability of the findings of the ITC Netherlands Survey and (iii) whether replenishment samples compensate for these possible effects.

Methods

The data

The ITC Netherlands Survey employs a probability-based web panel and consists of yearly survey waves with replenishment. The replenishment samples compensate for both the decrease in sample size and bias on demographic characteristics caused by attrition.

Wave 1 was conducted in April 2008 followed by a subwave in November and December 2008. Wave 2 was conducted in April and May 2009, wave 3 in May and June of 2010, wave 4 in May and June of 2011 and wave 5 in May and June 2012. The number of respondents surveyed in each wave and the number successfully followed from each wave is presented in figure 1. The total sample size ranged from 1717 respondents to 2022 respondents in any wave. Overall, 3437 respondents were included in this analysis. For the purpose of this analysis, we defined respondents who skipped a wave but later returned as dropouts once they missed one wave of data collection ($n = 305$).¹⁶ Information obtained from these participants after they returned was not used.

An extra survey wave was conducted after the first wave to evaluate the short-term effects of a media campaign. Only a subset of respondents from the baseline survey was approached and the timing between the first wave and this one did not follow the regular interval of one year. Therefore, we do not examine attrition in this extra wave, but we adjust the analyses for participation in this wave. In this study, this survey wave is called the 'subwave' to ease interpretation for readers, while the actual third wave is called the second wave and so on.

Respondents for the ITC Netherlands Survey were recruited from TNS NIPObase, a large probability-based database.¹⁵ Respondents are recruited into TNS NIPObase by phone or mail. Applying for participation is not permitted, which reduces the proportion of professional and inattentive respondents.¹⁷ A representative sample of Dutch smokers was obtained from TNS NIPObase with quotas on gender, age, geographic region and household size determined from the Dutch Continuous Survey of Smoking Habits,¹⁸ a national surveillance survey on smoking that has weekly measurements and, in 2007, had over 4000 respondents who were smokers.¹⁵ Younger respondents (15–30 years) were oversampled in order to increase

estimation precision for this specific subgroup.¹⁵ This procedure was also used for replenishment samples to ensure survey results were representative of the Dutch population of adult smokers. A previous study showed that there were only small differences in sample composition between the first ITC Netherlands survey and the Dutch population of smokers.¹⁵ Respondents were selected on the basis of three inclusion criteria: being 15 years or older, having smoked at least 100 cigarettes in their lifetime and smoking manufactured or roll-your-own cigarettes at least monthly. Respondents received points for questions they answered; points which could be exchanged for 5–7 euros per survey.

Ethics

The ITC Netherlands Surveys received ethics clearance from the University of Waterloo's Office of Research Ethics.

Measurements

The demographic characteristics gender, age, minority status, marital status, education, income, province and frequency of internet use were included in the analyses because previous studies indicated they were correlated with attrition.^{7,13,14,19} The smoking-related characteristics heaviness of smoking and ever made a quit attempt were included because they are key smoking-related characteristics within the ITC Project.²⁰ Descriptive statistics on these characteristics of the original sample are provided in the Supplementary Material (Supplementary Table S1).

Demographic characteristics

- Gender
- Age: categorized as 15–24, 25–39, 40–54 and 55+ years.
- Minority status: both parents born in The Netherlands versus otherwise.
- Marital status: not married, married, widowed and divorced.
- Highest achieved education: low (primary education and lower pre-vocational secondary education), moderate (middle pre-vocational secondary education and secondary vocational education) and high (senior general secondary education, [pre-]university education and higher professional education).
- Gross household income per month: low (less than 2000 Euro), middle (2000–3000 Euro) and high (more than 3000 Euro). An explicit category for nonresponse was also used (27.3%).
- Province of residence: 12 categories representing the provinces of The Netherlands.
- Frequency of internet use: low (<30 min/day), moderate (30–89 min/day) and high (90 min or more per day).

Smoking-related characteristics

- The Heaviness of Smoking Index had three categories indicative of nicotine dependence: low, moderate and high.²¹ It was calculated by summing two categorized measurements: number of cigarettes per day and time till first cigarette of the day, reverse scored.²² It had a fourth category indicating nonsmoking (quitter).
- Ever made a quit attempt: 'Yes' and 'No'.

Analyses

The effect of attrition on demographic and smoking-related characteristics of respondents was assessed using attrition ratios (ARs). ARs were calculated as:

$$AR = P_{fi}/P_{ri}$$

where P_{fi} is a proportion of a subgroup in the wave 1 sample and P_{ri} is the proportion of the same subgroup in the sample in wave 5. For example, if 20% of the wave 1 sample was married and 19% of this




Figure 1 Number of respondents of the ITC Netherlands Survey per wave and their retention rates. *In the subwave, 643 responded from 744 randomly selected participants from wave 1

sample retained in wave 5 was married, the AR would be $0.20/0.19 = 1.05$. This AR could be interpreted as a measure of change in the distribution of marital status between survey waves. The AR was calculated twice for every subgroup; once excluding the replenishment samples at wave 5 and once including them.

The ARs reported here represent the weight sizes needed to compensate for the effects of attrition on specific categories in the final wave.⁷ Although these ARs are similar to weight sizes, they are not the actual sampling weights used in the ITC Netherlands surveys. Those weights only account for gender, age, geographic region and household size.²³

A multivariable survival analysis was then conducted on the original unweighted data using a discrete-time proportional hazards regression model.²⁴ This specific multivariable survival analysis method can be performed with standard logistic regression^{25,26} and allowed us to disentangle the effects of correlated characteristics on attrition.⁷ The analysis was conducted in SPSS and examined participants that started in any wave and assessed which factors predicted whether they participated in all subsequent waves. The logit link was chosen over the complementary log–log link because it is easier to interpret, allowed us to report odds ratios and was used in similar previous studies.^{7,13,27} The main advantage of the complementary log–log link, that it is preferable for models of continuous time,²⁴ has no bearing on this analysis because our underlying metric of time was discrete: respondents were approached at the same period every year.

The same measures used in the analysis of ARs were included as covariates in the survival model. Because these variables were measured in every wave, they allowed the model to control for time varying characteristics. ‘Presence in subwave’ was added to control for whether participants were interviewed in the subwave. An interaction between ‘recruitment wave’ and ‘duration’ was also included. This was done in place of the intercept and allowed us to control for time.²⁴ ‘Recruitment wave’ indicated the wave in which a participant was first interviewed. ‘Duration’ indicated the number of interviews a respondent had completed including the current wave. An interaction between ‘recruitment wave’ and ‘duration’ controlled for the different time paths participants

followed. In other words, participants in their second wave (‘duration’ = 2) who were recruited for the cohort at wave 2 (‘recruitment wave’ = 2) were interviewed in 2010 (wave 3). Participants who were also in their second wave (‘duration’ = 2) but were recruited for the cohort at wave 1 (‘recruitment wave’ = 1) were interviewed in 2009 (wave 2).

The model was estimated with and without the three-way interactions of ‘recruitment wave’, ‘duration’ and each of the predictor variables using the deviance statistic.²⁴ None of the three-way interactions significantly improved model fit and they are therefore not reported here.

Results

Attrition ratios

Figures 2 and 3 illustrate the change in the proportion of respondents in each category, expressed as ARs. This is displayed in the figures for the original first wave sample present in wave 5 compared to the original first wave sample present in wave 1 (triangles) and for the total sample (i.e. with all replenishment samples) present in wave 5 compared to the first wave sample present in wave 1 (dots).

Figure 2 shows a relatively high AR of the original sample in wave 5 for the 15–24-year age group (AR = 1.46). The replenishment samples decreased the size of the ARs for the total sample in wave 5, from 1.46 to 1.12 in the 15–24-year age group and from 1.11 to 1.04 in ethnic minorities. However, the total sample also shows some small increases in the AR in a few demographic groups, most notably the low education group (AR increased from 1.00 to 1.23).

Figure 3 shows a large AR for participants who reported never having made a quit attempt in the original sample (AR = 1.94). Relatively high ARs were found for unmarried participants (1.29) and participants with low and medium nicotine dependence (1.38 and 1.29, respectively). ARs decreased towards 1.00 in these same categories when the total sample in wave 5 was used to calculate ARs. The largest difference in the AR occurred in the group of respondents who reported never having made a quit attempt (from 1.94 to 1.07). Other notable changes in the AR were seen for participants




Figure 2 ARs (weight sizes) for demographic characteristics of the original sample and the total sample. Legend: Δ Original sample ● Total sample




Figure 3 ARs (weight sizes) for demographic and smoking-related characteristics of the original sample and the total sample. Legend: Δ Original sample ● Total sample

with low nicotine dependence (from 1.38 to 1.25) and who were not married (from 1.29 to 1.15).

respondents aged 25–39 years (OR = 1.770) and moderate effects for respondents aged 15–24 years (OR = 2.377).

Multivariable attrition model

Table 1 reports the results of the multivariable survival analysis. It assesses the participant characteristics associated with dropout and includes the odds ratios for the independent variables. Respondents not participating in the subwave had slightly higher odds of attrition (OR = 1.755). There was a weak but significant effect of gender such that males had greater odds of dropping out than females (OR = 1.251). Younger respondents dropped out more often than the oldest respondents (aged 55+ years) with a weak effect among

Discussion

More than half of the ITC Netherlands sample that was recruited in 2008 was lost to follow-up by 2012. Despite this, only small (unweighted) ARs were found, suggesting the total sample present in wave 5 might not be biased by attrition. The survival analysis showed that the odds of attrition were somewhat higher for males, moderately higher for the youngest age group and moderately lower for those who participated in the subwave. This concurs with findings from the European Community Household Panel for 14

Table 1 Multivariable survival analysis: logistic regression of attrition between waves 1 and 5

Variables	Factors	OR ^a	95% CI ^b		Significance	Likelihood ratio test		
			Lower	Higher				
Present in subwave	Yes	Ref.				$\chi_{(1)}=52.73, P < 0.001$		
	No	1.755	1.504	2.051	***			
Gender	Female	Ref.				$\chi_{(1)}=12.65, P < 0.001$		
	Male	1.251	1.107	1.416	***			
Age (years)	55 and older	Ref.				$\chi_{(3)}=68.30, P < 0.001$		
	40–54	1.065	0.855	1.330				
	25–39	1.770	1.420	2.214	***			
	15–24	2.377	1.844	3.075	***			
Ethnicity	Dutch	Ref.				$\chi_{(1)}=0.35, P = 0.552$		
	Not Dutch	1.066	0.862	1.311				
Marital status	Married	Ref.				$\chi_{(3)}=4.98, P = 0.173$		
	Divorced	0.987	0.696	1.373				
	Widowed	1.365	0.798	2.233				
	Not married	1.167	0.996	1.367	†			
Highest education	High education	Ref.				$\chi_{(2)}=1.25, P = 0.535$		
	Moderate education	1.009	0.864	1.179				
	Low education	0.929	0.778	1.111				
Household income	3.000 euros and higher	Ref.				$\chi_{(3)}=1.20, P = 0.752$		
	Between 2.000 and 3.000 euros	0.985	0.828	1.172				
	Less than 2.000 euros	0.926	0.775	1.106				
	Unanswered	0.926	0.783	1.095				
Province	Limburg	Ref.				$\chi_{(11)}=12.55, P = 0.324$		
	Groningen	1.039	0.724	1.483				
	Friesland	0.912	0.630	1.312				
	Drenthe	0.829	0.545	1.242				
	Overijssel	1.050	0.768	1.435				
	Gelderland	0.975	0.731	1.305				
	Flevoland	0.983	0.606	1.557				
	Utrecht	0.906	0.640	1.278				
	Noord-Holland	1.135	0.871	1.486				
	Zuid-Holland	0.987	0.765	1.179				
	Zeeland	1.431	0.904	2.232				
	Noord-Brabant	0.844	0.644	1.109				
	Frequency of internet use	90 min or more per day	Ref.					$\chi_{(2)}=3.40, P = 0.183$
		30–89 min/day	0.969	0.837	1.123			
<30 min/day		1.110	0.945	1.303				
Heaviness of smoking	High nicotine dependence	Ref.				$\chi_{(3)}=1.50, P = 0.683$		
	Moderate nicotine dependence	0.933	0.719	1.224				
	Low nicotine dependence	1.016	0.770	1.352				
	Quit smoking	0.941	0.691	1.290				
Ever made a quit attempt	No	Ref.				$\chi_{(1)}=.74, P = 0.390$		
	Yes	1.068	0.920	1.242				
Interaction of recruitment wave and duration	1st wave, 1 wave	0.091	0.057	0.145	***	$\chi_{(3)}=38.86, P < 0.001$		
	1st wave, 2 waves	0.083	0.052	0.133	***			
	1st wave, 3 waves	0.118	0.073	0.189	***			
	1st wave, 4 waves	0.077	0.047	0.124	***			
	2nd wave, 1 wave	0.079	0.045	0.138	***			
	2nd wave, 2 waves	0.090	0.050	0.159	***			
	2nd wave, 3 waves	0.069	0.037	0.128	***			
	3rd wave, 1 wave	0.116	0.070	0.189	***			
	3rd wave, 2 waves	0.056	0.032	0.094	***			
	4th wave, 1 wave	0.070	0.041	0.118	***			

a: Ref., reference level.

b: CI, confidence interval.

* $P < 0.05$.** $P < 0.01$.*** $P < 0.001$.† $P < 0.10$.

countries, which found higher ARs for lower age groups⁷ and with findings from other ITC Europe Surveys.^{28,29} Other studies also found attrition effects for both gender and age,^{13,27} including ITC surveys conducted in other countries.³⁰

The lower risk of attrition for participants present in the subwave may indicate a greater probability of remaining in the survey for seven months (i.e. presence in the subwave) which would then predict remaining in the survey for one year (i.e. presence in the second wave). However, loyalty to the survey may also be a

factor: respondents who participated more often in the past are more likely to participate in subsequent waves. Survey loyalty may be due to the financial incentives provided for participation³¹ or because subwave participation encouraged respondent rapport, making them more devoted to future participation.

The tailored replenishment samples used in the ITC Netherlands Survey were designed to compensate for sample attrition and to ensure the cohort of smokers included remained representative of the population of Dutch smokers on specific variables (gender, age,

geographic region and household size). In doing so, the replenishment samples alleviated attrition bias on some other variables also (e.g. low nicotine dependence and never made a quit attempt). Additionally, the use of sampling weights increases the extent to which attrition effects are taken into account. It is therefore likely that the results from the ITC Netherlands survey are robust against deviations in sample compositions that result from respondent attrition.

Limitations and strengths

The analytic approach used in this study provides insight into the representativeness of the ITC Netherlands Survey and respondent attrition. ARs do not establish definitively whether a characteristic is significantly related to attrition or even whether participants dropped out of the cohort at all; the characteristics of a participant may just have changed (e.g. from never having made a quit attempt to having made a quit attempt). The sample is constantly changing and some of the observed changes in composition may be due to population changes. These changing characteristics may have inflated the ARs and they may also have an informing role on patterns of attrition.¹³ The survival model provides more insight into factors associated with attrition because it accounts for changing characteristics by way of time varying covariates. The model also makes it possible to account for the effects of correlated characteristics on attrition⁷ and includes replenishment samples in the analysis as potential sources of dropout, thus utilizing an extra source of information on attrition. However, our model does not contain data about characteristics of participants that changed since their last wave of participation before dropping out. Therefore, the model cannot account for such changes.⁷

Conclusion

Less than half of the ITC Netherlands sample that was recruited in 2008 remained in the cohort in 2012. However, the ARs calculated to measure attrition bias were small. Larger deviations, where participants with specific characteristics dropped out more often, were alleviated by replenishment samples, thus improving the sample structure. Our study revealed that only two variables out of ten were significant predictors of attrition: gender and age. Men and younger respondents dropped out more often. The replenishment samples in the ITC Netherlands Survey account for this because they are tailored to gender and age. These results support that sampling weights and tailored replenishment samples can compensate for attrition effects on the sample composition. Attrition prevention methods could also be aimed at participants who fall in categories that predict attrition.

Supplementary data

Supplementary data are available at *EURPUB* online.

Acknowledgements

Several members of the ITC Project team at the University of Waterloo have assisted in all stages of conducting the ITC Netherlands Surveys, which we gratefully acknowledge. In particular, we thank Thomas Agar, Project Manager of the ITC Europe countries.

Funding

The ITC Netherlands Surveys were supported by grants from the Netherlands Organization for Health Research and Development (ZonMw) (121010008, 70000001). The ITC Project was supported by the US National Cancer Institute (P50 CA111236, R01 CA100362 and P01 CA138389) and the Canadian Institutes of Health Research

(57897, 79551, 115016). G.T.F. was supported by a Senior Investigator Award from the Ontario Institute for Cancer Research and by a Prevention Scientist Award from the Canadian Cancer Society Research Institute.

Conflicts of interest: None declared.

Key points

- Replenishment samples tailored to keep the cohort representative of the population do compensate for attrition bias, even for variables other than the ones used to tailor the replenishment samples.
- Male gender and younger age were found to be the only characteristics that predicted attrition.
- Multivariable survival analyses across waves give a more exhaustive and therefore different picture of attrition than sampling weights do.
- Results suggest large dropout rates (>50%) in longitudinal surveys do not necessarily create problems with attrition bias.

References

- 1 Fitzgerald J, Gottschalk P, Moffitt R. An analysis of the impact of sample attrition on the second generation of respondents in the Michigan panel study of income dynamics. *J Hum Resour* 1998;33:300–44.
- 2 Leyland A, White I, Harding S, et al. Maximizing the return from cohort studies [published abstract]. *J Epidemiol Community Health* 2011;65 (Suppl 1):A21.
- 3 Menard S. Handbook of longitudinal research: design, measurement, and analysis. Burlington: Elsevier, 2008.
- 4 Poole WK, Perritt R, Shah KB, et al. A characterization of patient drop outs in a cohort of HIV positive/bisexual men and intravenous drug users. *J Epidemiol Community Health* 2001;55:66–7.
- 5 Fong GT, Cummings M, Borland R, et al. The conceptual framework of the International Tobacco Control (ITC) policy evaluation project. *Tob Control* 2006;15:3–11.
- 6 International Agency for Research on Cancer. IARC handbooks of cancer prevention, Vol. 12: methods for evaluating tobacco control policies. Lyon: World Health Organization, International Agency for Research on Cancer, 2008.
- 7 Watson D. Sample attrition between waves 1 and 5 in the European community household panel. *Eur Sociol Rev* 2003;19:361–78.
- 8 Deng Y, Hilligus DS, Reiter JP, et al. Handling attrition in longitudinal studies: the case for refreshment samples. *Stat Sci* 2013;28:238–56.
- 9 Alderman H, Behrman JR, Kohler HP, et al. Attrition in longitudinal household survey data: some tests for three developing-country samples. *Demogr Res* 2001;5:79–124.
- 10 Lohse GL, Bellman S, Johnson EJ. Consumer buying behavior on the internet: findings from panel data. *J Interact Mark* 2000;14:15–29.
- 11 Dennis JM, Li R. Effects of panel attrition on survey estimates. Paper presented at: 2003 Annual Meeting of the American Association for Public Opinion Research, Nashville, 2003. Available at: <http://www.knowledgenetworks.com/ganp/docs/kn%20panel%20attrition%20paper.pdf> (January 9 2016, date last accessed)
- 12 Mirowsky J, Reynolds JR. Age, depression, and attrition in the national survey of families and households. *Sociol Methods Res* 2000;28:476–504.
- 13 Bellón JA, de Dios Luna J, Moreno B, et al. Psychosocial and sociodemographic predictors of attrition in a longitudinal study of major depression in primary care: the predictD-Spain study. *J Epidemiol Community Health* 2010;64:874–84.
- 14 Burkam DT, Lee VE. Effects of monotone and nonmonotone attrition on parameter estimates in regression models with educational data. *J Hum Resour* 1998;33:556–74.
- 15 Nagelhout GE, Willemsen MC, Thompson ME, et al. Is web interviewing a good alternative to telephone interviewing? Findings from the International Tobacco Control (ITC) Netherlands Survey. *BMC Public Health* 2010;10:351.

- 16 Kruse Y, Callegaro M, Michael Dennis J, et al. Panel conditioning and attrition in the AP-Yahoo! news election panel study, 2009. Available at: http://www.knowledgenetworks.com/ganp/docs/jsm2009/Panel%20Conditioning%20and%20Attrition_JSM_2009_submitted.pdf (January 9 2016, date last accessed).
- 17 Willems P, Brown M, Van Ossenbruggen R, Vonk T. The effects of panel recruitment and management on research results. Dutch Online Panel Study NOPVO. Barcelona: ESOMAR, 2006.
- 18 Willemsen MC. The new EU cigarette health warnings benefit smokers who want to quit the habit: results from the Dutch continuous survey of smoking habits. *Eur J Public Health* 2005;15:389–92.
- 19 Vicente P, Reis E. “Frequency divide” on web surveys: differences of opinion, behaviour and demographics among internet users. Paper presented at: 58th World Statistical Congress, Dublin, 2011. Available at: <http://2011.isiproceedings.org/papers/950629.pdf> (January 9 2016, date last accessed).
- 20 Hyland A, Borland R, Li Q, et al. Individual-level predictors of cessation behaviours among participants in the International Tobacco Control (ITC) four country survey. *Tob control* 2006;15 (suppl 3):83–94.
- 21 Chaiton MO, Cohen JE, McDonald PW, Bondy SJ. The heaviness of smoking index as a predictor of smoking cessation in Canada. *Addict Behav* 2007;32:1031–42.
- 22 Heatherton TF, Kozlowski LT, Frecker RC, et al. Measuring the heaviness of smoking: using self-reported time to the first cigarette of the day and number of cigarettes smoked per day. *Br J Addict* 1989;84:791–800.
- 23 Boudreau C. Construction and use of sampling weights for the International Tobacco Control (ITC) Netherlands Survey. Waterloo, Ontario: Dept. of Statistics & Actuarial Science and Data Management Core (DMC) of ITC Project, 2010. Available at: http://www.itcproject.org/files/Report_Publications/Technical_Report/nl_w13_techreport_july62010rev.pdf (January 9 2016, date last accessed)
- 24 Singer JD, Willet JB. Applied longitudinal data analysis: modeling change and event occurrence. Oxford: Oxford University Press, 2003.
- 25 Efron B. Logistic regression, survival analysis, and the Kaplan-Meier curve. *J Am Stat Assoc* 1988;83:414–25.
- 26 Singer JD, Willet JB. It’s about time: using discrete-time survival analysis to study duration and the timing of events. *J Educ Behav Stat* 1993;18:155–95.
- 27 Behr A, Bellgardt E, Rendtel U. Extent and determinants of panel attrition in the European community household panel. *Eur Sociol Rev* 2005;21:489–512.
- 28 Brown A, Moodie C, Hastings G. A longitudinal study of policy effect (smoke-free legislation) on smoking norms: ITC Scotland/United Kingdom. *Nicotine Tob Res* 2009;11:924–32.
- 29 Mons U, Nagelhout GE, Guignard R, et al. Comprehensive smoke-free policies attract more support from smokers in Europe than partial policies. *Eur J Public Health* 2012;22:10–6.
- 30 Thompson ME, Fong GT, Hammond D, et al. Methods of the International Tobacco Control (ITC) four country survey. *Tob Control* 2006;15 (Suppl 3):12–8.
- 31 Olsen RJ. The problem of respondent attrition: survey methodology is key. *Mon Labor* 2005;128:63–70.