



UvA-DARE (Digital Academic Repository)

How did the treatment work for Robin? And for Dylan? Studying individual youth treatment mediators using single-case experimental designs

Maric, M.; Kok, S.I.

DOI

[10.1007/s10567-023-00442-7](https://doi.org/10.1007/s10567-023-00442-7)

Publication date

2023

Document Version

Final published version

Published in

Clinical Child and Family Psychology Review

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Maric, M., & Kok, S. I. (2023). How did the treatment work for Robin? And for Dylan? Studying individual youth treatment mediators using single-case experimental designs. *Clinical Child and Family Psychology Review*, 26(3), 721-726. <https://doi.org/10.1007/s10567-023-00442-7>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)



How did the Treatment Work for Robin? And for Dylan? Studying Individual Youth Treatment Mediators Using Single-Case Experimental Designs

Marija Maric^{1,2,3} · Shawn I. Kok¹

Accepted: 24 June 2023 / Published online: 5 July 2023
© The Author(s) 2023

Abstract

Study of individual treatment mechanisms in youth interventions facilitates evidence-based development, selection and implementation of treatment components that are most effective for each individual child. This position paper aims to bring together two important topics from the area of youth intervention research: mediators of treatment outcomes and single-case experimental design methodology. We start by outlining the benefits of studying within-person mechanisms and propose how statistical mediation analysis and single-case methods can be integrated to enable this type of research. Further, we review existing methodology for the study of individual youth treatment mechanisms and provide recommendations for the clinical practice research.

Keywords Treatment mediators · Single-case methods · Individual mechanisms · Youth interventions

In the 2012 volume of this journal, Maric et al. published a review on the importance of studying treatment mediation in youth intervention research. It was therewith emphasized that studying treatment mediators forms a crucial part in determining whether an intervention affects the intended symptom, risk factor or underlying process and hence whether it is effective (see also MacKinnon, 2008). Treatment mediators can thereby be seen as ‘mechanisms or processes through which a treatment achieves its effect’ (Kraemer et al., 2002). Furthermore, identifying treatment mediators can help to determine the most and least potent components of a treatment and thus further enhance its effectiveness and utility. Ultimately, this may decrease the mental burden that many children today are experiencing

by allowing for treatments to become more adapted to their unique situation.

One section of the earlier review by Maric et al. (2012) was devoted to drawing attention to studying treatment mediators in single-case experimental designs (SCEDs). At that time there was, however, not much to be mentioned except that we found it important to study mediating variables not only on the level of a group (as in randomized controlled trials [RCTs]), but also on the level of an individual client. Now, over a decade later, the topic of assessing mediators in SCEDs has become more relevant than ever. The aims of this position paper are therefore to: (i) define the relevance of studying youth treatment mediators on the level of a single client; (ii) discuss how statistical mediation analysis and single-case methods can be integrated to facilitate this research; and (iii) review existing methodology for the study of individual youth treatment mechanisms and provide recommendations for the clinical practice research.

✉ Marija Maric
m.maric@uva.nl

¹ Research Institute of Child Development and Education, University of Amsterdam, Amsterdam, The Netherlands

² Department of Child Development and Education, Faculty of Social and Behavioral Sciences, University of Amsterdam, Postbus 15776, 1001 NG Amsterdam, The Netherlands

³ University of Amsterdam, Nieuwe Achtergracht 127, 1018 WS Amsterdam, The Netherlands

Why is Attention for Individual Treatment Mediators Necessary?

There are at least three, somewhat interrelated, reasons to study treatment mediation in individual young clients. First, mental health problems in youth are not only highly

prevalent (Merikangas et al., 2010; World Health Organization, 2022), but are also hard to treat due to complex risk factor interplay. Various (combinations of) child, parent, family, and social context mechanisms can cause and maintain mental health problems in children and adolescents. In child A., for example, an interaction between negative self-esteem, changing schools, and cannabis use leads to depression, while in child B. an inhibited temperament in combination with poor social skills and parental psychopathology leads to the same problem. This heterogeneity in risk factors and mechanisms is not only causing extreme suffering in young people, but also points to the need to utilize personalized youth interventions. Discovering individual treatment mediators facilitates the development, selection and implementation of treatment components that are most effective for each individual client.

Second, our field has made a tremendous progress in the past 60 years regarding developing and testing interventions for various youth and family conditions. An empirical base of these interventions was also extensively established so that we have now, for example, learned that different modalities of Cognitive Behavioural Treatment (CBT) are effective for youth anxiety, major depression and trauma (e.g., Ewing et al., 2015; Kreuze et al., 2018; Oud et al., 2019; Sigurvinsdóttir et al., 2020; Weems & Neill, 2020). At the same time, however, we have also learned that not all youth profits equally when following treatment (Robinson et al., 2020). Disentangling reasons for this has turned out to be a difficult task considering the designs used to study efficacy and effectiveness of these interventions.

RCTs have long been the gold standard to evaluate the effects of youth treatments. However, to determine the effects of an intervention in RCTs, average symptom scores of treatment groups are compared to average symptom scores of a control group. Next to this, meta-analyses of RCTs are often used to underpin a treatment evidence-base, averaging these effects over different groups even further. On the one hand, RCTs are therewith suitable to study intervention effects in large-scale (e.g., prevention or school-based programs), answering questions such as ‘is intervention A *on average* more effective than intervention B?’. On the other hand, however, averaging participants’ scores and effects sizes (as in meta-analyses) is problematic and leads to loss of valuable information about individual variations in treatment effects. As mentioned, youth mental health problems are heterogeneous in nature and treatment effects that evolve within-persons cannot be captured by between-persons comparisons (Maric et al., 2023; Schuurman, 2023).

The third and final reason we want to mention in the context of this paper is that studying mediators on an individual level can aid the research in real-world settings and populations. It can assist clinicians and other mental health professionals in learning how the treatment works for their

individual clients and thus also other clients with similar profiles in similar circumstances. Furthermore, it provides a relatively easy way for researchers to test interventions and theories in concrete practical situations. All in all, this will certainly contribute to more evidence-based work of the therapists and more evidence-based clinical practice.

How SCEDs may Help in Assessing Individual Treatment Effects?


In the past decades, SCED methods have regained in popularity as a way to test treatment effects in youth populations (Kazdin, 2019). SCEDs therewith provide within-individual comparisons in which symptoms of interest of one or several participants are tested regularly (e.g., monthly, weekly, daily, and/or hourly) over a period of time. Examples of SCED designs include (1) an AB design (baseline phase A followed by an intervention phase B), (2) $A_1B_1A_2B_2$ design (in which an intervention is withdrawn during A_2 and again introduced during the B_2 phase), and (3) the multiple baseline SCEDs in which clients are randomized to different lengths of a baseline phase A before introducing an intervention phase B, making it possible to account for maturity effects in clients or passage of time (Barlow et al., 2009; Tate et al., 2016).

An important advantage of SCEDs is that, through the introduction of different phases of the study, individual clients serve as their own control. Therefore, a matching control group is not necessary. In Fig. 1B, for example, anxiety symptoms and coping of a single participant were monitored during the baseline and treatment phase. This design helps to answer the question whether there were changes in anxiety after the introduction of the treatment compared to the baseline phase. Likewise, we can assess whether the individual’s coping changed after the introduction of the treatment. To a certain degree, this will allow us to make some preliminary predictions regarding the potential of coping as a treatment mediator. Nonetheless, as we will demonstrate below, a proper evaluation of such a relation requires a more sophisticated approach.


Integrating Treatment Mediation in Youth Single-Case Research: Current Developments

The assessment of treatment mediators in SCEDs requires that two (or more) variables are tested simultaneously throughout each phase. Similar to group-level studies, the treatment mediator is therewith seen as a variable that lies within the causal chain between the intervention and (dependent) outcome (MacKinnon, 2008). An example of this relational model in SCEDs is depicted in Fig. 1A, in

(A) Basic mediation model with coping as a mediator



(B) Trends of coping and anxiety over time



(C) Mediation depicted over time assuming a direct (lag=1) relation




Fig. 1 Basic display of SCED mediation analysis. Panel **a** depicts the single mediator model. Applying this methodology on the example depicted in panel **(b)**, panel **(c)** shows a graphical representation of a

mediation effect of coping between a treatment and anxiety assuming a direct (lag = 1) effect

which a treatment (X) is hypothesized to produce changes in coping abilities (M), which in turn lead to changes in anxiety (Y). The effect of the treatment mediator can thereby be evaluated by statistically testing the effect of the treatment through paths a and b . It is important to note that the requirements of statistical mediation analysis which have previously been outlined for group studies (e.g., temporal precedence; Kazdin & Nock, 2003; MacKinnon, 2008; Maric et al., 2012) also apply when assessing mediators in SCEDs. Unlike group studies, however, SCEDs provide a within-subjects structure of data whereby each variable is repeatedly measured. This makes that the traditional mediation techniques may not be directly applied, and hence other data-analytic techniques are warranted.

While in the past decades attempts have been made to develop methodology to evaluate treatment mediators in SCEDs, each of them presented serious shortcomings. Gaynor and Harris (2008) designed a study to investigate whether changes in activation mediated changes in depressive symptoms in adolescents. The theoretical framework and the design were developed well, but the data were investigated visually. Although providing useful insights, visual inspections are nonetheless prone to bias and do not allow for a systematic evaluation of the extent to which the treatment mediator affects the outcome. In another study, Borckardt et al. (2008) tested changes in two processes over time, social engagement and mood, using cross-correlations. While correlations can be used to test associations in two variables over time, they are not a formal test of mediational pathways and treatment mediation. Finally, and more recent, Geuke et al. (2019) investigated the utility of combining previous SCED methodology to test for mediation and found that none of the techniques could be adapted to evaluate the b (the effect of the mediator on the outcome) and c' paths (the remaining effect of the treatment that is not accounted for by the mediator). Hence, further urging the need for new methodology to be developed.

Fueled by the Lorentz Workshop (2019), organized by Maric and colleagues in the city of Leiden in the Netherlands, preliminary efforts were made by a select group of clinical researchers and methodologists to develop methods to overcome these aforementioned shortcomings. Most of the proposed techniques consisted of piecewise linear regression analyses combined with interval testing for the obtained estimate of the mediated (or indirect) effect (Loeys & Rodenburg, 2022; Valente et al., 2022). In these techniques, each of the paths depicted in Fig. 1C are determined by assessing the change between phases in the averages (i.e., level), trends within the phase, or both. Furthermore, MacKinnon et al. (2022) described a randomized permutation test for single subject mediation using alternating treatment designs in which residuals are reassigned in many (i.e., thousands) possible combinations. This to acquire a

sampling distribution which allows to test the likelihood of the mediated effect found based on the actual data. In a study published in another journal, piecewise regression analysis was again applied, and it was emphasized that, similar to large group treatment mediation studies, it is important to study variables next to the mediator and outcome that may confound this relation (Valente et al., 2022).

Simultaneous to this development of new techniques, attention was also provided to the handling of autocorrelation, or the tendency of repeated measures to be correlated with one another (Busk & Marascuilo, 1988). For example, if a child shows a decline in symptoms at the start of the treatment phase, it is more likely that these symptoms will keep decreasing in subsequent measures. In other words, a value at one time point is predictive for the subsequent measures. This may have serious implications, however, for the underlying statistical assumptions and may in turn lead to biased estimates of standard error and hence inaccurate findings (Kazdin, 2011). In mediation analysis specifically, the computation of the intervals may be significantly influenced as these are dependent on the acquired standard errors. Therefore, methods have been proposed to take autocorrelation into account when assessing treatment mediation in SCEDs.

One common practise is to add lagged values of the mediator and outcome as covariates when determining the path values (MacKinnon et al., 2022). These lagged values can either be directly preceding (i.e., the previous measure) or more delayed (e.g., the measure from two or more time points ago) depending on the expected autocorrelation. Another method has been proposed by Loeys & Rodenburg (2022) in which Generalised Least Squares (GLS) models are used to compute an autocorrelation structure similar to multilevel approaches. Like with the lagged covariates, different orders of autocorrelation can be accounted for by fitting a corresponding structure. Moreover, the GLS method allows for the assessment of a small set of individuals at the same time. Thus, providing a means to compare individual trends with the total image on a statistical level.

Recommendations for the Youth Clinical Practice Research

Applying statistical mediation analysis in youth intervention research can be, in general, considered a challenging endeavor. Fortunately, established guidelines exist (Kazdin & Nock, 2003; MacKinnon, 2008; Maric et al., 2012) that can also be applied to design and conduct a study on treatment mediators in SCEDs. Because of the repeated structure of SCED data, existing statistical mediation data-analysis techniques do not apply and in this position paper we shared recent knowledge on the more suitable techniques.

We conclude by proposing guidelines for a set of issues that emerged from youth clinical practice research. One challenge, for instance, comes with the number of observations that is required to be able to obtain valid results. It has previously been estimated that between 30 and 100 observations are required to detect large effects depending on the type of technique used (Loeys & Rodenburg, 2022; Valente et al., 2022). This, however, is difficult to achieve within the clinical research field, as most treatments do not lend for such an extensive treatment window or the burden of repeated assessments for the clients may be too extensive. Our advice is to use short, personalized, assessments of hypothesized mediators and outcomes in the study and to choose smaller intervals (e.g., daily or twice-weekly assessments). Another issue concerns temporal precedence requirement for mediation, i.e., changes in the mediator should precede and influence changes in the outcomes. Temporal precedence can be achieved through shifting outcome and mediator measurements one (or more) lags to the left (Fig. 1C) or through choosing to conduct measurements directly after each other (e.g., as in the Ollendick et al., 1995 study). Finally, it should be considered that different SCED designs may be more or less suitable for the study of different types of mediators, and that we should keep account of the potential interplay between different SCED designs, types of interventions, and mediators. For now, multiple baseline SCEDs seem as the most feasible and experimental single-case design that can be implemented in youth intervention research. This design lends itself also for studying the changes in mediators and outcomes in the intervention phase as opposed to the baseline phase.

Conflict of Interest The authors have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Barlow, D. H., Nock, M., & Hersen, M. (2009). *Single case experimental designs: Strategies for studying behavior for change* (3rd ed.). Pearson Education.
- Borckardt, J. J., Nash, M. A., Murphy, M. J., Moore, M. H., Shaw, D., & O'Neil, P. M. (2008). Clinical practice as natural laboratory for psychotherapy research: A guide to case-based time-series analysis. *American Psychologist*, 63(2), 77–95. <https://doi.org/10.1037/0003-066x.63.2.77>
- Busk, P. L., & Marascuilo, L. A. (1988). APA PsycNet: A counterargument to the myth of no autocorrelation. *Behavioral Assessment*, 10(3), 229–242.
- Ewing, D. L., Monsen, J. J., Thompson, E. J., Cartwright-Hatton, S., & Field, A. P. (2015). A meta-analysis of transdiagnostic cognitive behavioural therapy in the treatment of child and young person anxiety disorders. *Behavioural and Cognitive Psychotherapy*, 43(5), 562–577. <https://doi.org/10.1017/s1352465813001094>
- Gaynor, S. G., & Harris, A. (2008). Single-participant assessment of treatment mediators. *Behavior Modification*, 32(3), 372–402. <https://doi.org/10.1177/0145445507309028>
- Geuke, G., Maric, M., Miočević, M., Wolters, L. H., & De Haan, E. (2019). Testing mediators of youth intervention outcomes using single-case experimental designs. *New Directions for Child and Adolescent Development*, 2019(167), 39–64. <https://doi.org/10.1002/cad.20310>
- Kazdin, A. E. (2011). *Single-case research designs: Method for clinical and applied settings* (2nd ed.). Oxford University Press.
- Kazdin, A. E. (2019). Single-case experimental designs. Evaluating interventions in research and clinical practice. *Behaviour Research and Therapy*, 117, 3–17. <https://doi.org/10.1016/j.brat.2018.11.015>
- Kazdin, A. E., & Nock, M. K. (2003). Delineating mechanisms of change in child and adolescent therapy: Methodological issues and research recommendations. *Journal of Child Psychology and Psychiatry*, 44(8), 1116–1129. <https://doi.org/10.1111/1469-7610.00195>
- Kraemer, H. C., Wilson, G., Fairburn, C. G., & Agras, W. S. (2002). Mediators and moderators of treatment effects in randomized clinical trials. *Archives of General Psychiatry*, 59(10), 877–883. <https://doi.org/10.1001/archpsyc.59.10.877>
- Kreuze, L., Pijnenborg, G. H. M., De Jonge, Y. B., & Nauta, M. (2018). Cognitive-behavior therapy for children and adolescents with anxiety disorders: A meta-analysis of secondary outcomes. *Journal of Anxiety Disorders*, 60, 43–57. <https://doi.org/10.1016/j.janxdis.2018.10.005>
- Loeys, T., & Rodenburg, L. A. (2022). Dealing with autocorrelation in the single case AB-design: A study assessing the mediating effect of infant crying on the relation between responsive soothing and infant sleep. *Evaluation & the Health Professions*, 45(1), 22–35. <https://doi.org/10.1177/01632787211069678>
- MacKinnon, D. P. (2008). Introduction to statistical mediation analysis. *Routledge*. <https://doi.org/10.4324/9780203809556>
- MacKinnon, D. P., Smyth, H. E., Somers, J., Ho, E., Norget, J., & Miočević, M. (2022). A randomization permutation test for single subject mediation. *Evaluation & the Health Professions*, 45(1), 54–65. <https://doi.org/10.1177/01632787211070811>
- Maric, M., Schumacher, L., Van den Noortgate, W., et al. (2023). A multilevel meta-analysis of single-case research on interventions for internalizing disorders in children and adolescents. *Clinical Child and Family Psychology Review*, 26, 416–429. <https://doi.org/10.1007/s10567-023-00432-9>
- Maric, M., Wiers, R. W., & Prins, P. J. M. (2012). Ten ways to improve the use of statistical mediation analysis in the practice of child and adolescent treatment research. *Clinical Child and Family Psychology Review*, 15(3), 177–191. <https://doi.org/10.1007/s10567-012-0114-y>
- Merikangas, K. R., He, J., Burstein, M., Swanson, S. A., Avenevoli, S., Cui, L., Benjet, C., Georgiades, K., & Swendsen, J. (2010). Lifetime prevalence of mental disorders in US adolescents: results from the national comorbidity survey replication-adolescent supplement (NCS-A). *Journal of the American Academy of Child*

- and *Adolescent Psychiatry*, 49(10), 980–989. <https://doi.org/10.1016/j.jaac.2010.05.017>
- Ollendick, T. H. (1995). Cognitive behavioral treatment of panic disorder with agoraphobia in adolescents: a multiple baseline design analysis. *Behavior Therapy*, 26(3), 517–531. [https://doi.org/10.1016/s0005-7894\(05\)80098-x](https://doi.org/10.1016/s0005-7894(05)80098-x)
- Oud, M., De Winter, L., Vermeulen-Smit, E., Bodden, D. H. M., Nauta, M. H., Stone, L. L., Van Den Heuvel, M. W. H., Taher, R. A., De Graaf, I., Kendall, T., Engels, R. C. M. E., & Stikkelbroek, Y. (2019). Effectiveness of CBT for children and adolescents with depression: A systematic review and meta-regression analysis. *European Psychiatry*, 57, 33–45. <https://doi.org/10.1016/j.eurpsy.2018.12.008>
- Robinson, L., Kellett, S., & Delgado, J. (2020). Dose-response patterns in low and high intensity cognitive behavioral therapy for common mental health problems. *Depression and Anxiety*, 37(3), 285–294. <https://doi.org/10.1002/da.22999>
- Schuurman, N. K. (2023). A “Within/Between Problem” Primer: *about (not) separating within-person variance and between-person variance in psychology*. Dynamic Modelling Lab, Utrecht University. <https://doi.org/10.31234/osf.io/7zgkx>
- Sigurvinsdóttir, A. L., Jensínudóttir, K. B., Baldvinsdóttir, K. D., Smáráson, O., & Skarphedinnsson, G. (2020). Effectiveness of cognitive behavioral therapy (CBT) for child and adolescent anxiety disorders across different CBT modalities and comparisons: A systematic review and meta-analysis. *Nordic J Psychiat*, 74(3), 168–180. <https://doi.org/10.1080/08039488.2019.1686653>
- Tate, R. L., Perdices, M., Rosenkoetter, U., Shadish, W., Vohra, S., Barlow, D. H., Horner, R., Kazdin, A., Kratochwill, T., McDonald, S., Sampson, M., Shamseer, L., Togher, L., Albin, R., Backman, C., Douglas, J., Evans, J. J., Gast, D., Manolov, R., ... Wilson, B. A. (2016). The single-case reporting guideline in behavioural interventions (SCRIBE) 2016 statement. *Physical Therapy*, 96(7), e1–e10. <https://doi.org/10.2522/ptj.2016.96.7.e1>
- Valente, M. J., Rijnhart, J. J., & Miočević, M. (2022). Causal mediation effects in single case experimental designs. *Psychological Methods*, 28(2), 488–506. <https://doi.org/10.1037/met0000497>
- Weems, C. F., & Neill, E. L. (2020). Empirically supported treatment options for children and adolescents with posttraumatic stress disorder: integrating network models and treatment components. *Current Treatment Options in Psychiatry*, 7(2), 103–119. <https://doi.org/10.1007/s40501-020-00206-y>
- World Health Organization. (2022). *Mental Health and COVID-19: Early evidence of the pandemic's impact* (WHO/2019-nCoV/Sci_Brief/Mental_health/2022.1). <https://apps.who.int/iris/handle/10665/352189>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.