UvA-DARE (Digital Academic Repository)

Post-burst quasi-periodic oscillations from GRO J1744-28 and from the rapid burster

Kommers, J.M.; Fox, D.W.; Rutledge, R.E.; Lewin, W.H.G.; van Paradijs, J.A.; Kouveliotou, C.

DOI

[10.1086/310669](https://doi.org/10.1086/310669)

Publication date

1997

Published in

Astrophysical Journal

[Link to publication](#)

Citation for published version (APA):

Kommers, J. M., Fox, D. W., Rutledge, R. E., Lewin, W. H. G., van Paradijs, J. A., & Kouveliotou, C. (1997). Post-burst quasi-periodic oscillations from GRO J1744-28 and from the rapid burster. *Astrophysical Journal*, 482, L53-L56. <https://doi.org/10.1086/310669>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

POSTBURST QUASI-PERIODIC OSCILLATIONS FROM GRO J1744–28 AND FROM THE RAPID BURSTER

JEFFERSON M. KOMMERS, DEREK W. FOX, AND WALTER H. G. LEWIN

Department of Physics and Center for Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139

ROBERT E. RUTLEDGE

Max-Planck-Institut für Extraterrestrische Physik, D-85740 Garching, Germany

JAN VAN PARADIJS

University of Alabama in Huntsville, Huntsville, AL 35812, and University of Amsterdam, Amsterdam, Netherlands

AND

CHRYSSA KOUVELIOTOU

Universities Space Research Association, Huntsville, AL 35800

Received 1997 January 10; accepted 1997 March 13

ABSTRACT

The repetitive X-ray bursts from the accretion-powered pulsar GRO J1744–28 show similarities to the type II X-ray bursts from the Rapid Burster. Several authors (notably, Lewin et al.) have suggested that the bursts from GRO J1744–28 are type II bursts (which arise from the sudden release of gravitational potential energy). In this paper, we present another similarity between these sources. *Rossi X-ray Timing Explorer* observations of GRO J1744–28 show that at least 10 out of 94 bursts are followed by quasi-periodic oscillations (QPOs) with frequencies of ~ 0.4 Hz. The period of the oscillations decreases over their ~ 30 – 80 s lifetime, and they occur during a spectrally hard “shoulder” (or “plateau”) that follows the burst. In one case, the QPOs show a modulation envelope that resembles simple beating between two narrow-band oscillations at ~ 0.325 and ~ 0.375 Hz. Using *EXOSAT* observations, Lubin et al. found QPOs with frequencies of 0.039 – 0.056 Hz following 10 out of 95 type II bursts from the Rapid Burster. As in GRO J1744–28, the period of these oscillations decreased over their ~ 100 s lifetime, and they occurred only during spectrally hard “humps” in the persistent emission. Even though the QPO frequencies differ by a factor of ~ 10 , we believe that this is further evidence that a similar accretion disk instability is responsible for the type II bursts from these two sources.

Subject headings: X-rays: bursts — stars: neutron — X-rays: general

1. INTRODUCTION

GRO J1744–28 is a recently discovered accretion-powered pulsar that shows repetitive X-ray bursts (Kouveliotou et al. 1996; Strickman et al. 1996; Giles et al. 1996; and references therein). Its unusual bursting behavior has been compared (see below) to that of the Rapid Burster (MXB 1730–335), which was discovered 20 years ago by Lewin et al. (1976). Analysis of the X-ray bursts from the Rapid Burster showed that it produced two distinct types of burst: type I, attributed to thermonuclear flashes on the surface of a neutron star; and type II, attributed to the release of gravitational potential energy due to spasmodic accretion. The mechanism responsible for the type II bursts has not been fully understood, although it is almost certainly related to an accretion disk instability (for reviews, see Lewin, van Paradijs, & Taam 1993, 1995). Rapidly repetitive type II bursts had previously been observed only from the Rapid Burster, so if the bursts from GRO J1744–28 are also of type II—as convincingly argued by Lewin et al. (1996)—then a comparison of these sources may constrain theories of the burst mechanism.

The pulse period of GRO J1744–28 is 467 ms, and the neutron star is in a 11.8 day binary orbit about a low-mass donor star (Finger et al. 1996). During the first 12 hr of bursting observed with BATSE on 1995 December 2 bursts occurred every 3–8 minutes, and for one 3 hr period the burst intervals clustered around 172 ± 15 s (Kouveliotou et al. 1996). Subsequently, the burst intervals became longer and

more erratic, and the burst rate settled at about 30–40 per day (corrected for Earth occultation and live time; Fishman et al. 1996). Burst durations were initially ~ 20 – 30 s and settled down to ~ 5 – 10 s. Unlike the type II bursts from the Rapid Burster, no relationship between the burst fluence and the time to the next (or previous) burst has been reported¹ (Kouveliotou et al. 1996; Strickman et al. 1996).

Using the first observations of GRO J1744–28 with the *Rossi X-ray Timing Explorer* (*RXTE*), Swank (1996) noted that the bursts were followed by a characteristic “dip” in the persistent emission level that took a few minutes to recover. No such dips were seen before the bursts. Subsequent *RXTE* observations showed that the bursts were sometimes preceded by steadily increasing variability in the persistent emission, including “micro” and “mini” bursts (Giles & Strohmayer 1996; Lewin et al. 1996; Giles et al. 1996).

Extensive reviews of the Rapid Burster and its complex behavior can be found in Lewin et al. (1993, 1995); see also references therein. Here we discuss the features of the type II bursts from this source that are relevant to our comparison with GRO J1744–28. The time intervals between type II bursts from the Rapid Burster range from ~ 10 s to ~ 1 hr, with the shorter intervals being more common. Burst durations range from ~ 2 s to ~ 680 s. The burst repetition pattern is that

¹ A preliminary analysis of the BATSE catalog of over 3000 bursts from GRO J1744–28 shows a statistically significant but weak correlation between the burst fluence scaled to the persistent emission level and the time to the next burst (Kommers et al. 1997).

of a relaxation oscillator: the fluence of a type II burst is roughly proportional to the time to the next burst (Lewin et al. 1976; Lewin et al. 1995). Persistent X-ray emission between the type II bursts is observed following long (duration > 30 s) bursts. The persistent flux emerges gradually after high-fluence bursts and decreases prior to the next burst (Marshall et al. 1979; Van Paradijs, Cominsky, & Lewin 1979; Stella et al. 1988a). These pre- and postburst features are referred to as “dips” in the persistent emission. The spectrum of the persistent emission is relatively soft during the dip just after a burst. It then rapidly increases in hardness, remaining hard for ~ 1 –2 minutes before gradually decreasing to again become very soft during the dip preceding the next burst (Stella et al. 1988b). The 1–2 minute period of spectrally hard emission corresponds to a “hump” in the persistent emission light curve.

Lubin et al. (1992) found “naked eye” quasi-periodic oscillations (QPOs) following 10 of 95 type II bursts from the Rapid Burster observed with *EXOSAT* in 1985 August. The oscillations occurred *only* during the spectrally hard humps (which immediately followed the postburst dips). The frequency of the oscillations ranged from 0.039 to 0.056 Hz, with a period decrease of 30%–50% observed over the ~ 100 s lifetime of the oscillations. The fractional root mean square (rms) variation in the oscillations was 5%–15%. In eight of the 10 cases, the “naked eye” oscillations were accompanied by ~ 4 Hz QPOs with rms variations of 6%–19% (Lubin et al. 1992).

Several authors have already noted similarities between the bursts from GRO J1744–28 and the Rapid Burster. After the discovery of the bursts from GRO J1744–28, Kouveliotou et al. (1996) suggested that the release of gravitational potential energy following an accretion instability might be responsible for the bursts. Lewin et al. (1996) made a detailed comparison between the two systems. Noting that both are transient low-mass X-ray binaries in which the accretor is a neutron star, they concluded that the bursts from GRO J1744–28 must be type II based on (1) the hardness of the burst spectra, (2) the lack of spectral evolution during the bursts, (3) the fact that the ratio of integrated energy in the persistent emission to that in the bursts was initially too small to allow for a thermonuclear burst mechanism, and (4) the presence of dips in the persistent emission following bursts (Lewin et al. 1996). Subsequently, Sturmer & Dermer (1996) reached the same conclusion. The possibility of thermonuclear burning in GRO J1744–28 has been discussed by Bildsten & Brown (1997).

In this paper, we present another similarity between the Rapid Burster and GRO J1744–28. Both sources show transient QPOs during spectrally hard emission intervals following bursts.

2. OBSERVATIONS

Since 1996 January 18, *RXTE* has performed numerous observations of GRO J1744–28. The data discussed here were taken by the Proportional Counter Array (PCA) instrument and are publicly available. Between 1996 January 18 and April 26, there were 94 main bursts observed (Giles et al. 1996). To refer to these bursts individually, we number them sequentially from 1 to 94.

The light curves of the bursts reveal a rich phenomenology in the burst profiles. Figure 1a shows the “dip” in persistent emission that follows some bursts, as first noted by Swank (1996). A broad “shoulder” (or “plateau”) of emission above the mean preburst level immediately follows some bursts, as

FIG. 1.—Light curves of three bursts from GRO J1744–28. The horizontal and vertical scales are each different to highlight the feature of interest. (a) The “dip” following burst 33. (b) A “shoulder” after burst 65; it occurs before a slight “dip.” QPOs with a mean frequency of $\sim 0.35 \pm 0.02$ Hz are present in the shoulder but are not apparent at this scale and time binning. (c) Shows an example of strong QPO with mean frequency 0.37 ± 0.03 Hz occurring during the shoulder after burst 68. The termination of the shoulder and the “dip” occur after the plotted time interval.

shown in Figure 1b. The shoulder occurs *before* the dip whenever both features are present. Some bursts show large-amplitude oscillations with frequencies ~ 0.4 Hz during their shoulder, as shown in Figure 1c (Kommers et al. 1996).

Oscillations can be seen during the shoulders of the light curves immediately following at least 10 of the 94 bursts. These 10 bursts were numbers 8, 12, 14, 18, 43, 53, 65, 68, 77, and 94. Typically 5–15 cycles of the QPOs are apparent, with as many as ~ 25 cycles seen in the case of burst 65. The low frequency of the oscillations and the low numbers of cycles make it difficult to use Fourier power spectra to study these QPOs. We have instead found the mean periods of the oscillations by estimating the time intervals between successive maxima in the count rates.

The mean frequency of the oscillations over the ensemble of bursts is 0.38 ± 0.04 Hz. The mean frequency of the oscillations after individual bursts varies from 0.35 ± 0.02 Hz in burst 65 to 0.49 ± 0.03 Hz in burst 12. (The uncertainty in these figures represents the uncertainty in the mean frequency.) The period of the QPOs following a given burst typically wanders nonmonotonically about a mean period by

FIG. 2.—Hardness, power spectrum, and intensity profile of burst 65. The top panel shows the ratio of the 3–5.6 keV count rate to the 2–3 keV count rate. (No instrumental dead time corrections have been applied, which may affect the hardness during the burst. The hardness during the shoulder is not significantly affected by dead time.) The middle panel shows a power spectrum with higher powers represented by darker shades. The lower panel shows the intensity profile. Notice that the 0.35 ± 0.02 Hz QPOs occur after the burst, and the spectrum during their presence is relatively hard. The inset shows the portion of the intensity profile during which the QPOs were observed. The data in the inset have been smoothed with a boxcar average to highlight the apparent beating between oscillations at ~ 0.325 Hz and ~ 0.375 Hz.

$\sim 20\%$ over the lifetime of the oscillations. Using the χ^2 statistic, we exclude (at the 2σ level or better in each of the 10 bursts) the null hypothesis that the period between successive QPO pulses is constant.

To identify any overall tendency toward increasing or decreasing QPO periods, we looked for bursts where the Spearman rank-order correlation coefficient (r_s) indicated that the time intervals between QPO pulses were correlated (or anti-correlated) with arrival time. In no case did we find a positive r_s , which would have indicated an overall trend toward increasing periods. For bursts 18, 53, 65 (see Fig. 2), and 94, we found an anticorrelation at the 2σ level or greater. In these bursts the QPO period tends to decrease by 10%–20% over the lifetimes (~ 50 – 80 s) of the oscillations.

The fractional rms variations of the oscillations range from $\sim 5\%$ – 13% . Using the counts-to-flux conversion 4.1×10^{-12} ergs $\text{cm}^{-2} \text{s}^{-1}$ per count (total PCA band; Giles et al. 1996) the rms variations in flux units range from 1.2×10^{-9} ergs $\text{cm}^{-2} \text{s}^{-1}$ to 1.4×10^{-8} ergs $\text{cm}^{-2} \text{s}^{-1}$.

The shoulders during which the postburst oscillations occur are spectrally harder than the persistent emission. Figure 2 (top panel) shows a hardness ratio for burst 65. The hardness is defined as the count rate (averaged over 1.0 s) in the 3–5.6 keV range divided by that in the 2–3 keV range. The middle panel shows the power spectrum as a function of time. Higher power levels are shown with darker shades. The presence of the QPO is shown by a dark band at 0.4 Hz lasting from the end of the burst until ~ 100 s after the peak. The bottom panel shows the burst intensity profile. The spectrum of the burst counts is clearly harder than that of the persistent emission, but even after the main part of the burst subsides the hardness ratios during the presence of the QPO (20–100 s after the burst) exceed those of the preburst persistent emission.

The QPOs following burst 65 show a modulation envelope reminiscent of “beating” between oscillations at separate frequencies; see the bottom panel of Figure 2. The count rates shown in the inset have been smoothed with a boxcar average to highlight the envelope of the (roughly) 0.05 Hz “beats.” If this modulation envelope can be attributed to simple beating, the two narrow-band oscillations must occur concurrently and have roughly comparable amplitudes. An estimate of the two frequencies involved can be obtained from the mean frequency $f_{\text{ave}} = (f_1 + f_2)/2$ and the beat frequency $f_{\text{beat}} = |f_1 - f_2|$. The mean frequency for the ~ 25 cycles of QPOs following this burst is 0.35 ± 0.02 Hz. The two frequencies that are beating must then be roughly $f_1 = f_{\text{ave}} - f_{\text{beat}}/2 = 0.325$ Hz and $f_2 = f_{\text{ave}} + f_{\text{beat}}/2 = 0.375$ Hz. The average frequency wanders by about 20%, so if beating is responsible for the apparent modulation envelope then the frequencies of the two narrow-band oscillations wander as well. The presence of a similar phenomenon following other bursts is difficult to determine because fewer cycles are available.

Over the course of the 1996 January–May *RXTE* observations, the burst fluence, peak flux, and persistent emission level each decreased (approximately linearly) by a factor of ~ 4 – 5 (Giles et al. 1996). The average frequency and rms amplitude of the postburst QPOs show no significant ($>2\sigma$) correlations with any of these quantities. QPOs with nearly the same frequency are seen in bursts for which the persistent emission level differs by a factor of ~ 4 .

Although our analysis focused on the 1995 December–1996 May outburst of GRO J1744–28, we note that postburst oscillations were also detected following a burst in 1996 June. This burst occurred when the source temporarily resumed bursting activity at a lower level than the main 1995 December–1996 May outburst (Jahoda, Strohmayer, & Corbet 1996). Kommers et al. (1996) reported 0.4 Hz oscillations during a ~ 25 s “shoulder” following the sixth of 7 bursts observed with the PCA on 1996 June 4. The fractional rms amplitude of these oscillations was $25\% \pm 5\%$. A second large outburst from GRO J1744–28 has been in progress since 1996 December 2, but we have not yet analyzed these observations (Kouveliotou et al. 1997; Stark et al. 1997).

3. DISCUSSION

The ~ 0.4 Hz oscillations following some bursts from GRO J1744–28 are reminiscent of oscillations that have been seen following some type II bursts from the Rapid Burster. This likeness between the two sources complements the comparisons made by Lewin et al. (1996). Although it is not certain that the postburst oscillations are the same phenomenon in

both sources, the following similarities are worth consideration. (1) The postburst oscillations follow bursts that do *not* show the profiles or spectral evolution characteristic of type I (thermonuclear flash) bursts. (2) The oscillations occur during a period of spectrally hard emission. (3) When an overall change in QPO period is observed, it is a decrease: 10%–20% for GRO J1744–28, and 30%–50% for the Rapid Burster. (4) The fractional rms amplitude of the oscillations is roughly 5%–15%.

There are some differences, however. (1) The frequency of the postburst oscillations in GRO J1744–28 is roughly 0.4 Hz, while in the Rapid Burster, the postburst oscillations have frequencies of 0.04 Hz and are in some cases accompanied by 4 Hz QPOs. (2) In GRO J1744–28, the oscillations occur *before* the dip in the persistent emission; but in the Rapid Burster, they appear *after* the dip (Lubin et al. 1992).

Cannizzo (1996a, 1996b) has shown that a simple numerical model of a thermal-viscous accretion disk instability can reproduce some features of the GRO J1744–28 bursts. In his model, interplay between radial and vertical energy transport in the disk causes oscillations in the ratio of gas pressure to total pressure (denoted by β). These oscillations increase in amplitude and eventually become nonlinear, leading to a Lightman-Eardley instability (Lightman & Eardley 1974). For specific choices of the inner and outer disk radii ($r_{\text{inner}} = 10^{7.5}$ cm, $r_{\text{outer}} = 10^9$ cm) and the form of the viscous stress, integration of the time-dependent model reproduces the ~ 1000 s recurrence times and the ~ 10 s durations of the bursts, as well as the postburst dips. Cannizzo (1996b) also points out that the increased variability *before* bursts may be related to the oscillations in β that precede the instability. These oscillations may occur in a variety of modes, but in his idealized case appear to have frequencies of about 0.05 Hz (Cannizzo 1996b). It remains to be seen whether this model can explain the oscillations reported here.

Abramowicz, Chen, & Taam (1995) have shown that low-frequency oscillations in mass flow can arise in accretion disk–corona systems. In their model, the development of a Lightman-Eardley instability is moderated by energy dissipation in the coronal region, which lies above the thin disk. They propose that the strong ~ 0.042 Hz oscillations seen in the Rapid Burster may arise from such a mechanism. The frequency of these QPOs may be relatively insensitive to the

accretion rate, since it is determined by several competing factors including the radius of the inner accretion disk. The spectrum of the QPOs is expected to be hard whenever the QPOs originate from the hot inner region (Abramowicz et al. 1995).

The common feature responsible for oscillatory behavior in these models is the presence of a mechanism that couples the vertical energy transport to the radial energy flux. In Cannizzo's model, this mechanism also acts to produce the bursts. To make a quantitative comparison of these models with the characteristics of the postburst oscillations in GRO J1744–28 and the Rapid Burster, one would have to know what QPO frequencies are expected based on the physical parameters that distinguish the two sources. The magnetic field strength and inner disk radius in particular should be quite different (Lewin et al. 1996; Finger et al. 1996; Sturmer & Dermer 1996), which might account for the factor of ~ 10 difference in postburst QPO frequency. Detailed information of this kind has not yet been presented for the models of Cannizzo (1996a, 1996b) or Abramowicz et al. (1995).

The presence of transient postburst QPOs is another similarity between the bursts from GRO J1744–28 and the type II bursts from the Rapid Burster. Combined with the comparison of Lewin et al. (1996), this similarity is further evidence that the bursts from GRO J1744–28 are of type II. If the same disk instability is responsible for the type II bursts from each of these sources, then the postburst oscillations provide another observational benchmark for models of the burst mechanism as well as the QPOs.

J. M. K. and D. W. F. acknowledge support from National Science Foundation Graduate Research Fellowships during the preliminary phase of this research. J. M. K. acknowledges subsequent support from a NASA Graduate Student Researchers Program Fellowship NGT8-52816. R. R. was supported by the NASA Graduate Student Researchers Program under grant NGT-51368. W. H. G. L. acknowledges support from NASA under grant NAG5-2046. J. v. P. acknowledges support from NASA under grant NAG5-2755. C. K. acknowledges support from NASA under grant NAG5-2560. We thank Dr. Ed Morgan for helpful discussions and assistance with the *RXTE* data archive at MIT.

REFERENCES

- Abramowicz, M. A., Chen, X., & Taam, R. E. 1995, *ApJ*, 452, 379
 Bildsten, L., & Brown, E. F. 1997, *ApJ*, 477, 897
 Cannizzo, J. K. 1996a, *ApJ*, 466, L31
 ———. 1996b, preprint
 Finger, M. H., Koh, D. T., Nelson, R. W., Prince, T. A., Vaughan, B. A., & Wilson, R. B. 1996, *Nature*, 381, 291
 Fishman, G. J., et al. 1996, *IAU Circ.*, No. 6290
 Giles, A. B., & Strohmayer, T. 1996, *IAU Circ.*, No. 6338
 Giles, A. B., Swank, J. H., Jahoda, K., Zhang, W., Strohmayer, T., Stark, M. H., & Morgan, E. H. 1996, *ApJ*, 469, L25
 Jahoda, K., Strohmayer, T., & Corbet, R. 1996, *IAU Circ.*, No. 6414
 Kommers, J. M., et al. 1997, in preparation
 Kommers, J. M., Rutledge, R. E., Fox, D. W., Lewin, W. H. G., Morgan, E. H., Kouveliotou, C., & Van Paradijs, J. 1996, *IAU Circ.*, No. 6415
 Kouveliotou, C., Deal, K., Richardson, G., Briggs, M., Fishman, G., & van Paradijs, J. 1997, *IAU Circ.*, No. 6530
 Kouveliotou, C., Van Paradijs, J., Fishman, G. J., Briggs, M. S., Kommers, J. M., Harmon, B. A., Meegan, C. A., & Lewin, W. H. G. 1996, *Nature*, 379, 799
 Lewin, W. H. G., et al. 1976, *ApJ*, 207, L95
 Lewin, W. H. G., Rutledge, R. E., Kommers, J. M., Van Paradijs, J., & Kouveliotou, C. 1996, *ApJ*, 462, L39
 Lewin, W. H. G., Van Paradijs, J., & Taam, R. E. 1993, *Space Sci. Rev.*, 62, 223
 ———. 1995, in *X-ray Binaries*, ed. W. H. G. Lewin, J. Van Paradijs, & E. P. J. Van den Heuvel (Cambridge: Cambridge Univ. Press), 175
 Lightman, A. P., & Eardley, D. M. 1974, *ApJ*, 187, L1
 Lubin, L. M., Lewin, W. H. G., Rutledge, R. E., Van Paradijs, J., Van der Klis, M., & Stella, L. 1992, *MNRAS*, 258, 759
 Marshall, H. L., Ulmer, M. P., Hoffman, J. A., Doty, J., & Lewin, W. H. G. 1979, *ApJ*, 227, 555
 Stark, M. J., Jahoda, K., Swank, J., & Strohmayer, T. 1997, *IAU Circ.*, No. 6548
 Stella, L., Haberl, F., Lewin, W. H. G., Parmar, A. N., Van der Klis, M., & Van Paradijs, J. 1988a, *ApJ*, 327, L13
 Stella, L., Haberl, F., Lewin, W. H. G., Parmar, A. N., Van Paradijs, J., & White, N. E. 1988b, *ApJ*, 324, 379
 Strickman, M. S., et al. 1996, *ApJ*, 464, L131
 Sturmer, S. J., & Dermer, C. D. 1996, *ApJ*, 465, L31
 Swank, J. 1996, *IAU Circ.*, No. 6291
 Van Paradijs, J., Cominsky, L., & Lewin, W. H. G. 1979, *MNRAS*, 189, 387