



## UvA-DARE (Digital Academic Repository)

### Alterations in the programming of energy metabolism in adolescents with background exposure to dioxins, dl-PCBs and PBDEs

Leijs, M.M.; Koppe, J.G.; Vulsma, T.; Olie, K.; van Aalderen, W.M.C.; de Voogt, W.P.; Legler, J.; ten Tusscher, G.W.

**DOI**

[10.1371/journal.pone.0184006](https://doi.org/10.1371/journal.pone.0184006)

**Publication date**

2017

**Document Version**

Final published version

**Published in**

PLoS ONE

**License**

CC BY

[Link to publication](#)

**Citation for published version (APA):**

Leijs, M. M., Koppe, J. G., Vulsma, T., Olie, K., van Aalderen, W. M. C., de Voogt, W. P., Legler, J., & ten Tusscher, G. W. (2017). Alterations in the programming of energy metabolism in adolescents with background exposure to dioxins, dl-PCBs and PBDEs. *PLoS ONE*, 12(9), [e0184006]. <https://doi.org/10.1371/journal.pone.0184006>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)*

RESEARCH ARTICLE

# Alterations in the programming of energy metabolism in adolescents with background exposure to dioxins, dl-PCBs and PBDEs

Marika M. Leijns<sup>1,2#a\*</sup>, Janna G. Koppe<sup>1,3</sup>, Thomas Vulmsa<sup>1</sup>, Kees Olie<sup>2</sup>, Wim M. C. van Aalderen<sup>1</sup>, Pim de Voogt<sup>2,4</sup>, Juliette Legler<sup>5#b</sup>, Gavin W. ten Tusscher<sup>6\*</sup>

**1** Department of Paediatrics and Neonatology, Emma Children's Hospital Academic Medical Center, Amsterdam, The Netherlands, **2** IBED/AEE, University of Amsterdam, Amsterdam, The Netherlands, **3** Ecobaby Foundation, Loenersloot, The Netherlands, **4** KWR Watercycle Research Institute, Nieuwegein, The Netherlands, **5** Institute for Environmental Studies (IVM) VU University, Amsterdam, The Netherlands, **6** Department of Paediatrics and Neonatology, Westfriesgasthuis, Hoorn, The Netherlands

#a Current address: Department of Dermatology and Allergology, RTWH University Aachen, Aachen, Germany

#b Current address: College of Health and Life Sciences, Brunel University, London, United Kingdom

\* [g.tentusscher@westfriesgasthuis.nl](mailto:g.tentusscher@westfriesgasthuis.nl) (GWTT); [mleijns@ukaachen.de](mailto:mleijns@ukaachen.de) (MML)



**OPEN ACCESS**

**Citation:** Leijns MM, Koppe JG, Vulmsa T, Olie K, van Aalderen WMC, de Voogt P, et al. (2017) Alterations in the programming of energy metabolism in adolescents with background exposure to dioxins, dl-PCBs and PBDEs. *PLoS ONE* 12(9): e0184006. <https://doi.org/10.1371/journal.pone.0184006>

**Editor:** Umberto Simeoni, Centre Hospitalier Universitaire Vaudois, FRANCE

**Received:** May 18, 2017

**Accepted:** August 16, 2017

**Published:** September 12, 2017

**Copyright:** © 2017 Leijns et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data is contained within the paper and Supporting Information files.

**Funding:** The study was financially supported by the Netherlands Ministry of Housing, Spatial Planning and the Environment. The paper was supported by E.C. grant: OBELIX nr: 227391.

**Competing interests:** The authors have declared that no competing interests exist.

## Abstract

### Objectives

Dioxins and PCBs are highly toxic and persistent environmental pollutants that are measurable in humans worldwide. These persistent organic pollutants are associated with a higher incidence of diabetes mellitus. We hypothesise that perinatal (background) exposure to industrial pollutants like dioxins also influences body mass development and energy metabolism in later life.

### Study design

In The Netherlands, the perinatal exposure (prenatal exposure and postnatal lactational intake) to dioxins has been studied prospectively since 1987. Fasting glucose, insulin, HbA1c and leptin were analysed in 33 children of the original cohort of 60. BMI, glucose:insulin and BMI:leptin ratios were calculated. Prenatal exposure, lactational intake and current serum levels of dioxins (PCDD/F), dl-PCBs and PBDE concentrations were determined using (HR)GC-MS.

### Results

Prenatal dioxin (PCDD/F) exposure was positively correlated to the glucose:insulin ratio ( $p = 0.024$ ) and negatively correlated to the fasting insulin concentration ( $p = 0.017$ ) in adolescence. Postnatal lactational PCDD/F intake was also negatively correlated to fasting insulin concentration ( $p = 0.028$ ). Current serum levels of PCDD/Fs and total TEQ (dl-PCBs + PCDD/Fs) were positively correlated to the fasting serum glucose concentration ( $p = 0.015$  and  $p = 0.037$ , respectively). No metabolic effects were seen in association with current serum levels of PBDEs. A positive correlation between the insulin and leptin concentrations

**Abbreviations:** PCDD, polychlorinated dibenzo-*p*-dioxins; PCDF, polychlorinated dibenzofurans; TCDD, 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin; PCB, polychlorinated biphenyls; PBDE, polybrominated diphenylethers; AhR, aryl hydrocarbon receptor; BMI, body mass index; HR, high resolution; GC/MS, gas chromatography/mass spectrometry; HbA1c, hemoglobin A1c; glycosylated hemoglobin; DDE, dichlorodiphenyldichloroethene; DDT, dichlorodiphenyltrichloroethane; PBB, polybrominated biphenyls; I-TEQ, International Toxic Equivalents; dl, dioxin-like; RA, retinoic acid.

( $p = 0.034$ ) was observed. No effects were found on leptin levels, BMI:leptin ratio, HbA1c levels or BMI.

## Discussion/Conclusion

This study indicates that prenatal and lactational exposure influences glucose metabolism in adolescents, presumably through a negative effect on insulin secretion by pancreatic beta cells. Additionally, the very low recent background exposure to dioxins in puberty possibly has an effect on the glucose level.

## Introduction

Polychlorinated dibenzo-*p*-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs) and polychlorinated biphenyls (PCBs) belong to the group of most hazardous environmental toxicants. PCDDs and PCDFs are unintentionally formed by-products of the incineration of waste, metallurgic processes, the production of chlorinated phenols, and bleaching of paper pulp [1, 2]. PCBs were intentionally produced mainly because of their extreme stability and resistance to acids, bases, hydrolysis and heat [3]. As a result of emissions during these processes, these compounds are widespread in nature and bio-accumulate, resulting in the contamination of human food and breast milk. As a result, all European children are exposed to detectable levels [4]. PCDDs, PCDFs and planar PCBs are grouped together as 'dioxins' or 'dioxin-like compounds', because of planar structure which gives the ability to bind the aryl hydrocarbon (Ah) receptor. Genomic and non-genomic pathways have been identified after binding the Ah-receptor. The genomic (or classical action) pathway is interrelated with the retinoic acid (RA) signalling pathway [5]. Binding with this receptor mediates many pathological processes like teratogenesis and tumor promotion as well as disturbances in many metabolic processes in humans and animals [6]. The non-genomic pathway has been identified to play an important role in the inflammatory action of dioxins (intracellular  $\text{Ca}^{2+}$  concentration, enzymatic activation of phospholipase A2 and Cox2) [7] as well as other processes like insulin secretion, fat storage, liver damage and tumorigenesis, and is potentially regulated by miRNAs [8]. Dioxin-like compounds as well as PCBs have been proven to be endocrine disruptors by a variety of mechanisms [9–11].

Other, more recently produced persistent toxic compounds are the flame retardants, polybrominated diphenylethers (PBDEs) [12]. An increase in serum concentrations of these compounds began in the nineties [13]. Over the last three decades these compounds have been used more and more frequently in various materials such as electronic equipment, plastics, carpet liners and furniture textiles. Human uptake is mostly through ingestion by eating polluted food. Inhalation of dust is also an important exposure route, especially in younger children [14].

Diabetes and obesity are currently epidemics throughout the Western World. Besides eating habits and genetic factors, environmental factors such as smoking during pregnancy and famine during the first three months of pregnancy, followed by an abundance of food in utero and later, play a role in later body mass index (BMI) and energy metabolism [15]. Various correlations between persistent organic pollutants, especially dioxins, and insulin resistance and diabetes mellitus have been published [16].

The environment in early (perinatal) life sets the stage for lifelong health. There is empirical support from the Dutch famine study that early-life environmental conditions can cause epigenetic changes that persist throughout life [17].

Most studies on dioxins in relation to diabetes are based on cohorts of highly exposed populations, such as veterans exposed to dioxins through contact with the defoliant Agent Orange

[18;19], and the highly exposed Yucheng cohort, exposed to PCBs and PCDFs through contaminated rice oil [20]. A higher incidence of diabetes was also seen in the Italian Seveso cohort, exposed to dioxins following an explosion at a herbicide manufacturing plant [21] and in a recent German occupational study [22]. Studies of lower exposed subjects include studies from Japan [23], the National Health and Nutrition Survey from the US [24], and Belgium [25]. Most of these studies, however, were not prospective but cross sectional and the studies were performed in adults. Limited prospective studies have focussed on developmental exposures to dioxin, and to our knowledge, no epidemiological studies have been carried out which examine possible effects of PBDE exposure on diabetes and obesity.

Our prospective study of mother-baby pairs was started in 1987-1990/1991 to study the effects of dioxins in the prenatal and nursing period [26]. Previous results linked to metabolism include a significant decrease in serum retinol binding protein (RBP) found during the eleventh postnatal week in relation to a higher lactational exposure to dioxins and furans (PCDD/Fs). This finding is suggestive of an inhibiting effect of dioxins on adipocyte differentiation, that normally takes place after birth, since RBP is a protein produced by mature adipocytes [27]. Interestingly, the inhibition of adipocyte differentiation by TCDD has been shown *in vitro* [28].

These results indicate that dioxin exposure has biological effects early in life. Given the associations mentioned in the aforementioned studies, we analysed energy metabolism parameters in the cohort, including fasting glucose, fasting insulin, glucose:insulin ratio, glycosylated haemoglobin (HbA1c), and leptin, in relation to prenatal and lactational dioxin exposure as well as current dioxin, dioxin-like PCB and PBDE concentrations.

Leptin is a hormone produced by white adipose tissue. It has an important function in the regulation of appetite in the hypothalamic appetite centre, giving a feeling of satiation in order to limit caloric surplus. It also plays a role in the energy production from fatty acids in skeletal muscle cells [29].

We hypothesise that prenatal, lactational and current dioxin and PBDE exposure affects metabolic processes. Theoretically, this may lead to an increase in the risk of developing diabetes and obesity.

To our knowledge this is the first human study evaluating metabolic parameters after perinatal and current exposure in puberty to dioxins. In addition, effects of the current exposure to PBDEs were studied.

## Methods and materials

### Study population

In 1987 in the Amsterdam-Zaandam region a longitudinal cohort study on the effects of background exposure to dioxins was started. Selection criteria of the cohort were an optimal pregnancy, birth weight above 2500 grams, gestational age between 37 and 41 weeks and all the children were to be breast fed. This study is part of the longitudinal cohort study of 14–19 year old children, studied previously during their neonatal ( $n = 60$ ) [30], toddler ( $n = 31$ ) [31] and pre-pubertal period ( $n = 41$ ) [32]. In the current (pubertal) follow-up all 33 children (18 girls and 15 boys) participating were born in the Amsterdam/Zaandam region and 25 of them are still inhabitants of this region. Prenatal PCDD/F exposure and lactational PCDD/F intake (together the perinatal exposure) were determined in breast milk soon after birth [33].

The participants and their parents were interviewed to obtain data on individual characteristics including residential histories, social economic status, smoking habits, past history of diseases and treatments as well as present current illnesses, medication usage and allergies. A physical examination by one and the same physician took place to assess the height and weight

as well as the pubertal development [9]. Body mass index was calculated as weight (in kilograms) divided by height (in meters) squared.

Of the total cohort of 41 subjects who participated in the pre-pubertal study, one subject was excluded from the current follow-up because of an Ewing sarcoma, one was partly excluded because of an extra Y-chromosome, and one had passed away due to leukaemia. Five subjects declined to participate in the follow-up, two could not be traced. One of the children, who did not participate in the pre-pubertal follow-up, consented to the current follow up. Of the 33 examined adolescents, 3 refused blood sampling.

The study was approved by the Medical Ethics Committee of the Academic Medical Centre, Meibergdreef 9, Amsterdam, and was conducted according to the principles expressed in the Declaration of Helsinki, and its amendments. Following verbal consent, written informed consent was obtained from all the children and their parents, prior to inclusion in the previous and current study.

## Laboratory analyses

For measuring fasting glucose, HbA1c, insulin, leptin, and PCDD/Fs, dioxin-like-PCBs (dl-PCB) and PBDE serum concentrations, 30 subjects underwent vena puncture in 2005/2006, following overnight fasting for at least six hours. Serum was obtained and stored at -20° C until analysis.

The serum leptin concentrations were analysed using radioimmunoassay (RIA) from Millipore in the department for clinical chemistry in the VUMC in Amsterdam. Insulin samples were measured using the Immunoradiometric assay (IRMA) technique from Biosense in the Algemeen Medisch Laboratorium (AML) in Antwerpen. Glucose was determined using the hexokinase method and HbA1c using HPLC in the department for clinical chemistry in the Zaans Medisch Centrum in Zaandam. Perinatal PCDD/F levels and recent serum levels of PCDD/Fs, dl-PCBs and PBDEs were measured in an uncontaminated laboratory dedicated to low-level dioxin sample treatment, at the Environmental Chemistry Section of IBED/ESS of the University of Amsterdam. Levels of the 19 most toxic PCDD/F congeners (seven PCDDs and twelve PCDFs) and 3 dl-PCBs (77, 126, 169) as well as 8 PBDEs (28, 47, 85, 99, 100, 153, 154 and 183) were determined. The perinatal dioxin levels are expressed as International Toxic Equivalents (I-TEQ). The concentration of PCDD/F and dl-PCB congeners are expressed in TEQ pg/g lipid using WHO-TEF values [34].

For group separation of the compounds we used an activated carbon column (Carbo-sphere). The PCDD/F and dl-PCB fraction was isolated and a clean-up was performed using a column of AgNO<sub>3</sub>-impregnated silica gel and a column of activated Al<sub>2</sub>O<sub>3</sub> on top of silica gel. The PBDE fraction was purified using activated Al<sub>2</sub>O<sub>3</sub> on silica gel and an activated alumina column. After concentrating the sample, quantification of dioxins and dl-PCBs was performed using HR-GC/HR-MS. PBDEs were determined using HR-GC/LR-MS. As an internal standard, a mixture of <sup>13</sup>C-labelled PCDD/Fs, dl-PCBs and PBDEs was used. More detailed information about the analysis has been published previously [4].

PCDD/F concentrations were previously determined in mothers' milk 3–4 weeks after birth, which is indicative of the *prenatal* exposure. The cumulative total *postnatal*/lactational intake was calculated as the measured PCDD/F concentration in breast milk multiplied by the total breast milk intake [33].

## Statistical analyses

For statistical analyses, simple linear regression, and Spearman's correlation coefficient (when the correlation was not typically linear in the scatter diagram or when there were outliers) was used using SPSS®.

As dependent values we used serum glucose, serum insulin, glucose:insulin ratio, leptin, BMI and BMI:leptin ratio and HbA1c. The prenatal, lactational, serum PCDD/Fs and the serum dl-PCBs, the total dioxin WHO-TEQ (PCDD/F+dl-PCB) and PBDE concentrations were the independent values using linear regression and Spearman’s correlation coefficient.

To quantify the relationship of two variables, for linear regression the regression coefficient B was used and for Spearman’s rho the Spearman’s correlation coefficient. To adjust for possible confounding factors the partial correlation coefficient was used. The level of significance was 5% (p = 0.05).

**Confounders.** Evaluation of variables or confounding factors (age, sex and BMI) was performed.

Gender, age and BMI were considered possible confounders. Due to the limited size of the cohort, ANOVA was not viable. We therefore assessed the dependency according to the formula:

$$y = N0 + N1x + N2z + N3xz$$

whereby x is the independent, y is the dependent and z the possible confounder.

Gender versus insulin (p = 0.815), versus glucose (p = 0.133) and versus the glucose:insulin ratio (p = 0.460) were not significant. In other words, gender was not a confounder in the outcomes. Similarly, age (p = 0.063, p = 0.054 and p = 0.336, respectively) and BMI (p = 0.640, p = 0.035 and p = 0.066, respectively) were analysed as possible confounders.

## Results

Descriptive statistics of the cohort (age, BMI, prenatal dioxin exposure, lactational dioxin intake and, current serum dioxin, dl-PCB and PBDE levels) are presented in Table 1. The measured serum dioxin (PCDD/Fs) levels in the adolescents (current serum levels) were much lower than the prenatal exposure (levels in the mothers). There was no significant correlation between the current serum levels and the prenatal exposure levels nor with the lactational intake. Descriptive statistics of the measured metabolic parameters are given in Table 2. A summary of the results of the statistical analyses is given in Table 3.

All clinical laboratory measurements in the adolescents were within normal limits. Our population of 33 adolescents had heights, weights and BMIs within the normal range for Dutch adolescents. The mean BMI was 21 kg/m<sup>2</sup>. There were no children with obesity, but 3 children (9%) were overweight (BMI 25–29.9). Life style (diets, consumption of alcohol, cigarettes) as well as socio economic status had no influence on the dioxin, dl-PCB or PBDE levels. None of the participants had a known exposure to other toxicants.

**Table 1. Descriptive statistics of the cohort and the cohort’s exposure to PCDD/Fs, dl- PCBs and PBDEs.**

N = 33	Median	Mean	Range	95 <sup>th</sup> —percentile
Age (years)	14.3	15.0	14.0–18.7	18.5
Prenatal PCDD/F exposure I-TEQ (pg/g lipid)	29.8	32.6	9.05–88.8	74.8
LactationalPCDD/Fexposure I-TEQ (ng)	45.9	66.9	4.34–279	239.1
BMI (kg/m <sup>2</sup> )	19.7	21.0	17.4–30.9	29.9
Current serum PCDD/F WHO I-TEQ (pg/g lipid)	1.6	2.2	0.4–6.1	6.1
Current serum dl-PCBs WHO I-TEQ (pg/g lipid)	1.8	2.2	0.04–7.8	7.3
Current serum PBDEs(ng/g lipid)	9.9	14	4.9–73.6	22.1

<https://doi.org/10.1371/journal.pone.0184006.t001>



**Table 2. Descriptive statistics of the cohort's measured metabolic parameters.**

N = 33	Median	Mean	Range	Standard Deviation
HbA1c (%)	5.6	5.5	5–6.1	0.24
Leptin (µg/l)	5.3	6.9	0.5–27.6	7.1
Glucose (mmol/l)	4.0	4.0	3.2–5.0	0.4
Insulin (mU/l)	9.2	9.7	4.4–16.1	3.2

<https://doi.org/10.1371/journal.pone.0184006.t002>

### Glucose concentrations, HbA1c levels

Serum samples were taken to measure the fasting glucose in the adolescents. After statistical analysis using linear regression we found a significant correlation of glucose levels with current serum PCDD/Fs ( $p = 0.015$ , correlation coefficient ( $r$ ) = 0.46, std. error = 0.040, see Fig 1). Neither gender, nor age or BMI were identified as confounding factors in this case.

We also found a significant positive correlation between the current total dioxin WHO-TEQ (PCDD/Fs+dl-PCBs) serum levels and fasting glucose concentrations ( $p = 0.037$ , correlation coefficient ( $r$ ) = 0.43, std. error = 0.028). No correlation between dl-PCBs ( $p = 0.994$ ), PBDEs ( $p = 0.564$ ), lactational intake ( $p = 0.287$ ) or prenatal ( $p = 0.631$ ) PCDD/F exposures and fasting glucose levels were observed. Similarly, we found no correlation between HbA1c levels and PCDD/F, dl-PCB or PBDE serum levels, prenatal PCDD/F concentrations or lactational intake. HbA1c gives an indication of the average serum glucose concentration of the previous 2–3 months.

### Insulin concentration and glucose:Insulin ratio

We observed a significant negative correlation between insulin levels and prenatal PCDD/F exposure ( $p = 0.017$ , correlation coefficient ( $r$ ) = - 0.44, std. error = 0.031) (Fig 2), as well as with postnatal lactational intake ( $p = 0.028$ ,  $r = -0.41$ , std. error = 0.008). For the serum PCDD/F ( $p = 0.114$ ) serum dl-PCBs ( $p = 0.916$ ), and total WHO-TEQ in serum ( $p = 0.592$ ), no correlation was seen with insulin concentrations. Age, sex nor BMI were identified as confounding factors in this model.

For the glucose:insulin ratio a significant positive correlation was seen with the prenatal PCDD/F exposure using linear regression ( $p = 0.024$ ,  $r = 0.42$ , std. error = 0.002) (Fig 3). We found no correlation with postnatal lactational PCDD/F intake ( $p = 0.095$ ) or current serum PCDD/F ( $p = 0.983$ ), dl-PCB ( $p = 0.811$ ) and PBDE ( $p = 0.633$ ) levels.

### Body mass index

No significant correlation was observed between the current BMI and the prenatal exposure ( $p = 0.920$ ) or lactational intake ( $p = 0.177$ ) of dioxins, nor with the serum dioxin ( $p = 0.951$ ), dl-PCB ( $p = 0.788$ ) and PBDE ( $p = 0.735$ ) levels. After separation of the group for boys and girls, no significant correlation was found for either group.

We found no correlation between the age of the group ( $p = 0.772$ ) and the BMI.

### Leptin concentration and BMI:Leptin ratio

A clear positive correlation between BMI and serum leptin was found ( $p < 0.001$ ) using Pearson's correlation coefficient. As expected, a positive correlation between leptin and insulin was also seen ( $p = 0.034$ ). Using the partial correlation coefficient showed that gender and age did not influence these correlations.


**Table 3. Summary of the statistical significance (p-values) of relationships between exposure and metabolic parameters of the cohort calculated with linear regression/multiple regression and Spearman's 2-tailed correlation.**

	Prenatal PCDD/F exposure I-TEQ (pg/g lipid)			Postnatal Lactational PCDD/F intake I-TEQ (ng)			Current total dioxins in serum WHO I-TEQ(PCDD/F +dl-PCB) (pg/g lipid)			Current serum PCDD/F WHO I-TEQ (pg/g lipid)			Current serum dl-PCBs WHO I-TEQ (pg/g lipid)			Current serum PBDEs (ng/g lipid)		
	Slope (neg/pos)	Linear regression p-value	Spearman 2-tailed correlation	Slope (neg/pos)	Linear regression p-value	Spearman 2-tailed correlation	Slope (neg/pos)	Linear regression p-value	Spearman 2-tailed correlation	Slope (neg/pos)	Linear regression p-value	Spearman 2-tailed correlation	Slope (neg/pos)	Linear regression p-value	Spearman 2-tailed correlation	Slope (neg/pos)	Linear regression p-value	Spearman 2-tailed correlation
Glucose	+	0.631	0.631	+	0.278	0.287	+	0.037 <sup>a</sup>	0.037 <sup>a</sup>	+	0.015 <sup>a</sup>	0.015 <sup>a</sup>	+	0.994	0.994	+	0.564	0.564
Insulin	-	0.017 <sup>a</sup>	0.017 <sup>a</sup>	-	0.028 <sup>a</sup>	0.028 <sup>a</sup>	-	0.592	0.592	-	0.114	0.114	-	0.916	0.916	-	0.494	0.494
Glucose: insulin ratio	+	0.024 <sup>a</sup>	0.023 <sup>a</sup>	+	0.095	0.095	+	0.577	0.588	+	0.983	0.998	+	0.811	0.819	+	0.633	0.640
Leptine (lin. regression)	-	0.138	0.138	-	0.228	0.228	-	0.337	0.337	-	0.139	0.139	-	0.630	0.630	-	0.584	0.584
BMI	-	0.920	0.650	+	0.177	0.983	+	0.916	0.916	+	0.951	0.914	-	0.788	0.474	-	0.735	0.852
HbA1c	+	0.427	0.427	-	0.857	0.857	+	0.577	0.916	+	0.426	0.426	-	0.185	0.185	-	0.471	0.471
Age	+	Not relevant	Not relevant	+	Not relevant	Not relevant	-	0.191	0.191	-	0.391	0.391	-	0.354	0.354	+	0.838	0.838

<sup>a</sup> values correspond to statistically significant correlations

<https://doi.org/10.1371/journal.pone.0184006.t003>





**Fig 1. Current fasting glucose and current serum PCDD/Fs.** ( $p = 0.015$  correlation coefficient  $r = 0.46$ , std. error = 0.040).

<https://doi.org/10.1371/journal.pone.0184006.g001>


We found no correlation between leptin and the measured compounds, statistical information is found in [Table 3](#).

The statistical analysis of leptin concentrations in relation to the environmental pollutants was strongly influenced by the confounding factor gender. Analysis stratified by gender showed no significant correlations with the measured compounds ( $p = 0.979$  for boys and  $p = 0.230$  for girls). Similarly, no significant correlations were found for the BMI:leptin ratio and the measured compounds when using gender stratified analysis.

## Discussion

### Glucose metabolism

The main finding of this study is a negative correlation between (perinatal) dioxin exposure on glucose metabolism in adolescence. Lower insulin secretion in adolescence was seen in association with both prenatal exposure and lactational dioxin intake. The glucose:insulin ratio is



**Fig 2. Fasting insulin levels and prenatal PCDD/F exposure.** ( $p = 0.017$ , correlation coefficient  $r = -0.44$ , std. error = 0.031).

<https://doi.org/10.1371/journal.pone.0184006.g002>

positively associated with prenatal dioxin exposure. The very low serum PCDD/F levels (mean 2.2 pg/g fat) and total WHO- TEQ serum levels (PCDD/F+dl-PCB levels (mean 4.4 pg/g fat)) were positively correlated with serum fasting glucose levels.


We found no effects on serum glucose, insulin or the glucose:insulin ratio with serum PBDE levels in the adolescents.

Effects however, were seen in an animal study. One study found a higher glucose:insulin ratio in PBDE (DE-71) treated rats compared to controls. Fasting plasma glucose, insulin, and C-peptide levels were not markedly affected in this study [35]

We have to bear in mind that PBDE congeners have a much shorter half-life than dioxins. This means that a single measurement of serum levels might be insufficient to give an indication of long term exposure to these compounds.

Our results suggest an influence of dioxins in the perinatal period on pancreas development.

The significant correlation between current serum dioxin exposure and fasting serum glucose, confirms findings seen in a study with higher dioxin exposure [36], and a study of Agent Orange veterans exposed to the TCDD-containing defoliant [37], as well as the Seveso study



**Fig 3. Glucose:insulin ratio and prenatal PCDD/F exposure.** ( $p = 0.024$ , correlation coefficient  $r = 0.42$ , std. error = 0.002).

<https://doi.org/10.1371/journal.pone.0184006.g003>

[21]. Lower insulin levels were also found in a study of Faroese residents in relation to PCBs [38]. These findings in adult populations are also seen in our adolescent cohort, with the current very low background exposure. However contrasting studies have also been published. In a study on rat insulin-secreting beta cell lines, higher levels of insulin were observed after stimulation with TCDD. In this study a continuous insulin secretion was observed by stimulation of exocytosis for secretory vesicles containing insulin followed by beta cell exhaustion [39]. In a mice study, chronic exposure to Aroclor 1254 (a mixture of PCB congeners) induced hyperinsulinemia with an elevation of glucose and glucagon levels. Hereby it was observed, that Aroclor 1254 inhibited the expression of the insulin receptor signaling cascade [40].

In a cross-sectional study on Danish children, a reduced serum insulin was seen with higher PCB levels. However, in the study no separate statistical analyses were made on dl- and non-dl-PCB congeners, though different congeners may exert different effects. No effect on fasting glucose was seen in the study [41].

We found no correlation between HbA1c levels and PCDD/Fs, dl/PCBs or PBDEs. This could be due to the limited number of studied subjects or the low exposure status of the cohort. One cross-sectional study with 1374 (15–73 y) participants (mean total TEQ 24.08) in Japan found a correlation between HbA1c and the total TEQ of PCDD/Fs and dl-PCBs [23].

Direct effects of dioxins on insulin secretion have been described in animal studies. The disruptive effect of dioxins on beta cell functioning and lower insulin secretion of pancreatic islets was seen in a rat study [42]. In a study in mice, the insulin secretion was found to be significantly decreased by TCDD exposure, probably via the AhR signaling pathway [43].

Another example of disturbances of the energy homeostasis after dioxin exposure is the commonly seen wasting syndrome. This is known as the acute state of (permanent) reduced food intake and wasting after acute dioxin poisoning. Although the exact mechanism of this syndrome is unknown, there are signs of derangement of the energy balance [44].

Another explanation for our findings, the prenatal and lactational exposure to dioxins and later effects on insulin secretion might be caused by epigenetic changes. There is empirical support that early-life environmental conditions can cause epigenetic changes that persist throughout life [17].

Dioxins have been shown to exert epigenetic effects in sperm cells [45;46].

Possible epigenetic changes could involve the SOX9 gene [47]. Its homolog in zebra fish is downregulated by dioxins. [48]. Both SOX 9 and PDX1 genes, important for the development of the beta cells, are two of the earliest genes expressed in pancreatic tissue [49]. An association between increased methylation of the promoter of the RXRA gene in the cord blood of children, and later obesity was seen in 2 cohorts in the UK [50].

Theoretically, the perinatal dioxin exposure, that was quite high in Western Europe when our cohort was born (1987–1991), could have left modifications in the form of epigenetic changes resulting in effects on glucose metabolism in later life.

## BMI and leptin

No correlation was seen between BMI and the perinatal dioxin exposure, including after correction for confounding factors (age, gender). No correction was made for the confounding factor parental BMI, however the mothers' BMI was divided equally in two different groups and there were no obese women who participated the study [51]. It must be borne in mind that the number of subjects in the study is limited, and the children at the time of blood sampling, were still in their puberty, a period with changing BMIs. The adolescents now studied had no history of severe illnesses, and no abnormal weight gain was observed.

We could find one animal TCDD study that reports on serum concentrations of leptin, glucose, insulin and triglyceride in rats. In this study, similarly to our study, no effects on leptin concentrations were observed [44]. In a recent Japanese study, serum levels of leptin were significantly lower in the highly (PCB/PCDF) exposed Yusho victims [52].

A limitation of our present study is the rather small number of participants. Furthermore, the fact that since only breast fed participants were included, we cannot exclude modification by breast feeding. The HOMA-IR and HOMA- $\beta$  was not calculated in this study. The effect of exposure to other compounds, like DDT (dichlorodiphenyltrichloroethane), hexachlorobenzene, non-dioxin-like PCBs, PBB (polybrominated biphenyls), perfluorooctane sulfonate, phthalates and bisphenol-A cannot be excluded.

Further research on the possible mechanism and interaction of dioxins with the developing pancreas and effects at older age is warranted.

## Conclusion

Effects seen in this prospective long-term study of mother-baby pairs, selected on the basis of an optimal pregnancy, delivery and birth weight, indicate the influence of prenatal dioxin exposure on the metabolic parameter insulin secretion in later life. Since the levels of dioxin exposure are so-called background levels in Western Europe in the period 1987–1991, these

findings warrant concern. Further follow-up needs to elucidate if these disturbances predict pathology, such as diabetes mellitus, in later life.

## Supporting information

**S1 File. Dataset SPSS.**  
(SAV)

## Acknowledgments

We are grateful to the late Dr. Hans Oosting, for statistical support. We are indebted to Matthijs Westra, paediatrician, for facilitating performing the study in the Zaans Medical Centre, Dr. Richard Wang, of The Centers for Disease and Control, Atlanta, USA, Dr. Joantine van Esterik, University of Utrecht, Faculty of Veterinary Medicine, Utrecht, Dr. Rebecca Simmons, neonatologist Perelman School of Medicine, University of Pennsylvania, USA, Dr. Eric Fliers, neuro-endocrinologist and Dr. Elisabeth Mathus-Vliegen, gastro-enterologist of the AMC, University of Amsterdam, Dr. Richard Peterson, University of Wisconsin, and the OBELIX consortium for their advice and critique.

## Author Contributions

**Conceptualization:** Marike M. Leijs, Janna G. Koppe, Thomas Vulmsma, Kees Olie, Wim M. C. van Aalderen, Pim de Voogt, Gavin W. ten Tusscher.

**Data curation:** Marike M. Leijs, Janna G. Koppe, Gavin W. ten Tusscher.

**Formal analysis:** Marike M. Leijs, Gavin W. ten Tusscher.

**Funding acquisition:** Janna G. Koppe.

**Investigation:** Marike M. Leijs, Janna G. Koppe, Kees Olie, Gavin W. ten Tusscher.

**Methodology:** Marike M. Leijs, Janna G. Koppe, Thomas Vulmsma, Kees Olie, Wim M. C. van Aalderen, Pim de Voogt, Gavin W. ten Tusscher.

**Project administration:** Marike M. Leijs, Janna G. Koppe, Gavin W. ten Tusscher.

**Resources:** Janna G. Koppe.

**Supervision:** Janna G. Koppe, Thomas Vulmsma, Kees Olie, Wim M. C. van Aalderen, Pim de Voogt, Gavin W. ten Tusscher.

**Validation:** Marike M. Leijs, Janna G. Koppe, Gavin W. ten Tusscher.

**Writing – original draft:** Marike M. Leijs.

**Writing – review & editing:** Marike M. Leijs, Janna G. Koppe, Thomas Vulmsma, Kees Olie, Wim M. C. van Aalderen, Pim de Voogt, Juliette Legler, Gavin W. ten Tusscher.

## References

1. Liem AK, Furst P, Rappe C. Exposure of populations to dioxins and related compounds. *Food Addit Contam* 2000 Apr; 17(4):241–59. <https://doi.org/10.1080/026520300283324> PMID: 10912239
2. Olie K, Vermeulen PL, Hutzinger O. Chloro-dibenzo-p-dioxins and chlorodibenzofurans are trace components of fly ash and flue gas of some municipal incinerators in the Netherlands. *Chemosphere* 1977 Jan 1; 6:455–9.
3. de Voogt P, Brinkman UAT. Production, properties and usage of polychlorinated biphenyls. In: Kimbrough RD, Jensen AA, editors. *Halogenated biphenyls, terphenyls, naphthalenes, dibenzodioxins and related products*. Second ed. Amsterdam: Elsevier; 1989. p. 3–45.

4. Leijs MM, Teunenbroek TV, Olie K, Koppe JG, Tusscher GW, Aalderen WM, et al. Assessment of current serum levels of PCDD/Fs, dl-PCBs and PBDEs in a Dutch cohort with known perinatal PCDD/F exposure. *Chemosphere* 2008; 73:176–81. <https://doi.org/10.1016/j.chemosphere.2008.05.056> PMID: [18599107](https://pubmed.ncbi.nlm.nih.gov/18599107/)
5. Schmidt JV, Bradfield CA. Ah receptor signaling pathways. *Annu Rev Cell Dev Biol* 1996; 12:55–89. <https://doi.org/10.1146/annurev.cellbio.12.1.55> PMID: [8970722](https://pubmed.ncbi.nlm.nih.gov/8970722/)
6. Mimura J, Yamashita K, Nakamura K, Morita M, Takagi TN, Nakao K. Loss of teratogenic response to 2,3,7,8-tetrachlorodibenzo-p-dioxin and furans. *Gene Cells* 1997; 2:645–54.
7. Matsumura F. The significance of the nongenomic pathway in mediating inflammatory signaling of the dioxin-activated Ah receptor to cause toxic effects. *Biochem Pharmacol* 2009; 77:608–26. <https://doi.org/10.1016/j.bcp.2008.10.013> PMID: [18996358](https://pubmed.ncbi.nlm.nih.gov/18996358/)
8. Yoshioka W, Higashiyama W, Tohyama C. Involvement of MicroRNAs in Dioxin-induced Liver Damage in the Mouse. *Toxicol Sci* 2011; 122(2):457–65. <https://doi.org/10.1093/toxsci/kfr130> PMID: [21602190](https://pubmed.ncbi.nlm.nih.gov/21602190/)
9. Leijs MM, Koppe JG, Olie K, van Aalderen WM, Voogt P, Vulsma T, et al. Delayed initiation of breast development in girls with higher prenatal dioxin exposure; a longitudinal cohort study. *Chemosphere* 2008 Oct; 73(6):999–1004. <https://doi.org/10.1016/j.chemosphere.2008.05.053> PMID: [18707752](https://pubmed.ncbi.nlm.nih.gov/18707752/)
10. Leijs M, van der Linden L, Koppe JG, Olie K, van Aalderen W, ten Tusscher GW. The influence of perinatal and current dioxin and PCB exposure on reproductive parameters (sex-ratio, menstrual cycle characteristics, endometriosis, semen quality, and prematurity): a review. *Biomonitoring* 2014; 1(1):1–15.
11. Schoeters G, Den Hond E., Dhooge W, van Larebeke N., Leijs M. Endocrine disruptors and abnormalities of pubertal development. *Basic Clin Pharmacol Toxicol* 2008 Feb; 102(2):168–75. <https://doi.org/10.1111/j.1742-7843.2007.00180.x> PMID: [18226071](https://pubmed.ncbi.nlm.nih.gov/18226071/)
12. Damerud PO, Eriksen GS, Johannessen T, Larsen PB, Viluksela M. Polybrominated diphenylethers: occurrence, dietary exposure, and toxicology. *Environ Health Perspect* 2001; 109 Suppl. 1:49–68.
13. Koizumi A, Yoshinaga T, Harada K, Inoue K, Morikawa A, Muroi J, et al. Assessment of human exposure to polychlorinated biphenyls and polybrominated diphenyl ethers in Japan using archived samples from the early 1980s and mid-1990s. *Environ Res* 2005 Sep; 99(1):31–9. <https://doi.org/10.1016/j.envres.2004.12.002> PMID: [16053925](https://pubmed.ncbi.nlm.nih.gov/16053925/)
14. Stapleton HM, Eagle S, Sjodin A, Webster TF. Serum PBDEs in a North Carolina Toddler Cohort: Associations with Wipes, House Dust and Socioeconomic Variables. *Environ Health Perspect* 2012; 120:1049–54. <https://doi.org/10.1289/ehp.1104802> PMID: [22763040](https://pubmed.ncbi.nlm.nih.gov/22763040/)
15. Ravelli ACI, van der Meulen JH, Osmond C., Barker DJP, Bleker OP. Obesity after prenatal exposure to famine in men and women at the age of 50. *Int J Obesity* 1998; 22:S18.
16. Jones OAH, Maguire ML, Griffin JL. Environmental pollution and diabetes: a neglected association. *Lancet* 2008; 371:287–8. [https://doi.org/10.1016/S0140-6736\(08\)60147-6](https://doi.org/10.1016/S0140-6736(08)60147-6) PMID: [18294985](https://pubmed.ncbi.nlm.nih.gov/18294985/)
17. Heijmans BT, Tobi EW, Stein AD, Putter H, Blauw GJ, Susser ES. Persistent epigenetic differences associated with prenatal exposure to famine in humans. *Proc Natl Acad Sci U S A* 2008; 105:17046–9. <https://doi.org/10.1073/pnas.0806560105> PMID: [18955703](https://pubmed.ncbi.nlm.nih.gov/18955703/)
18. Michalek JE, Ketchum NS, Tripathi RC. Diabetes mellitus and 2,3,7,8-tetrachlorodibenzo-p-dioxin elimination in veterans of Operation Ranch Hand. *J Toxicol Environ Health A* 2003 Feb 14; 66(3):211–21. <https://doi.org/10.1080/15287390306373> PMID: [12521668](https://pubmed.ncbi.nlm.nih.gov/12521668/)
19. Michalek JE, Pavuk M. Diabetes and cancer in veterans of Operation Ranch Hand after adjustment for calendar period, days of spraying, and time spent in Southeast Asia. *J Occup Environ Med* 2008 Mar; 50(3):330–40. <https://doi.org/10.1097/JOM.0b013e31815f889b> PMID: [18332783](https://pubmed.ncbi.nlm.nih.gov/18332783/)
20. Wang SL, Tsai PC, Yang CY, Leon GY. Increased risk of diabetes and polychlorinated biphenyls and dioxins: a 24-year follow-up study of the Yucheng cohort. *Diabetes Care* 2008 Aug; 31(8):1574–9. <https://doi.org/10.2337/dc07-2449> PMID: [18487481](https://pubmed.ncbi.nlm.nih.gov/18487481/)
21. Bertazzi PA, Consonni D, Bachetti S, Rubagotti M, Baccarelli A, Zocchetti C. Health effects of dioxin exposure: a 20-year mortality study. *Am J Epidemiol* 2001; 153:1031–44. PMID: [11390319](https://pubmed.ncbi.nlm.nih.gov/11390319/)
22. Esser A, Schettgen T, Gube M, Koch J, Kraus T. Association between polychlorinated biphenyls and diabetes mellitus in the German HELPCB cohort. *Int J Hyg Environ Health* 2016; 219(6):557–65. <https://doi.org/10.1016/j.ijheh.2016.06.001> PMID: [27397874](https://pubmed.ncbi.nlm.nih.gov/27397874/)
23. Uemura H, Arisawa K, Hiyoshi M, Satoh H, Sumiyoshi Y, Morinaga K, et al. Associations of environmental exposure to dioxins with prevalent diabetes among general inhabitants in Japan. *Environ Res* 2008 Sep; 108(1):63–8. <https://doi.org/10.1016/j.envres.2008.06.002> PMID: [18649880](https://pubmed.ncbi.nlm.nih.gov/18649880/)
24. Lee DH, Lee IK, Song K, Steffes M, Toscano W, Baker BA, et al. A strong dose-response relation between serum concentrations of persistent organic pollutants and diabetes: results from the National



- Health and Examination Survey 1999–2002. *Diabetes Care* 2006 Jul; 29(7):1638–44. <https://doi.org/10.2337/dc06-0543> PMID: 16801591
25. Fierens S, Mairesse H, Heilier JF, De Burbure C., Focant JF, Eppe G, et al. Dioxin/polychlorinated biphenyl body burden, diabetes and endometriosis: findings in a population-based study in Belgium. *Biomarkers* 2003 Nov; 8(6):529–34. <https://doi.org/10.1080/1354750032000158420> PMID: 15195683
  26. ten Tusscher GW, Koppe JG. Perinatal dioxin exposure and later effects—a review. *Chemosphere* 2004 Mar; 54(9):1329–36. [https://doi.org/10.1016/S0045-6535\(03\)00254-6](https://doi.org/10.1016/S0045-6535(03)00254-6) PMID: 14659426
  27. ten Tusscher GW, Pluim HJ, Olie K., Koppe JG. Clinical laboratory manifestations of perinatal exposure to background levels of dioxins in Dutch children. *Organohalogen Compounds* 1999; 42:213.
  28. Bastos SL, Kamstra JH, Cenijn PH, van Rijt LS, Hamers T, Legler J. Effects of endocrine disrupting chemicals on in vitro global DNA methylation and adipocyte differentiation. *Toxicol In Vitro* 2013 Sep; 27(6):1634–43. <https://doi.org/10.1016/j.tiv.2013.04.005> PMID: 23603478
  29. Bouret SG, Simerly RB. Leptin and development of hypothalamic feeding circuits. Minireview. *Endocrinology* 2004; 145(6):2621–6. <https://doi.org/10.1210/en.2004-0231> PMID: 15044371
  30. Pluim HJ. pre- and postnatal exposure in the human newborn University of Amsterdam; 2003.
  31. Ilsen A, Briet JM, Koppe JG, Pluim HJ, Oosting J. Signs of enhanced neuromotor maturation in children due to perinatal load with background levels of dioxins. Follow-up until age 2 years and 7 months. *Chemosphere* 1996 Oct; 33(7):1317–26. PMID: 8799996
  32. ten Tusscher GW. Later childhood effects of perinatal exposure to background levels of dioxins in the Netherlands. Ph.D Thesis, University of Amsterdam. 2002.
  33. Pluim HJ, Koppe JG, Olie K, van der Slikke JW, Slot PC, van Boxtel CJ. Clinical laboratory manifestations of exposure to background levels of dioxins in the perinatal period. *Acta Paediatr* 1994 Jun; 83(6):583–7. PMID: 7919753
  34. van den Berg M., Birnbaum LS, Denison M, De VM, Farland W, Feeley M, et al. The 2005 World Health Organization reevaluation of human and Mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 2006 Oct; 93(2):223–41. <https://doi.org/10.1093/toxsci/kfl055> PMID: 16829543
  35. Nash J, Szabo D, Carey G. Brominated diphenyl ethers alter hepatic phosphoenolpyruvate carboxylase enzyme kinetics in male Wistar rats: implications for lipid and glucose metabolism. *J Toxicol Environ Health A*. 2013; 76(2):142–56 <https://doi.org/10.1080/15287394.2012.738457> PMID: 23294302
  36. Remillard RB, Bunce NJ. Linking dioxins to diabetes: epidemiology and biologic plausibility. *Environ Health Perspect* 2002 Sep; 110(9):853–8. PMID: 12204817
  37. Michalek J, Longnecker MP. Veterans and Agent Orange: Herbicide/Dioxin exposure and type 2 diabetes. National Academy Press Washington D.C; 2000.
  38. Grandjean P, Henriksen JE, Choi AL, Petersen MS, Dalgard C, Nielsen F, et al. Marine food pollutants as a risk factor for hypoinsulinemia and type 2 diabetes. *Epidemiology* 2011 May; 22(3):410–7. <https://doi.org/10.1097/EDE.0b013e318212fab9> PMID: 21364465
  39. Kim YH, Shim YJ, Shin YJ, Sul D, Lee E, Min BH. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) induces calcium influx through T-type calcium channel and enhances lysosomal exocytosis and insulin secretion in INS-1 cells. *Int J Toxicol* 2009 May; 28(3):151–61. <https://doi.org/10.1177/1091581809336885> PMID: 19546254
  40. Zhang S, Wu T, Chen M, Guo Z, Yang Z, Zuo Z, et al. Chronic exposure to Aroclor 1254 disrupts glucose Homeostasis in male mice via inhibition of the insulin receptor signal pathway. *Environ Sci Technol* 2015; 49(16):84–92.
  41. Jensen TK, Timmermann AG, Rossing LI, Ried-Larsen M, Grontved A, Andersen LB, et al. Polychlorinated biphenyl exposure and glucose metabolism in 9-year-old Danish children. *J Clin Endocrinol Metab* 2014 Dec; 99(12):E2643–E2651. <https://doi.org/10.1210/jc.2014-1683> PMID: 25093617
  42. Novelli M, Piaggi S, De Tata V. 2,3,7,8-Tetrachlorodibenzo-p-dioxin-induced impairment of glucose-stimulated insulin secretion in isolated rat pancreatic islets. *Toxicol Lett* 2005 Apr 10; 156(2):307–14. <https://doi.org/10.1016/j.toxlet.2004.12.004> PMID: 15737493
  43. Kurita H, Yoshioka W, Nishimura N, Kubota N, Kadowaki T, Tohyama C. Aryl hydrocarbon receptor-mediated effects of 2,3,7,8-dibenzo-p-dioxin on glucose-stimulated insulin secretion in mice. *J Appl Toxicol* 2009; 29(8):689–94. <https://doi.org/10.1002/jat.1459> PMID: 19623578
  44. Linden J, Korkalainen M, Lensu S, Tuomisto J, Pohjanvirta R. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and leptin on hypothalamic mRNA expression of factors participating in food intake regulation in a TCDD-sensitive and a TCDD-resistant rat strain. *J Biochem Mol Toxicol* 2005; 19:139–48. <https://doi.org/10.1002/jbt.20065> PMID: 15977195

45. Manikkam m, Guerrero-Bosagna C, Tracey R, Haque M, Skinner M. Transgenerational actions of environmental compounds on reproductive disease and identification of epigenetic biomarkers of ancestral exposures. *PLoS One* 2012; 7(2):1–12.
46. Pilsner J, Parker M, Sergejev O, Suvorov A. Spermatogenesis disruption by dioxins: Epigenetic reprogramming and windows of susceptibility. *Reprod Toxicol* 2017; 69:221–9. <https://doi.org/10.1016/j.reprotox.2017.03.002> PMID: [28286111](https://pubmed.ncbi.nlm.nih.gov/28286111/)
47. McDonald E, Li J, Krishnamurthy M, Fellows GF, Goodyer CG, Wang R. SOX9 regulates endocrine cell differentiation during human fetal pancreas development. *Int J Biochem Cell Biol* 2012; 44:72–83. <https://doi.org/10.1016/j.biocel.2011.09.008> PMID: [21983268](https://pubmed.ncbi.nlm.nih.gov/21983268/)
48. Villano CM, White LA. The aryl hydrocarbon receptor-signaling pathway and tissue remodeling: Insights from the zebrafish (*Danio rerio*) model system. *Toxicol Sci* 2006; 92(1):1–4. PMID: [16848026](https://pubmed.ncbi.nlm.nih.gov/16848026/)
49. Lee Young-Hoon, Saint-Jeannet JP. Sox9, a novel pancreatic marker in *Xenopus*. *Int J Dev Biol* 2003; 47:459–62. PMID: [14584783](https://pubmed.ncbi.nlm.nih.gov/14584783/)
50. Godfrey KM, Sheppard A, Gluckman PD, Lillycrop KA, Burdge GC, McLean C, et al. Epigenetic Gene Promotor Methylation at Birth is Associated With Child's Later Adiposity. *Diabetes* 2011; 60:1528–34. <https://doi.org/10.2337/db10-0979> PMID: [21471513](https://pubmed.ncbi.nlm.nih.gov/21471513/)
51. Pluim HJ, Boersma ER, Kramer I, Olie K, van der Slikke JW, Koppe JG. Influence of short-term dietary measures on dioxin concentrations in human milk. *Environ Health Perspect* 1994 Nov; 102(11):968–71. PMID: [9738212](https://pubmed.ncbi.nlm.nih.gov/9738212/)
52. Uchi H, Furue M. [Adipokine profile of Yusho patients]. *Fukuoka Igaku Zasshi* 2013 Apr; 104(4):85–7. PMID: [23858783](https://pubmed.ncbi.nlm.nih.gov/23858783/)