



UvA-DARE (Digital Academic Repository)

Asymptotic results in nonparametric Bayesian function estimation

Kirichenko, A.

Publication date

2017

Document Version

Other version

License

Other

[Link to publication](#)

Citation for published version (APA):

Kirichenko, A. (2017). *Asymptotic results in nonparametric Bayesian function estimation*.

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Asymptotic results in nonparametric Bayesian function estimation

Alisa Kirichenko

Bayesian nonparametric methods are widely used in practical applications. They have numerous attractive features such as their philosophical appeal, conceptual simplicity, and ability to easily incorporate prior knowledge about the model. However, putting a prior on a large function space might result in erroneous estimates or suboptimal performance. Therefore, it is essential to study Bayesian procedures to gain insight about which priors to use and how their tuning affects the performance of these procedures. One way to do it is to take an asymptotic approach and to analyse Bayesian methods from the frequentist point of view by assuming that there exists a true underlying function and studying how fast a particular Bayesian procedure captures the truth as the number of observations goes to infinity.

In this thesis we consider function estimation problems in two different statistical settings. First, we discuss regression and binary classification problems on large graphs, where the goal is to estimate a smooth function defined on the vertices of a graph. In the second setting we aim to estimate the intensity of an inhomogeneous Poisson process from a realised point pattern. For both problems we develop adaptive Bayesian procedures and study their asymptotic behaviour from the frequentist perspective. In particular, we derive contraction rates for our procedures and show that they are optimal in a minimax sense.

Asymptotic results in nonparametric Bayesian function estimation

Alisa Kirichenko

*Asymptotic results in nonparametric
Bayesian function estimation*

PROEFSCHRIFT

ter verkrijging van de graad van doctor
aan de Universiteit van Amsterdam
op gezag van de Rector Magnificus, prof.dr.ir. K.L.J. Maex
ten overstaan van een door het College voor Promoties ingestelde commissie,
in het openbaar te verdedigen in de Agnietenkapel
op 5 oktober 2017, te 12:00 uur

door

Alisa Kirichenko

geboren te Sint-Petersburg, Rusland

Dit proefschrift is goedgekeurd door de promotoren en de samenstelling van de promotiecommissie is als volgt:

Promotor:	prof.dr. J.H. van Zanten	Universiteit van Amsterdam
Copromotor:	dr. B.J.K. Kleijn	Universiteit van Amsterdam
Overige leden:	dr. A.J. van Es	Universiteit van Amsterdam
	prof.dr. S. Ghoshal	North Carolina State University
	prof.dr. M.C.M. de Gunst	Vrije Universiteit Amsterdam
	prof.dr. M.R.H. Mandjes	Universiteit van Amsterdam
	prof.dr. R. Nunez Queija	Universiteit van Amsterdam

Faculteit der Natuurwetenschappen, Wiskunde en Informatica

Contents

Introduction	5
1 Bayesian nonparametric statistics	11
1.1 Statistical inference	11
1.2 Nonparametric models	12
1.3 Frequentist asymptotics	13
1.4 Bayesian approach	15
1.5 Bayesian asymptotics	16
1.6 Adaptation	19
2 The problem of function estimation on large graphs	21
2.1 Introduction	21
2.2 Asymptotic geometry assumption on graphs	23
2.3 Smoothness assumption	29
2.4 Minimax rates for function estimation problems on graphs	30
2.5 Proofs	33
3 Bayesian function estimation on large graphs	43
3.1 Introduction	43
3.2 General results on prior concentration	45
3.3 Function estimation on graphs	46
3.4 Function estimation on graphs with missing observations	48
3.5 Proofs	55
4 Poisson intensity estimation	71
4.1 Introduction	71
4.2 The SGCP model	73
4.3 Main result	75
4.4 Proofs	76
References	81
Summary	87

Samenvatting	89
Author contributions	91
Acknowledgments	93

Introduction

This thesis is dedicated to the scientific field of mathematical statistics. In order to understand what kind of problems we are dealing with consider the following example. Figure 1 shows a component of the social network graph from the Framingham Heart Study conducted in 2000. The vertices of the graph represent people and the edges show ties among siblings, spouses, and friends. Vertex size is made proportional to people's body mass index (BMI), and the vertex is coloured yellow if the BMI of the corresponding person is above 30 (which indicates obesity) and green otherwise. Christakis and Fowler [2007] show that the chance of a person becoming obese increases if his graph-neighbours (siblings/spouses or friends) became obese. In real life one wouldn't have the value of BMI for every person, hence some of the vertices are going to be blank. Using statistical analysis we can try to fill in that data by predicting a person's BMI based on the geometry of the social network and the information about other people in the graph.

The problem above is an example of function estimation problem, where the function of interest is the function on the vertices of the graph describing the BMI index. Of course, there are plenty of other scenarios in which the target function can be defined on different domains and take values in different spaces. In order to be able to make predictions a statistician chooses a model that incorporates his knowledge and assumptions about the underlying function. A fundamental problem in statistics is to develop models based on a sample of observations so that further analyses can be carried out. If the set of functions in the model is not very large (can be indexed by a subset of \mathbb{R}^d) such a model is called parametric. If the collection of functions considered is infinite dimensional the model is nonparametric. When there is enough information about the underlying function, parametric models are favourable. However, such models lack flexibility and are sometimes not robust enough in the sense that even slight contamination of the data by observations not following the particular parametric model might lead to incorrect conclusions. Nonparametric models are more flexible in that sense, and more robust against misspecification.

The study of nonparametric models started in the mid twentieth century. According to Prakasa Rao [1983], the first paper in the area of nonparametric function estimation is Rosenblatt [1956]. Since then nonparametric models have greatly evolved and gained tremendous popularity. A good introduction to the topic is the