UvA-DARE (Digital Academic Repository)

Early-life stress diminishes the increase in neurogenesis after exercise in adult female mice

Abbink, M.R.; Naninck, E.F.G.; Lucassen, P.J.; Korosi, A.

DOI

[10.1002/hipo.22745](https://doi.org/10.1002/hipo.22745)

Publication date

2017

Document Version

Final published version

Published in

Hippocampus

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Abbink, M. R., Naninck, E. F. G., Lucassen, P. J., & Korosi, A. (2017). Early-life stress diminishes the increase in neurogenesis after exercise in adult female mice. *Hippocampus*, 27(8), 839-844. <https://doi.org/10.1002/hipo.22745>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

RAPID COMMUNICATION

Early-life stress diminishes the increase in neurogenesis after exercise in adult female mice

M. R. Abbink* | E. F. G. Naninck* | P. J. Lucassen | A. Korosi

Swammerdam Institute for Life Sciences,
Center for Neuroscience, Brain plasticity
group, University of Amsterdam,
Amsterdam, The Netherlands

Correspondence

E.F.G. Naninck, Swammerdam Institute for
Life Sciences, Center for Neuroscience,
University of Amsterdam, Brain plasticity
group, Science Park 904, 1098 XH
Amsterdam, The Netherlands.
Email: e.f.g.naninck@uva.nl

Funding information

NWO and ISAO, Grant/Award numbers:
NWO Meervoud 2013/15445/ALW; NWO
FCB 14-48; ISAO grant #12536. Alzheimer
Nederland WE-03-2012-41

Abstract

Exposure to early-life stress (ES) has long-lasting consequences for later cognition and hippocampal plasticity, including adult hippocampal neurogenesis (AHN), i.e., the generation of new neurons from stem/progenitor cells in the adult hippocampal dentate gyrus. We had previously demonstrated a sex-specific vulnerability to ES exposure; female mice exposed to ES from P2-P9 exhibited only very mild cognitive changes and no reductions in AHN as adult, whereas ES-exposed male mice showed impaired cognition closely associated with reductions in AHN. Given the apparent resilience of AHN to ES in females, we here questioned whether ES has also altered the capacity to respond to positive stimuli for neurogenesis. We therefore investigated whether exercise, known for its strong pro-neurogenic effects, can still stimulate AHN in adult female mice that had been earlier exposed to ES. We confirm a strong pro-neurogenic effect of exercise in the dorsal hippocampus of 8-month-old control female mice, but this positive neurogenic response is less apparent in female ES mice. These data provide novel insights in the lasting consequences of ES on hippocampal plasticity in females and also indicate that ES might lastingly reduce the responsiveness of the hippocampal stem cell pool, to exercise, in female mice.

KEYWORDS

dentate gyrus, doublecortin, Ki67, running, sex-specific

1 | INTRODUCTION

Exposure to early-life stress (ES) lastingly affects brain structure and function. ES has e.g., been associated with later hippocampal volume reductions and cognitive impairments (Maccari, Krugers, Morley-Fletcher, Szyf, & Brunton, 2014; Nemeroff, 2016; Teicher, Anderson, & Polcari, 2012). Interestingly, vulnerability to ES appears to be sex-specific and human and rodent female offspring seem often resilient to ES (Frodl, Reinhold, Koutsouleris, Reiser, & Meisenzahl, 2010; Loi et al., 2017; Oomen et al., 2009).

One form of hippocampal plasticity consistently affected by ES in male rodents, is adult hippocampal neurogenesis (AHN) (Korosi et al., 2011; Lajud & Torner, 2015; Loi, Koricka, Lucassen, & Joels, 2014). Neuronal progenitors in the dentate gyrus (DG) undergo proliferation, migrate into the granular cell layer and eventually differentiate into functional neurons. These adult-generated neurons have been implicated in various hippocampus-dependent cognitive functions (Clark

et al., 2012; Oomen, Bekinschtein, Kent, Saksida, & Bussey, 2014) and their level is strongly regulated by various factors (Lucassen et al., 2015; Schoenfeld & Gould, 2012), including stress (Schoenfeld & Gould, 2012), circulating sex hormones (Pawluski et al., 2009), and physical exercise (Duzel, van Praag, & Sendtner, 2016).

Previously we have shown that, whereas male mice exposed to chronic ES exhibited a reduced survival of adult-born neurons, associated with learning and memory impairments, female mice, exposed to the same ES, did not show such changes in AHN and exhibited less prominent cognitive deficits (Naninck et al., 2015). In male rodents, exercise has been able to rescue various measures of neuroplasticity that were altered by different types of adverse early-life events (Maniam & Morris, 2010; Naylor et al., 2008; Lajud & Torner, 2015). Despite the apparent resilience of female mice to ES, it is unknown whether the neurogenic capacity to respond to a positive stimulus is also affected in this sex. We hypothesize that in female mice, ES occurring during a period of ongoing hippocampal development, may lastingly program the stem cells or the neurogenic niche. Because exercise has strong pro-neurogenic effects, we here studied whether prolonged exercise can still stimulate AHN in female ES-exposed mice during late adulthood.

*Shared first authors.

2 | MATERIALS AND METHODS

All animal procedures were performed as previously described in (Naninck et al., 2015). In short, a total of 9 litters with six C57Bl/6J mouse pups each, including both males and females were bred in house. At postnatal day (P) 2, litters were randomly assigned to the control (CTL) or ES condition, consisting of exposure to the limited

nesting/bedding-material model from P2–9. At P21, animals were weaned and housed in groups of the same sex with 2–3 animals/cage. At 8 months of age, female mice were moved to a larger cage (type III, Technilab-BMI, Someren, the Netherlands) and housed with 2–3 animals/cage to avoid isolation stress (Leasure & Decker, 2009; Stranahan, Khalil, & Gould, 2006). For six weeks (P237–P280) the cages were equipped with either a functional or a locked running wheel

FIGURE 1.

(160 mm diameter, Ferplast, Italy) to control for cage enrichment (see Figure 1a) (Dostes et al., 2016). For the experiments described here, a total of 23 females was used (CTL-RUN $N = 6$ (from 3 litters), ES-RUN $N = 6$ (from 2 litters), CTL-SED $N = 4$ (from 2 litters), ES-SED $N = 7$ (from 4 litters)). The relatively low power of some of the experimental groups calls for caution when interpreting the data from this study. Engagement in exercise was monitored and confirmed daily by the researcher (30 minute observations/day for 6 weeks). No differences were observed in the engagement in exercise between control and ES-exposed animals. At the end of the running period (P280) animals were sacrificed. Estrous cycle stage at P280 was determined by vaginal smear cytology (Caligioni, 2009). All experimental procedures were conducted under Dutch national law and EU directives and approved by the animal welfare committee of the University of Amsterdam.

Immunocytochemistry on perfusion-fixed brain sections was performed to identify proliferating (Ki67⁺) cells and newborn neurons (Doublecortin, DCX⁺ cells) in the hippocampus according to previously described methods (Naninck et al., 2015). Antibodies used were: polyclonal rabbit-anti-Ki67 (Novocastra NCL-L-Ki67_MM1, 1:20,000), goat anti-rabbit biotinylated secondary antibody (Vector Laboratories, 1:500), polyclonal goat-anti-DCX (SantaCruz Biotechnology sc-8066; 1:800) and donkey-anti-goat biotinylated secondary antibody (Jackson Laboratories, 1:500). All Ki67⁺ and DCX⁺ immune-reactive cell counts were performed manually by means of a modified stereological procedure, using a 20x objective (200x magnification), as described in detail in (Naninck et al., 2015). Immuno-reactive cells were counted in the subgranular zone of the dentate gyrus and cell numbers are expressed unilaterally. For each hippocampus, eight coronal, 40 μ m thick sections were stereologically sampled along the rostro-caudal axis (corresponding with Bregma: -1.34 , -1.70 , -2.06 , -2.46 , -2.92 , -3.16 , -3.52 , -3.80). A distinction was made between the rostral DG (i.e., the first four sections, (Vivar, Peterson, & van Praag, 2016)) versus caudal DG (the last four sections, (Vivar et al., 2016)), and the supra- versus infrapyramidal blade. Morphological subtypes of DCX⁺ cells (i.e., proliferative, intermediate, or post mitotic stage) were analyzed as described before (Oomen et al., 2010).

Data were analyzed using SPSS 24.0 (IBM software) and Graphpad Prism 5 (Graphpad software). To compare body weight gain (BWG) between CTL and ES animals, independent t-tests were used. To compare all four groups, we used mixed models with fixed factors: condition (CTL vs. ES) and exercise (RUN vs SED). The number of newborn cells in the left and the right hippocampus were analyzed separately for each animal and nested under the factor "animal" (Aarts, Verhage, Veenvliet, Dolan, & van der Sluis, 2014). Random factors "animal," "litter" and/or "estrous cycle stage" were included only when a factor significantly influenced the outcome variable. For the final Ki67 analysis, none of the random factors were included, while for DCX-data, the effect of the factor "animal" was significant (likelihood ratio = 9.935 $p = .001$) and therefore included in the model. Post-hoc analyses were performed using LSD multiple comparison tests when appropriate.

3 | RESULTS AND DISCUSSION

We confirm that ES-exposure induced physiological signs of chronic ES in the pups, including a reduced body weight gain from P2-P9 (CTL: 3.49 ± 0.11 , ES: 2.76 ± 0.16 , independent t-test $t(21) = 3.58$, $p = .002^*$) (Naninck et al., 2015). Bodyweight was no longer different from weaning (P21) onwards (bodyweight at P21: 9.69 ± 0.19 grams in CTL vs. 9.61 ± 0.51 grams in ES, independent t-test $t(21) = 0.153$, $p = .88$).

We questioned whether ES affected neurogenic capacity in response to physical exercise and compared AHN in various anatomical and functional subregions of the dentate gyrus (i.e., rostral and caudal) between all four experimental groups. The number of proliferating (Ki67⁺) cells was reduced by ES in the rostral hippocampus ($F_{\text{condition}}(1,40) = 5.103$ $p = .03^*$; $F_{\text{running}}(1,40) = 0.999$ $p = .32$; $F_{\text{interaction}}(1,40) = 3.789$ $p = .058$, trend, Figure 1b). Because of the strong trend in the interaction, we performed exploratory *posthoc* testing to further study the relationships between condition and running. This revealed that there was no significant difference between CTL-SED and ES-SED (LSD, $p = .28$) under basal conditions, while after running, the number of Ki67⁺ cells was higher in the CTL-RUN females compared to ES-RUN (LSD, $p = .04^*$) suggesting that ES might reduce the

FIGURE 1 Early-life stress diminishes the increase in neurogenesis after exercise in adult female mice. (a) Experimental design. (b) The number of proliferating (Ki67⁺) cells in the rostral hippocampus was reduced by ES (two-way ANOVA, main effect of condition $F_{1,40} = 5.103$ $p = .03$, no effect of running $F_{1,40} = 0.999$ $p = .32$, trend towards interaction effect $F_{1,40} = 3.789$ $p = .058$). The effect of condition was mainly attributable to the difference between the running wheel exposed CTL and ES females, as exploratory post hoc testing revealed no differences in proliferating (Ki67⁺) cells between CTL-SED and ES-SED animals (post hoc comparison, LSD, $p = .28$), but showed a significant lower number of proliferating (Ki67⁺) cells in ES-RUN females compared to CTL-RUN females (post hoc comparison, LSD, $p = .04$). (c) In the caudal hippocampus, no differences in the number of proliferating (Ki67⁺) cells were observed (two-way ANOVA no effect of condition: $F_{1,40} = 0.358$ $p = .55$, no effect of running: $F_{1,40} = 0.865$ $p = .36$, no interaction effect: $F_{1,40} = 0.273$ $p = .60$) (d) Representative images of a nuclear Ki67 immunostaining, indicating proliferating cells in a CTL-RUN animal (upper panel), versus a ES-RUN animal (lower panel). (e) The number of differentiating, immature neurons (DCX⁺ cells) in the rostral hippocampus is increased by running (main effect of running $F_{1,23} = 4.236$ $p = .05$, no effect of condition $F_{1,23} = 0.082$ $p = .78$, no interaction effect $F_{1,23} = 1.775$ $p = .20$). (f) In the caudal hippocampus, no differences in the number of DCX⁺ cells were observed (two-way ANOVA, no effect of condition: $F_{1,23} = 0.125$ $p = .73$, no effect of running: $F_{1,23} = 0.315$ $p = .58$, no interaction effect: $F_{1,23} = 0.414$ $p = .53$). (g) Representative image of a cytoplasmic DCX immunostaining, indicating newborn immature neurons in a CTL-RUN animal (upper panel), versus a ES-RUN animal (lower panel). (h) Voluntary wheel running particularly increases the number of postmitotic DCX⁺ cells in the rostral hippocampus (two-way ANOVA, main effect of running $F_{1,23} = 5.758$ $p = 0.03$, no effect of condition $F_{1,23} = 0.370$ $p = .55$, no interaction effect $F_{1,23} = 2.447$ $p = .13$). * = Main effect of running, # = significant post hoc comparison [Color figure can be viewed at wileyonlinelibrary.com]

neurogenic response to physical exercise. In the *caudal* hippocampus, there were no significant differences among the groups ($F_{\text{condition}}(1,40) = 0.358$ $p = .55$; $F_{\text{running}}(1,40) = 0.865$ $p = .36$; $F_{\text{interaction}}(1,40) = 0.273$ $p = .60$, Figure 1c).

Running increased the number of young (DCX^+) neurons in the *rostral*, but not caudal hippocampus. In the rostral hippocampus, DCX^+ cell numbers were significantly increased after exercise ($F_{\text{running}}(1,23) = 4.236$ $p = .05^*$, Figure 1e), without a main effect of condition ($F_{\text{condition}}(1,23) = 0.082$ $p = .78$, $F_{\text{interaction}}(1,23) = 1.775$ $p = .20$, Figure 1e). Even though no significant interaction was observed in this study, it is interesting to note that DCX expression is modulated by ES and running in a similar fashion as Ki67^+ : Rostral-hippocampal DCX^+ cells are 39.0% higher in CTL-RUN females compared to CTL-SED females, while there is only a minor (5.4%) increase between ES-SED and ES-RUN females. In the caudal hippocampus, no differences were observed ($F_{\text{condition}}(1,23) = 0.125$ $p = .73$, $F_{\text{running}}(1,23) = 0.315$ $p = .58$, $F_{\text{interaction}}(1,23) = 0.414$ $p = .53$, Figure 1f). For neither Ki67 nor DCX were differences observed between the supra- and infra-pyramidal blade.

Interestingly, detailed analyses of the three different morphological types of DCX^+ cells (i.e., proliferative, intermediate and post-mitotic) (Figure 1h) revealed that exercise particularly enhanced the number of postmitotic DCX^+ cells in the rostral hippocampus (mixed model analysis for postmitotic cells $F_{\text{running}}(1,23) = 5.758$ $p = .03^*$, $F_{\text{condition}}(1,23) = 0.370$ $p = .55$, $F_{\text{interaction}}(1,23) = 2.447$ $p = .131$). Exercise increased the number of postmitotic neurons by 66.2% in CTL-RUN vs CTL-SED females and only by 7.9% in ES-RUN vs ES-SED females (Figure 1h). Thus, suggesting that exercise selectively enhanced DCX^+ cell numbers in control mice, but that previous exposure to ES diminished this pro-neurogenic effect.

To summarize our findings, prolonged voluntary exercise in eight-month old female mice (i) increases AHN selectively in the rostral hippocampus, and (ii) previous ES-exposure seems to reduce this exercise-induced neurogenic response. Our data further indicate that exercise in late adulthood has a survival promoting effect, specifically for the post-mitotic DCX^+ cells, an effect that is less pronounced for proliferation. The reason for the modest effect of running on the number of proliferating cells might be explained by the fact that control mice were exposed to a locked running wheel, a form of cage enrichment which alone can already increase proliferation (Dostes et al., 2016). In addition, our paradigm of 6 weeks of running might have contributed as well, as effects of exercise on cell proliferation typically peak at 3–10 days after running onset and return to baseline after 32 days of running (Kronenberg et al., 2006; Naylor, Persson, Eriksson, Jonsdottir, & Thorlin, 2005). In contrast, in line with our observations, the number of DCX^+ cells is stably increased when longer exercise periods are involved (Kronenberg et al., 2006). Finally, we housed our animals together to prevent isolation stress, which could also have interfered with the stimulatory effect of running on AHN (Leasure & Decker, 2009; Stranahan et al., 2006).

Our findings also highlight regional differences in the neurogenic response to voluntary exercise with a pronounced increase found in DCX^+ numbers in the rostral/dorsal but not caudal/ventral hippocampus. This is consistent with previous studies showing similar rostral-

specific changes in AHN upon running. Three weeks of voluntary exercise e.g., increased DCX numbers in young (8-week old) C57Bl/6 female mice to a greater extent in the rostral than in the caudal hippocampus (Bolz, Heigele, & Bischofberger, 2015), while retroviral- and BrdU-based approaches in male mice also revealed selective increases after exercise in the dorsal hippocampus (Vivar et al., 2016). These AHN effects of exercise mainly in the dorsal hippocampus suggest that exercise may preferentially benefit spatial aspects of cognition (Wu & Hen, 2014). The mechanisms underlying these topographical differences remain unclear, but differences in input, mossy cell activity and GABAergic or glutamatergic cells density likely contribute.

Up to date, little was known as to how ES exposure affects the later neurogenic response to a positive stimulus in female mice. As mentioned above, our study has a relatively low power, which calls for caution in its interpretation. Despite this limitation, our data are the first to suggest that the running-induced increase in newborn cells in the rostral hippocampus, as occurs in control animals, is less pronounced in ES animals. Intriguingly, while AHN in female mice appeared to be resilient to the negative effects of ES (Naninck et al., 2015; Loi et al., 2017), this indicates that ES-exposed females might also be less responsive to a potent positive stimulus for AHN. This concept of ES-induced unresponsiveness at the level of this structural plasticity has been suggested before (Korosi et al., 2011) and is in line with previous studies showing that learning in prenatally stressed rats could not upregulate AHN (Lemaire, Koehl, Le Moal, & Abrous, 2000). The reasons why later exercise fails to stimulate AHN in ES-exposed females remain elusive, but as the dentate gyrus still undergoes active development from P2-P9 (Navarro-Quiroga, Hernandez-Valdes, Lin, & Naegel, 2006), a substantial proportion of the developmentally born granule cells have likely been migratory and/or dividing during the period of ES, and may thus have been particularly sensitive to programming effects induced by the elevated stress hormone levels, that could have altered neurogenic properties, e.g., through epigenetic modifications (Singh-Taylor et al., 2017).

Various (growth) factors have been implicated in the pro-neurogenic effects of exercise, including catepsin B (Moon et al., 2016), BDNF (Marlatt, Potter, Lucassen, & van Praag, 2012), cytokines and various neurotransmitters, a.o. (Bolinj & Lucassen, 2015). Hence, to explain the current results, it will be interesting to study in the future whether ES differentially affects the induction of these factors by exercise. Another candidate is an altered programming of the hypothalamic-pituitary-adrenal-axis by ES. Running reduces stress (Kannangara et al., 2011) and ES-exposed male rats subjected to exercise display e.g., a lower corticosterone response (Maniam & Morris, 2010), but whether such effects also apply to female mice awaits future study.

4 | CONCLUSION

In conclusion, while seemingly resilient to the negative effects of ES, AHN in adult female ES mice is less responsive to the positive stimulus of exercise at a later age, suggesting that ES could induce lasting effects on the responsiveness and properties of the hippocampal stem

cell pool in females. These data underscore the importance of studies that assess the sex-dependent vulnerability to ES, and highlight that further research is needed to better understand the mechanisms that underlie the programming of stem cell properties, and of hippocampal plasticity in general, by ES.

ACKNOWLEDGMENTS

AK is supported by NWO (Meervoud grant). PJL is supported by Alzheimer Nederland. We thank L.P. de Vries for practical assistance.

REFERENCES

- Aarts, E., Verhage, M., Veenliet, J. V., Dolan, C. V., & van der Sluis, S. (2014). A solution to dependency: using multilevel analysis to accommodate nested data. *Nature Neuroscience*, 17, 491–496.
- Bolijn, S., & Lucassen, P. J. (2015). How the body talks to the brain; peripheral mediators of physical activity-induced proliferation in the adult hippocampus. *Brain Plasticity*, 1, 5–27.
- Bolz, L., Heigele, S., & Bischofberger, J. (2015). Running improves pattern separation during novel object recognition. *Brain Plasticity*, 1, 129–141.
- Caligioni, C. S. (2009). Assessing reproductive status/stages in mice. *Current Protocols in Neuroscience*, Appendix 4:Appendix 4I.
- Clark, P. J., Bhattacharya, T. K., Miller, D. S., Kohman, R. A., DeYoung, E. K., & Rhodes, J. S. (2012). New neurons generated from running are broadly recruited into neuronal activation associated with three different hippocampus-involved tasks. *Hippocampus*, 22, 1860–1867.
- Dostes, S., Dubreucq, S., Ladevèze, E., Marsicano, G., Abrous, D. N., Chaouloff, F., & Koehl, M. (2016). Running per se stimulates the dendritic arbor of newborn dentate granule cells in mouse hippocampus in a duration-dependent manner. *Hippocampus*, 26, 282–288.
- Duzel, E., van Praag, H., & Sendtner, M. (2016). Can physical exercise in old age improve memory and hippocampal function? *Brain*, 139, 662–673.
- Frodl, T., Reinhold, E., Koutsouleris, N., Reiser, M., & Meisenzahl, E. M. (2010). Interaction of childhood stress with hippocampus and prefrontal cortex volume reduction in major depression. *Journal of Psychiatric Research*, 44, 799–807.
- Kannangara, T. S., Lucero, M. J., Gil-Mohapel, J., Drapala, R. J., Simpson, J. M., Christie, B. R., & van Praag, H. (2011). Running reduces stress and enhances cell genesis in aged mice. *Neurobiology of Aging*, 32, 2279–2286.
- Korosi, A., Naninck, E. F. G., Oomen, C. A., Schouten, M., Krugers, H., Fitzsimons, C., & Lucassen, P. J. (2011). Early-life stress mediated modulation of adult neurogenesis and behavior. *Behavioural Brain Research*, 227, 400–409.
- Kronenberg, G., Bick-Sander, A., Bunk, E., Wolf, C., Ehninger, D., & Kempermann, G. (2006). Physical exercise prevents age-related decline in precursor cell activity in the mouse dentate gyrus. *Neurobiology of Aging*, 27, 1505–1513.
- Lajud, N., & Torner, L. (2015). Early life stress and hippocampal neurogenesis in the neonate: sexual dimorphism, long term consequences and possible mediators. *Frontiers in Molecular Neuroscience*, 8, 3.
- Leasure, J. L., & Decker, L. (2009). Social isolation prevents exercise-induced proliferation of hippocampal progenitor cells in female rats. *Hippocampus*, 19, 907–912.
- Lemaire, V., Koehl, M., Le Moal, M., & Abrous, D. N. (2000). Prenatal stress produces learning deficits associated with an inhibition of neurogenesis in the hippocampus. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 11032–11037.
- Loi, M., Koricka, S., Lucassen, P. J., & Joels, M. (2014). Age- and sex-dependent effects of early life stress on hippocampal neurogenesis. *Frontiers in Endocrinology (Lausanne)*, 5, 13.
- Loi, M., Mossink, J. C., Meerhoff, G. F., Den Blaauwen, J. L., Lucassen, P. J., & Joels, M. (2017). Effect of early-stress on cognitive function and hippocampal structure in female rodents. *Neuroscience*, 342, 101–119.
- Lucassen, P. J., Naninck, E. F. G., van Goudoever, J. B., Fitzsimons, C., Joels, M., & Korosi, A. (2013). Perinatal programming of adult hippocampal structure and function; emerging roles of stress, nutrition and epigenetics. *Trends in Neuroscience*, 36, 621–631.
- Lucassen, P. J., Oomen, C. A., Naninck, E. F. G., Fitzsimons, C. P., Van Dam, A. M., Czéh, B., & Korosi, A. (2015). Regulation of adult neurogenesis and plasticity by (early) stress, glucocorticoids, and inflammation. *Cold Spring Harbor Perspectives in Biology*, 7, a021303.
- Maccari, S., Krugers, H. J., Morley-Fletcher, S., Szyf, M., & Brunton, P. J. (2014). The consequences of early-life adversity: neurobiological, behavioural and epigenetic adaptations. *Journal of Neuroendocrinology*, 26, 707–723.
- Maniam, J., & Morris, M. J. (2010). Voluntary exercise and palatable high-fat diet both improve behavioural profile and stress responses in male rats exposed to early life stress: role of hippocampus. *Psychoneuroendocrinology*, 35, 1553–1564.
- Marlatt, M. W., Potter, M. C., Lucassen, P. J., & van Praag, H. (2012). Running throughout middle-age improves memory function, hippocampal neurogenesis and BDNF levels in female C57Bl/6J mice. *Developmental Neurobiology*, 72, 943–952.
- Moon, H. Y., Becke, A., Berron, D., Becker, B., Sah, N., Benoni, G., ... van Praag, H. (2016). Running-induced systemic cathepsin B secretion is associated with memory function. *Cell Metabolism*, 24, 332–340.
- Naninck, E. F. G., Hoeijmakers, L., Kakava-Georgiadou, N., Meesters, A., Lazic, S. E., Lucassen, P. J., & Korosi, A. (2015). Chronic early-life stress alters developmental and adult neurogenesis and impairs cognitive function in mice. *Hippocampus*, 25, 309–328.
- Navarro-Quiroga, I., Hernandez-Valdes, M., Lin, S. L., & Naegele, J. R. (2006). Postnatal cellular contributions of the hippocampus subventricular zone to the dentate gyrus, corpus callosum, fimbria, and cerebral cortex. *Journal of Comparative Neurology*, 497, 833–845.
- Naylor, A. S., Bull, C., Nilsson, M. K. L., Zhu, C., Bjork-Eriksson, T., Eriksson, P. S., ... Kuhn, H. G. (2008). Voluntary running rescues adult hippocampal neurogenesis after irradiation of the young mouse brain. *Proceedings of the National Academy of Science of the United States of America*, 105, 14632–14637.
- Naylor, A. S., Persson, A. I., Eriksson, P. S., Jonsdottir, I. H., & Thorlin, T. (2005). Extended voluntary running inhibits exercise-induced adult hippocampal progenitor proliferation in the spontaneously hypertensive rat. *Journal of Neurophysiology*, 93, 2406–2414.
- Nemeroff, C. B. (2016). Paradise lost: The neurobiological and clinical consequences of child abuse and neglect. *Neuron*, 89, 892–909.
- Oomen, C. A., Bekinschtein, P., Kent, B. A., Saksida, L. M., & Bussey, T. J. (2014). Adult hippocampal neurogenesis and its role in cognition. *WIREs Cognitive Science*, 5, 573–587.
- Oomen, C. A., Girardi, C. E. N., Cahyadi, R., Verbeek, E. C., Krugers, H., Joels, M., & Lucassen, P. J. (2009). Opposite effects of early maternal deprivation on neurogenesis in male versus female rats. *PLoS One*, 4, e3675.
- Oomen, C. A., Soeters, H., Audureau, N., Vermunt, L., van Hasselt, F. N., Manders, E. M., ... Krugers, H. (2010). Severe early life stress

- hampers spatial learning and neurogenesis, but improves hippocampal synaptic plasticity and emotional learning under high-stress conditions in adulthood. *Journal of Neuroscience*, 30, 6635–6645.
- Pawluski, J. L., Brummelte, S., Barha, C. K., Crozier, T. M., & Galea, L. A. (2009). Effects of steroid hormones on neurogenesis in the hippocampus of the adult female rodent during the estrous cycle, pregnancy, lactation and aging. *Frontiers in Neuroendocrinology*, 30, 343–357.
- Singh-Taylor, A., Molet, J., Jiang, S., Korosi, A., Bolton, J. L., Noam, Y., ... Baram, T. Z. (2017). NRSF-dependent epigenetic mechanisms contribute to programming of stress-sensitive neurons by neonatal experience, promoting resilience. *Molecular Psychiatry*. doi:10.1038/mp.2016.240
- Schoenfeld, T. J., & Gould, E. (2012). Stress, stress hormones, and adult neurogenesis. *Experimental Neurology*, 233, 12–21.
- Stranahan, A. M., Khalil, D., & Gould, E. (2006). Social isolation delays the positive effects of running on adult neurogenesis. *Nature Neuroscience*, 9, 526–533.
- Teicher, M. H., Anderson, C. M., & Polcari, A. (2012). Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 563–572.
- Vivar, C., Peterson, B. D., & van Praag, H. (2016). Running rewires the neuronal network of adult-born dentate granule cells. *Neuroimage*, 131, 29–41.
- Wu, M. V., & Hen, R. (2014). Functional dissociation of adult-born neurons along the dorsoventral axis of the dentate gyrus. *Hippocampus*, 24, 751–761.

How to cite this article: Abbink MR, Naninck EFG, Lucassen PJ, Korosi A. Early-life stress diminishes the increase in neurogenesis after exercise in adult female mice. *Hippocampus*. 2017;27:839–844. <https://doi.org/10.1002/hipo.22745>