



UvA-DARE (Digital Academic Repository)

Soil networks become more connected and take up more carbon as nature restoration progresses

Morriën, E.; Hannula, S.E.; Snoek, L.B.; Helmsing, N.R.; Zweers, H.; de Hollander, M.; Luján Soto, R.; Bouffaud, M.-L.; Buée, M.; Dimmers, W.; Duyts, H.; Geisen, S.; Girlanda, M.; Griffiths, R.I.; Jørgensen, H.-B.; Jensen, J.; Plassart, P.; Redecker, D.; Schmelz, R.M.; Schmidt, O.; Thomson, B.C.; Tisserant, E.; Uroz, S.; Winding, A.; Bailey, M.J.; Bonkowski, M.; Faber, J.H.; Martin, F.; Lemanceau, P.; de Boer, W.; van Veen, J.A.; van der Putten, W.H.

DOI

[10.1038/ncomms14349](https://doi.org/10.1038/ncomms14349)

Publication date

2017

Document Version

Other version

Published in

Nature Communications

[Link to publication](#)

Citation for published version (APA):

Morriën, E., Hannula, S. E., Snoek, L. B., Helmsing, N. R., Zweers, H., de Hollander, M., Luján Soto, R., Bouffaud, M.-L., Buée, M., Dimmers, W., Duyts, H., Geisen, S., Girlanda, M., Griffiths, R. I., Jørgensen, H.-B., Jensen, J., Plassart, P., Redecker, D., Schmelz, R. M., ... van der Putten, W. H. (2017). Soil networks become more connected and take up more carbon as nature restoration progresses. *Nature Communications*, 8, Article 14349. <https://doi.org/10.1038/ncomms14349>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Download date:29 Apr 2025

1 **Supplementary Information**

2 **Supplementary Methods**

3 Sequencing

4

5 Bacterial 16S rRNA gene fragments were amplified using the primer pair 530F (5'-
6 ACTCCTACGGGAGGCAGCAG)¹ and 803R (5'-CTACCGGGTATCTAAT-3'). The fungal ITS2 region was
7 amplified using primers ITS4 (5'-TCC TCC GCT TAT TGA TAT GC-3') and ITS9 (5'- GAA CGC AGC RAA IIG
8 YGA-3')². Finally, bacterial and fungal amplicons were pooled at equimolar concentrations to provide
9 a total of 2 ng DNA in each library.

10

11 Bacterial sequences were analysed using a Snakemake workflow³ that follows the SOP for 454-
12 sequencing data in mothur version 1.33.2⁴. The flowgrams were demultiplexed with a mismatch of 2
13 to the barcode, 3 mismatches to the primer, and flowgrams were trimmed to a size of 575 flows.
14 Flowgrams were corrected using the shhh.flows command, which is the mothur implementation of
15 the original PyroNoise algorithm⁵. Afterwards the results of the different sff files were combined for
16 further analysis. The merged sequences were aligned and classified with SINA⁶ against the SILVA 119
17 database⁷. Only reads that starts at position 9878 and stop at a position so that 90% of the sequence
18 fall into this region were kept. To reduce sequence errors even more the pre.cluster command is
19 used to merge sequences that are within 2 mismatches of each other. Chimeric sequences were
20 identified and removed using the chimera.uchime command⁸. OTUs were formed at maximum
21 distance of 0.03 using the dist.seqs command and average neighbour clustering. For each OTU a
22 consensus taxonomy was determined using the classify.otu command. Representative sequences for
23 each OTU were re-aligned to the Silva reference alignment and a neighbour joining tree was created
24 using the clearcut program⁹. Taxonomic classification and OTU clustering data were combined into
25 the BIOM format¹⁰ for further downstream statistical analysis with the Phyloseq¹¹ package for R.

26

27 Fungal sequences were initially demultiplexed according to their multiplex identifier (MID) using the
28 sffinfo command of Mothur v.1.22.2⁴. The raw flowgrams were filtered using the trim.flows
29 command to a minimum flowgram length of 360 cycles before the first noisy signal (signal ranging
30 from 0.5-0.7). All flowgrams were then truncated at 720 cycles. The internal transcribed spacer 2
31 (ITS2) region was extracted using Fungal ITS extractor v 2¹² and sequences shorter than 100 bp were
removed. See for more details Thomson *et al.*¹³.

32

33 For each microbial group, all sequences were pooled and sorted by decreasing length as
34 recommended for Usearch clustering¹⁴. Operational taxonomic units (OTUs) were generated from
35 these reads using the cluster_smallmem command at a 97 % similarity threshold. Phylogenetic
36 assignment was done for each OTU consensus sequence using the Basic Local Alignment Search Tool
37 (BLAST) algorithm v 2.2.23¹⁵ against the Ribosomal Data Project (RDP) database release 10.3¹⁶ for
38 bacterial and archaeal sequences. For fungi, the assignment was performed using the UNITE
39 database release 5.0¹⁷. All phylogenies were determined using a minimum e-value cut-off of 1e-5.


40

41 For bacteria and fungi, matrices containing the sequence abundances of different OTUs in each soil
42 sample were created. To obtain comparable samples with different number of sequences, data were
43 rarefied to least amount of reads using the 'rrarefy' comment in the vegan package in R¹⁸. We
44 confirmed that the selected amount of reads was approaching the saturation end of the rarefaction
45 curve. This rarefaction procedure resulted in 2800 sequences for bacteria and 635 for fungi. The
46 abundance of a certain OTU in a sample was used as an abundance measure for network analysis.
47 The fungal reads were assigned into 15 ecological functional guilds (nr.31 till nr. 45) according to the
48 Supplementary Table 5.


49 **Supplementary References**

- 50 1 Acosta-Martinez, V., Dowd, S., Sun, Y. & Allen, V. Tag-encoded pyrosequencing analysis of
51 bacterial diversity in a single soil type as affected by management and land use. *Soil Biol*
52 *Biochem* **40**, 2762-2770, doi:10.1016/j.soilbio.2008.07.022 (2008).
- 53 2 Ihrmark, K. *et al.* New primers to amplify the fungal ITS2 region - evaluation by 454-
54 sequencing of artificial and natural communities. *FEMS Microbiol. Ecol.* **82**, 666-677,
55 doi:10.1111/j.1574-6941.2012.01437.x (2012).
- 56 3 Koster, J. & Rahmann, S. Snakemake-a scalable bioinformatics workflow engine.
57 *Bioinformatics* **28**, 2520-2522, doi:10.1093/bioinformatics/bts480 (2012).
- 58 4 Schloss, P. D. *et al.* Introducing mothur: Open-Source, Platform-Independent, Community-
59 Supported Software for Describing and Comparing Microbial Communities. *Appl Environ*
60 *Microbiol* **75**, 7537-7541, doi:10.1128/aem.01541-09 (2009).
- 61 5 Quince, C., Lanzen, A., Davenport, R. J. & Turnbaugh, P. J. Removing Noise From
62 Pyrosequenced Amplicons. *BMC Bioinformatics* **12**, doi:10.1186/1471-2105-12-38 (2011).
- 63 6 Pruesse, E., Peplies, J. & Glöckner, F. O. SINA: Accurate high-throughput multiple sequence
64 alignment of ribosomal RNA genes. *Bioinformatics* **28**, 1823-1829,
65 doi:10.1093/bioinformatics/bts252 (2012).
- 66 7 Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing
67 and web-based tools. *Nucleic Acids Res* **41**, D590-D596, doi:10.1093/nar/gks1219 (2013).
- 68 8 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. UCHIME improves sensitivity
69 and speed of chimera detection. *Bioinformatics* **27**, 2194-2200,
70 doi:10.1093/bioinformatics/btr381 (2011).
- 71 9 Sheneman, L., Evans, J. & Foster, J. A. Clearcut: a fast implementation of relaxed neighbor
72 joining. *Bioinformatics* **22**, 2823-2824, doi:10.1093/bioinformatics/btl478 (2006).
- 73 10 McDonald, D. *et al.* An improved Greengenes taxonomy with explicit ranks for ecological and
74 evolutionary analyses of bacteria and archaea. *Isme Journal* **6**, 610-618,
75 doi:10.1038/ismej.2011.139 (2012).
- 76 11 McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis
77 and Graphics of Microbiome Census Data. *Plos One* **8**, doi:10.1371/journal.pone.0061217
78 (2013).
- 79 12 Nilsson, R. H. *et al.* An open source software package for automated extraction of ITS1 and
80 ITS2 from fungal ITS sequences for use in high-throughput community assays and molecular
81 ecology. *Fungal Ecology* **3**, 284-287, doi:10.1016/j.funeco.2010.05.002 (2010).
- 82 13 Thomson, B. C. *et al.* Soil conditions and land use intensification effects on soil microbial
83 communities across a range of European field sites. *Soil Biology and Biochemistry* **88**, 403-
84 413, doi:http://dx.doi.org/10.1016/j.soilbio.2015.06.012 (2015).
- 85 14 Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**,
86 2460-2461, doi:10.1093/bioinformatics/btq461 (2010).
- 87 15 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic Local Alignment Search
88 Tool. *J Mol Biol* **215**, 403-410, doi:10.1006/jmbi.1990.9999 (1990).
- 89 16 Cole, J. R. *et al.* The Ribosomal Database Project: improved alignments and new tools for
90 rRNA analysis. *Nucleic Acids Res* **37**, D141-D145, doi:10.1093/nar/gkn879 (2009).
- 91 17 Koljalg, U. *et al.* Towards a unified paradigm for sequence-based identification of fungi.
92 *Molec. Ecol.* **22**, 5271-5277, doi:10.1111/mec.12481 (2013).
- 93 18 R: A language and environment for statistical computing. R foundation for statistical
94 computing, Vienna, <http://www.R-project.org/>. (2013).

97 Supplementary Figures




98




99

100 Supplementary Figure 1: Co-occurrence network visualisation sub-groups. Network visualisation of the interaction strengths between the species sub-groups in recently, mid-
 101 term and long-term abandoned agricultural fields. Spearman correlations between the co-occurrences of all individual species of different fields were calculated. The proportion of
 102 correlations > 0.9 was divided by the total number of possible interactions in order to obtain the interaction strength between two groups of species. Line width is proportional to the
 103 absolute number of correlations > 0.9. Line colour and transparency is proportional to the interactions strength as indicated in the legend. The size of the circles is proportional to the number
 104 of species/taxa in that group. Red circles are bacterial groups, blue circles are fungal groups. Abbreviations: H.=Herbivorous R.F.=Root-feeding S.=Saprotrophic F.=Fungivorous
 105 B.=Bacterivorous H.F.=Herbofungivorous N.=Nematophagous O.=Omnivorous O.C.=Omni-carnivorous P.=Predaceous.



106



107

108

109

110

111


112

113

114


115

Supplementary Figure 2: Co-occurrence network visualisation main groups. Network visualisation of the interaction strengths between the main species groups in recently, mid-term and long-term abandoned agricultural fields. Spearman correlations between the co-occurrence of all individual species of different fields were calculated. The proportion of correlations > 0.9 was divided by the total number of possible interactions to obtain the interaction strength between two groups of species. Line width is proportional to the absolute number of correlations > 0.9. Line colour and transparency is proportional to the interactions strength as indicated in the legend. The size of the circles is proportional to the number of species/taxa in that group. Red circles are bacterial groups, blue circles are fungal groups. Abbreviations: H.=Herbivorous R.F.=Root-feeding S.=Saprotrophic F.=Fungivorous B.=Bacterivorous H.F.=Herbofungivorous N.=Nematophagous O.=Omnivorous O.C.=Omni-carnivorous P.=Predaceous.



116


117 Supplementary Figure 3: Graphical representation of the Supplementary Table 5. Showing number of taxa in each stage of abandonment per sampling site.
 118 Abbreviations of field sites see site description Supplementary Table 1. Additional to Supplementary Table 5 the ectomycorrhizal fungi and predaceous mesostigmata are displayed since they
 119 show significant patterns along the successional gradient. Total amount of species within the different species groups were plotted among field sites (each bar), categories of sites (recent-
 120 (blue), mid- (red) and long-term (green) abandoned) and in years since abandonment (left to right). Line represents the median. The boxes represent the upper- and under-quartile of the
 121 data range. The whiskers represent 1.5 times the distance between the first and third quartile. Data beyond the end of the whiskers are outliers and plotted as points (as specified by Tukey).
 122 Abbreviations: E. M. fungi=Ectomycorrhizal fungi, F. C. mites=Fungivorous cryptostigmatic mites, P. mesostigmata=Predaceous mesostigmatic mites, R.F. nematodes=Root-feeding
 123 nematodes, B. nematodes=Bacterivorous nematodes, OC. nematodes=Omni-carnivorous nematodes, F. nematodes=Fungivorous nematodes, P. mites= Predaceous mites, F.NC.
 124 Mites=Fungivorous non-cryptostigmatic mites, O. mites=Omnivorous mites H. mites=Herbivorous mite.



125
126

127 Supplementary Figure 4: Results of Analysis of similarities. ANOSIM (Euclidian, R- and p-values given) on
 128 soil properties at recent, mid- and long-term abandoned fields. The variation of the similarities between groups (recent-,
 129 mid-term-, long-term abandoned since agricultural practice, first boxplot left) is larger than the variation within groups 1, 2
 130 and 3 representing recent-, mid-term and long-term abandonment (R=0.2333; p=0.0041). Within groups the variation of
 131 the recently abandoned fields is significantly higher than of the mid-term abandoned fields (p=0.0033), however, not
 132 significantly higher than of the long-term abandoned fields (p=0.0544).

133



134

135 Supplementary Figure 5: NMDS of the rank abundance of plants in cores. Non-metric Multidimensional
 136 Scaling (NMDS) of the rank abundance of the three most dominated plant species (*Agrostis capillaris*, *Holcus lanatus*,
 137 *Plantago lanceolata*) in each of the experimental cores, which shows strong overlap in the plant community between the
 138 cores.


139

140


141

142

143




146 Supplementary Figure 6: Carbon and nitrogen flow in the soil food-web during 2 weeks. The root
 147 root derived carbon is marked with grey, shoot carbon with green, bacterial-based channel with red,
 148 fungal-based channel with blue and the higher trophic level interactions with purple and pink. The plots with standard deviation represent the
 149 amount of ¹³C excess in pmol per gram soil (bacteria, fungi, AMF), or δ^{13} C and δ^{15} N excess (all other groups) compared to
 150 natural values measured from non-labelled controls in recently (light grey), mid-term (dark grey) and long-term (black)
 151 abandoned agricultural fields. The statistical analyses of the treatment effects with field site as a factor nested in
 152 successional stage are presented in boxes next to the figure. The darker coloured arrows depict the carbon flow in the food
 153 web while the lighter coloured (dashed) arrows depict the nitrogen flow. The width of the arrows between groups reflects the
 154 percentage of correlation >0.9 between groups in all the fields as in Fig. 1. The pie charts above the arrows are also
 155 calculated from Fig. 1, and represent the proportion of significant interactions in recently (light grey), mid-term (dark grey)
 156 and long-term (black) abandoned fields. The (rho)p values at the arrows represent the Spearman-rank correlation values
 157 between the groups that are connected by the arrow, which is calculated from the labelling data. Significant correlations
 158 are bold and marked with darker arrow colour. By correlation analysis of pulse labelling data we wanted to analyse the
 159 potential of feeding relationships that emerged on the basis of correlations in the network analysis.



164 Supplementary Figure 7: Map of sampling sites in The Netherlands

174


175



176

177 Supplementary Figure 8: Shoot + root biomass & C/N ratio of the plants. A. Shoot + root biomass in grams
 178 of the plants from the cores at the time of harvest. Abbreviations of field sites see site description Supplementary Table 1.
 179 B. C/N ratio of the plant shoots and roots from the cores at the time of harvest. Abbreviations of field sites see site

180 description Supplementary Table 1.
 181 Error bars represent standard errors
 182 of the mean (s.e.m.).




184

185 Supplementary Figure 9: PCA of site variation explained by the physiochemical data. Principal
 186 Component Analysis of site variation explained by the physiochemical data collected in October 2011. Abbreviations of
 187 field sites have been provided in the site description of Supplementary Table 4. Phosphorus (P_2O_5) is measured as P-Olsen
 188 in g/kg, Cation Exchange Capacity (Cobaltihexamine) and all ions indicated with a '+' are measured as cmol/kg, Residual
 189 water content is the g/kg water remaining after drying at 105°C, metals have mg/kg as a unit, Organic Matter, Organic
 190 Carbon and Total N are measured as g/kg.

191

192

193



194

195 Supplementary Figure 10: Core harvesting scheme.

196

197

198

199

200

201

202

203

204

205

206

207

208

209 **Supplementary Tables**

210 Supplementary Table 1: Nested ANOVA results for Supplementary Fig. 4. Nested ANOVA results for each of
 211 the graphs from Supplementary Fig. 4. We analysed the number of species per aggregated group in three ways: the effect
 212 of site, succession, and time since abandonment. The sites OR, REY and TW (S-Table 1) were categorized as recently
 213 abandoned fields, MO, NR and WV as mid-term abandoned fields and MV, DK and BB as long-term abandoned fields. These
 214 categories mark the factor 'succession'. We also analysed the effect as a regression taking 'time since abandonment' as a
 215 continuous variable (S-Table 1). For the other factors we used a nested ANOVA approach: when testing 'site' as a factor,
 216 subplots were nested in 'site' and when testing 'succession' as a factor, sites were nested in 'succession'.

217

	Site		Succession		Time since abandonment	
	<i>F-values</i>	<i>P-values</i>	<i>F-values</i>	<i>P-values</i>	<i>F-values</i>	<i>P-values</i>
Bacteria	12.4	<0.0001	44.4	<0.0001	1.28	0.274
Archaea	1.11	0.412	0.348	0.712	1.06	0.316
Collembola	1.03	0.449	0.225	0.801	<0.001	0.984
Earthworms	1.96	0.112	1.86	0.185	<0.001	0.978
Enchytraeids	0.407	0.902	0.215	0.808	0.399	0.536
AMF	1.85	0.136	2.71	0.095	5.93	0.026
Fungi	1.3	<0.0001	33.8	<0.0001	0.156	0.698
Predaceous mites	2.2	0.078	7.51	0.0043	14.4	0.001
Fungivorous cryptostigmatic mites	8.82	<0.0001	14.4	<0.001	36.3	<0.0001
Omnivorous mites	0.763	0.639	0.684	0.517	1.58	0.225
Herbivorous mites	6.38	<0.001	9.75	0.0014	27.3	<0.0001
Fungivorous non-cryptostigmatic mites	1.4	0.262	1.4	0.272	0.072	0.792
Root-feeding nematodes	4.85	0.003	10.7	<0.001	10.5	0.005
Bacterivorous nematodes	2.52	0.049	4.84	0.021	9.07	0.007
Fungivorous nematodes	4.67	0.003	1.17	0.334	1.98	0.176
Omni-carnivorous nematodes	1.02	0.458	0.557	0.583	0.173	0.683
Plants	9.48	<0.0001	11.3	<0.001	31.4	<0.0001
Ectomycorrhiza	2.6	0.047	4.95	0.02	0.144	0.709
Predaceous mesostigmata	2.06	0.096	5.48	0.014	9.54	0.006

218

219

220 Supplementary Table 2: Amount of variation explained by all measured soil properties via constrained analysis. Linear (RDA) or unimodal (CCA), dependent on the
 221 length of the gradient. Then via a forward selection procedure the most explaining soil properties were selected and used to re-analyse the constrained analysis. The variation explained by
 222 each of the significant contributing soil properties is displayed below. Phosphorus (P₂O₅) is measured as P-Olsen in g/kg, Cation Exchange Capacity (Cobaltihexamine) and all ions indicated
 223 with a '+' are measured as cmol+/kg, Residual water content is the g/kg water remaining after drying at 105°C, metals have mg/kg as a unit, Organic Matter, Organic Carbon and Total N are
 224 measured as g/kg.

Group	% var. expl. by all soil properties	Most expl. soil property 1			Most expl. soil property 2			Most expl. soil property 3			Most expl. soil property 4			Most expl. soil property 5			Most expl. soil property 6		
		Env. Var.	Var. expl.	P	Env. Var.	Var. expl.	P	Env. Var.	Var. expl.	P	Env. Var.	Var. expl.	P	Env. Var.	Var. expl.	P	Env. Var.	Var. expl.	P
Bacteria	38.9% (CCA)	Fe ⁺	10.5%	0.002	Water	6.7%	0.002	Fe	6%	0.002	CEC	5.8%	0.01						
Archaea	46.1% (RDA)	Cu	12.4%	0.042															
Fungi (incl. AMF)	38.4% (CCA)	Fe ⁺	10.9%	0.002	C/N	6.2%	0.002	Fe	5.7%	0.006	P	5.3	0.036	pH	5.4%	0.022			
Protists	28.4% (CCA)	Fe ⁺	17.1%	0.002	Fe	7.9%	0.004	Cu	7.3%	0.004	P	6.1%	0.004						
Nematodes	34.6% (RDA)	Na ⁺	12.6%	0.004	Org. C	7%	0.062	Mg ⁺	7.1%	0.013	Ca ⁺	9.5%	0.036						
Enchytraeids	54.4% (CCA)	Fe ⁺	17%	0.002	Fe	7.9%	0.004	Cu	7.3%	0.004	P	6.1%	0.01	Org C	6.4%	0.004	Ca	5.0%	0.038
Mites	29.4% (RDA)	P	13%	0.016	Al ⁺	12.3%	0.018												
Collembola	48.5% (RDA)	Fe ⁺	16.6%	0.024	K ⁺	11.5%	0.05	C/N	14.4%	0.0.28									
Plants	68.4% (CCA)	pH	13.2%	0.002	Org. C	11.1%	0.002	P	7.9%	0.008	Al ⁺	6.8%	0.002	C/N	6.9%	0.006	Tot N	6.9%	0.002

225

226
227
228
229
230
231

Supplementary Table 3: Average label in each of the measured groups at all harvest points: 1 day, 1 week and 2 weeks after pulse labelling in atom% C. The table shows the significance levels of the three successional stages in an ANOVA with a nested design. The sites OR, REY and TW (S-Table 1) were categorized as recently abandoned fields, MO, NR and WV as mid-term abandoned fields and MV, DK and BB as long-term abandoned fields. These categories mark the factor 'succession'. Where sites are nested in succession time.

Generalized Linear Model with Nested design. Field nested in succession time

			Average	Stdev	Stats	
1 day	Plant shoots	Recent	1.62	0.334	F(2,17) = 0.18 , p=0.83	
		Mid	1039,00	0.482		
		Long	1098,00	0.363		
	Plant roots	Recent	0.119	0.086	F(2,15) = 0.42 , p=0.67	
		Mid	0.080	0.052		
		Long	0.038	0.017		
	Soil	Recent	0.000	0.000	F(2,16) = 2.49 , p=0.11	
		Mid	0.001	0.001		
		Long	0.000	0.000		
1 week	Earthworms	Recent	0.008	0.005	F(2,5) = 0.70 , p=0.54	
		Mid	0.001	0.000		
		Long	0.003	nd		
	Root-feeding nematodes	Recent	0.175	0.062	F(2,14) = 0.20 , p=0.82	
		Mid	0.164	0.072		
		Long	0.157	0.055		
	Fungal-feeding nematodes	Recent	0.045	0.040	F(2,6) = 0.35 , p=0.72	
		Mid	0.003	0.004		
		Long	0.009	0.003		
	Fungivorous collembola	Recent	0.125	0.056	F(2,15) = 2.54 , p=0.11	
		Mid	0.204	0.107		
		Long	0.011	0.010		
	Fungivorous cryptostigmatic mites	Recent	0.007	0.004	F(2,8) = 6.62 , p=0.02	
		Mid	0.056	0.067		
		Long	0.025	0.022		
	Fungivorous prostigmatic mites	Recent	0.000	0.000	F(4,5) = 9.42 , p=0.02	
		Mid	0.012	0.010		
		Long	0.001	0.001		
	Herbivorous bugs	Recent	0.340	0.233	F(1,6) = 0.48 , p=0.52	
		Mid	0.293	0.131		
		Long	nd	nd		
	2 weeks	Bacterial feeding nematodes	Recent	0.009	0.003	F(2,16) = 2.16 , p=0.15
			Mid	0.013	0.007	
			Long	0.021	0.008	
		Fungivorous astigmatic mites	Recent	0.004	0.003	F(2,6) = 5.79 , p=0.04
			Mid	0.013	0.012	
			Long	0.005	0.002	
Predaceous mites		Recent	0.008	0.007	F(2,38) = 0.79 , p=0.46	
		Mid	0.01	0.006		
		Long	0.059	0.085		
Predaceous spiders		Recent	0.126	0.012	F(2,10) = 0.56 , p=0.59	
		Mid	0.216	0.035		
		Long	0.177	nd		
Omnivorous nematodes		Recent	0.014	0.006	F(2,18) = 0.90 , p=0.43	
		Mid	0.026	0.009		
		Long	0.016	0.008		

232

233 Supplementary Table 4: Averages and standard deviations of delta ¹³C values. Averages and standard
 234 deviations of delta ¹³C values of unlabelled controls and labelled material of 1 day, 1 week and two weeks after pulse
 235 labelling. Numbers in bold indicate the highest numbers, the moment at which the pulse has in incorporated most ¹³C in
 236 the tissue of the measured group of soil biota. That point was chosen to represent the label incorporated in that specific
 237 group of soil organisms.

238

	Non-labeled		1 day		1 week		2 weeks	
	Average	Stdev	Average	Stdev	Average	Stdev	Average	Stdev
Plant shoots	-29.54	1.28	1002.33	713.03	959.07	363.87	563.79	358.55
Plant roots	-29.78	0.57	42.65	108.56	23.42	72.20	53.36	99.01
Soil	-27.92	0.96	-27.42	1.18	-27.54	1.45	-27.08	1.38
Earthworms	-25.82	1.12	-22.94	6.60	-22.72	5.31	-22.07	3.38
Root-feeding nematodes	-27.43	5.78	80.54	96.50	125.39	111.64	125.95	172.38
Fungal-feeding nematodes	-27.94	2.35	-25.47	3.10	-7.33	49.83	-9.42	25.23
Fungivorous collembola	-23.42	1.84	57.19	62.80	93.05	149.33	81.15	106.62
Fungivorous cryptostigmatic mites	-24.63	0.27	-17.22	15.17	-11.72	27.60	-20.58	6.00
Fungivorous prostigmatic mites	-23.99	0.32	-16.20	13.21	-18.68	13.20	-19.34	8.14
Herbivorous bugs	-21.65	-	541.14	963.12	294.63	358.55	135.46	130.44
Bacterial feeding nematodes	-26.66	2.59	-16.60	27.08	-19.68	7.18	-15.11	11827.00
Fungivorous astigmatic mites	-22.45	3.76	-22.94	4.47	-23.03	3.73	-17.38	13.47
Predaceous mites	-23.84	2.52	-16.65	9.22	-14.31	18.28	-1.66	86.77
Predaceous spiders	-24.21	1.82	-24.83	1.72	167.54	262.84	125.70	76.11
Omnivorous nematodes	-24.06	9.11	-20.71	9.32	-1.656	36.39	-6.84	14.56
Enchytraeds	-27.67	-	-17.21	10.78	-39.79	-	-28.94	5.12

239

240 Supplementary Table 5: Description of field sites coordinates and time since abandonment.

Field name	Geocoordinates	Abandoned since
Oud Reemst (OR)	N 52°2'27 E 5°48'34	2005
Reijerskamp (REY)	N 52°1'0 E 5°46'21	2005
Telefoonweg (TW)	N 52°00'9 E 5°45'8	2002
De Mossel (MO)	N 52°3'40 E 5°45'8	1995
Nieuw Reemst (NR)	N 52°2'33 E 5°46'29	1990
Wolfhezer Veld (WV)	N 51°59'43 E 5°47'24	1988
Mosselse Veld (MV)	N 52°4'23 E 5°44'13	1985
Dennenkamp (DK)	N 52°1'43 E 5°48'2	1982
Boersbos (BB)	N 52°3'44 E 5°59'57	1982

241

242 Supplementary Table 6: Number of taxa within sub-group. Number of taxa within sub-group in each stage of
 243 abandonment (occurrence > 1 sample), for recent, mid-term and long-term abandoned fields, used in the Spearman
 244 correlation interaction strength network visualization (Fig. 1A)

Nr.	Names	Recent	Mid-term	Long-term	Nr.	Names	Recent	Mid-term	Long-term
1	Proteobacteria	398	403	346	31	AMF	17	22	30
2	Chloroflexi	54	69	51	32	Ascomycota	70	86	70
3	Actinobacteria	280	241	230	33	Basidiomycota	7	8	9
4	Firmicutes	38	47	34	34	Chytridiomycota	5	2	0
5	Acidobacteria	254	213	245	35	Endophytes	11	10	13
6	Verrucomicrobia	4	99	3	36	Molds	17	41	27
7	Gemmatimonadetes	40	35	36	37	Ectomycorrhiza	5	2	7
8	Nitrospirae	12	5	15	38	Nematophagous fungi	3	9	7
9	Unclassified (bacteria)	258	274	214	39	Other (fungi)	15	17	13
10	Bacteroidetes	64	91	44	40	Potential plant-pathogen	10	14	9
11	WD272	6	7	8	41	Saprotrophic fungi	30	36	31
12	Candidate_division_WS3	11	4	6	42	Unknown fungi	128	136	106
13	Planctomycetes	30	104	39	43	Wood decomposer or parasite	8	9	5
14	Candidate_division_TM7	10	12	6	44	Yeasts	13	14	13
15	Fibrobacteres	3	1	3	45	Zygomycota	4	3	8
16	SHA-109	2	2	0	46	Predaceous mesostigmata	10	14	16
17	Elusimicrobia	1	1	2	47	Herbo-fungivorous cryptostigmata	3	3	3
18	WCHB1-60	1	2	1	48	Predaceous prostigmata	0	3	5
19	Thermotogae	0	0	1	49	Fungivorous cryptostigmata	3	7	10
20	Cyanobacteria	0	2	0	50	Omnivorous prostigmata	1	1	1
21	Chlamydiae	0	3	0	51	Bacterivorous astigmata	0	1	0
22	Armatimonadetes	1	2	1	52	Herbivorous prostigmata	4	4	4
23	Chlorobi	0	1	0	53	Herbivorous cryptostigmata	0	1	2
24	TM6	0	1	1	54	Fungivorous prostigmata	2	3	2
25	SM2F11	0	0	0	55	Fungivorous astigmata	0	1	0
26	Archaea	54	54	54	56	Root-feeding nematodes	6	7	10
27	Fungivorous collembola	12	12	12	57	Bacterivorous nematodes	11	11	13
28	Predaceous collembola	2	2	1	58	Fungivorous nematodes	3	4	3
29	Earthworms	2	2	2	59	Omni-carnivorous nematodes	10	8	10
30	Enchytraeids	15	18	20	60	Plants	29	29	24

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259 Supplementary Table 7: Number of taxa within main group. Number of taxa within main group in each stage
 260 of abandonment (occurrence > 1 sample), for recent, mid-term and long-term abandoned fields, used in the Spearman
 261 correlation interaction strength network visualization (Fig. 1B).

Nr.	Names	Recent	Mid-term	Long-term
1	Bacteria	1467	1620	1286
2	Archaea	54	54	54
3	Collembola	14	14	13
4	Earthworms	2	2	2
5	Enchytraeids	15	18	20
6	AMF	17	22	30
7	Fungi	326	387	318
8	Predaceous mites	10	17	21
9	Fungivorous cryptostigmatic mites	3	7	10
10	Omnivorous mites	4	5	4
11	Herbivorous mites	4	5	6
12	Fungivorous non-cryptostigmatic mites	2	4	2
13	Root-feeding nematodes	6	7	10
14	Bacterivorous nematodes	11	11	13
15	Fungivorous nematodes	3	4	3
16	Omni-carnivorous nematodes	10	8	10
17	Plants	29	29	24
	Subtotals	1977	2214	1826

262