



UvA-DARE (Digital Academic Repository)

Lateral Occipital cortex responsive to local correlation structure of natural images

Scholte, H.S.; Ghebreab, S.; Smeulders, A.; Lamme, V.

DOI

[10.3389/conf.fnins.2010.03.00162](https://doi.org/10.3389/conf.fnins.2010.03.00162)

Publication date

2010

Document Version

Final published version

Published in

Frontiers in Neuroscience

[Link to publication](#)

Citation for published version (APA):

Scholte, H. S., Ghebreab, S., Smeulders, A., & Lamme, V. (2010). Lateral Occipital cortex responsive to local correlation structure of natural images. *Frontiers in Neuroscience, Conference abstract*(Computational and Systems Neuroscience 2010). <https://doi.org/10.3389/conf.fnins.2010.03.00162>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

EVENT ABSTRACT

[« Back to Event](#)

Lateral Occipital cortex responsive to local correlation structure of natural images

 H.Steven Scholte^{1*}, Sennay Ghebream¹, Arnold Smeulders² and Victor Lamme¹
¹ University of Amsterdam, Department of Psychology, Netherlands

² University of Amsterdam, Department of Informatics, Netherlands

It is clear from behavioral experiments that subjects can rapidly access information about visual scenes (Potter, 1976) and that different types of scenes (such as beaches and mountains) differ in terms of low level image statistics (Torralba & Oliva, 2003). Furthermore, the distribution of local contrasts in natural images adheres to the Weibull distribution (Geusebroek & Smeulders, 2005), which is a family of distribution deforming from power-law to normal with two free parameters, beta and gamma. The beta parameter indicates the scale of the distribution, whereas the gamma parameter represents its shape. Spatially coherent scenes with one or a few objects tend to have a low gamma value, i.e. the distribution of their local contrast values approximates a power-law. In contrast, cluttered scenes with many uncorrelated visual structures typically have a high gamma values corresponding with a Gaussian distribution. We recently showed that the brain is capable of estimating the beta and gamma value of a scene by summarizing the X and Y cell populations of the LGN (Scholte et al., 2009). Here we investigated to what degree the brain is sensitive to differences in the global correlation (gamma) of a scene by presenting subjects with a wide range of natural images while measuring BOLD-MRI. Covariance analysis of the single-trial BOLD-MRI data with the gamma parameter showed that only the lateral occipital cortex (LO), and no other areas, responds stronger to low gamma values (corresponding to images with a power-law distribution) than high gamma values (corresponding to images with a normal distribution). The analysis of the covariance matrix of the voxel-pattern cross-correlated single-trial data further revealed that responses to images containing clear objects are more similar in their spatial structure than images that do not contain objects. This data is consistent with a wide range of literature on object perception and area LO (Grill-Spector et al., 2001), and extends our understanding of object recognition by showing that the global correlation structure of a scene is (part of) the diagnostics that are used by the brain to detect objects.

Conference: Computational and Systems Neuroscience 2010, Salt Lake City, UT, United States, 25 Feb - 2 Mar, 2010.

Presentation Type: Poster Presentation Topic: Poster session I

 Citation: Scholte H, Ghebream S, Smeulders A and Lamme V (2010). Lateral Occipital cortex responsive to local correlation structure of natural images. *Front. Neurosci. Conference Abstract: Computational and Systems Neuroscience 2010*. doi: 10.3389/conf.fnins.2010.03.00162

Received: 02 Mar 2010; Published Online: 02 Mar 2010.

* Correspondence: H.Steven Scholte, University of Amsterdam, Department of Psychology, Amsterdam, Netherlands, h.s.scholte@uva.nl

Abstract Info

Abstract

Export Citation ▼

The Authors in

Frontiers ▼

Google ▼

Google Scholar ▼

PubMed ▼

Related Article

in Frontiers

Google Scholar

PubMed

[Back to top](#) ▲

[Home](#)
[Contact](#)
[Submit](#)
[Newsletters](#)
[About Frontiers](#)
[Media Relations](#)
[FAQs](#)
[RSS/Twitter](#)