UNIVERSITY OF AMSTERDAM

UvA-DARE (Digital Academic Repository)

Limits to Gauge Coupling in the Dark Sector Set by the Nonobservation of Instanton-Induced Decay of Super-Heavy Dark Matter in the Pierre Auger Observatory data

Abreu, P.; de Jong, S.J.; Vink, J.; Pierre Auger Collaboration

DOI
10.1103/PhysRevLett.130.061001

Publication date
2023

Document Version
Final published version

Published in
Physical Review Letters

Link to publication

Citation for published version (APA):
Abreu, P., de Jong, S. J., Vink, J., & Pierre Auger Collaboration (2023). Limits to Gauge Coupling in the Dark Sector Set by the Nonobservation of Instanton-Induced Decay of Super-Heavy Dark Matter in the Pierre Auger Observatory data. *Physical Review Letters*, *130*(6), Article 061001. https://doi.org/10.1103/PhysRevLett.130.061001

General rights
It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations
If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: https://uba.uva.nl/en/contact, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (https://dare.uva.nl)

Download date:08 Oct 2026

Limits to Gauge Coupling in the Dark Sector Set by the Nonobservation of Instanton-Induced Decay of Super-Heavy Dark Matter in the Pierre Auger Observatory Data

P. Abreu,[1] M. Aglietta,[2,3] J. M. Albury,[4] I. Allekotte,[5] K. Almeida Cheminant,[6] A. Almela,[7,8] R. Aloisio,[9,10] J. Alvarez-Muñiz,[11] R. Alves Batista,[12] J. Ammerman Yebra,[11] G. A. Anastasi,[2,3] L. Anchordoqui,[13] B. Andrada,[7] S. Andringa,[1] C. Aramo,[14] P. R. Araújo Ferreira,[15] E. Arnone,[16,3] J. C. Arteaga Velázquez,[17] H. Asorey,[7] P. Assis,[1] G. Avila,[18] E. Avocone,[19,9] A. M. Badescu,[20] A. Bakalova,[21] A. Balaceanu,[22] F. Barbato,[9,10] J. A. Bellido,[4,23] C. Berat,[24] M. E. Bertaina,[16,3] G. Bhatta,[6] P. L. Biermann,[25] V. Binet,[26] K. Bismark,[27,7] T. Bister,[15] J. Biteau,[28] J. Blazek,[21] C. Bleve,[24] J. Blümer,[29] M. Boháčová,[21] D. Boncioli,[19,10] C. Bonifazi,[30,31] L. Bonneau Arbeletche,[32] N. Borodai,[6] A. M. Botti,[7] J. Brack,[33] T. Bretz,[15] P. G. Brichetto Orchera,[7] F. L. Briechle,[15] P. Buchholz,[34] A. Bueno,[35] S. Buitink,[36] M. Buscemi,[37] M. Büsken,[27,7] K. S. Caballero-Mora,[38] L. Caccianiga,[39,40] F. Canfora,[12,41] I. Caracas,[42] R. Caruso,[43,37] A. Castellina,[2,3] F. Catalani,[44] G. Cataldi,[45] L. Cazon,[11] M. Cerda,[46] J. A. Chinellato,[32] J. Chudoba,[21] L. Chytka,[47] R. W. Clay,[4] A. C. Cobos Cerutti,[48] R. Colalillo,[49,14] A. Coleman,[50] M. R. Coluccia,[45] R. Conceição,[1] A. Condorelli,[9,10] G. Consolati,[40,51] F. Contreras,[18] F. Convenga,[29] D. Correia dos Santos,[52] C. E. Covault,[53] S. Dasso,[54,55] K. Daumiller,[29] B. R. Dawson,[4] J. A. Day,[4] R. M. de Almeida,[52] J. de Jesús,[7,29] S. J. de Jong,[12,41] J. R. T. de Mello Neto,[31,56] I. De Mitri,[9,10] J. de Oliveira,[57] D. de Oliveira Franco,[32] F. de Palma,[58,45] V. de Souza,[59] E. De Vito,[58,45] A. Del Popolo,[43,37] M. del Río,[18] O. Deligny,[28] L. Deval,[29,7] A. di Matteo,[3] M. Dobre,[22] C. Dobrigkeit,[32] J. C. D'Olivo,[60] L. M. Domingues Mendes,[1] R. C. dos Anjos,[61] M. T. Dova,[62] J. Ebr,[21] R. Engel,[27,29] I. Epicoco,[58,45] M. Erdmann,[15] C. O. Escobar,[63] A. Etchegoyen,[7,8] H. Falcke,[12,64,41] J. Farmer,[65] G. Farrar,[66] A. C. Fauth,[32] N. Fazzini,[63] F. Feldbusch,[67] F. Fenu,[16,3] B. Fick,[68] J. M. Figueira,[7] A. Filipčič,[69,70] T. Fitoussi,[29] T. Fodran,[12] T. Fujii,[65,71] A. Fuster,[7,8] C. Galea,[12] C. Galelli,[39,40] B. García,[48] A. L. Garcia Vegas,[15] H. Gemmeke,[67] F. Gesualdi,[7,29] A. Gherghel-Lascu,[22] P. L. Ghia,[28] U. Giaccari,[12] M. Giammarchi,[40] J. Glombitza,[15] F. Gobbi,[46] F. Gollan,[7] G. Golup,[5] M. Gómez Berisso,[5] P. F. Gómez Vitale,[18] J. P. Gongora,[18] J. M. González,[5] N. González,[72] I. Goos,[5,29] D. Góra,[6] A. Gorgi,[2,3] M. Gottowik,[42] T. D. Grubb,[4] F. Guarino,[49,14] G. P. Guedes,[73] E. Guido,[3,16] S. Hahn,[29,7] P. Hamal,[21] M. R. Hampel,[7] P. Hansen,[62] D. Harari,[5] V. M. Harvey,[4] A. Haungs,[29] T. Hebbeker,[15] D. Heck,[29] G. C. Hill,[4] C. Hojvat,[63] J. R. Hörandel,[12,41] P. Horvath,[47] M. Hrabovský,[47] T. Huege,[29,36] A. Insolia,[43,37] P. G. Isar,[74] P. Janecek,[21] J. A. Johnsen,[75] J. Jurysek,[21] A. Kääpä,[42] K. H. Kampert,[42] B. Keilhauer,[29] A. Khakurdikar,[12] V. V. Kizakke Covilakam,[7,29] H. O. Klages,[29] M. Kleifges,[67] J. Kleinfeller,[46] F. Knapp,[27] N. Kunka,[67] B. L. Lago,[76] N. Langner,[15] M. A. Leigui de Oliveira,[77] V. Lenok,[29] A. Letessier-Selvon,[78] I. Lhenry-Yvon,[28] D. Lo Presti,[43,37] L. Lopes,[1] R. López,[79] L. Lu,[80] Q. Luce,[27] J. P. Lundquist,[70] A. Machado Payeras,[32] G. Mancarella,[58,45] D. Mandat,[21] B. C. Manning,[4] J. Manshanden,[81] P. Mantsch,[63] S. Marafico,[28] F. M. Mariani,[39,40] A. G. Mariazzi,[62] I. C. Mariş,[72] G. Marsella,[82,37] D. Martello,[58,45] S. Martinelli,[29,7] O. Martínez Bravo,[79] M. Mastrodicasa,[19,10] H. J. Mathes,[29] J. Matthews,[83] G. Matthiae,[84,85] E. Mayotte,[75,42] S. Mayotte,[75] P. O. Mazur,[63] G. Medina-Tanco,[60] D. Melo,[7] A. Menshikov,[67] S. Michal,[47] M. I. Micheletti,[26] L. Miramonti,[39,40] S. Mollerach,[5] F. Montanet,[24] L. Morejon,[42] C. Morello,[2,3] M. Mostafá,[86] A. L. Müller,[21] M. A. Muller,[32] K. Mulrey,[12,41] R. Mussa,[3] M. Muzio,[66] W. M. Namasaka,[42] A. Nasr-Esfahani,[42] L. Nellen,[60] G. Nicora,[87] M. Niculescu-Oglinzanu,[22] M. Niechciol,[34] D. Nitz,[68] I. Norwood,[68] D. Nosek,[88] V. Novotny,[88] L. Nožka,[47] A. Nucita,[58,45] L. A. Núñez,[89] C. Oliveira,[59] M. Palatka,[21] J. Pallotta,[87] P. Papenbreer,[42] G. Parente,[11] A. Parra,[79] J. Pawlowsky,[42] M. Pech,[21] J. Pękala,[6] R. Pelayo,[90] J. Peña-Rodriguez,[89] E. E. Pereira Martins,[27,7] J. Perez Armand,[91] C. Pérez Bertolli,[7,29] L. Perrone,[58,45] S. Petrera,[9,10] C. Petrucci,[19,10] T. Pierog,[29] M. Pimenta,[1] V. Pirronello,[43,37] M. Platino,[7] B. Pont,[12] M. Pothast,[41,12] P. Privitera,[65] M. Prouza,[21] A. Puyleart,[68] S. Querchfeld,[42] J. Rautenberg,[42] D. Ravignani,[7] M. Reininghaus,[29,7] J. Ridky,[21] F. Riehn,[1] M. Risse,[34] V. Rizi,[19,10] W. Rodrigues de Carvalho,[12] J. Rodriguez Rojo,[18] M. J. Roncoroni,[7] S. Rossoni,[81] M. Roth,[29] E. Roulet,[5] A. C. Rovero,[54] P. Ruehl,[34] A. Saftoiu,[22] M. Saharan,[12] F. Salamida,[19,10] H. Salazar,[79] G. Salina,[85] J. D. Sanabria Gomez,[89] F. Sánchez,[7] E. M. Santos,[91] E. Santos,[21] F. Sarazin,[75] R. Sarmento,[1] C. Sarmiento-Cano,[7] R. Sato,[18] P. Savina,[80] C. M. Schäfer,[29] V. Scherini,[58,45] H. Schieler,[29] M. Schimassek,[27,7] M. Schimp,[42] F. Schlüter,[29,7] D. Schmidt,[27] O. Scholten,[36] H. Schoorlemmer,[12,41] P. Schovánek,[21] F. G. Schröder,[50,29] J. Schulte,[15] T. Schulz,[29] S. J. Sciutto,[62] M. Scornavacche,[7,29] A. Segreto,[92,37] S. Sehgal,[42] R. C. Shellard,[93] G. Sigl,[81] G. Silli,[7,29] O. Sima,[22,94] R. Smau,[22] R. Šmída,[65] P. Sommers,[86] J. F. Soriano,[13] R. Squartini,[46] M. Stadelmaier,[29,7] D. Stanca,[22] S. Stanič,[70] J. Stasielak,[6] P. Stassi,[24] A. Streich,[27,7] M. Suárez-Durán,[72] T. Sudholz,[4] T. Suomijärvi,[28] A. D. Supanitsky,[7] Z. Szadkowski,[95] A. Tapia,[96] C. Taricco,[16,3]

 © 2023 American Physical Society

C. Timmermans,[41,12] O. Tkachenko,[29] P. Tobiska,[21] C. J. Todero Peixoto,[44] B. Tomé,[1] Z. Torrès,[24] A. Travaini,[46] P. Travnicek,[21] C. Trimarelli,[19,10] M. Tueros,[62] R. Ulrich,[29] M. Unger,[29] L. Vaclavek,[47] M. Vacula,[47] J. F. Valdés Galicia,[60] L. Valore,[49,14] E. Varela,[79] A. Vásquez-Ramírez,[89] D. Veberič,[29] C. Ventura,[56] I. D. Vergara Quispe,[62] V. Verzi,[85] J. Vicha,[21] J. Vink,[97] S. Vorobiov,[70] H. Wahlberg,[62] C. Watanabe,[31] A. A. Watson,[98] A. Weindl,[29] L. Wiencke,[75] H. Wilczyński,[6] D. Wittkowski,[42] B. Wundheiler,[7] A. Yushkov,[21] O. Zapparrata,[72] E. Zas,[11] D. Zavrtanik,[70,69] M. Zavrtanik,[69,70] and L. Zehrer[70]

(Pierre Auger Collaboration)[*]

[1]*Laboratório de Instrumentação e Física Experimental de Partículas—LIP and Instituto Superior Técnico—IST, Universidade de Lisboa—UL, Lisboa, Portugal*
[2]*Osservatorio Astrofisico di Torino (INAF), Torino, Italy*
[3]*INFN, Sezione di Torino, Torino, Italy*
[4]*University of Adelaide, Adelaide, S.A., Australia*
[5]*Centro Atómico Bariloche and Instituto Balseiro (CNEA-UNCuyo-CONICET), San Carlos de Bariloche, Argentina*
[6]*Institute of Nuclear Physics PAN, Krakow, Poland*
[7]*Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), Buenos Aires, Argentina*
[8]*Universidad Tecnológica Nacional—Facultad Regional Buenos Aires, Buenos Aires, Argentina*
[9]*Gran Sasso Science Institute, L'Aquila, Italy*
[10]*INFN Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila), Italy*
[11]*Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
[12]*IMAPP, Radboud University Nijmegen, Nijmegen, Netherlands*
[13]*Department of Physics and Astronomy, Lehman College, City University of New York, Bronx, New York, USA*
[14]*INFN, Sezione di Napoli, Napoli, Italy*
[15]*RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
[16]*Università Torino, Dipartimento di Fisica, Torino, Italy*
[17]*Universidad Michoacana de San Nicolás de Hidalgo, Morelia, Michoacán, México*
[18]*Observatorio Pierre Auger and Comisión Nacional de Energía Atómica, Malargüe, Argentina*
[19]*Università dell'Aquila, Dipartimento di Scienze Fisiche e Chimiche, L'Aquila, Italy*
[20]*University Politehnica of Bucharest, Bucharest, Romania*
[21]*Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic*
[22]*"Horia Hulubei" National Institute for Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
[23]*Universidad Nacional de San Agustin de Arequipa, Facultad de Ciencias Naturales y Formales, Arequipa, Peru*
[24]*Université Grenoble Alpes, CNRS, Grenoble Institute of Engineering Université Grenoble Alpes, LPSC-IN2P3, 38000 Grenoble, France*
[25]*Max-Planck-Institut für Radioastronomie, Bonn, Germany*
[26]*Instituto de Física de Rosario (IFIR)—CONICET/U.N.R. and Facultad de Ciencias Bioquímicas y Farmacéuticas U.N.R., Rosario, Argentina*
[27]*Karlsruhe Institute of Technology (KIT), Institute for Experimental Particle Physics, Karlsruhe, Germany*
[28]*Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*
[29]*Karlsruhe Institute of Technology (KIT), Institute for Astroparticle Physics, Karlsruhe, Germany*
[30]*International Center of Advanced Studies and Instituto de Ciencias Físicas, ECyT-UNSAM and CONICET, Campus Miguelete—San Martín, Buenos Aires, Argentina*
[31]*Universidade Federal do Rio de Janeiro, Instituto de Física, Rio de Janeiro, RJ, Brazil*
[32]*Universidade Estadual de Campinas, IFGW, Campinas, SP, Brazil*
[33]*Colorado State University, Fort Collins, Colorado, USA*
[34]*Universität Siegen, Department Physik—Experimentelle Teilchenphysik, Siegen, Germany*
[35]*Universidad de Granada and C.A.F.P.E., Granada, Spain*
[36]*Vrije Universiteit Brussels, Brussels, Belgium*
[37]*INFN, Sezione di Catania, Catania, Italy*
[38]*Universidad Autónoma de Chiapas, Tuxtla Gutiérrez, Chiapas, México*
[39]*Università di Milano, Dipartimento di Fisica, Milano, Italy*
[40]*INFN, Sezione di Milano, Milano, Italy*
[41]*Nationaal Instituut voor Kernfysica en Hoge Energie Fysica (NIKHEF), Science Park, Amsterdam, Netherlands*
[42]*Bergische Universität Wuppertal, Department of Physics, Wuppertal, Germany*
[43]*Università di Catania, Dipartimento di Fisica e Astronomia "Ettore Majorana," Catania, Italy*
[44]*Universidade de São Paulo, Escola de Engenharia de Lorena, Lorena, SP, Brazil*
[45]*INFN, Sezione di Lecce, Lecce, Italy*
[46]*Observatorio Pierre Auger, Malargüe, Argentina*

[47]Palacky University, RCPTM, Olomouc, Czech Republic
[48]Instituto de Tecnologías en Detección y Astropartículas (CNEA, CONICET, UNSAM), and Universidad Tecnológica Nacional—Facultad Regional Mendoza (CONICET/CNEA), Mendoza, Argentina
[49]Università di Napoli "Federico II," Dipartimento di Fisica "Ettore Pancini," Napoli, Italy
[50]University of Delaware, Department of Physics and Astronomy, Bartol Research Institute, Newark, Delaware, USA
[51]Politecnico di Milano, Dipartimento di Scienze e Tecnologie Aerospaziali, Milano, Italy
[52]Universidade Federal Fluminense, EEIMVR, Volta Redonda, RJ, Brazil
[53]Case Western Reserve University, Cleveland, Ohio, USA
[54]Instituto de Astronomía y Física del Espacio (IAFE, CONICET-UBA), Buenos Aires, Argentina
[55]Departamento de Física and Departamento de Ciencias de la Atmósfera y los Océanos, FCEyN, Universidad de Buenos Aires and CONICET, Buenos Aires, Argentina
[56]Universidade Federal do Rio de Janeiro (UFRJ), Observatório do Valongo, Rio de Janeiro, RJ, Brazil
[57]Instituto Federal de Educação, Ciência e Tecnologia do Rio de Janeiro (IFRJ), Brazil
[58]Università del Salento, Dipartimento di Matematica e Fisica "E. De Giorgi," Lecce, Italy
[59]Universidade de São Paulo, Instituto de Física de São Carlos, São Carlos, SP, Brazil
[60]Universidad Nacional Autónoma de México, México, D.F., México
[61]Universidade Federal do Paraná, Setor Palotina, Palotina, Brazil
[62]IFLP, Universidad Nacional de La Plata and CONICET, La Plata, Argentina
[63]Fermi National Accelerator Laboratory, Fermilab, Batavia, Illinois, USA
[64]Stichting Astronomisch Onderzoek in Nederland (ASTRON), Dwingeloo, Netherlands
[65]University of Chicago, Enrico Fermi Institute, Chicago, Illinois, USA
[66]New York University, New York, New York, USA
[67]Karlsruhe Institute of Technology (KIT), Institut für Prozessdatenverarbeitung und Elektronik, Karlsruhe, Germany
[68]Michigan Technological University, Houghton, Michigan, USA
[69]Experimental Particle Physics Department, J. Stefan Institute, Ljubljana, Slovenia
[70]Center for Astrophysics and Cosmology (CAC), University of Nova Gorica, Nova Gorica, Slovenia
[71]Present address: Hakubi Center for Advanced Research and Graduate School of Science, Kyoto University, Kyoto, Japan
[72]Université Libre de Bruxelles (ULB), Brussels, Belgium
[73]Universidade Estadual de Feira de Santana, Feira de Santana, Brazil
[74]Institute of Space Science, Bucharest-Magurele, Romania
[75]Colorado School of Mines, Golden, Colorado, USA
[76]Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Nova Friburgo, Brazil
[77]Universidade Federal do ABC, Santo André, SP, Brazil
[78]Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE), Sorbonne Université, Université de Paris, CNRS-IN2P3, Paris, France
[79]Benemérita Universidad Autónoma de Puebla, Puebla, México
[80]University of Wisconsin-Madison, Department of Physics and WIPAC, Madison, Wisconsin, USA
[81]Universität Hamburg, II. Institut für Theoretische Physik, Hamburg, Germany
[82]Università di Palermo, Dipartimento di Fisica e Chimica "E. Segrè," Palermo, Italy
[83]Louisiana State University, Baton Rouge, Louisiana, USA
[84]Università di Roma "Tor Vergata," Dipartimento di Fisica, Roma, Italy
[85]INFN, Sezione di Roma "Tor Vergata," Roma, Italy
[86]Pennsylvania State University, University Park, Pennsylvania, USA
[87]Centro de Investigaciones en Láseres y Aplicaciones, CITEDEF and CONICET, Villa Martelli, Argentina
[88]Charles University, Faculty of Mathematics and Physics, Institute of Particle and Nuclear Physics, Prague, Czech Republic
[89]Universidad Industrial de Santander, Bucaramanga, Colombia
[90]Unidad Profesional Interdisciplinaria en Ingeniería y Tecnologías Avanzadas del Instituto Politécnico Nacional (UPIITA-IPN), México, D.F., México
[91]Universidade de São Paulo, Instituto de Física, São Paulo, SP, Brazil
[92]Istituto di Astrofisica Spaziale e Fisica Cosmica di Palermo (INAF), Palermo, Italy
[93]Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, RJ, Brazil
[94]Also at University of Bucharest, Physics Department, Bucharest, Romania
[95]University of Łódź, Faculty of High-Energy Astrophysics, Łódź, Poland
[96]Universidad de Medellín, Medellín, Colombia
[97]Universiteit van Amsterdam, Faculty of Science, Amsterdam, Netherlands
[98]School of Physics and Astronomy, University of Leeds, Leeds, United Kingdom

(Received 9 August 2022; revised 30 November 2022; accepted 14 December 2022; published 7 February 2023)

Instantons, which are nonperturbative solutions to Yang-Mills equations, provide a signal for the occurrence of quantum tunneling between distinct classes of vacua. They can give rise to decays of particles

otherwise forbidden. Using data collected at the Pierre Auger Observatory, we search for signatures of such instanton-induced processes that would be suggestive of super-heavy particles decaying in the Galactic halo. These particles could have been produced during the post-inflationary epoch and match the relic abundance of dark matter inferred today. The nonobservation of the signatures searched for allows us to derive a bound on the reduced coupling constant of gauge interactions in the dark sector: $\alpha_X \lesssim 0.09$, for $10^9 \lesssim M_X/\mathrm{GeV} < 10^{19}$. Conversely, we obtain that, for instance, a reduced coupling constant $\alpha_X = 0.09$ excludes masses $M_X \gtrsim 3 \times 10^{13}$ GeV. In the context of dark matter production from gravitational interactions alone, we illustrate how these bounds are complementary to those obtained on the Hubble rate at the end of inflation from the nonobservation of tensor modes in the cosmological microwave background.

DOI: 10.1103/PhysRevLett.130.061001

Should a flux of astrophysical photons with energies in excess of $\simeq 10^8$ GeV be detected, it could be compelling evidence for the decay of super-heavy relics dating from the early universe [1,2]. Possible mechanisms taking place during or at the end of the inflationary era in big bang cosmology have been shown to be capable of producing such particles [3–14]. The abundance of the stable super-heavy particles could then evolve to match the relic abundance of dark matter (DM) inferred today, for viable parameters governing the thermal history and the geometry of the universe, such as the reheating temperature or the Hubble expansion rate at the end of inflation. Stability for super-heavy particles is more easily achieved for a dark sector interacting with the standard model (SM) sector only via gravity. The absence of other DM-SM couplings is consistent with the extensive observational evidence for the existence of DM based on gravitational effects alone. However, even particles protected from decay by a symmetry can eventually disintegrate due to nonperturbative effects in non-Abelian gauge theories and produce ultra-high energy (UHE) photons. In this Letter, we show that the absence of such photons in the data of the Pierre Auger Observatory provides constraints on the coupling constant of a hidden sector pertaining to super-heavy dark matter (SHDM), possibly unified with SM interactions at a high scale. The constraints are illustrated in Fig. 1 in terms of the reduced coupling constant of a hidden gauge interaction and the mass of the SHDM candidate. Our results show that the coupling should be less than $\simeq 0.09$ for a wide range of masses. After explaining how these constraints are obtained, we briefly discuss their relevance for delineating viable regions of cosmological parameters, in a manner complementary to the constraints provided by the non-detection so far of tensor modes in the cosmological microwave background anisotropies [15,16].

Contemporary motivations for SHDM.—Technical naturalness has provided an important motive for the emergence of new physics at the TeV scale [18], but the corresponding new particles have escaped detection so far [19–21]. An alternative tool to infer the energy scale of new physics relies on assessing the scale Λ_I at which the Higgs potential develops an instability at large field values. Its estimation at the two-loop level was made possible by the precise measurements of the Higgs mass and the top Yukawa coupling [22–24]. It turns out to result in a very high energy scale, $\Lambda_I = 10^{10}$ to 10^{12} GeV. Moreover, the particular slow running of the Callan-Symanzik β_λ function relative to the self-Higgs coupling makes it possible to extrapolate the SM up to M_{Pl} without encountering any instabilities [22]. Renouncing naturalness to solve the problem of the mass hierarchy, new degrees of freedom could thus appear only in the range between Λ_I and M_{Pl}, motivating searches for SHDM. We note also that while some have argued that the properties of nuclei and atoms would not allow complex chemistry if the electroweak scale were too far from the confinement scale of QCD [25], there is no such anthropic requirement for the mass scale of DM.

Independent of the intrinsic consistency of the SM up to a very high scale, models of gravitational production of



FIG. 1. Upper limits at 95% C.L. (Confidence Level) on the coupling constant α_X of a hidden gauge interaction as a function of the mass M_X of a dark matter particle decaying into a dozen $q\bar{q}$ pairs. For reference, the unification of the three SM gauge couplings is shown as the blue dashed line in the framework of supersymmetric grand unified theory [17].

DM provide another motivation for a spectrum of superheavy particles. While no coupling between the SM and DM sectors is introduced in the concordance model of cosmology, most DM models invoke some weak couplings, or new feeble couplings, to explain DM production during the postinflationary reheating period. It turns out, however, that the introduction of such couplings is not a compelling necessity if one considers the minimal assumption of graviton exchange to act as the only portal. Recent studies have indeed shown that, on the condition that DM is super heavy, the relic abundance observed today can be reproduced for tenable ranges of quantities governing the inflationary and reheating eras in the early Universe [9,14]. In addition, while structure formation constrains the mass density of DM, it does not preclude SHDM models as it leaves a *carte blanche* for the mass spectrum of the particles.

SHDM particles interacting with SM particles through gravitons alone have been dubbed as Planckian-interacting massive particles (PIDM) [9], and we shall use this term when we need to be specific. There are only a few possible signatures to test this scenario for DM. We show that if instanton effects are strong enough, PIDM particles could decay and their by-products could be detected in ultra-high energy cosmic ray (UHECR) data. Conversely, the nonobservation of these by-products allows us to set upper bounds on the dark sector coupling constant. We note that these limits are, to date, the best ever obtained from instanton-mediated processes; they are an indirect probe of the instanton strength.

Decay mechanisms of SHDM particles.—Some SHDM models postulate the existence of super-weak couplings between the dark and SM sectors. The lifetime τ_X of the particles is then governed by the strength of the couplings g_X and by the dimension n of the operator standing for the SM fields in the effective interaction [26]. This results in lifetimes that are in general far too short for DM to be stable enough, unless a practically untenable fine tuning between g_X and n holds [3,26,27]. Stability of super-heavy particles is thus preferentially calling for a new quantum number conserved in the dark sector so as to protect the particles from decaying. Nevertheless, as we have already pointed out in the study motivation, even stable particles in the perturbative domain will in general eventually decay due to nonperturbative effects in non-Abelian gauge theories [28–30].

Instanton-induced decay can thus make observable a dark sector that would otherwise be totally hidden by the conservation of a quantum number [31]. Assuming quarks and leptons carry this quantum number and so contribute to anomaly relationships with contributions from the dark sector, they will be by-products of decays together with the lightest hidden fermion. The lifetime of the decaying particle follows from Ref. [32],

$$\tau_X \simeq M_X^{-1} \exp\left(4\pi/\alpha_X\right), \tag{1}$$

with α_X being the reduced coupling constant of the hidden gauge interaction. In this expression, we retained only the exponential dependency in α_X^{-1}, dropping the functional determinants arising from the exact content of fields of the underlying theory. The constraints inferred using Eq. (1) are indeed barely destabilized for a wide range of numerical factors given the exponential dependency in α_X^{-1}. Equation (1) provides us with a relationship connecting the lifetime τ_X, which is shown below to be constrained by the absence of UHE photons, to the coupling constant α_X.

Production of ultrahigh energy photons.—In most SHDM models, the production of quark-antiquark pairs is expected in the decay by-products, giving rise to large fluxes of UHE particles such as nucleons, photons, and neutrinos. This is because each pair triggers a QCD cascade until the hadronization of the partons occurs and the unstable hadrons eventually decay. Various computational schemes have been applied to predict the energy spectra of the UHE particles [33–37]. The fragmentation of a parton into a hadron is determined from the fragmentation functions of partons convolved with hadronization functions, which do not depend on the scale M_X and can therefore be calculated from the available data. The fragmentation functions, on the other hand, are evolved starting from measurements at the electroweak scale up to the energy scale fixed by M_X using the Dokshitzer–Gribov–Lipatov–Altarelli–Parisi equation. We use the scheme detailed in Ref. [33] in this study. Overall, the spectra of the UHE particles is of the form $E^{-1.9}$ in the $q\bar{q}$ channel, and is barely softened by kinematical effects in large-multiplicity final states [34,38].

As shown below, it turns out to be more efficient to search for decaying super-heavy particles via UHE-photon by-products. Because of their attenuation over intergalactic distances, only those emitted in the Milky Way can survive on their way to Earth. The emission rate per unit volume and unit energy q_γ from any point labeled by its Galactic coordinates is shaped by the density of DM particles, $n_{\rm DM}$,

$$q_\gamma(E, \mathbf{x}_\odot + s\mathbf{n}) = \frac{1}{\tau_X}\frac{dN_\gamma}{dE} n_{\rm DM}(\mathbf{x}_\odot + s\mathbf{n}), \tag{2}$$

where τ_X is the lifetime of the particle, $\mathbf{x}_\odot$ is the position of the Solar System in the Galaxy, s is the distance from $\mathbf{x}_\odot$ to the emission point, and $\mathbf{n} \equiv \mathbf{n}(\ell, b)$ is a unit vector on the sphere pointing to the Galactic longitude ℓ and latitude b. Hereafter, the density is more conveniently expressed in terms of energy density $\rho_{\rm DM}(\mathbf{x}) = M_X n_{\rm DM}(\mathbf{x})$, normalized to $\rho(\mathbf{x}_\odot) = 0.3\ {\rm GeV\,cm^{-3}}$. There are uncertainties in the determination of this profile. We take as reference the traditional Navarro–Frenk–White profile [39]. We have checked that other profiles such as those from Einasto [40],

Burkert [41], or Moore [42] would lead to differences within 2% in the determination of α_X.

The expected flux (per steradian) of UHE photons produced by the decay of super-heavy particles, $J_{\mathrm{DM},\gamma}(E,\mathbf{n})$, is obtained by integrating the position-dependent emission rate q_γ along the path of the photons in the direction $\mathbf{n}$,

$$J_{\mathrm{DM},\gamma}(E,\mathbf{n}) = \frac{1}{4\pi}\int_0^\infty ds\, q_\gamma(E,\mathbf{x}_\odot + s\mathbf{n}), \qquad (3)$$

where the 4π normalization factor accounts for the isotropy of the decay processes. While the peak value of the flux is inversely proportional to the unknown M_X and τ_X parameters, the energy and directional dependencies are determined by Eq. (2). The exact content of quarks and leptons depends on the specific underlying model. Yet, instanton-induced decays obey selection rules that involve necessarily large multiplicities. As a proxy, we consider a dozen $q\bar{q}$ pairs and adapt dN_γ/dE in Eq. (2) accordingly [27]. The flux pattern on the sky is more intense in a hot-spot region around the Galactic center; it provides therefore clear signatures. On the other hand, the nonobservation of UHE photons enables one to constrain the all-sky flux observed over the solid angle $\Delta\Omega$, $\langle J_{\mathrm{DM},\gamma}(E,\mathbf{n})\rangle = \int_{\Delta\Omega} d\mathbf{n} J_{\mathrm{DM},\gamma}(E,\mathbf{n})/\Delta\Omega$, and thus to constrain the unknown M_X and τ_X parameters.

Constraints on dark-sector coupling constant from instanton-induced decays.—Of particular interest would thus be the detection of UHE photons from regions of denser DM density such as the center of our Galaxy. Because of the spectral steepness of the expected flux, this search can presently only be done through large ground-based detectors that exploit the phenomenon of extensive air showers. The identification of photon primaries relies on the ability to distinguish showers generated by photons from those initiated by the overwhelming background of protons and heavier nuclei. Since the radiation length in the atmosphere is more than 2 orders of magnitude smaller than the mean free path for photonuclear interactions, the transfer of energy to the hadron-muon channel is reduced in photon showers with respect to the bulk of hadron-induced showers, resulting in a lower number of secondary muons. Additionally, as the development of photon showers is delayed by the typically small multiplicity of electromagnetic interactions, they reach the maximum development of the shower, X_{max}, deeper in the atmosphere with respect to showers initiated by hadrons.

Both the ground signal and X_{max} can be measured at the Pierre Auger Observatory [43], where a hybrid detection technique is employed for the observation of extensive air showers by combining a fluorescence detector (FD) with a ground array of particle detectors [surface detector, (SD)] separated by 1500 m. The FD provides direct observation of the longitudinal shower profile, which allows for the measurement of the energy and the X_{max} of a shower, while the SD samples the secondary particles at ground level. Although showers are observed at a fixed slice in depth with the SD, the longitudinal development is embedded in the signals detected. The FD and SD are complemented with the low-energy enhancements of the Observatory, namely three additional fluorescence telescopes with an elevated field of view, overlooking a denser SD array, in which the stations are separated by 750 m. The combination of these instruments allows showers to be measured in the energy range above 10^8 GeV.



FIG. 2. Flux upper limits of UHE photons, neutrinos, and cosmic rays as a function of energy thresholds.

Three different analyses, differing in the detector used, have been developed to cover the wide energy range probed at the Observatory [44–46]. No photons with energies above 2×10^8 GeV have been unambiguously identified so far, leading to the 95% C.L. flux upper limits displayed in Fig. 2. The limit above $10^{11.2}$ GeV, stemming from the nondetection so far of any UHECR [47], including photons, is also constraining [37,48]. For comparison purposes, neutrino limits obtained at the Observatory [49] are also displayed as the continuous line. Indeed, neutrinos constitute in the figure another emblematic signature of decays of super-heavy particles. Except at the lowest energies, these limits are seen to be superseded by photon limits, as are those from anisotropy signatures searched for in the bulk of UHECR data shown as red-filled circles [27].

Assuming that the relic abundance of DM is saturated by super-heavy particles, constraints can be inferred in the plane (τ_X, M_X) by requiring the all-sky flux integrated above some energy threshold E to be less than the limits, $\int_E^\infty dE' \langle J_{\mathrm{DM},\gamma}(E',\mathbf{n})\rangle \leq J_\gamma^{95\%}(\geq E)$. For a specific upper limit at one energy threshold, a scan of the value of the mass M_X is carried out so as to infer a lower limit of the τ_X parameter, which is subsequently transformed into an upper limit on α_X by means of Eq. (1). This defines a curve. By repeating the procedure for each upper limit on $J_\gamma^{95\%}(\geq E)$,

a set of curves is obtained, reflecting the sensitivity of a specific energy threshold to some range of mass. The union of the excluded regions finally provides the constraints in the plane (α_X, M_X). In this manner the shaded red area is obtained in Fig. 1. As already noted, additional model-dependent factors could be at play in the vacuum transition amplitude [50] and thus in Eq. (1). Explicit constructions of the dark sector are required to calculate these factors. Such constructions are well beyond the scope of this Letter. Although the limits presented in Fig. 1 are hardly destabilized due to the exponential dependence in α_X^{-1}, we note that a shift of $\pm 0.0013k$ would arise for factors $10^{\pm k}$. We limit ourselves to showing with dotted and dashed lines the bounds that would be obtained for $k = 2$ and $k = 4$, respectively. These factors are by far the dominant systematic uncertainties.

Connection to cosmological scenarios.—We now briefly mention how the results shown in Fig. 1 can be connected to cosmological scenarios. Further details can be found in a companion paper [27]. In inflationary cosmologies, the inflaton field responsible for the rapid expansion of the Universe, ϕ, slowly rolls down to its minimum of potential before starting to oscillate about this minimum. This marks the end of the inflation era at time $H_{\rm inf}^{-1}$, with $H_{\rm inf}$ the Hubble rate at this time, and the beginning of a matter-dominated era during which the production of SM particles accompanying the decay of coherent oscillations of the inflaton field reheats the Universe. The temperature rises rapidly to a maximum before decreasing slowly until the reheating era ends at time Γ_ϕ^{-1}, marking the beginning of the radiation-dominated era when the temperature decreases more rapidly as a^{-1}, with a being the cosmological scale factor. The temperature at the end of reheating, $T_{\rm rh}$, is, together with $H_{\rm inf}$, an important parameter governing the dynamics of the reheating era summarized here—see Ref. [51] for details. A relevant combination of these parameters is the reheating efficiency ε, which, defined as $\varepsilon = (\Gamma_\phi/H_{\rm inf})^{1/2}$ [52], measures the duration of the reheating period. It can be related to $T_{\rm rh}$ and $H_{\rm inf}$ through $\varepsilon \simeq T_{\rm rh}/[0.25(M_{\rm Pl}H_{\rm inf})^{1/2}]$ [9].

PIDM particles can be produced during reheating by annihilation of SM particles [9] or inflaton particles [14] through gravitational interaction. The energy density of the Universe is then in the form of unstable inflaton particles, SM radiation, and stable massive particles, the time evolution of which is governed by a set of coupled Boltzmann equations [53]. However, because the energy density of the massive particles is always subdominant, the evolution of the inflationary and radiation energy densities largely decouple from the time evolution of the PIDM-particle density n_X. In addition, because PIDM particles interact through gravitation only, they never come to thermal equilibrium. In this case, the collision term in the Boltzmann equation can be approximated as a source term only,



FIG. 3. Constraints in the $(H_{\rm inf}, M_X)$ plane. The red region is excluded by the nonobservation of tensor modes in the cosmic microwave background [9,16]. The regions of viable $(H_{\rm inf}, M_X)$ values needed to set the right abundance of DM are delineated by the blue lines for different values of reheating efficiency ε [54]. Additional constraints from the nonobservation of instanton-induced decay of SHDM particles allow for excluding the mass ranges in the regions to the right of the vertical lines, for the specified values of the dark-sector gauge coupling.

$$\frac{dn_X(t)}{dt} + 3H(t)n_X(t) \simeq \sum_i \gamma_i \bar{n}_i^2(t). \tag{4}$$

Here, the sum on the right hand side represents the contributions from the SM and inflationary sectors. Using, on the one hand, the evolution of the SM matter and inflaton densities derived in Ref. [5] and Ref. [14] respectively, and, on the other hand, the SM + SM → PIDM + PIDM and $\phi + \phi \to$ PIDM + PIDM reaction rates derived for fermionic DM in Ref. [54] and Ref. [14] respectively, the present-day relic abundance of DM, Ω_X, can be related to M_X, $H_{\rm inf}$, and ε through

$$\Omega_X h^2 = \frac{1.4 \times 10^{23} \varepsilon M_X}{M_{\rm Pl}^{5/2} H_{\rm inf}^{3/2}} \int_{a_{\rm inf}}^{\infty} \frac{da}{H(a)} \sum_i \gamma_i a^2 \bar{n}_i^2(a), \tag{5}$$

where h is the dimensionless expansion rate and $a_{\rm inf}$ the scale factor at the end of inflation.

The viable $(H_{\rm inf}, M_X)$ parameter space satisfying Eq. (5) is delineated by the blue curves corresponding to different values of ε in Fig. 3. Values for $(H_{\rm inf}, M_X)$ above (below) the lines lead to overabundance of (negligible quantity of) DM. Arbitrarily large values of $H_{\rm inf}$ are however not permitted because of the 95% C.L. on the tensor-to-scalar ratio in the cosmic microwave background anisotropies, which, once converted into limits on the energy scale of inflation when the pivot scale exits the Hubble radius, yield $H_{\rm inf} \leq 4.9 \times 10^{-6} M_{\rm Pl}$ [9,16]. For efficiencies larger than a few percent, PIDM particles are dominantly produced by

the thermal bath of SM particles. A wide range of masses M_X is then allowed, including the Grand Unified scale, provided that the energy scale of the inflation ($H_{\rm inf}$ being the proxy) is high enough [9] and that the dark-sector gauge coupling α_X is less than $\simeq 0.085$. Larger values of α_X shrink the allowed range of M_X, with, for instance, $M_X \lesssim 2 \times 10^9$ GeV for $\alpha_X = 0.1$. For efficiencies below the percent level, the production of PIDM particles from the inflaton condensate dominates, allowing smaller values of $T_{\rm rh}$ to be viable. The allowed region for M_X shrinks around 10^{13} GeV, close to the inflaton mass adopted here (3×10^{13} GeV). We see that the scenario is then tenable for $\alpha_X \lesssim 0.09$.

In summary, we have illustrated here the power of upper limits on the flux of UHE photons obtained at the Pierre Auger Observatory to place constraints on physics in the reheating epoch that could be related to Grand Unified models. The minimal setup to produce DM is from gravitational effects alone, consistent with the concordance model of cosmology. This production mechanism could lead to high values of the Hubble rate at the end of inflation that could be revealed by future measurements of primordial tensor-to-scalar ratio provided that $H_{\rm inf} \gtrsim 6 \times 10^{12}$ GeV [55,56]. However, the only unambiguous signature to capture the existence of PIDM is through the detection of UHE photons produced by the instanton-induced decay. The nonobservation of such fluxes has allowed us to probe in a unique way to date the instanton strength through the dark-sector gauge coupling. It is likely that the use of limits on UHE photon fluxes made in this Letter only scratches the surface of the power of these limits to constrain physics otherwise beyond the reach of laboratory experiments.

The successful installation, commissioning, and operation of the Pierre Auger Observatory would not have been possible without the strong commitment and effort from the technical and administrative staff in Malargüe. We are very grateful to the following agencies and organizations for financial support: Argentina—Comisión Nacional de Energía Atómica; Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET); Gobierno de la Provincia de Mendoza; Municipalidad de Malargüe; NDM Holdings and Valle Las Leñas; in gratitude for their continuing cooperation over land access; Australia—the Australian Research Council; Belgium—Fonds de la Recherche Scientifique (FNRS); Research Foundation Flanders (FWO); Brazil—Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); Financiadora de Estudos e Projetos (FINEP); Fundação de Amparo à Pesquisa do Estado de Rio de Janeiro (FAPERJ); São Paulo Research Foundation (FAPESP) Grants No. 2019/10151-2, No. 2010/07359-6, and No. 1999/05404-3; Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTIC); Czech Republic—Grants No. MSMT CR LTT18004, No. LM2015038, No. LM2018102, No. CZ.02.1.01/0.0/0.0/16_013/0001402, No. CZ.02.1.01/0.0/0.0/18_046/0016010, and No. CZ.02.1.01/0.0/0.0/17_049/0008422; France—Centre de Calcul IN2P3/CNRS; Centre National de la Recherche Scientifique (CNRS); Conseil Régional Ile-de-France; Département Physique Nucléaire et Corpusculaire (PNC-IN2P3/CNRS); Département Sciences de l'Univers (SDU-INSU/CNRS); Institut Lagrange de Paris (ILP) Grant No. LABEX ANR-10-LABX-63 within the Investissements d'Avenir Programme Grant No. ANR-11-IDEX-0004-02; Germany—Bundesministerium für Bildung und Forschung (BMBF); Deutsche Forschungsgemeinschaft (DFG); Finanzministerium Baden-Württemberg; Helmholtz Alliance for Astroparticle Physics (HAP); Helmholtz-Gemeinschaft Deutscher Forschungszentren (HGF); Ministerium für Innovation, Wissenschaft und Forschung des Landes Nordrhein-Westfalen; Ministerium für Wissenschaft, Forschung und Kunst des Landes Baden-Württemberg; Italy—Istituto Nazionale di Fisica Nucleare (INFN); Istituto Nazionale di Astrofisica (INAF); Ministero dell'Istruzione, dell'Universitá e della Ricerca (MIUR); CETEMPS Center of Excellence; Ministero degli Affari Esteri (MAE); México—Consejo Nacional de Ciencia y Tecnología (CONACYT) No. 167733; Universidad Nacional Autónoma de México (UNAM); PAPIIT DGAPA-UNAM; The Netherlands—Ministry of Education, Culture and Science; Netherlands Organisation for Scientific Research (NWO); Dutch national e-infrastructure with the support of SURF Cooperative; Poland—Ministry of Education and Science, Grant No. DIR/WK/2018/11; National Science Centre, Grants No. 2016/22/M/ST9/00198, No. 2016/23/B/ST9/01635, and No. 2020/39/B/ST9/01398; Portugal—Portuguese national funds and FEDER funds within Programa Operacional Factores de Competitividade through Fundação para a Ciência e a Tecnologia (COMPETE); Romania—Ministry of Research, Innovation and Digitization, CNCS/CCCDI—UEFISCDI, Projects No. PN19150201/16N/2019, No. PN1906010, No. TE128, and No. PED289, within PNCDI III; Slovenia—Slovenian Research Agency, Grants No. P1-0031, No. P1-0385, I0-0033, No. N1-0111; Spain—Ministerio de Economía, Industria y Competitividad (FPA2017-85114-P and PID2019–104676 GB-C32), Xunta de Galicia (ED431C 2017/07), Junta de Andalucía (SOMM17/6104/UGR, P18-FR-4314) Feder Funds, RENATA Red Nacional Temática de Astropartículas (FPA2015-68783-REDT) and María de Maeztu Unit of Excellence (MDM-2016-0692); USA—Department of Energy, Contracts No. DE-AC02-07CH11359, No. DE-FR02-04ER41300, No. DE-FG02-99ER41107, and No. DE-SC0011689; National Science

Foundation, Grant No. 0450696; The Grainger Foundation; Marie Curie-IRSES/EPLANET; European Particle Physics Latin American Network; and UNESCO. We acknowledge for this work the support of the Institut Pascal at Université Paris-Saclay during the Paris-Saclay Astroparticle Symposium 2021, with the support of the P2IO Laboratory of Excellence (program "Investissements d'avenir" ANR-11-IDEX-0003-01 Paris-Saclay and ANR-10-LABX-0038), the P2I axis of the Graduate School Physics of Université Paris-Saclay, as well as IJCLab, CEA, IPhT, APPEC, the IN2P3 master projet UCMN and EuCAPT ANR-11-IDEX-0003-01 Paris-Saclay and ANR-10-LABX-0038).

*spokespersons@auger.org

[1] P. Bhattacharjee and G. Sigl, Phys. Rep. **327**, 109 (2000).
[2] L. A. Anchordoqui, Phys. Rep. **801**, 1 (2019).
[3] J. R. Ellis, G. Gelmini, J. L. Lopez, D. V. Nanopoulos, and S. Sarkar, Nucl. Phys. **B373**, 399 (1992).
[4] V. Berezinsky, M. Kachelrieß, and A. Vilenkin, Phys. Rev. Lett. **79**, 4302 (1997).
[5] D. J. H. Chung, E. W. Kolb, and A. Riotto, Phys. Rev. D **59**, 023501 (1998).
[6] V. Kuzmin and I. Tkachev, Phys. Rev. D **59**, 123006 (1999).
[7] D. J. H. Chung, E. W. Kolb, A. Riotto, and I. I. Tkachev, Phys. Rev. D **62**, 043508 (2000).
[8] M. A. Fedderke, E. W. Kolb, and M. Wyman, Phys. Rev. D **91**, 063505 (2015).
[9] M. Garny, M. C. Sandora, and M. S. Sloth, Phys. Rev. Lett. **116**, 101302 (2016).
[10] J. Ellis, M. A. G. Garcia, D. V. Nanopoulos, K. A. Olive, and M. Peloso, J. Cosmol. Astropart. Phys. 03 (2016) 008.
[11] E. W. Kolb and A. J. Long, Phys. Rev. D **96**, 103540 (2017).
[12] E. Dudas, Y. Mambrini, and K. A. Olive, Phys. Rev. Lett. **119**, 051801 (2017).
[13] K. Kaneta, Y. Mambrini, and K. A. Olive, Phys. Rev. D **99**, 063508 (2019).
[14] Y. Mambrini and K. A. Olive, Phys. Rev. D **103**, 115009 (2021).
[15] P. A. R. Ade *et al.* (BICEP2 and Planck Collaborations), Phys. Rev. Lett. **114**, 101301 (2015).
[16] P. A. R. Ade *et al.* (Planck Collaboration), Astron. Astrophys. **594**, A20 (2016).
[17] P. A. Zyla *et al.* (Particle Data Group), Prog. Theor. Exp. Phys. **2020**, 083C01 (2020).
[18] G. 't Hooft, NATO Sci. Ser. B **59**, 135 (1980).
[19] T. Marrodán Undagoitia and L. Rauch, J. Phys. G **43**, 013001 (2016).
[20] S. Rappoccio, Rev. Phys. **4**, 100027 (2019).
[21] J. M. Gaskins, Contemp. Phys. **57**, 496 (2016).
[22] D. Buttazzo, G. Degrassi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio, and A. Strumia, J. High Energy Phys. 12 (2013) 089.
[23] S. Alekhin, A. Djouadi, and S. Moch, Phys. Lett. B **716**, 214 (2012).
[24] A. V. Bednyakov, B. A. Kniehl, A. F. Pikelner, and O. L. Veretin, Phys. Rev. Lett. **115**, 201802 (2015).
[25] T. Damour and J. F. Donoghue, Phys. Rev. D **78**, 014014 (2008).
[26] H. J. de Vega and N. G. Sanchez, Phys. Rev. D **67**, 125019 (2003).
[27] P. Abreu *et al.* (Pierre Auger Collaboration), companion paper, Phys. Rev. D **107**, 042002 (2023).
[28] A. A. Belavin, A. M. Polyakov, A. S. Schwartz, and Y. S. Tyupkin, Phys. Lett. B **59**, 85 (1975).
[29] S. R. Coleman, Subnucl. Ser. **15**, 805 (1979).
[30] A. I. Vainshtein, V. I. Zakharov, V. A. Novikov, and M. A. Shifman, Sov. Phys. Usp. **25**, 195 (1982).
[31] V. A. Kuzmin and V. A. Rubakov, Phys. At. Nucl. **61**, 1028 (1998).
[32] G. 't Hooft, Phys. Rev. Lett. **37**, 8 (1976).
[33] R. Aloisio, V. Berezinsky, and M. Kachelriess, Phys. Rev. D **69**, 094023 (2004).
[34] S. Sarkar and R. Toldra, Nucl. Phys. **B621**, 495 (2002).
[35] C. Barbot and M. Drees, Astropart. Phys. **20**, 5 (2003).
[36] M. Kachelriess, O. E. Kalashev, and M. Y. Kuznetsov, Phys. Rev. D **98**, 083016 (2018).
[37] E. Alcantara, L. A. Anchordoqui, and J. F. Soriano, Phys. Rev. D **99**, 103016 (2019).
[38] C. Barbot, M. Drees, F. Halzen, and D. Hooper, Phys. Lett. B **563**, 132 (2003).
[39] J. F. Navarro, C. S. Frenk, and S. D. M. White, Astrophys. J. **462**, 563 (1996).
[40] J. Einasto, Tr. Astrofiz. Inst. Alma-Ata **5**, 87 (1965).
[41] A. Burkert, Astrophys. J. Lett. **447**, L25 (1995).
[42] B. Moore, S. Ghigna, F. Governato, G. Lake, T. R. Quinn, J. Stadel, and P. Tozzi, Astrophys. J. Lett. **524**, L19 (1999).
[43] A. Aab *et al.* (Pierre Auger Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **798**, 172 (2015).
[44] P. Abreu *et al.* (Pierre Auger Collaboration), Astrophys. J. **933**, 125 (2022).
[45] P. Savina (Pierre Auger Collaboration), Proc. Sci., (ICRC2021) (**2021**) 373.
[46] P. Abreu *et al.* (Pierre Auger Collaboration), arXiv: 2209.05926.
[47] A. Aab *et al.* (Pierre Auger Collaboration), Phys. Rev. D **102**, 062005 (2020).
[48] L. A. Anchordoqui *et al.*, Astropart. Phys. **132**, 102614 (2021).
[49] A. Aab *et al.* (Pierre Auger Collaboration), J. Cosmol. Astropart. Phys. 10 (2019) 022.
[50] G. 't Hooft, Phys. Rev. D **14**, 3432 (1976); **18**, 2199(E) (1978).
[51] G. F. Giudice, E. W. Kolb, and A. Riotto, Phys. Rev. D **64**, 023508 (2001).
[52] E. W. Kolb and M. S. Turner, *The Early Universe* (CRC Press, Boca Raton, 1990), Vol. 69.
[53] D. J. H. Chung, E. W. Kolb, and A. Riotto, Phys. Rev. D **60**, 063504 (1999).
[54] M. Garny, A. Palessandro, M. Sandora, and M. S. Sloth, J. Cosmol. Astropart. Phys. 02 (2018) 027.
[55] K. N. Abazajian *et al.* (CMB-S4 Collaboration), arXiv:1610 .02743.
[56] H. Ishino, Int. J. Mod. Phys. Conf. Ser. **43**, 1660192 (2016).