



UvA-DARE (Digital Academic Repository)

Interventions aimed at automatic processes in addiction: considering necessary conditions for efficacy

Gladwin, T.E.; Wiers, C.E.; Wiers, R.W.

DOI

[10.1016/j.cobeha.2016.08.001](https://doi.org/10.1016/j.cobeha.2016.08.001)

Publication date

2017

Document Version

Final published version

Published in

Current Opinion in Behavioral Sciences

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Gladwin, T. E., Wiers, C. E., & Wiers, R. W. (2017). Interventions aimed at automatic processes in addiction: considering necessary conditions for efficacy. *Current Opinion in Behavioral Sciences*, 13, 19-24. <https://doi.org/10.1016/j.cobeha.2016.08.001>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Interventions aimed at automatic processes in addiction: considering necessary conditions for efficacy

Thomas E Gladwin^{1,2}, Corinde E Wiers³ and Reinout W Wiers¹



Automatic processes related to addiction can be directly targeted in novel training paradigms. First studies have demonstrated that Cognitive Bias Modification (CBM) targeting approach biases can enhance treatment outcomes when added to regular treatment. However, the overall efficacy of CBM is debated. We argue that considering the modulating role of motivation and the mediating role of actual bias change are essential to drawing valid conclusions. Findings on mediating cognitive and neural mechanisms underlying clinical effects provide further sources of evidence on CBM. Taken together the literature supports the claim that cognitive bias change can improve clinical outcome, but that there are necessary conditions that must be met. Improved theoretical understanding of changing biases and new techniques such as neuromodulation may be needed to optimally apply CBM to help patients overcome addictive behavior.

Addresses

¹ Addiction, Development and Psychopathology (ADAPT)-lab, Department of Psychology, Amsterdam Brain & Cognition (ABC), Research Priority Area Yield, University of Amsterdam, The Netherlands
² Research Centre – Military Mental Health, Ministry of Defense, Utrecht, The Netherlands
³ National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda, MD, USA

Corresponding author: Wiers, Reinout W (r.wiers@uva.nl)

Current Opinion in Behavioral Sciences 2017, 13:19–24

This review comes from a themed issue on **Addiction**

Edited by **Scott Edwards** and **Karen D Ersche**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 18th August 2016

<http://dx.doi.org/10.1016/j.cobeha.2016.08.001>

2352-1546/© 2016 Elsevier Ltd. All rights reserved.

Introduction

Theories of addiction range from considering it a moral failure [1,2*] to a brain disease [3,4]. Essentially, the role of agency varies in these accounts [5]. Does one choose to use drugs, despite personally and socially harmful consequences, for selfish reasons? Or is the brain of an addict compromised, rendering them unable to control their drug seeking and consumption? Such contradictory views are a problem, as they suggest quite different kinds of intervention. Dual-process models of addiction [6,7] resolve this conflict by understanding choice itself as an

emergent brain function — the outcome of interactions between fast automatic processes underlying impulsive behavior versus slow reflective processing allowing self-regulation [8,9]. Neurobiological consequences of drug use [10] may well lead to involuntary automatic responses involving drug cues, but the ultimate impact of these effects on behavior is modulated by higher-order reflective cognition. Such models also have implications for interventions. First and foremost, they suggest the use of techniques collectively termed Cognitive Bias Modification (CBM).

CBM for addiction

A cognitive bias is a difference in the automatic processes evoked by a certain class of stimuli. Such biases can be related to a psychiatric disorder, such as anxiety [11] or addiction [12]; motivational states such as hunger [13]; training [14]; or expertise and interest (e.g., car-lovers quickly spot a rare sports-car). Importantly, in relation to psychiatric disorders, these biases contribute to the etiology and maintenance of the disorder [15]. An illustration of this is a problem drinker who may have good intentions to cut down drinking for various long-term reasons, such as health and relationship problems, but a conditioned alcohol-related stimulus is likely to catch the eye (attentional bias), bring to mind tempting memories of fun or reduced stress (association activation), and increase the likelihood of actions aimed at alcohol consumption (approach bias) [16]. Hence, these biases may contribute to continued drinking or other addictive behaviors despite explicit intentions. Where such automatic processes play a role, CBM may help patients to remain abstinent [17].

The bias that has shown the most promising results in CBM is the tendency to approach addiction-related stimuli. Wiers, Rinck and colleagues first developed a task to indirectly assess this bias [18], in which heavy drinkers showed an approach bias for alcohol-related stimuli, moderated by the presence of a g-allele in the OPRM1 gene (previously related to cue-induced craving). Participants were requested to pull or push a joystick in response to the format of a picture (e.g., push landscape pictures, pull portrait pictures), which was followed by a zoom-mechanism contingent upon the movement (zoom in after pushing the joystick, zoom out after pulling), creating a sense of avoidance or approach, respectively. A first proof-of-concept study in students demonstrated that this bias could be changed, with generalization to untrained pictures and to drinking behavior in a taste-test [19]. In a first randomized clinical trial (RCT), patients

received four sessions of approach-bias retraining or placebo-training (continued assessment) or no training. Training changed the approach-bias more strongly than the control conditions, with generalization to untrained pictures and related verbal memory-associations. Moreover, trained patients were less likely to have relapsed a year after treatment discharge [20^{••}]. In a recent large RCT, this effect on relapse was replicated, and was found to be mediated by the change in approach-tendencies [21[•]]. A very recent study has reported positive findings for approach bias retraining during detox [22].

Another common target for CBM is the attentional bias, which can be targeted for retraining using similar variations of assessment tasks such as the dot-probe task [23]. A first small clinical Randomized Controlled Trial has provided preliminary evidence that this may be effective in alcoholic patients [24] and a larger replication is underway. Other methods have shown promise in changing biases in heavy drinking students; for example, evaluative conditioning [25], interpretation bias modification [26], and response inhibition training [27]. However, clinical effects of these variants of CBM remain to be established. Further, many published results in healthy subjects involve exploratory analyses, or some degree of uncontrolled multiple testing [28], and require replication.

Working memory training for addiction

Although CBM for addiction is aimed at changing automatic processes, dual-process models suggest that improving cognitive control might also help individuals to regulate their behavior in the face of dysfunctional biases [29]. Working memory training has been applied with some success to addiction [30,31,32[•]]. The first study [30] found reductions in alcohol consumption in problem drinkers, but only in those who demonstrated relatively strong positive alcohol associations. Bickel and colleagues [31,32[•]] found that working memory training in stimulant addicted patients has an effect on delay discounting, a facet of impulsivity associated with addiction. However, results of cognitive control training in addiction are quite preliminary at this stage. We briefly note that game-versions are being developed to reduce the problem of working memory training being extensive and boring, but it is not yet clear how well such approaches will work [33[•]].

The efficacy debate: motivation, mediation and variations

Various papers have argued that the evidence for CBM, or specific CBM variants, is weak [34,35]. Responses to this claim [17,36^{••}] have made two particular important counter-arguments, both of which are in line with our theoretical basis in dual-process models.

First is that CBM can only be evaluated in the context of effective motivation: Without adequate motivation-to-change and associated goals and strategies at the level

of reflective processing, no effect of CBM can be expected. Note that it is a crucial ingredient of many interventions in addiction to help people develop a long-term perspective, including personal goals that conflict with continuation of the addictive behavior [37,38]. This is exemplified in the motivational interviewing approach, a therapeutic technique originally developed by Bill Miller in the context of addiction treatment, and now applied more widely [39]. For example, a recent study found initial evidence that change talk — speech expressing motivation to reduce usage — in adolescents (elicited with motivational interviewing techniques) and its neural correlates predicted prospective marijuana use [40]. Note that in the RCTs for CBM discussed above [20^{••},21[•],24], training was added to regular treatment, in which motivation to change is enhanced and the patient is helped to develop an alternative long-term perspective. In line with the idea that motivation is a necessary condition for a positive outcome, CBM has demonstrated no effect in participants who were not motivated to change [41], self-selected problem drinkers with uncertain motivation to change [42] (although all groups including the control group did reduce their drinking), or community smokers not wanting to quit smoking [43[•]]. Promisingly, however, a recent online study on smoking only required smokers to make an actual quit-attempt to be included and found increased chance of abstinence half a year later (in heavy smokers only) in the training group compared to the control group [44[•]].

Second, if the rationale for CBM is correct then clinical change is mediated by changes in bias. If an intervention fails to change the targeted bias, then a lack of clinical effects cannot be taken as evidence against CBM. The only conclusion that can be drawn is that the particular method did not, in fact, modify a cognitive bias. The pattern of results over studies shows that studies with clinically relevant effects almost always show a change in cognitive bias, while negative studies do not [36^{••},45]. This pattern is just what would be expected if cognitive biases play a causal role in outcome. Note that various mediating biases have been found for the approach bias retraining [20^{••},21[•],46[•]].

Analyses that fail to consider these factors might lead to invalid or less relevant conclusions. A more general point about pooling results is that the existence of some effective forms of CBM does not imply that every attempt is likely to be successful. It may be more useful at this stage to try to answer less generalizing questions such as: What makes which methods more or less effective; which conditions are necessary; and what are the mechanisms underlying effects?

New developments in theory and methods

We are only beginning to learn about the underlying neurocognitive mechanisms of CBM [47]. Regarding

cognitive mechanisms, clinical effects in approach-bias retraining are mediated by alcohol-related action tendencies and avoidance associations [21*,46*], and it was also found that patients with a strong approach bias profited most from CBM as a supplement to therapy [21*]. Only a few studies have been published on the neural mechanisms of CBM in addiction. Corinde Wiers and colleagues [48**] found that alcohol approach-bias retraining CBM reduced alcohol cue-evoked activation in amygdala and nucleus accumbens in a passive viewing task. Reduction in amygdala activation was associated with reduction in craving within the CBM group. In an approach-bias assessment task in the scanner, CBM was found to decrease medial prefrontal cortex activation [49*], an area previously related to an alcohol-approach bias [50,51]. Note that such results provide convergent evidence that CBM can affect at least certain neurocognitive processes.

Understanding these mechanisms depends on our theoretical foundations. Recent theoretical criticisms of dual-process models are therefore highly relevant to CBM. Models that posit or imply dual systems have been argued to have theoretical problems, as arguments for clearly distinguishable systems or types of process appear to be flawed [52]. For instance, there does not appear to be a neurobiological basis for dual systems despite interesting early models [53]; for example, subcortical regions attributed to the reflexive system also play a role in working memory [54]. Further, systems with disjunctive sets of functions run into difficulties when, logically, these functions must be integrated and work together somehow. If 'hot' motivational and emotional functions are assigned to an impulsive system, for example, then a problem termed the motivational homunculus occurs: Why does the reflective system exert control in line with the individual's goals and motivation? One solution to such conceptual problems was proposed via the R³ model [8,55], which focuses on generalized response selection mechanisms, emphasizing iterative processes with varying temporal dynamics [56], consideration of the levels of description of different concepts, and the reinforcement of covert cognitive functions [57]. The model supplies a proof-of-principle definition of a continuum of automatic versus reflective processing, in terms of how much evaluation precedes selection. This avoids the above theoretical problems while retaining the useful heuristic of automatic versus controlled processes and relatively impulsive versus reflective behavior. Further, more specific research continues on automatic processes and implicit measures involved in addiction. For instance, temporal dynamics [58,59] and trial-to-trial fluctuations [60,61] appear to play a role in attentional biases. Such research is particularly relevant in terms of finding novel targets for CBM, although current methods appear to remain focused on simple biases.

An improved grasp of neurocognitive mechanisms in CBM may also inform the use of neuromodulation

methods such as transcranial Direct Current Stimulation (tDCS) or repetitive Transcranial Magnetic Stimulation (rTMS) [62]. Although generalizations about its efficacy are debated [63*,64], prefrontal tDCS has been found to improve working memory [65,66], facilitate the training of cognitive skills [66,67] and reduce cue-evoked craving for various addictive substances [68–70]. This suggests various mechanisms by which tDCS could interact with CBM. The cognitive state of participants could be made more amenable to addiction-related training, either via reflective (improved working memory) or impulsive (reduced craving) pathways, or general neural learning mechanisms could be modulated [71]. Two attentional bias modification studies have indeed shown such enhancement concerning attention to threatening stimuli [72*,73]. Analogous applications of tDCS in addiction CBM are to our knowledge yet to be published, but in our lab studies are underway with promising results. Note that the essential point here is that tDCS by itself is not expected to have persistent effects: It is expected to enhance training effects.

Finally, it is an interesting question to what extent treatment can be provided with a minimal face-to-face protocol [74], or perhaps even automated and online, which has proved to reduce problem drinking by itself [75]. Such interventions could easily be supplemented with online training [76]. Rapidly developing technological possibilities will permit further tailoring of cognitive training to the individual, making it more personally salient, and permit mobile interventions that occur in situations relevant to the bias being trained [28].

Conclusions

Targeting automatic processes may be an important piece of the puzzle of effective treatment for addiction, and complementary methods focused specifically on these processes could improve outcome for some patients. Training and neuromodulation may reduce impulsive tendencies or help patients to overcome them, but alternative choices and long-term perspectives that are incompatible with continuation of the addictive behavior appear necessary to profit from training effects [77*,78]. Further, a CBM intervention must actually change a proximal bias to be expected to have distal effects on clinical outcome. Much work remains to be done to determine the most relevant biases to target, and the most effective training tasks for those biases. Research therefore needs a healthy combination of exploratory studies, aimed at finding relationships between biases and symptoms and between training variants and changes in biases; confirmatory RCTs testing clinical effects of the most promising variants; and meta-analyses on appropriate subsets of confirmatory studies, taking into account theoretical considerations.

Conflicts of interest

Nothing declared.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Dalrymple T: **Doctors, lies and the addiction bureaucracy [Internet]**. *J Health Serv Res Policy* 2008, **13**:124-126.
 2. Hellman M, Berridge V, Duke KL, Mold A: *Concepts of Addictive Substances and Behaviours Across Time and Place*. Oxford University Press; 2015.
- A recent, comprehensive historical and cross-cultural overview of concepts related to addiction.
3. Leshner AI: **Addiction is a brain disease, and it matters [Internet]**. *Science* 1997, **278**:45-47.
 4. Volkow ND, Koob GF, McLellan AT: **Neurobiologic advances from the brain disease model of addiction**. *N Engl J Med* 2016, **374**:363-371.
 5. Levy N: **Addiction is not a brain disease (and it matters) [Internet]**. *Front Psychiatry* 2013, **4**:24.
 6. Bechara A: **Decision making, impulse control and loss of willpower to resist drugs: a neurocognitive perspective [Internet]**. *Nat Neurosci* 2005, **8**:1458-1463.
 7. Wiers RW, Bartholow BD, van Den Wildenberg E, Thush C, Engels RC, Sher KJ, Grenard J, Ames SL, Stacy AW: **Automatic and controlled processes and the development of addictive behaviors in adolescents: a review and a model [Internet]**. *Pharmacol Biochem Behav* 2007, **86**:263-283.
 8. Gladwin TE, Figner B, Crone EA, Wiers RW: **Addiction, adolescence, and the integration of control and motivation [Internet]**. *Dev Cogn Neurosci* 2011, **1**:364-376.
 9. Smith E, DeCoster J: **Dual-process models in social and cognitive psychology: conceptual integration and links to underlying memory systems [Internet]**. *Personal Soc Psychol Rev* 2000, **4**:108-131.
 10. Robinson TE, Berridge KC: **Review. The incentive sensitization theory of addiction: some current issues. [Internet]**. *Philos Trans R Soc Lond B Biol Sci* 2008, **363**:3137-3146.
 11. Cisler JM, Koster EHW: **Mechanisms of attentional biases towards threat in anxiety disorders: an integrative review [Internet]**. *Clin Psychol Rev* 2010, **30**:203-216.
 12. Field M, Cox WM: **Attentional bias in addictive behaviors: a review of its development, causes, and consequences [Internet]**. *Drug Alcohol Depend* 2008, **97**:1-20.
 13. Loeber S, Grosshans M, Herpertz S, Kiefer F, Herpertz SC: **Hunger modulates behavioral disinhibition and attention allocation to food-associated cues in normal-weight controls**. *Appetite* 2013, **71**:32-39.
 14. Shiffrin RM, Schneider W: **Controlled and automatic human information processing: II. Perceptual learning, automatic attending and a general theory. [Internet]**. *Psychol Rev* 1977, **84**:127-190.
 15. Stacy AW, Wiers RW: **Implicit cognition and addiction: a tool for explaining paradoxical behavior [Internet]**. *Annu Rev Clin Psychol* 2010, **6**:551-575.
 16. Wiers RW, Stacy AW: **Implicit cognition and addiction**. *Curr Dir Psychol Sci* 2006, **15**:292-296.
 17. Gladwin TE, Wiers CE, Wiers RW: **Cognitive neuroscience of cognitive retraining for addiction medicine: from mediating mechanisms to questions of efficacy [Internet]**. *Progr Brain Res* 2016, **224**:323-344.
 18. Wiers RW, Rinck M, Dictus M, van Den Wildenberg E: **Relatively strong automatic appetitive action-tendencies in male carriers of the OPRM1 G-allele [Internet]**. *Genes Brain Behav* 2009, **8**:101-106.
 19. Wiers RW, Rinck M, Kordts R, Houben K, Strack F: **Retraining automatic action-tendencies to approach alcohol in hazardous drinkers [Internet]**. *Addiction* 2010, **105**:279-287.
 20. Wiers RW, Eberl C, Rinck M, Becker ES, Lindenmeyer J: **Retraining automatic action tendencies changes alcoholic patients' approach bias for alcohol and improves treatment outcome**. *Psychol Sci* 2011, **22**:490-497.
- First RCT showing that CBM can have long-term clinical effects for alcohol-dependent patients when added to regular treatment.
21. Eberl C, Wiers RW, Pawelczack S, Rinck M, Becker ES, Lindenmeyer J: **Approach bias modification in alcohol dependence: do clinical effects replicate and for whom does it work best? Dev Cogn Neurosci** 2013, **4**:38-51.
- Large replication study of [25] showing that the long-term clinical effect is mediated by the change in approach-bias.
22. Manning V, Staiger P, Hall K, Garfield J, Flaks G, Leung D, Hughes L, Lum J, Lubman D, Verdejo-García A: **Cognitive bias modification training during inpatient alcohol detoxification reduces early relapse: a randomized controlled trial**. *Alcohol Clin Exp Res* 2016 <http://dx.doi.org/10.1111/acer.13163>. in press (<http://onlinelibrary.wiley.com/doi/10.1111/acer.13163/abstract>; jsessionid=2673094FE272AAB367B273B7827E9515.f04t04).
 23. Mogg K, Bradley BP: **Orienting of attention to threatening facial expressions presented under conditions of restricted awareness [Internet]**. *Cogn Emot* 1999, **13**:713-740.
 24. Schoenmakers TM, de Bruin M, Lux IFM, Goertz AG, Van Kerkhof DHAT, Wiers RW: **Clinical effectiveness of attentional bias modification training in abstinent alcoholic patients [Internet]**. *Drug Alcohol Depend* 2010, **109**:30-36.
 25. Houben K, Havermans RC, Wiers RW: **Learning to dislike alcohol: conditioning negative implicit attitudes toward alcohol and its effect on drinking behavior**. *Psychopharmacology (Berl)* 2010, **211**:79-86.
 26. Woud ML, Postma P, Holmes EA, Mackintosh B: **Reducing analogue trauma symptoms by computerized reappraisal training – considering a cognitive prophylaxis? [Internet]**. *J Behav Ther Exp Psychiatry* 2013, **44**:312-315.
 27. Houben K, Nederkoorn C, Wiers RW, Jansen A: **training response inhibition**. *Drug Alcohol Depend* 2011 <http://dx.doi.org/10.1016/j.drugalcdep.2010.12.011>.
 28. Christiansen P, Schoenmakers TM, Field M: **Less than meets the eye: reappraising the clinical relevance of attentional bias in addiction**. *Addict Behav* 2015, **44**:43-50.
 29. Hofmann W, Gschwendner T, Friese M, Wiers RW, Schmitt M: **Working memory capacity and self-regulatory behavior: toward an individual differences perspective on behavior determination by automatic versus controlled processes [Internet]**. *J Pers Soc Psychol* 2008, **95**:962-977.
 30. Houben K, Wiers RW, Jansen A: **Getting a grip on drinking behavior: training working memory to reduce alcohol abuse [Internet]**. *Psychol Sci* 2011, **22**:968-975.
 31. Bickel WK, Yi R, Landes RD, Hill PF, Baxter C: **Remember the future: working memory training decreases delay discounting among stimulant addicts [Internet]**. *Biol Psychiatry* 2011, **69**:260-265.
 32. Bickel WK, Landes RD, Kurth-Nelson Z, Redish AD: **A quantitative signature of self-control repair: rate-dependent effects of successful addiction treatment**. *Clin Psychol Sci* 2014, **2**:685-695.
- An interesting study on the 'control' side of the dual-process equation, linking delay discounting to treatment outcome and effects of working memory training, considering individual change trajectories.
33. Boendermaker WJ, Prins PJM, Wiers RW: **Cognitive Bias Modification for adolescents with substance use problems – can serious games help? [Internet]**. *J Behav Ther Exp Psychiatry* 2015, **49**:13-20.
- One way to counter the rather boring nature of cognitive training is to use gamification. However, this review argued that this changes motivation to perform training rather than motivation to change the addictive behavior.
34. Cristea IA, Kok RN, Cuijpers P: **Efficacy of cognitive bias modification interventions in anxiety and depression: meta-analysis [Internet]**. *Br J Psychiatry* 2015, **206**:7-16.

35. Emmelkamp PM: **Attention bias modification: the Emperor's new suit?** [Internet]. *BMC Med* 2012, **10**:63.
36. Clarke PJF, Notebaert L, MacLeod CM: **Absence of evidence or evidence of absence: reflecting on therapeutic implementations of attentional bias modification** [Internet]. *BMC Psychiatry* 2014, **14**:8.
Provides important nuancing counterarguments to criticisms of CBM.
37. Kopetz CE, Lejuez CW, Wiers RW, Kruglanski AW: **Motivation and self-regulation in addiction: a call for convergence.** *Perspect Psychol Sci* 2013, **8**:3-24.
38. Lewis M: *The Biology of Desire: Why Addiction Is Not a Disease.* Public Affairs; 2015.
39. Miller W, Rollnick G: *Motivational Interviewing: Helping People Change.* Guilford; 2013.
40. Feldstein Ewing SW, McEachern AD, Yezhuvath U, Bryan AD, Hutchison KE, Filbey FM: **Integrating brain and behavior: evaluating adolescents' response to a cannabis intervention** [Internet]. *Psychol Addict Behav* 2012, **27**:510-525.
41. Lindgren KP, Wiers RW, Teachman BA, Gasser ML, Westgate EC, Cousijn J, Enkema MC, Neighbors C: **Attempted training of alcohol approach and drinking identity associations in US undergraduate drinkers: null results from two studies** [Internet]. *PLoS One* 2015, **10**:e0134642.
42. Wiers RW, Houben K, Fardard J, van Beek P, Rhemtulla M, Cox WM: **Alcohol Cognitive Bias Modification training for problem drinkers over the web** [Internet]. *Addict Behav* 2015, **40**:21-26.
43. Kerst WF, Waters AJ: **Attentional retraining administered in the field reduces smokers' attentional bias and craving** [Internet]. *Health Psychol* 2014, **33**:1232-1240.
Study showing that cognitive biases in addiction can also be changed on a mobile device. No effects on smoking behavior were found, which is not surprising given that participants were not motivated to quit smoking.
44. Elfeddali I, de Vries H, Bolman C, Pronk T, Wiers RW: **A randomized controlled trial of Web-based Attentional Bias Modification to help smokers quit.** *Health Psychol* 2016, **35**:870-880 <http://dx.doi.org/10.1037/hea0000346>.
Study showing that internet-delivered CBM can work in heavy smokers, if they are selected to make a quit attempt.
45. MacLeod C, Clarke PJF: **The attentional bias modification approach to anxiety intervention** [Internet]. *Clin Psychol Sci* 2015, **3**:58-78.
46. Gladwin TE, Rinck M, Eberl C, Becker ES, Lindenmeyer J, Wiers RW: **Mediation of cognitive bias modification for alcohol addiction via stimulus-specific alcohol avoidance association** [Internet]. *Alcohol Clin Exp Res* 2015, **39**:101-107.
Exploratory re-analysis suggesting that the clinical result of CBM, as reported in the first large RCT for approach-retraining for alcohol addiction was mediated by effects on automatic alcohol-approach associations. Such results are important because they support the central claim of CBM. That the method works by changing automatic processes and that these changes have clinical effects.
47. Verdejo-García A: **Cognitive training for substance use disorders: neuroscientific mechanisms** [Internet]. *Neurosci Biobehav Rev* 2016 <http://dx.doi.org/10.1016/j.neubiorev.2016.05.018>.
48. Wiers CE, Stelzel C, Gladwin TE, Park SQ, Pawelczack S, Gawron CK, Stuke H, Heinz A, Wiers RW, Rinck M *et al.*: **Effects of cognitive bias modification training on neural alcohol cue reactivity in alcohol dependence.** *Am J Psychiatry* 2015, **172**:335-343.
One of the first studies to show effects of CBM on cue-evoked meso-limbic brain activity. Importantly, changes in amygdala activation appeared to mediate effects on craving. This is a first step in understanding which neural changes should be targeted to achieve clinically relevant outcomes.
49. Wiers CE, Ludwig VU, Gladwin TE, Park SQ, Heinz A, Wiers RW, Rinck M, Lindenmeyer J, Walter H, Bermphol F: **Effects of cognitive bias modification training on neural signatures of alcohol approach tendencies in male alcohol-dependent patients** [Internet]. *Addict Biol* 2015, **20**:990-999.
Showed that CBM affects brain activation related to alcohol approach biases.
50. Ernst LH, Plichta MM, Dresler T, Zesewitz AK, Tupak SV, Haeussinger FB, Fischer M, Polak T, Fallgatter AJ, Ehlis A-C: **Prefrontal correlates of approach preferences for alcohol stimuli in alcohol dependence** [Internet]. *Addict Biol* 2014, **19**:497-508.
51. Wiers CE, Stelzel C, Park SQ, Gawron CK, Ludwig VU, Gutwinski S, Heinz A, Lindenmeyer J, Wiers RW, Walter H *et al.*: **Neural correlates of alcohol-approach bias in alcohol addiction: the spirit is willing but the flesh is weak for addicts** [Internet]. *Neuropsychopharmacology* 2014, **39**:688-697.
52. Keren G, Schul Y: **Two is not always better than one: a critical evaluation of two-system theories** [Internet]. *Perspect Psychol Sci* 2009, **4**:533-550.
53. Lieberman MD: **The X- and C-systems: the neural basis of automatic and controlled social cognition** [Internet]. In *Social Neuroscience: Integrating Biological and Psychological Explanations of Social Behavior.* Edited by Harmon-Jones E, Winkielman P. Guilford Press; 2007:290-315.
54. Van Hecke J, Gladwin TE, Coremans J, Destoop M, Hulstijn W, Sabbe B: **Prefrontal, parietal and basal activation associated with the reordering of a two-element list held in working memory** [Internet]. *Biol Psychol* 2010, **85**:143-148.
55. Gladwin TE, Figner B: **"Hot" cognition and dual systems: introduction, criticisms, and ways forward.** In *Frontiers of Cognitive Psychology Series: Neuroeconomics, Judgment and Decision Making.* Edited by Wilhelms E, Reyna VF. Psychology Press; 2014:157-180.
56. Cunningham WA, Zelazo PD, Packer DJ, Van Bavel JJ: **The iterative reprocessing model: a multilevel framework for attitudes and evaluation** [Internet]. *Soc Cogn* 2007, **25**:736-760.
57. Bunge SA: **How we use rules to select actions: a review of evidence from cognitive neuroscience** [Internet]. *Cogn Affect Behav Neurosci* 2004, **4**:564-579.
58. Noël X, Colmant M, Van Der Linden M, Bechara A, Bullens Q, Hanak C, Verbanck P: **Time course of attention for alcohol cues in abstinent alcoholic patients: the role of initial orienting** [Internet]. *Alcohol Clin Exp Res* 2006, **30**:1871-1877.
59. Mogg K, Bradley BP, Miles F, Dixon R: **BRIEF REPORT Time course of attentional bias for threat scenes: testing the vigilance-avoidance hypothesis** [Internet]. *Cogn Emot* 2004, **18**:689-700.
60. Zvielli A, Bernstein A, Koster EHW: **Dynamics of attentional bias to threat in anxious adults: bias towards and/or away?** [Internet]. *PLoS One* 2014, **9**:e104025.
61. Gladwin TE: **Attentional bias variability and cued attentional bias for alcohol stimuli.** *Addict Res Theory* 2016 <http://dx.doi.org/10.1080/16066359.2016.1196674>.
62. Jansen JM, Daams JG, Koeter MWJ, Veltman DJ, van den Brink W, Goudriaan AE: **Effects of non-invasive neurostimulation on craving: a meta-analysis** [Internet]. *Neurosci Biobehav Rev* 2013, **37**:2472-2480.
63. Antal A, Keiser D, Priori A, Padberg F, Nitsche M: **Conceptual and procedural shortcomings of the systematic review' evidence that transcranial direct current stimulation (tDCS) generates little-to-no reliable neurophysiologic effect beyond MEP amplitude modulation in healthy human subjects: a systematic R.** *Brain Stimul* 2015, **8**:846-849.
Provides arguments against the conclusions of a prominent meta-analysis of tDCS. Although there may in fact be little evidence against the ability of tDCS to affect cognition, caution is needed when making generalized claims about its efficacy, similarly to the case for CBM.
64. Horvath JC, Forte JD, Carter O: **Evidence that transcranial direct current stimulation (tDCS) Generates little-to-no reliable neurophysiologic effect beyond MEP amplitude modulation in healthy Human subjects: a systematic review** [Internet]. *Neuropsychologia* 2014, **66C**:213-236.
65. Gladwin TE, den Uyl TE, Fregni FF, Wiers RW: **Enhancement of selective attention by tDCS: interaction with interference in a Sternberg task** [Internet]. *Neurosci Lett* 2012, **512**:33-37.

66. Mancuso L, Lieva I, Hamilton R, Farah M: **Does transcranial direct current stimulation improve healthy working memory? A meta-analytic review.** *J Cogn Neurosci* 2016. [no volume].
67. Ditye T, Jacobson L, Walsh V, Lavidor M: **Modulating behavioral inhibition by tDCS combined with cognitive training [Internet].** *Exp Brain Res* 2012, **219**:363-368.
68. Boggio PS, Sultani N, Fecteau S, Merabet L, Mecca T, Pascual-Leone A, Basaglia A, Fregni FF: **Prefrontal cortex modulation using transcranial DC stimulation reduces alcohol craving: a double-blind, sham-controlled study [Internet].** *Drug Alcohol Depend* 2008, **92**:55-60.
69. Fregni FF, Orsati F, Pedrosa W, Fecteau S, Tome FAM, Nitsche MA, Mecca T, Macedo EC, Pascual-Leone A, Boggio PS: **Transcranial direct current stimulation of the prefrontal cortex modulates the desire for specific foods [Internet].** *Appetite* 2008, **51**:34-41.
70. den Uyl TE, Gladwin TE, Wiers RW: **Transcranial direct current stimulation, implicit alcohol associations and craving [Internet].** *Biol Psychol* 2014, **105C**:37-42.
71. Nitsche MA, Kuo M-F, Karrasch R, Wächter B, Liebetanz D, Paulus W: **Serotonin affects transcranial direct current-induced neuroplasticity in humans [Internet].** *Biol Psychiatry* 2009, **66**:503-508.
72. Clarke PJF, Browning M, Hammond G, Notebaert L, MacLeod CM: **The causal role of the dorsolateral prefrontal cortex in the modification of attentional bias: evidence from transcranial direct current stimulation [Internet].** *Biol Psychiatry* 2014, **76**:946-952.
73. Heeren A, Baeken C, Vanderhasselt M-A, Philippot P, de Raedt R: **Impact of anodal and cathodal transcranial direct current stimulation over the left dorsolateral prefrontal cortex during attention bias modification: an eye-tracking study [Internet].** *PLoS One* 2015, **10**:e0124182.
74. Boffo M, Pronk T, Wiers RW, Mannarini S: **Combining cognitive bias modification training with motivational support in alcohol dependent outpatients: study protocol for a randomised controlled trial [Internet].** *Trials* 2015, **16**:63.
75. Riper H, Blankers M, Hadiwijaya H, Cunningham J, Clarke S, Wiers RW, Ebert D, Cuijpers P: **Effectiveness of guided and unguided low-intensity internet interventions for adult alcohol misuse: a meta-analysis [Internet].** *PLoS One* 2014, **9**:e99912.
76. van Deursen DS, Saleminck E, Smit F, Kramer J, Wiers RW: **Web-based cognitive bias modification for problem drinkers: protocol of a randomised controlled trial with a 2×2×2 factorial design [Internet].** *BMC Public Health* 2013, **13**:674.
77. Wiers RW, Becker D, Holland R, Moggi F, Lejuez CW, Yield A: **Cognitive motivational processes underlying addiction treatment.** In *Addiction*. Edited by Kopetz CE, Lejuez CW.. Psychology Press; 2016:201-236.
- Argues for the need to integrate CBM and regular treatment.
78. Wiers RW, Gladwin TE: **Reflective and impulsive processes in addiction and the role of motivation.** In *Reflective and Impulsive Determinants of Human Behavior*. Edited by Deutsch R, Gawronski B, Hofmann W. Psychology Press; 2016:173-188.