



UvA-DARE (Digital Academic Repository)

Brain training improves recovery after stroke but waiting list improves equally: A multicenter randomized controlled trial of a computer-based cognitive flexibility training

van de Ven, R.M.; Buitenweg, J.I.V.; Schmand, B.; Veltman, D.J.; Aaronson, J.A.; Nijboer, T.C.W.; Kruiper-Doesborgh, S.J.C.; van Bennekom, C.A.M.; Rasquin, S.M.C.; Ridderinkhof, K.R.; Murre, J.M.J.

DOI

[10.1371/journal.pone.0172993](https://doi.org/10.1371/journal.pone.0172993)

Publication date

2017

Document Version

Other version

Published in

PLoS ONE

[Link to publication](#)

Citation for published version (APA):

van de Ven, R. M., Buitenweg, J. I. V., Schmand, B., Veltman, D. J., Aaronson, J. A., Nijboer, T. C. W., Kruiper-Doesborgh, S. J. C., van Bennekom, C. A. M., Rasquin, S. M. C., Ridderinkhof, K. R., & Murre, J. M. J. (2017). Brain training improves recovery after stroke but waiting list improves equally: A multicenter randomized controlled trial of a computer-based cognitive flexibility training. *PLoS ONE*, *12*, [e0172993]. <https://doi.org/10.1371/journal.pone.0172993>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the library <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

S3 Table. Mean (standard deviation) and repeated-measures MANOVA of the outcome measures at follow-up.

measure	Intervention group (n = 38)			Active control group (n = 35)			Comparison											
	Follow-up	Pre-training		Post-training		Follow-up	Time (T0T3)			Time*group			Time (T2T3)			Time*group		
		F ^α	p-value	η_p^2	F ^α		p-value	η_p^2	F ^α	p-value	η_p^2	F ^α	p-value	η_p^2	F ^α	p-value	η_p^2	
ToL (optimal - moves)	-26.9 (17.8)	-34.0 (26.0)	-29.0 (21.5)	-25.6 (19.4)	9.2	<.001*	.61	F _(10 , 62)	.82	.09	F _(10 , 62)	<.001*	.39	F _(10 , 62)	.11	.21		
Switch RT (ms)	-321 (218)	-418 (328)	-416 (316)	-378 (293)	6.8	.01*	.09				0.7	.41	.01	4.3	.04	.06		
Switch Acc (trials)	-3.0 (3.4)	-2.2 (7.3)	-3.0 (4.9)	-3.4 (5.5)	0.8	.36	.01				0.2	.65	.00	0.2	.65	.00		
TMT B (sec) ^b	74.8 (37.6)	110.5 (47.8)	102.1 (42.3)	87.8 (41.9)	28.1	<.001*	.28				11.4	<.01*	.14					
TMT A (sec) ^b	41.4 (15.1)	57.3 (20.7)	52.5 (24.3)	48.4 (20.2)	33.3	<.001*	.32				1.8	.18	.03					
DSC online (correct)	38.5 (8.3)	31.7 (7.4)	35.0 (7.8)	34.7 (7.7)	31.1	<.001*	.30				0.2	.69	.00					
Corsi (span)	7.2 (1.2)	6.8 (1.1)	7.0 (1.0)	6.9 (1.1)	0.4	.52	.01				0.2	.70	.00					
Click (sec) ^b	28.7 (11.8)	37.6 (22.1)	30.9 (17.2)	31.3 (17.1)	12.7	<.001*	.15				0.1	.73	.00					
Drag (sec) ^b	2.1 (0.7)	3.4 (2.1)	2.7 (1.5)	2.6 (1.3)	29.3	<.001*	.29				3.8	.06	.05					
Peg (sec) ^b	66.2 (17.1)	83.5 (35.9)	71.0 (21.1)	73.6 (22.0)	20.5	<.001*	.22				2.9	.09	.04					

Note. Follow-up scores were only available for a selection of the tasks. Bold values are considered significant. Pre- and post-training scores of the intervention group can be found in table 4. * = remains significant after Bonferroni-Holm adjustment; b = lower values represent better performance; T0 = pre-training; T2 = post-training; T3 = 4 weeks after training completion; TMT = Trail Making Test; ToL = Tower of London; Acc = accuracy; DSST = Digit-Symbol-Coding.