



UvA-DARE (Digital Academic Repository)

The role of vanadium haloperoxidases in the formation of volatile brominated compounds and their impact on the environment

Wever, R.; van der Horst, M.A.

DOI

[10.1039/c3dt50525a](https://doi.org/10.1039/c3dt50525a)

Publication date

2013

Document Version

Final published version

Published in

Dalton Transactions

[Link to publication](#)

Citation for published version (APA):

Wever, R., & van der Horst, M. A. (2013). The role of vanadium haloperoxidases in the formation of volatile brominated compounds and their impact on the environment. *Dalton Transactions*, 42(33), 11778. <https://doi.org/10.1039/c3dt50525a>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Cite this: *Dalton Trans.*, 2013, **42**, 11778

The role of vanadium haloperoxidases in the formation of volatile brominated compounds and their impact on the environment

Ron Wever* and Michael A. van der Horst

Vanadium haloperoxidases differ strongly from heme peroxidases in substrate specificity and stability and in contrast to a heme group they contain the bare metal oxide vanadate as a prosthetic group. These enzymes specifically oxidize halides in the presence of hydrogen peroxide into hypohalous acids. These reactive halogen intermediates will react rapidly and aspecifically with many organic molecules. Marine algae and diatoms containing these iodo- and bromoperoxidases produce short-lived brominated methanes (bromoform, CHBr_3 and dibromomethane CH_2Br_2) or iodinated compounds. Some seas and oceans are supersaturated with these compounds and they form an important source of bromine to the troposphere and lower stratosphere and contribute significantly to the global budget of halogenated hydrocarbons. This perspective focuses, in particular, on the biosynthesis of these volatile compounds and the direct or indirect involvement of vanadium haloperoxidases in the production of huge amounts of bromoform and dibromomethane. Some of the global sources are discussed and from the literature a picture emerges in which oxidized brominated species generated by phytoplankton, seaweeds and cyanobacteria react with dissolved organic matter in seawater, resulting in the formation of intermediate brominated compounds. These compounds are unstable and decay *via* a haloform reaction to form an array of volatile brominated compounds of which bromoform is the major component followed by dibromomethane.

Received 26th February 2013,

Accepted 12th April 2013

DOI: 10.1039/c3dt50525a

www.rsc.org/dalton

University of Amsterdam, Van't Hoff Institute for Molecular Sciences, Science Park
904, 1098 XH Amsterdam, The Netherlands. E-mail: r.wever@uva.nl



Ron Wever and Michael A. van der Horst

Ron Wever (right) (Amsterdam, The Netherlands, 1947) received his PhD in 1975 at the University of Amsterdam. In 2001 he was nominated as a professor in Biocatalysis at the same university. His research was focussed on metalloproteins and their reactions with oxygen or oxygen derivatives. His group discovered the new class of enzymes in which vanadium is the prosthetic group, the vanadium bromo- and chloroperoxidases. These enzymes are still being studied in his group. More recently he became interested in enzymes that are able to phosphorylate or sulfate compounds with the aim to replace chemical synthesis methods by biocatalytic procedures.


Michael Alois van der Horst (left) (Vienna, Austria, 1975) finished his chemistry studies at the University of Amsterdam in 1999. He received his PhD in Microbiology in 2004 working on bacterial signal transduction with Prof. K. J. Hellingwerf at the University of Amsterdam. During his PhD and post-doctoral (KU Leuven, Belgium)

positions, his research focused on the interface of microbiology, biochemistry and biophysics. During his most recent positions in the life-science consulting company InnoTact Consulting and in the Biocatalysis research group of Prof. R. Wever, Michael's interest shifted towards green chemistry and bio-based products.




Introduction


Vanadium haloperoxidases occur widely in nature and are found in a large number of brown, red and green seaweeds, in diatoms, in bacteria and in a group of fungi, the dematiaceous hyphomycetes. The prosthetic group in these enzymes consists of metal oxide vanadate,¹ and unlike the heme peroxidases, UV-VIS spectra of these vanadium enzymes show only a modest absorption in the optical spectra around 315 nm. The bromo and chloroperoxidases are remarkably stable,^{2–6} are highly resistant towards elevated temperatures and denature only at temperatures above 70 °C.^{7,8} Detergents such as SDS or water-miscible organic solvents *e.g.* ethanol or dioxane hardly denature these enzymes. Also oxidative agents such as H₂O₂ in high concentrations, HOCl and singlet oxygen have little effect on the activity of the chloroperoxidase.⁹ The enzymes are also resistant towards proteolytic degradation. This makes the vanadium haloperoxidases unique enzymes and also raises the question of their persistence in the environment after death and decay of the host organism. Haloperoxidases catalyze the two-electron oxidation of halides (Cl[−], Br[−], I[−]) by H₂O₂ to hypohalous acids:



Most enzymes during turnover release these hypohalous acids or related halogenating intermediates, such as OX[−], X₃[−], and X⁺ from the active site in the medium, although there are exceptions in which vanadium haloperoxidases regioselectively halogenate substrates.^{10–12} In most cases these halogenating intermediates react nonspecifically with a variety of organic compounds (RH) that are susceptible for electrophilic attack. This will result in the production of a wide array of halogenated compounds (RX).



When a nucleophilic acceptor is absent a reaction may also occur between HOX and H₂O₂ resulting in the formation of singlet oxygen:



The historical nomenclature convention of the vanadium haloperoxidases is based on the most electronegative halide which these enzymes are able to oxidize. Chloroperoxidases catalyze the oxidation of Cl[−], Br[−] and I[−], bromoperoxidases catalyze only the oxidation of Br[−] and I[−] and iodoperoxidases¹³ are specific for iodide oxidation. It should be noted however that vanadium chloroperoxidase has a much higher rate in the oxidation of bromide (*k*_{cat} about 250 s^{−1}) than in the oxidation of chloride (*k*_{cat} about 20 s^{−1}).¹

Many red, brown and green macro-algae and ice microalgae from the Arctic, Antarctic, shores of temperate zones and shores of the subtropics and tropics produce CH₂Br₂ and CHClBr₂ at very high rates.^{14–22} Also diatoms found in oceans produce these halogenated compounds.^{23–26} The short-lived brominated and volatile methanes bromoform (CHBr₃) and dibromomethane (CH₂Br₂) are a very important source of

bromine to the troposphere and lower stratosphere.^{27,28} These compounds represent the largest natural contributions to atmospheric organic bromine and 20–30% of stratospheric and tropospheric O₃ depletion is due to these compounds.²⁹ A recent review³⁰ highlights the role of reactive halogens species in the chemistry and oxidizing capacity of the troposphere. The depletion of ozone through efficient catalytic cycles is well known but these compounds also oxidize dimethyl sulphide, affect the oxidation of organic compounds and reduce the lifetime of the greenhouse gas methane.

Most of these seaweeds have been shown to contain vanadium bromo- and iodoperoxidases^{31–37} and the majority of these halogenated products are formed directly or indirectly by the enzymatic activity of the vanadium enzymes. However, some seaweeds also contain heme peroxidases and for diatoms both the presence of heme and vanadium peroxidase has been reported.^{23,26} The major source of atmospheric CHBr₃ in the marine environment^{21,38,39} is the sea-to-air flux originating from macroalgal and planktonic sources, and is therefore of importance as a source of reactive halogens to the troposphere and lower stratosphere. However, this flux of CHBr₃ alone cannot account⁴⁰ for the high atmospheric mixing ratios observed in the tropical Atlantic Ocean.^{38–42} Possibly terrestrial contributions for CHBr₃ are involved and it is conceivable that soils in the tropical area produce brominated compounds. A global three-dimensional chemical model used to simulate atmospheric bromoform also indicates that emissions are likely to be significantly larger than previously assumed.⁴⁴

The vanadium iodoperoxidases in seaweeds, their role in the formation of I₂ and iodinated compounds and the impact on tropospheric photochemistry, *via* the intermediate formation of iodine radicals, will not be discussed here. For this the reader is referred to recent reviews.^{45–47} Similarly the vanadium chloroperoxidases and their involvement in the formation of chlorinated^{1,48} and brominated compounds in the terrestrial environment will be discussed elsewhere.

Formation of HOBr by seaweeds

It has been shown that in some brown³³ and a green macro-alga²² the bromoperoxidases are located at or near the surface of the seaweed. This was demonstrated for the brown seaweed *Ascophyllum nodosum* (knotted wrack) by exposing the intact plant in seawater to hydrogen peroxide. Under these conditions the plants produce HOBr by oxidation of bromide (1 mM) present in seawater resulting in the bromination of phenol red to bromophenol blue. This phenomenon was also observed for the brown seaweeds *Laminaria digitata*,⁴⁹ *L. saccharina*, *Pelvetia canaliculata*, *Fucus vesiculosus*³³ and more recently the marine red seaweed *Delisea pulchra*.¹¹ However, seaweeds are very different in their morphology and physiology and are difficult to compare, *e.g.* in some red seaweeds the peroxidases are present in glands.

The possible role of the extracellular enzymatic system may be to control the colonization of the surfaces of the seaweed



by generating HOBr which is directly bactericidal.^{50,51} In addition, HOBr in very low concentrations reacts with bacterial homoserine lactones, resulting in inactivated brominated lactones.⁴⁹ Many bacteria secrete these lactones (Fig. 1) and they play an important role in bacterial signalling systems and interference with these systems inhibits quorum sensing and bacterial biofilm formation, a first step in the fouling of surfaces.⁴⁹

Also these lactones are coregulatory ligands required for the control of the expression of genes encoding virulence traits in many Gram-negative bacterial species. Thus, HOBr is not only directly bactericidal but the oxidized halogen species also inactivate homoserine lactones and disturb communication signals between bacteria. Some red macro-algae produced halogenated furanones^{11,52–54} (Fig. 2) and the halogenated furanones inhibit *N*-acyl-L-homoserine lactones-dependent gene expression by bacteria.

The furanones are encapsulated in gland cells in the seaweed and this provides a mechanism for the delivery of the metabolites to the surface of the algae. In the macroalga *Delisea pulchra* a vanadium bromoperoxidase¹¹ halogenates these furanones.

When the intact seaweed *A. nodosum* in seawater was exposed to direct sunlight the observation was made that this seaweed generated HOBr.³³ The amounts formed under these conditions were 68 nmol g⁻¹ h⁻¹. Although the biomass of this seaweed is not known exactly, it grows in large quantities in the North Atlantic and western Russian polar seas⁵⁵ and a

biomass of 10¹¹ is certainly an accepted value. From this biomass an input of 2 × 10⁹ g of HOBr in the biosphere, produced annually by this particular seaweed, can be estimated. From the rate of release of halometabolites by several algae and a global biomass⁵⁶ of 10¹³ g, Gschwend *et al.*¹⁴ estimated an annual global input in the biosphere of 10⁴ tons per year and concluded that the macroalgae are an important source of bromine-containing material released in the atmosphere. More recent data show that these quantities for coastal seawater are even higher. Carpenter and Liss²¹ arrived at values of 1.6 Gmol Br per year from macroalgae and Carpenter *et al.*⁴² estimated on the basis of air-sea fluxes and bulk seawater and atmospheric concentrations of CHBr₃ and CH₂Br₂ a coastal flux of 2.5 Gmol Br per year corresponding to annual fluxes of 1.3–2 × 10⁵ tons per year of bromine in brominated compounds, respectively.

Mechanisms of the formation of bromoform and dibromomethane by algae and diatoms

Many seaweeds that contain bromoperoxidases produce volatile brominated compounds like CHBr₃ or dibromomethane in very large amounts. These seaweeds produce other volatile and non-volatile halogenated compounds as well but in general to a lesser extent and they will not be considered in this review. For a detailed discussion on the occurrence of other halogenated metabolites see the review by Paul and Pohnert.⁵⁷ For a long time the biosynthetic pathways by which these compounds are produced were puzzling but that bromoperoxidases are involved is obvious. Two biosynthetic pathways are conceivable. The first possibility is based upon observations on the red seaweed *Bonnemaisonia hamifera* that ketones present in seaweeds are brominated by the peroxidase. These ketones are unstable and decay *via* the haloform reaction to form volatile brominated compounds.⁵⁸ This requires however that ketones are present near or at the thallus surface and this has not been studied.

A more likely possibility as originally proposed³³ is that HOBr in the case of *A. nodosum* and other brown seaweeds is just released in seawater and reacts with dissolved organic matter (DOM) to form unstable brominated compounds which decay to form CHBr₃ and other brominated compounds. This process is similar to the production of trihalomethanes (which include CHCl₃, CHBr₃, CHCl₂Br, and CHClBr₂) as by-products of drinking water disinfection using chlorination.⁵⁹ The reaction of HOBr and DOM is very rapid, with a half-life in the millisecond range in high-DOM-containing surface waters.⁶⁰ Not only polybromomethanes will be formed, but also as known from water disinfection procedures the released HOBr will react with reactive components of DOM to form non-volatile brominated organics.⁶¹

Substantial evidence that bromination of DOM by bromoperoxidase leads to polybrominated compounds comes from studies by Manley and coworkers. It was shown²² for example that the addition of H₂O₂ to the green seaweed *Ulva lactuca* that contains a bromoperoxidase located outside of the cell




Fig. 1 Chemical structures of some of the homoserine lactones produced by bacteria.






Fig. 2 Structures of halogenated furanones produced by *Delisea pulchra*. (a) 4-Bromo-3-butyl-5-(dibromomethylene)-2(5H)-furanone. (b) 4-Bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone.



membrane resulted in an enhanced production of CHBr_3 and that removal of DOM decreased the CHBr_3 production. CHBr_3 production was nearly tripled in the light compared to the dark. The results also show that H_2O_2 produced as a result of photosynthetic and respiratory electron transport is available to the bromoperoxidase for bromination leading to the production of volatile polybromomethanes. Similarly a study²⁶ on eleven polar, temperate and warm water marine diatoms showed that 6 species have an extracellular bromoperoxidase which is located in the apoplast, the space between the cell membrane and silicified frustule. These diatoms showed a significant release of reactive bromine and iodine (primarily HOBr and HOI) in the presence of hydrogen peroxide. This is an important phenomenon that has not been reported before for diatoms. The species *Porosira glacialis* which is known to produce CHBr_3 ²³ exhibits the highest bromoperoxidase activity.²⁶ Based on its resistance towards high concentrations of H_2O_2 the bromoperoxidase in this diatom is probably a vanadium enzyme. Most of the HOBr and HOI released by these diatoms into the sea will react with DOM to form polybromo- and poly-iodomethanes. Importantly the rate of HOBr release was much greater than the CHBr_3 emissions measured by others from laboratory cultures and sea ice algae in the field indicating that most of the HOBr released may react with dissolved organic matter to form non-volatile brominated compounds.²⁶ The release of the reactive brominating species may present a defence mechanism against pathogenic marine bacteria and viruses. The subsequent reaction of HOBr or HOI with DOM may be responsible for oceanic CHBr_3 production, and CHBr_3 , CH_2Br_2 , CHI_3 , and CH_2I_2 production in polar regions. Direct evidence that reactive bromine and iodine generated by a vanadium bromoperoxidase reacts with DOM, resulting in CHBr_3 and dibromomethane, was published recently.⁶² It was shown that coastal waters contain organic compounds that are susceptible to both bromination by HOBr formed enzymatically by a bromoperoxidase or HOBr from the bottle (bromine water). In both cases the same polybromomethanes were produced with CHBr_3 the dominant species. Thus it is very likely that the release of HOBr and HOI by phytoplankton is responsible for oceanic production of CHBr_3 , CH_2Br_2 , CH_3 and CH_2I_2 .^{29,41}

The equations below summarize the sequence of events leading to the formation of CHBr_3 and other polybromomethanes:



It should be noted that not all the brominated DOM will decay to CHBr_3 but other brominated compounds will be formed as well with unknown stability and fate. Some of the brominated material may actually end up in marine sediments. Indeed, it has been shown that marine sediments also contain organic brominated species associated with humic material.⁶³

If these other compounds are formed as well this raises an important question: is the variability and amount of the polybromomethanes produced by either phytoplankton or seaweeds not a function of the origin, nature and concentration of DOM in seawater? Indeed DOM in coastal waters may differ from ocean DOM in terms of chemical composition and it may react and decay differently when exposed to HOBr. Near coastal waters contain humic material that originates mostly from degradation products of leaves and this differs from DOM of marine ecosystems.⁶⁴ This suggests strongly that the source of the seawater used, the site at which water samples were taken, and in which phytoplankton and seaweeds were incubated determine to a large extent the nature and the amounts of volatile halogenated compounds formed. This may for example explain the differences in distribution of CH_2Br_2 and CHBr_3 in warm and nitrogen depleted surface waters and cold nitrogen enriched deeper waters in the Mauritanian upwelling observed in the study by Quack *et al.*⁶⁵ Similarly Carpenter *et al.*⁴² suggested that the surface seawater concentration ratio of CH_2Br_2 and CHBr_3 is dependent upon location, that is, open ocean and coastal regions.

Effect of light, H_2O_2 and stress on bromoform production by algae and diatoms

It is now clear why the addition of H_2O_2 to seaweeds resulted in the formation of brominated compounds. The vanadium bromoperoxidases that in general are extracellular are accessible to added hydrogen peroxide, start producing HOBr that will in turn react with DOM in seawater or in some cases with ketoacids from the seaweed. Consequently volatile halogenated compounds will be formed as shown by several authors.^{66–72}

It is well known that plant cells during photosynthesis and photorespiration generate H_2O_2 .⁷³ Also intra- and intercellular levels of H_2O_2 increase during environmental stresses experienced by plant cells. Several groups showed that illumination of seaweeds indeed results in the formation of volatile halogenated compounds.^{66,67,71,74,75} When six algal species (a diatom, a brown algae and four filamentous green algae) were illuminated a strong correlation was observed⁷² between the release of hydrogen peroxide and the release of brominated compounds to the medium. Surprisingly these species also generated substantial amounts of chloroform with the diatom *Pleurosira laevis* as one of the major producers. Nightingale *et al.*¹⁶ reported that several seaweed species also release chloroform, although substantially less than CHBr_3 . This suggests strongly that some of the seaweed species contain chloroperoxidase activity. This requires however that the reaction of enzymatically generated HOCl with DOM be faster than the reaction of HOCl with Br^- present in seawater; otherwise only CHBr_3 would be detected.

The highest production rates of halocarbons by macroalgae were observed when photosynthesis was maximal in a rock pool containing macroalgae⁷⁶ under near natural conditions. The concentrations of halocarbons corresponded well with the concentration of H_2O_2 . This again establishes a clear



relationship between photosynthesis, H_2O_2 and halocarbon formation. In line with this, bromination of externally added phenol red occurs^{33,66} when the brown macroalga *A. nodosum* and the red alga *Meristiella gelidium*, which contain (vanadium) bromoperoxidases, were exposed to light. This demonstrates again the direct release of HOBr by these algae.

Stress experienced by the red seaweed *Soliera chordalis* also affects the release of volatile halogenated compounds, e.g. hypo-osmotic conditions strongly enhanced the production rate of volatile halogenated compounds.⁷⁴

In line with this exposure of laboratory-grown and harvested wild individuals of *L. digitata* to oligoguluronates, carbohydrates which are released when the plant is wounded and stressed resulted in the release of H_2O_2 .⁷⁷ Further, concomitantly *L. digitata* produces a number of volatile brominated and iodinated compounds.

Thus, illumination and stressing of seaweeds result in the formation of hydrogen peroxide and as a consequence of the presence of the bromoperoxidases, halogenated compounds will be formed *via* the intermediate formation of HOBr, as has been demonstrated for several different algae under a variety of conditions.^{66–72,74–77} However, although Antarctic macroalgae release CHBr_3 and dibromomethane no clear light-dependent relationship was found.⁷⁸ Similarly the exposure of three species of marine microalgae that were selected to represent three different algal groups that are common to the marine environment, to light stress did not result in the release of iodinated or brominated volatile compounds.⁷⁹

Bromoform and dibromomethane in oceans and coastal areas

Tropical oceans. There is substantial evidence that the tropical ocean is a main source of bromoform produced by phytoplankton. The data by Quack *et al.*⁴¹ show that the sea-to-air flux is strongly localized, and includes intense emissions in tropical open ocean regions where bromoform can be transported rapidly to the troposphere–stratosphere boundary. The measurements show that there is also widespread production within the open ocean that can support elevated atmospheric CHBr_3 levels. Not only physical evidence is present but also the global three dimensional chemical model used to simulate atmospheric bromoform indicates that emissions from oceans are likely to be significantly larger than previously assumed.⁴⁴ Further, according to this model a large proportion of emissions is situated in tropical regions. Later data of Carpenter *et al.*⁴² also show that both bromoform and dibromomethane are supersaturated throughout these regions. In line with this, extensive halogen mediated ozone destruction was observed over the tropical Atlantic Ocean as measured at the Cabo Verde archipelago.^{43,80} The West Pacific Ocean also emits CHBr_3 and CH_2Br_2 but emissions are low to moderate.⁸¹ As observed previously⁴⁰ the concentration ratio of $\text{CH}_2\text{Br}_2/\text{CHBr}_3$ in seawater is a strong function of concentration (and location), with a lower $\text{CH}_2\text{Br}_2/\text{CHBr}_3$ ratio found in coastal regions near macroalgal sources.⁴² As pointed out this may relate to the nature of the DOM present at these locations,⁶⁴

its reactivity towards HOBr and its ability to undergo the haloform reaction. US coastal waters also are supersaturated with bromoform and values were found ranging from 2 to 1724 pmol L^{-1} .⁸² The highest concentrations along the coast-line were found north of 40°N partly originating from macroalgal sources.

Arctic ocean. Several studies have shown that in the Arctic Ocean halocarbons are present in the ng L^{-1} range. The early data from Dyrssen and Fogelqvist⁸³ demonstrate that the source of CHBr_3 is biogenic and that algal belts in the coastal area of Svalboard are partly responsible. Some sampling sites in and outside fjords showed the presence of very high concentrations of CHBr_3 (up to 300 ng L^{-1} or 1185 pmol L^{-1}). High surface water concentrations (up to 12 ng L^{-1}) were found for stations situated far from the coast. This biogenic production of CHBr_3 in the Arctic Ocean causes an oversaturation in the open sea surface waters when compared to the atmospheric background and this results in a flux of CHBr_3 to the atmosphere. Studies by Fogelqvist^{84,85} show that phytoplankton in the Arctic Ocean is involved in the production of CHBr_3 . At the Nansen basin in the Arctic Ocean there is a CHBr_3 maximum (3–7 ng L^{-1}) which is indicative of production very close to the surface and this suggests that pelagic marine algae are the source of CHBr_3 in the Northern Atlantic Ocean.⁸⁶ Indeed it has been shown^{23,24–26} that marine phytoplankton and sea-ice algae, especially diatoms, are producers of CHBr_3 . As discussed, bromo and iodoperoxidase activities have been detected in these diatoms.

The Gulf Stream may also play a role in this by transporting CHBr_3 -rich waters from coastal areas to the Arctic Ocean. For some time the CHBr_3 released from seaweed and sea-ice algae^{24,88–90} was taken as the major source of bromine responsible for the ozone loss in the arctic troposphere at polar sunrise.⁸⁷ Recent studies^{91,92} showed that the photolysis of CHBr_3 was too slow to account for the formation of reactive bromine species and subsequently the reactive gas phase bromine “Bromine explosion” has been explained by heterogeneous autocatalytic reactions taking place on salt-laden ice surfaces.^{93,94} Globally, however, the direct emission of brominated compounds and molecular iodine contributes significantly to halogen loading in coastal marine boundary layers^{95,96} and oceans and it may be a new link between climate change and the composition of the global atmosphere.^{28,29,38}

Interestingly a study⁹⁷ of the marine boundary layer at Roscoff, Brittany, where large macroalgal beds exposed to a large tidal range are present, showed in addition to significant I_2 emission the formation of BrO. This pointed to extremely localized emissions of reactive bromine. It is tempting to speculate that the brown algae, mainly Laminariaceae, are responsible for this since it is known that they produce HOBr directly. Overall these brominated compounds either abiotic or biotically produced play an important role in tropospheric and lower stratospheric ozone chemistry.

Antarctic. The annual ozone loss in the Antarctic has also triggered substantial research on biological halocarbon



formation²⁵ and fate in this area of the world. Carpenter *et al.*⁹⁸ showed, in the Antarctic Spring Summer, that at several locations in the Southern Ocean within 40 km of the Arctic, sea ice surface maxima of several dihalogenated methane and bromoform occurred in the water column (5–100 m depth). This is consistent with *in situ* surface halocarbon production by ice algae liberated from the sea ice. Concurrent measurements in air suggested a sea–air flux of bromoform near the Antarctic coast. Interestingly, in pelagic polar oceans the concentrations of bromoform are a factor of 2–4 lower³⁸ than observed (57 pmol) in the study by Carpenter *et al.*⁹⁸ close to sea ice.

Values of 50 pmol L⁻¹ have been reported by Moore *et al.*⁹⁹ at Resolute Bay (Arctic waters) during spring ice algal bloom and by Fogelqvist and Tunhua¹⁰⁰ of 54 pmol L⁻¹ in close proximity to an ice flow in the Weddel Sea in Antarctica. Similarly near the King George Island values of up to 190 pmol L⁻¹ (50 ng L⁻¹) were found.¹⁰¹ Strong seasonality in CHBr₃ and CH₂Br₂ concentrations was observed¹⁰² and the highest bromocarbon concentrations (up to 276.4 ± 13.0 pmol CHBr₃ L⁻¹ and 30.0 ± 0.4 pmol CH₂Br₂ L⁻¹) were found to coincide with the annual microalgal bloom during the austral summer. The timing of the initial increase in bromocarbon concentrations was related to the sea-ice retreat and onset of the microalgal bloom. The high sea–air flux rate of bromocarbon emissions observed at the seasonal ice edge zone bloom, which occurs each year over large areas of the Southern Ocean during the austral summer, could have an important impact on the chemistry of the Antarctic atmosphere.¹⁰² Similarly seawater CHBr₃ concentrations in samples collected in coastal waters of the western Antarctic Peninsula were found¹⁰³ to be significantly higher (122 pmol L⁻¹) in samples collected in diatom bloom compared to non-bloom waters (42.9 [12.0–126] pmol L⁻¹). A comparison of sea-to-air flux rates suggests that a switch from diatom bloom to non-bloom conditions results in a factor of 3–4 decrease in average CHBr₃ sea-to-air emission rates which will reduce the supply of biogenic bromine to the atmosphere. These concentrations of bromoform are still 1–2 orders of magnitude less than global near shore values influenced by macro-algae.³⁸

General remarks

From the studies described it is clear that in oceans marine phytoplankton is the main source of bromoform and dibromomethane. Similarly sea ice algae present in the Arctic or Antarctic also produce these brominated compounds in substantial amounts. In this case their formation correlates strongly with microalgal bloom in spring. Some tropical regions are even supersaturated with bromoform and dibromomethane which by sea–air flux leads to halogen-mediated ozone destruction in the atmosphere. The concentrations in the pelagic zone are in general lower than in the coastal waters^{38,42,82,104} and in going from the open ocean *via* shelf to coastal areas, saturations and sea–air fluxes of bromoform and

dibromomethane in general increase. Several studies have established that in these coastal waters marine macro algae are responsible for the high concentrations of the brominated compounds formed. Tidal cycles and day light seem to affect their formation as was shown for the Iberian upwelling off Portugal.¹⁰⁵

In the upwelling along the coast, an additional coastal source was present that further increased the bromocarbon concentration. This coastal source was correlated to the diurnal and tidal cycles, showing that intertidal macro-algae beds were involved in halocarbon production.¹⁰⁵ Indeed ample evidence exists that photosynthesis plays an important role in the formation of these brominated compounds as has been shown for many seaweed species in laboratory experiments^{66,67,71,73,75} but also under near natural conditions.⁷⁶ Phytoplankton and macro algae are not the only source of volatile brominated compounds. Data obtained from a cruise in the Baltic established¹⁰⁶ that cyanobacteria form also significant amounts of CHBr₃, CH₃I, CH₂Br₂ and CHBr₂Cl and in particular during summer, cyanobacteria become highly dominant in the Baltic Sea, forming extensive blooms. The calculated total production of bromoform, in the Baltic Sea, could be of the order of 110 Mg per year for a cyanobacterial bloom period of 1 month and a maximum bloom coverage of 130 000 km². For the other compounds a range of Mg per year is calculated. Interestingly the addition of hydrogen peroxide to a culture of the cyanobacterium *Nodularia spumigena* resulted in an increased production of halocarbons, indicating haloperoxidase activity.¹⁰⁶ Indeed it has been shown that in several sequenced genomes of marine cyanobacteria, putative vanadium bromoperoxidases are present.^{107,108} Protein extracts of cultures of *Synechococcus* sp. CC9311 show bromoperoxidase activity establishing a functional vanadium bromoperoxidase in this strain. Homologous bromoperoxidase sequences are also present in the genomes of *Acaryochloris marina*, *Crocospaera watsonii* and *Nodularia spumigena*.¹⁰⁸ As has also been observed for many seaweed species^{16,74,76} the presence of halocarbons in surface waters of the Baltic Sea showed a diurnal variation with the highest concentrations around noon.¹⁰⁶

Conclusions

Fig. 3 summarizes and simplifies the events occurring in the formation of CHBr₃ and its ventilation to the atmosphere. Seaweeds, phytoplankton and cyanobacteria triggered by light or stress start releasing HOBr in seawater. This will react with DOM, resulting in brominated DOM that partly decays *via* the haloform reaction to volatile brominated compounds, mainly CHBr₃ and CH₂Br₂. The brominated DOM that does not decay may end up in the marine sediment.⁶³ The volatile compounds are ventilated to the atmosphere and as a result of photolysis in the troposphere and lower stratosphere bromine radicals will be formed that cause depletion of ozone.³⁰






Fig. 3 The formation of bromine radicals results from the photolysis of brominated methanes formed by the reaction of DOM with HOBr produced by seaweed, phytoplankton and cyanobacteria. By a catalytic cycle these radicals will cause depletion of ozone.³⁰

References

- R. Wever and W. Hemrika, in *Handbook of Metalloproteins*, ed. A. Messerschmidt, R. Hubert, T. Poulos and K. Wieghardt, John Wiley & Sons, Ltd, Chichester, 2001, pp. 1417–1428.
- M. Almeida, S. Filipe, M. Humanes, M. F. Maia, R. Melo, N. Severino, J. A. L. da Silva, J. J. R. F. da Silva and R. Wever, *Phytochemistry*, 2001, **57**, 633–642.
- D. J. Sheffield, T. Harry, A. J. Smith and L. J. Rogers, *Phytochemistry*, 1993, **32**, 21–26.
- M. G. M. Tromp, G. Olafsson, B. E. Krenn and R. Wever, *Biochim. Biophys. Acta*, 1990, **1040**, 192–198.
- B. Zhang, X. Cao, X. Cheng, P. Wu, T. Xiao and W. Zhang, *Biotechnol. Lett.*, 2011, **33**, 545–548.
- E. Garcia-Rodriguez, T. Ohshiro, T. Aibara, Y. Izumi and J. Littlechild, *J. Biol. Inorg. Chem.*, 2005, **10**, 275–282.
- H. Plat, B. E. Krenn and R. Wever, *Biochem. J.*, 1987, **248**, 277–279.
- J. W. P. M. van Schijndel, P. Barnett, J. Roelse, E. G. M. Vollenbroek and R. Wever, *Eur. J. Biochem.*, 1994, **22**, 151–157.
- R. Renirie, C. Pierlot, J. M. Aubry, A. F. Hartog, H. E. Schoemaker, P. L. Alsters and R. Wever, *Adv. Synth. Catal.*, 2003, **345**, 849–858.
- A. Butler and M. Sandy, *Nature*, 2009, **460**, 848–854.
- M. Sandy, J. N. Carter-Franklin, J. D. Martiny and A. Butler, *Chem. Commun.*, 2011, **47**, 12086–12088.
- J. M. Winter, M. C. Moffit, E. Zazopolos, J. B. McAlpine, P. C. Dorrestein and B. S. Moore, *J. Biol. Chem.*, 2007, **282**, 16362–16368.
- C. Colin, C. Leblanc, E. Wagner, L. Delage, E. Leize-Wagner, A. van Dorsselaer, B. Kloareg and P. Potin, *J. Biol. Chem.*, 2003, **278**, 23545–23552.
- P. Gschwend, J. MacFarlane and K. Newman, *Science*, 1985, **227**, 1033–1035.
- S. L. Manley, K. Goodwin and W. J. North, *Limnol. Oceanogr.*, 1992, **37**, 1652–1659.
- P. D. Nightingale, G. Malin and P. S. Liss, *Limnol. Oceanogr.*, 1995, **40**, 680–689.
- F. Laturus, *Mar. Chem.*, 1996, **55**, 359–366.
- N. Itoh, M. Tsujita, T. Ando, G. Hisatomi and T. Higashi, *Phytochemistry*, 1996, **45**, 67–73.
- K. D. Goodwin, W. J. North and M. E. Lidstrom, *Limnol. Oceanogr.*, 1997, **42**, 1725–1734.
- F. Laturus, C. Wiencke and H. Klöser, *Mar. Environ. Res.*, 1996, **41**, 169–181.
- L. Carpenter and P. S. Liss, *J. Geophys. Res., [Atmos.]*, 2000, **105**, 539–548.
- S. L. Manley and P. E. Barbero, *Limnol. Oceanogr.*, 2001, **46**, 1392–1399.
- R. M. Moore, M. Webb, R. Tokarczyk and R. Wever, *J. Geophys. Res., [Atmos.]*, 1996, **101**, 20899–20908.
- W. T. Sturges, G. F. Cota and P. T. Buckley, *Nature*, 1992, **358**, 660–662.
- W. T. Sturges, C. W. Sullivan, R. C. Schnell, L. E. Heidt and W. H. Pollock, *Tellus Ser. B*, 1993, **45**, 120–126.
- V. L. Hill and S. L. Manley, *Limnol. Oceanogr.*, 2009, **54**, 812–822.
- World Meteorological Organization, *Scientific assessment of ozone depletion: 2006*, Global Ozone Res. Monit. Proj. 50, Geneva, Switzerland, 2007, p. 498.
- X. Yang, R. A. Cox, N. J. Warwick, J. A. Pyle, G. D. Carver and N. H. Savage, *J. Geophys. Res., [Atmos.]*, 2005, **110**, 1984–2012.
- R. Salawitch, *Nature*, 2006, **439**, 275–277.
- A. Saiz-Lopez and R. von Glasow, *Chem. Soc. Rev.*, 2012, **41**, 6448–6472.
- E. de Boer, M. G. M. Tromp, H. Plat, G. E. Krenn and R. Wever, *Biochim. Biophys. Acta*, 1986, **872**, 104–115.
- B. E. Krenn, Y. Izumi, H. Yamada and R. Wever, *Biochim. Biophys. Acta*, 1989, **998**, 63–68.
- R. Wever, M. G. M. Tromp, B. E. Krenn, A. Marjani and M. van Tol, *Environ. Sci. Technol.*, 1991, **25**, 446–449.
- R. Wever, M. G. M. Tromp, J. W. P. M. van Schijndel and E. Vollenbroek, in *Biogeochemistry of Global Change*, ed. R. D. Oremland, Chapman and Hall, New York, 1993, pp. 811–824.
- T. Ohshiro, S. Nakano, Y. Takahashi, M. Suzuki and Y. Izumi, *Phytochemistry*, 1999, **52**, 1211–1215.
- J. N. Carter-Franklin and A. Butler, *J. Am. Chem. Soc.*, 2004, **126**, 15060–15066.
- T. Suthiphongchai, P. Boonsiri and B. Panijpan, *J. Appl. Phycol.*, 2008, **20**, 271–278.
- B. Quack and D. W. R. Wallace, *Global Biogeochem. Cycles*, 2003, **17**, 1023.
- L. J. Carpenter, D. J. Wevill, J. R. Hopkins, R. M. Dunk, C. E. Jones, K. E. Hornsby and J. B. McQuaid, *Geophys. Res. Lett.*, 2007, **34**, L11810, DOI: 10.1029/2007gl029893.



- 40 B. Quack, I. Peeken, G. Petrick and K. Nachtigall, *J. Geophys. Res.*, 2007, **112**, C10006, DOI: 10.1029/2006jc003803.
- 41 B. Quack, E. Atlas, G. Petrick, V. Stroud, S. Schauffler and D. W. R. Wallace, *Geophys. Res. Lett.*, 2004, **31**, L23S05, DOI: 10.1029/2004gl020597.
- 42 L. J. Carpenter, C. E. Jones, R. M. Dunk, K. E. Hornsby and J. Woeltjen, *Atmos. Chem. Phys.*, 2009, **9**, 1805–1816.
- 43 K. A. Read, A. S. Mahajan, L. J. Carpenter, M. J. Evans, B. V. E. Faria, D. E. Heard, J. R. Hopkins, J. D. Lee, S. J. Moller, A. C. Lewis, L. Mendes, J. B. McQuaid, H. Oetjen, A. Saiz-Lopez, M. J. Pilling and J. M. C. Plane, *Nature*, 2008, **453**, 1232–1235.
- 44 N. J. Warwick, J. A. Pyle, G. D. Carver, X. Yang, N. H. Savage, F. M. O'Connor and R. A. Cox, *J. Geophys. Res.*, 2006, **111**, D24305, DOI: 10.1029/2006jd007264.
- 45 R. Wever, in *Vanadium Biochemical and Molecular Biological Approaches*, ed. H. Michibata, Springer, Dordrecht, Heidelberg, London, New York, 2012, pp. 95–125.
- 46 G. McFiggans, C. S. E. Bale, S. M. Ball, J. M. Beames, W. J. Bloss, L. J. Carpenter, J. Dorsey, R. Dunk, M. J. Flynn, K. L. Furneaux, M. W. Gallagher, D. E. Heard, A. M. Hollingsworth, K. Hornsby, T. Ingham, C. E. Jones, R. L. Jones, L. J. Kramer, J. M. Langridge, C. Leblanc, J.-P. LeCrane, J. D. Lee, R. J. Leigh, I. Longley, A. S. Mahajan, P. S. Monks, H. Oetjen, A. J. Orr-Ewing, J. M. C. Plane, P. Potin, A. J. L. Shillings, F. Thomas, R. von Glasow, R. Wada, L. K. Whalley and J. D. Whitehead, *Atmos. Chem. Phys.*, 2010, **10**, 2975–2999.
- 47 S. La Barre, P. Potin, C. Leblanc and L. Delage, *Mar. Drugs*, 2010, **8**, 988–1010.
- 48 P. Ortiz-Bermúdez, K. C. Hirth, E. Srebotnik and K. E. Hammel, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 3895–3900.
- 49 S. A. Borchardt, E. J. Allain, J. J. Michels, G. W. Stearns, R. F. Kelly and W. F. McCoy, *Appl. Environ. Microbiol.*, 2001, **67**, 3174–3179.
- 50 R. Renirie, A. Dewilde, C. Pierlot, R. Wever, D. Hober and J.-M. Aubry, *J. Appl. Microbiol.*, 2008, **105**, 264–270.
- 51 E. H. Hansen, L. Albertsen, T. Schafer, C. Johansen, J. C. Frisvad, S. Molin and L. Gram, *Appl. Environ. Microbiol.*, 2003, **69**, 4611–4617.
- 52 T. B. Rasmussen, M. Manefield, J. B. Andersen, L. Eberl, U. Anthoni, C. Christophersen, P. Steinberg, S. Kjelleberg and M. Givskov, *Microbiology*, 2000, **146**, 3237–3244.
- 53 S. Kjelleberg and P. Steinberg, *Microbiology Today*, 2001, **28**, 134–135.
- 54 M. Manefield, T. B. Rasmussen, M. Henzter, J. B. Andersen, P. Steinberg, S. Kjelleberg and M. Givskov, *Microbiology*, 2002, **148**, 1119–1127.
- 55 L. Zenkevitch, *Biology of the Seas of the USSR*, Interscience Publishers, New York, 1963, 955.
- 56 R. J. Waaland, in *The Biology of Seaweeds*, ed. C. S. Lobban and M. J. Wynne, University of California Press, Berkeley and Los Angeles, 1981, pp. 726–741.
- 57 C. Paul and G. Pohnert, *Nat. Prod. Rep.*, 2011, **28**, 186–195.
- 58 R. Theiler, J. C. Cook, L. P. Hager and J. F. Siuda, *Science*, 1978, **202**, 1094–1096.
- 59 G. R. Helz and R. Y. Hsu, *Limnol. Oceanogr.*, 1978, **23**, 858–869.
- 60 D. A. Jaworske and G. R. Helz, *Environ. Sci. Technol.*, 1985, **19**, 1188–1191.
- 61 S. D. Boyce and J. F. Hornig, *Environ. Sci. Technol.*, 1983, **17**, 202–211.
- 62 C.Y. Lin and S. L. Manley, *Limnol. Oceanogr.*, 2012, **56**, 1857–1866.
- 63 A. C. Leri, J. A. Hakala, M. A. Marcus, A. Lanzirrotti, C. M. Reddy and S. C. B. Myneni, *Global Biogeochem. Cycles*, 2010, **24**, GB4017, DOI: 10.1029/2010gb003794.
- 64 S. Opsahl and R. Benner, *Nature*, 1997, **6**, 480–482.
- 65 B. Quack, E. Atlas, G. Petrick and D. Wallace, *J. Geophys. Res.*, 2007, **112**, D09312, DOI: 10.1029/2006jd007614.
- 66 M. Pedersen, J. Collen, K. Abrahamsson and A. Ekdahl, *Sci. Mar.*, 1996, **60**, 257–263.
- 67 J. Collen, A. Ekdahl, K. Abrahamsson and M. Pedersen, *Phytochemistry*, 1994, **36**, 1197–1203.
- 68 J. Sundström, J. Collin, K. Abrahamsson and M. Pedersén, *Phytochemistry*, 1996, **42**, 1527–1530.
- 69 N. Ohsawa, Y. Ogata, N. Okada and N. Itoh, *Phytochemistry*, 2001, **58**, 683–692.
- 70 R. A. Marshall, D. B. Harper, W. McRoberts and M. J. Dring, *Limnol. Oceanogr.*, 1999, **44**, 1348–1352.
- 71 L. Mata, H. Gaspar, F. Justini and R. Santos, *J. Appl. Phycol.*, 2011, **23**, 827–832.
- 72 K. Abrahamsson, K.-S. Choo, M. Pedersen, G. Johansson and P. Snoeijs, *Phytochemistry*, 2003, **64**, 725–734.
- 73 I. Ślesak, M. Libik, B. Karpinska, S. Karpinski and Z. Miszalski, *Acta Biochim. Pol.*, 2007, **54**, 39–50.
- 74 S. Bondu, B. Cocquempot, E. Deslandes and P. Morin, *Bot. Mar.*, 2008, **51**, 485–492.
- 75 M. S. P. Mtolera, J. Collen, M. Pedersen, A. Ekdahl, K. Abrahamsson and A. K. Semesi, *Eur. J. Phycol.*, 1996, **31**, 89–95.
- 76 A. Ekdahl, M. Pedersen and K. Abrahamsson, *Mar. Chem.*, 1998, **63**, 1–8.
- 77 F. Thomas, A. Cosse, S. Goullitquer, S. Raimund, P. Morin, M. Valero, C. Leblanc and P. Potin, *PLoS One*, 2011, **6**, e21475.
- 78 F. Laturus, C. Wiencke and F. C. Adam, *Mar. Environ. Res.*, 1998, **45**, 285–294.
- 79 C. Hughes, G. Malin, P. D. Nightingale and P. S. Liss, *Limnol. Oceanogr.*, 2006, **51**, 2849–2854.
- 80 A. S. Mahajan, J. M. C. Plane, H. Oetjen, L. Mendes, R. W. Saunders, A. Saiz-Lopez, C. E. Jones, L. J. Carpenter and G. B. McFiggans, *Atmos. Chem. Phys.*, 2010, **10**, 4611–4624.
- 81 S. Tegtmeier, K. Krüger, B. Quack, E. L. Atlas, I. Pisso, A. Stohl and X. Yang, *Atmos. Chem. Phys.*, 2012, **12**, 10633–10648.
- 82 Y. Liu, S. A. Yvon-Lewis, L. Hu, J. E. Salisbury and J. E. O'Hern, *J. Geophys. Res.*, 2011, **116**, C10004, DOI: 10.1029/2010jc006729.



- 83 D. Dyrssen and E. Fogelqvist, *Ocean. Acta*, 1981, **43**, 313–317.
- 84 E. Fogelqvist, *J. Geophys. Res.*, [Atmos.], 1985, **90**, 9181–9193.
- 85 E. Fogelqvist, B. Josefsson and C. Roos, *Environ. Sci. Technol.*, 1982, **16**, 479–482.
- 86 M. Krysell, *Mar. Chem.*, 1991, **33**, 187–197.
- 87 L. A. Barrie, J. W. Bottenheim, R. C. Schnell, P. J. Crutzen and R. A. Rasmussen, *Nature*, 1988, **334**, 138–141.
- 88 R. Wever, *Nat. Correspondence*, 1988, **335**, 501.
- 89 R. Wever, in *Microbial growth on C1 Compounds*, ed. J. C. Murell and D. P. Kelly, Intercept Ltd, Andover, 1993, pp. 35–45.
- 90 N. Itoh and M. Shinya, *Mar. Chem.*, 1994, **45**, 95–103.
- 91 M. Martinez, T. Arnold and D. Perner, *Ann. Geophys.*, 1999, **17**, 941–956.
- 92 W. R. Simpson, R. von Glasow, K. Riedel, P. Anderson, P. Ariya, J. Bottenheim, J. Burrows, L. J. Carpenter, U. Friess, M. E. Goodsite, D. Heard, M. Hutterli, H. W. Jacobi, L. Kaleschke, B. Neff, J. Plane, U. Platt, A. Richter, H. Roscoe, R. Sander, P. Shepson, J. Sodeau, A. Steffen, T. Wagner and E. Wolff, *Atmos. Chem. Phys.*, 2007, **7**, 4375–4418.
- 93 K. L. Foster, R. A. Plastridge, J. W. Bottenheim, P. B. Shepson, B. J. Finlayson-Pitts and C. W. Spicer, *Science*, 2001, **291**, 471–474.
- 94 L. Kaleschke, A. Richter, J. Burrows, O. Afe, G. Heygster, J. Notholt, A. M. Rankin, H. K. Roscoe, J. Hollwedel, T. Wagner and H. W. Jacobi, *Geophys. Res. Lett.*, 2004, **31**, L16114, DOI: 10.1029/2004gl020655.
- 95 L. J. Carpenter, W. T. Sturges, S. A. Penkett, P. S. Liss, B. Alicke, K. Heibestreit and U. Platt, *J. Geophys. Res.*, [Atmos.], 1999, **104**, 1679–1689.
- 96 C. J. Palmer, T. L. Anders, L. J. Carpenter, F. C. Kupper and G. McFiggans, *Environ. Chem.*, 2005, **2**, 282–290.
- 97 D. Lowe, J. Ryder, R. Leigh, J. R. Dorsey and G. McFiggans, *Atmos. Chem. Phys.*, 2011, **11**, 979–994.
- 98 L. J. Carpenter, D. J. Wevill, C. J. Palmer and J. Michels, *Mar. Chem.*, 2007, **103**, 227–236.
- 99 R. M. Moore and R. Tokarczyk, *Geophys. Res. Lett.*, 1992, **19**, 1779–1782.
- 100 E. Fogelqvist and T. Tanhua, in *Naturally-Produced Organohalogens*, ed. A. Grimvall and E. W. B. de Leer, Kluwer Academic Publishers, Dordrecht, the Netherlands, 1995, pp. 295–306.
- 101 W. Reifenhäuser and K. G. Heumann, *Chemosphere*, 1992, **24**, 1293–1300.
- 102 C. Hughes, A. Chuck, H. Rossetti, P. J. Mann, S. M. Turner, A. Clarke, R. Chance and P. S. Liss, *Global Biogeochem. Cycles*, 2009, **23**, GB2024, DOI: 10.1029/2008gb003268.
- 103 C. Hughes, M. Johnson, R. von Glasow, R. Chance, H. Atkinson, T. Souster, G. A. Lee, A. Clarke, M. Meredith, H. J. Venables, S. M. Turner, G. Malin and P. S. Liss, *Global Biogeochem. Cycles*, 2012, **26**, GB3019, DOI: 10.1029/2012gb004295.
- 104 R. M. Moore and R. Tokarczyk, *Global Biogeochem. Cycles*, 1993, **7**, 195–210.
- 105 S. Raimund, B. Quack, Y. Bozec, M. Vernet, V. Rossi, V. Garçon, Y. Morel and P. Morin, *Biogeosciences*, 2011, **8**, 1551–1564.
- 106 A. Karlsson, N. Auer, D. Schulz-Bull and K. Abrahamsson, *Mar. Chem.*, 2008, **110**, 129–139.
- 107 B. Palenik, Q. Ren, C. Dupont, G. Myers, J. Heidelberg, J. Badger and R. Madupu, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 13555–13559.
- 108 T. L. Johnson, B. Palenik and B. Brahamsha, *J. Phycol.*, 2011, **47**, 792–801.

