



## UvA-DARE (Digital Academic Repository)

### IGR J18245-2452 is a new transient located in the core of the globular cluster M28

Heinke, C.O.; Bahramian, A.; Wijnands, R.; Altamirano, D.

**Publication date**

2013

**Document Version**

Final published version

**Published in**

The astronomer's telegram

[Link to publication](#)

**Citation for published version (APA):**

Heinke, C. O., Bahramian, A., Wijnands, R., & Altamirano, D. (2013). IGR J18245-2452 is a new transient located in the core of the globular cluster M28. *The astronomer's telegram*, 4927. <http://www.astronomerstelegam.org/?read=4927>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

24 Jun 2014; 15:22 UT

## Outside

GCN  
IAUCs

## Other

ATel on [Twitter](#) and [Facebook](#)  
ATELstream  
ATel Community Site  
MacOS: [Dashboard Widget](#)This space for free for your  
conference.IAU Symposium 305  
Polarimetry: From the Sun to  
Stars and Stellar  
Environments  
Costa Rica  
Nov 30-Dec 5 2014[ [Previous](#) | [Next](#) | [ADS](#) ]

## IGR J18245-2452 is a new transient located in the core of the globular cluster M28

ATel #4927; *C. O. Heinke, A. Bahramian (U. Alberta), R. Wijnands, D. Altamirano (U. Amsterdam)*

on 30 Mar 2013; 06:33 UT

Credential Certification: *Craig Heinke (cheinke@virginia.edu)*

Subjects: X-ray, Globular Cluster, Transient

Referred to by ATel #: [4929](#), [4934](#), [4959](#), [4960](#), [4961](#), [4964](#), [4981](#), [5003](#), [5031](#), [5045](#), [5068](#), [5069](#), [5086](#)

Eckert et al. (Atel #4925) report a new INTEGRAL ISGRI transient, IGR J18245-2452, consistent with the location of the globular cluster M28, but with a large (1.4', 90% conf) error circle. We used the UK Swift Science Centre XRT products tool, and our own processing, to analyze a Swift observation of IGR J18245-2452 (PI Ferrigno), which started on March 29, 2013 and lasted 1977 seconds. The transient is clearly detected and heavily piled-up, but the PSF is symmetrical, allowing a reliable centroid position. The centroid position is (J2000) 18:24:32.20,-24:52:05.5, with error radius 3.5" (at 90% confidence), inside the core of M28. We note that this position is inconsistent with the one known quiescent LMXB in the cluster, at position 18:24:32.821,-24:52:08.26 (Becker et al. 2003, ApJ, 594, 798), and is only consistent (within 90% confidence) with one source from Becker et al. (2003), their source 22. Note that source 22 is probably a combination of X-ray emission from millisecond pulsar L and another source (see Bogdanov et al. 2011, ApJ, 730, 81). It is quite possible that the true quiescent counterpart to IGR J18245-2452 is fainter than Becker et al.'s detection limit.

The Swift XRT observation shows substantial variability, by over a factor of two within less than 100 seconds. Spectral fitting (excluding 17 arcseconds around the central piled-up core) gives  $N_H=4.4\pm 0.4e21$  cm<sup>-2</sup> (slightly higher than the cluster  $N_H$  of  $2.4e21$ ), photon index=1.41±0.07, and 0.5-10 keV (unabsorbed) flux of  $5.7\pm 0.2e-10$  erg/cm<sup>2</sup>/s. For M28's distance of 5.5 kpc (Harris, 1996, AJ, 112, 1487; 2010 edition), this gives an inferred 0.5-10 keV luminosity of  $2.1\pm 0.1e36$  ergs/s.

## Related

- 5086 [Coherent Pulsations and Burst Oscillations in the Millisecond Pulsar IGR J18245-2452/PSR J1824-2452 in M28](#)
- 5069 [The transient low-mass X-ray binary IGR J18245-2452 is again active as a radio pulsar](#)
- 5068 [Burst Oscillations in the X-Ray Binary IGR J18245-2452 in M28](#)
- 5045 [A Chandra observation of the neutron-star transient IGR J18245-2452 in M28](#)
- 5031 [HST search of the region around IGR J18245-2452](#)
- 5003 [The optical counterpart to the transient IGR J18245-2452 in the globular cluster M28](#)
- 4981 [ATCA radio observation of the region around IGR J18245-2452](#)
- 4964 [Optical amateur observations of the field of IGR J18245-2452 in M28](#)
- 4961 [MAXI/GSC detected two X-ray bursts from IGR J18245-2452 in M28](#)
- 4960 [IGR J18245-2452: an accreting neutron star and thermonuclear burster in M28](#)
- 4959 [A type-I X-ray burst detected by Swift/XRT from the direction of IGR J18245-2452](#)
- 4934 [Possible 6.6-mHz modulation in IGR J18245-2452](#)
- 4929 [Swift observations of IGR J18245-2452](#)
- 4927 [IGR J18245-2452 is a new transient located in the core of the globular cluster M28](#)
- 4925 [IGR J18245-2452: a new hard X-ray transient discovered by INTEGRAL](#)
- 4924 [INTEGRAL/IBIS detects a new hard X-ray transient IGR J17446-2948 in the Galactic Center region](#)
- 4000 [INTEGRAL Galactic Bulge monitoring detects activity from XMMU J174445.5-295044 with JEM-X](#)

[ [Telegram Index](#) ]

R. E. Rutledge, Editor-in-Chief

Derek Fox, Editor

Mansi M. Kasliwal, Co-Editor

rrutledge@astronomerstelegram.org

dfox@astronomerstelegram.org

mansik@astronomerstelegram.org

ATel #4927: IGR J18245-2452 is a new transient located in the core of the globular cluster M28