



**UvA-DARE (Digital Academic Repository)**

**The biochemistry and genetics of floral scent production as part of the petunia pollination syndrome**

Shaipulah, N.F.M.

[Link to publication](#)

*Citation for published version (APA):*

Shaipulah, N. F. M. (2018). The biochemistry and genetics of floral scent production as part of the petunia pollination syndrome

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

# CHAPTER 6

## General discussion

1. Circadian rhythm of floral scent

*How does the earlier expression ODO1 affect the composition of the FVBP?*

*Does the EE in wild petunia contribute to late emission of FVBP?*

2. The metabolic connection between scent and color biosynthesis

3. Outlook

## GENERAL DISCUSSION

Different species of plants advertise unique floral features which specialize in attracting particular animals or group of animals. A suite of floral traits such as color, scent, shape and nectar volume have evolved to maximize pollination by animals. Ensembles of these floral traits demonstrate the adaptive value in pollination syndromes (Fenster *et al.*, 2004).

Among the traits related to a pollination syndrome, floral scent is known as an honest signal, that associates with the quality and quantity of nectar (Wright and Schiestl, 2009). Floral volatiles serve as signals to specific pollinators through emission at a specific time of the day when these pollinators are active. In petunia, floral scent production is temporally regulated by a few transcription factors. Although many studies have reported the rhythmic emission of floral volatiles in plants (Verdonk *et al.*, 2003, Effmert *et al.*, 2005, Kong *et al.*, 2012, Whitehead and Peakall, 2014), only for a few species the contribution of circadian clock in floral scent has been demonstrated (Kolossova *et al.*, 2001a, Fenske *et al.*, 2015, Yon *et al.*, 2016). In recent work, a clock gene, *LATE ELONGATED HYPOCOTYL (LHY)* has been shown to involve in regulating the timing of emission in petunia and *Nicotiana attenuata* flowers (Fenske *et al.*, 2015, Yon *et al.*, 2016).

### 1. Circadian rhythm of floral scent

The circadian clock has been shown to be involved in regulating volatile emission in petunia flowers (*P.hybrida* cv. Mitchell, *P.axillaris*) and snapdragon (*Antirrhinum majus*) (Kolossova *et al.*, 2001a, Oyama-Okubo *et al.*, 2005, Fenske *et al.*, 2015), but molecular components of this “scent” circadian clock have only been discovered recently (Fenske *et al.*, 2015). The emission of *P.hybrida* cv. Mitchell floral volatile benzenoids /phenylpropanoids (FVBP) is rhythmic and peaks in the dark (Verdonk *et al.*, 2003), whereas snapdragon flowers rhythmically emit methylbenzoate with its maximum peak during the day (Kolossova *et al.*, 2001a). The rhythmic emission by these two plant species during night and day period coincides with foraging activities of hawkmoths and bees, respectively. The question remains: how is the rhythm of floral scent emission achieved?

FVBP emission in *P.hybrida* cv. Mitchell shows a circadian oscillation in continuous dark, but not in continuous light, suggesting that the timing of FVBP genes expression is influenced by the circadian clock (Fenske *et al.*, 2015). In general, the expression of circadian clock genes maintain rhythmicity within periods of 24h in continuous light or darkness (McClung, 2006). This is the case for snapdragon flowers, for which the emission of methylbenzoate remains rhythmic in continuous light and dark, indicating that the circadian clock is more pronounced in these flowers (Kolossova *et al.*, 2001a). *Phenylacetaldehyde synthase (PAAS)*, *S-adenosyl-L-methionine:benzoic acid/salicylic acid carboxyl methyltransferase (BSMT)*, *benzoyl-CoA:benzylalcohol/2-phenylethanol benzoyltransferase (BPBT)* and *isoeugenol synthase (IGS)* are biosynthetic genes encoding the enzymes in the final steps for the production of phenylacetaldehyde, methylbenzoate, benzylbenzoate and isoeugenol respectively in *P.hybrida* cv. Mitchell (Negre *et al.*, 2003, Boatright *et al.*, 2004, Kaminaga *et al.*, 2006, Koeduka *et al.*, 2008). The expression of these genes oscillate in the light and dark (L/D) cycle, but *BSMT*, *IGS* and *PAAS* do not sustain this oscillation in continuous dark (Fenske *et al.*, 2015, Cheng *et al.*, 2016). In snapdragon flowers, *BAMT* activity does not oscillate during the L/D cycle (Kolossova *et al.*, 2001b), indicating these biosynthetic genes are not an oscillation-determining factors. Interestingly, the levels of benzoic acid (BA), the precursor of

methylbenzoate change rhythmically in L/D cycle in *P.hybrida* cv. Mitchell and snapdragon flowers, with maximum levels at night and day, respectively. The BA rhythmicity is sustained in continuous dark, correlating with the emission of methylbenzoate (Kolosova *et al.*, 2001a). The level of L-phenylalanine (Phe), the precursor from which all FVBPs are derived, in the *P.hybrida* cv. Mitchell and snapdragon flowers also oscillates in L/D cycle with maximum peaks at night and day, respectively, concurrent with the timing of FVBP emission (Kolosova *et al.*, 2001a, Maeda *et al.*, 2010). The Phe level also peaks at night in *Nicotiana attenuata* flowers, correlating with the night emission of benzyl acetone (Kim *et al.*, 2011). Together, this suggests that the “scent” circadian clock influences the precursor availability and additionally the expression of FVBP genes in the final step of benzenoid/phenylpropanoid pathway.

The genes encoding enzymes in the upstream pathways and the regulators that control the precursor genes may control the precursor levels at night in *P.hybrida* cv. Mitchell. The transcript levels of the shikimate/arogenate pathway genes and upstream genes of FVBP biosynthesis: 5-enolpyruvylshikimate-3-phosphate (*EPSPS*), 3-deoxy-D-arabino-heptulosonate-7-phosphate (*DAHPS*) and arogenase dehydratase (*ADT*), and phenylalanine ammonia lyase (*PAL*), cinnamate 4-hydroxylase (*C4H*), 4-coumarate CoA ligase (*4CL*) display oscillatory patterns in L/D cycle, in which the highest transcript levels are 1 to 3 hour before the onset of dark period. Importantly, these genes sustain their rhythmic expression in continuous dark, suggesting the circadian emission of FVBP at night is mainly regulated in the upstream pathway (Fenske *et al.*, 2015, Cheng *et al.*, 2016). For a few of the precursor genes, the transcription factors that regulate them have been identified. The R2R3-MYB ODORANT1 (*ODO1*) regulates the shikimate pathway in *P.hybrida* cv. Mitchell, by activating the *EPSPS* promoter (Verdonk *et al.*, 2005) (Fig. 1). Although a direct interaction was not shown in flowers, *ODO1* is able to activate the *EPSPS* promoter in petunia leaves (Verdonk *et al.*, 2005). Down-regulation of *ODO1* reduced the expression of shikimate pathway genes and emission of FVBP, as well as benzoic acid pools (Verdonk *et al.*, 2005). The role of *ODO1* in regulating petunia volatiles has been further supported by the identification of this gene as a major Quantitative Trait Loci (QTL) in scent production (Klahre *et al.*, 2011). Two MYB binding sites (MBS) in the *ODO1* promoter are necessary for high promoter activity in fragrant petunia (Van Moerkercke *et al.*, 2011). In addition, *P.hybrida* cv. Mitchell promoter contains two evening elements (EEs), which are located directly downstream of each MBS (Chapter 4). The EEs are potential binding sites for the circadian clock transcription factor, LATE ELONGATED HYPOCOTYL (*LHY*) in *P.hybrida* cv. Mitchell flower (Fenske *et al.*, 2015). Reduced expression of *LHY* leads to early production of scent, while constitutive expression of *LHY* leads to a strong reduction of scent production (Fenske *et al.*, 2015). Chapter 4 shows that transgenic Mitchell plants with an additional copy of *ODO1*, driven by its own promoter, but with the EEs deleted, expressed *ODO1* earlier and emitted FVBP earlier, but the expression of most biosynthetic genes was not affected. Furthermore, these transgenic flowers emitted more benzaldehyde and less benzylbenzoate and isoeugenol (Chapter 4). This phenotype has not been observed in suppression lines of *LHY*. Reduced expression of *LHY* not only shifted the peak expression of *ODO1*, but other FVBP genes as well (*EPSPS*, *chorismate mutase 1(CMI)*, *ADT*, *PAL*, *3-ketoacyl-CoA ketothiolase (KATI)* and *BPBT*) toward the morning (Fenske *et al.*, 2015). *LHY* binds to the EE sites in other promoters (*EPSPS* and *IGS*) as well, and possibly can bind to other promoters of genes in FVBP pathway, suggesting that *LHY* is likely setting the phase of *ODO1*, *EPSPS* and other FVBP genes for evening expression (Fenske *et al.*, 2015). Chapter 4 shows that lack of EEs did not lead to arrhythmic expression of *ODO1*, suggesting that other circadian clock factors than *LHY* are also involved regulating the rhythmic expression of


*ODO1*. Two clock homologs in *P.hybrida* cv. Mitchell, GIGANTEA (GI) and PSEUDO RESPONSE REGULATOR 5 (PRR5) showed same expression pattern as *ODO1*, peaking in the evening, suggesting LHY and these circadian clock genes may work in concert to regulate rhythmic FVBP production in *P.hybrida* cv. Mitchell (Fenske *et al.*, 2015). The R2R3-MYBs *EOBI* and *EOBII* that activate the *ODO1* promoter (Van Moerkercke *et al.*, 2011, Spitzer-Rimon *et al.*, 2012) do not oscillate in continuous dark (Fenske *et al.*, 2015), hence, we exclude their contribution. It seems that the interaction of LHY with the EEs in the *ODO1* promoter, determines the timing of *ODO1* expression and the downstream precursor genes in the hours preceding production of FVBPs.

*How does the earlier expression ODO1 affect the composition of the FVBP?*

As described above, benzoic acid (BA) levels in snapdragon and *P.hybrida* cv. Mitchell flowers correspond with the rhythmic emission of methylbenzoate during the day and night, respectively (Kolossova *et al.*, 2001a). However, it is still a question which pathways contribute to BA biosynthesis. Genetic evidence showed that BA production in *P.hybrida* cv. Mitchell flowers dominantly occurs via the  $\beta$ -oxidative pathway; cinnamoyl-CoA  $\rightarrow$  3-hydroxy-3-phenylpropanoyl-CoA (3H3PP-CoA)  $\rightarrow$  3-oxo-3-phenylpropanoyl-CoA (3O3PP-CoA)  $\rightarrow$  Benzoyl-CoA  $\rightarrow$  BA in peroxisome during the night (Orlova *et al.*, 2006, Van Moerkercke *et al.*, 2009, Colquhoun *et al.*, 2012, Qualley *et al.*, 2012) (Fig. 2-Chapter 1). However, BA can also be synthesized via non-  $\beta$ -oxidative pathway, with benzaldehyde as intermediate, which has been reported in snapdragon (Long *et al.*, 2009). Although there is an argument that conversion of exogenous supplied of benzaldehyde to BA cannot be taken as *in planta* evidence (Van Moerkercke *et al.*, 2009), different mechanism to synthesize BA during day and night might exist. Moreover, flux analysis showed that BA can be produced via both  $\beta$ -oxidative and non-  $\beta$ -oxidative pathway in *P.hybrida* cv. Mitchell flowers and can vary over L/D cycle (Boatright *et al.*, 2004, Orlova *et al.*, 2006).

In chapter 4, I showed that earlier expression of *ODO1* not only resulted in earlier emission of FVBP, but also elevated the emission levels of benzaldehyde in *P.hybrida* cv. Mitchell (Fig. 1). The question is what makes the change in flux to benzaldehyde production? I will discuss the possible mechanisms in this section.

Early expression of *ODO1* resulted in earlier expression of *EPSPS*, which thus may lead to the early synthesis of Phe in petals (not measured). Of note is the fact that *P.hybrida* cv. Mitchell is derived from *P.axillaris*, and that benzenoid/phenylpropanoid composition and timing of emission are similar to that of *P.axillaris* (Verdonk *et al.*, 2003). Rhythmic FVBP emission by *P.axillaris* also occurs in continuous light (Hoballah *et al.*, 2005), suggesting that circadian clock regulates the FVBP is more robust in *P.axillaris* than *P.hybrida* cv. Mitchell. The clock phenotype can be different among *Petunia* species. For instance, *P.integrifolia* flowers emits solely benzaldehyde during the day (PhD thesis M. Boersma, unpublished) which is light dependent (Hoballah *et al.*, 2005). In *P.hybrida* cv. Mitchell, a combination of both circadian and environmental factors may regulate the rhythmic emission of FVBP.



**Figure 1:** Differences in rhythmic pattern and composition of benzenoid/phenylpropanoid in *P.hybrida* cv. Mitchell and transgenic evening elements mutated (mEE) Mitchell lines. Transcript levels of *ODO1* and *EPSPS* are evening-phased in *P.hybrida* cv. Mitchell and midday-phased in mEE lines. Phe and BA pools in *P.hybrida* cv. Mitchell are night-phased, preceding the emission of volatile compounds (red letters-underlined). Blue and red vertical lines indicate MYB binding sites (MBS) and EEs motifs in promoter, respectively. White and grey boxes represent light and dark period, respectively. Abbreviation: *EPSPS*: 5-enolpyruvylshikimate-3-phosphate synthase; BA: Benzoic acid; *BPBT*: benzoyl-CoA:benzylalcohol/2-phenylethanol benzoyltransferase; *BSMT*: S-adenosyl-L-methionine:benzoic acid/salicylic acid carboxyl methyltransferase; *CCoAOMT*: Caffeoyl-CoA O-methyltransferase; *CNL*: Cinnamate-CoA ligase; *EOBII*: EMISSION OF BENZENOID II; *IGS*: Isoeugenol synthase; *LHY*: LATE ELONGATED HYPOCOTYL; *ODO1*: *ODORANT1*; Phe: Phenylalanine; *t*-CA: trans- cinnamic

It has been shown that light and temperature also contribute to rhythmic emission of *P.hybrida* cv. Mitchell (Cheng *et al.*, 2016). As Phe accumulates during the day, the FVBP genes are also expressed early in the light as shown in Chapter 4, including *PAL* in the *P.hybrida* cv. V26 line. High temperature and light exposure during the day have a negative effect on *PAL* transcript levels and enzymatic activity (Lo Piero *et al.*, 2005, Boo *et al.*, 2011). High temperatures lead to lower anthocyanin levels due to the lower *PAL* activity in plants (Shaked-Sachray *et al.*, 2002, Lo Piero *et al.*, 2005). A similar result has been reported for volatile biosynthesis in *P.hybrida* cv. Mitchell. For instance, increasing the ambient temperature from night to day can reduce the emission of floral volatiles produced by petunia (Cna'ani *et al.*, 2015a). Interestingly, the

negative effect of temperature was also robust at endogenous levels of volatiles during the day, indicating the expression of FVBP genes is also influenced by temperature (Ben Zvi *et al.*, 2008, Sagae *et al.*, 2008). PAL is the first committed enzyme that catalyzes the conversion of Phe to *t*-cinnamic acid (*t*-CA) (Ritter and Schulz, 2004) and *t*-CA must be in enough supply for benzenoid production in petunia petals via  $\beta$ -oxidative pathway. It seems that, during the day, light and temperature may take part in regulation of FVBP emission by controlling the flux distribution at benzenoid/phenylpropanoid pathway. Instead of importing the precursor into peroxisome, *t*-CA is redirected to the non- $\beta$ -oxidative pathway and resulting in more benzaldehyde and benzylalcohol production in the cytosol during the day, since formation of *t*-CA from Phe is a one-step conversion (Coquoz *et al.*, 1998). Moreover, the flux contribution from benzaldehyde to BA appeared to be higher in the presence of light than dark in *BPBT* silencing lines, supporting that the benzenoid/phenylpropanoid branch is also light dependent (Orlova *et al.*, 2006). Thus, Phe levels and flux distribution analysis during day and night should be performed to fully understand the molecular mechanisms of FVBP production in peroxisomal  $\beta$ -oxidative and non- $\beta$ -oxidative pathway in the cytosol. The internal volatiles are yet to be quantified in continuous light and dark, albeit the rhythm of both emission and endogenous pools were same in L/D cycle (Fenske *et al.*, 2015).

#### *Does the EE in wild petunia contribute to late emission of FVBP?*

A polymorphism in a single gene of a flower trait has been reported result in the attraction of different pollinators. QTL analysis in petunia suggests a shift in pollination syndromes can be explained by polymorphisms of transcription factors involved in floral color, UV absorbance or scent production (Hoballah *et al.*, 2007, Klahre *et al.*, 2011, Sheehan *et al.*, 2016). The mutation in the coding region of a R2R3-MYB transcription factor ANTHOCYANIN2 (AN2) of *P.axillaris* leads to a shift from purple to white flowers (Quattrocchio *et al.*, 1999, Hoballah *et al.*, 2007). In recent work, mutations in the *MYB-FL* promoter enhanced the expression of flavonol synthase (FLS), and in turn lead to enhanced UV absorbance in *P.axillaris* (Sheehan *et al.*, 2016). Another R2R3-MYB transcription factor, *ODO1*, localizes to a QTL on chromosome VII, which is responsible for scent production in petunia flowers (Klahre *et al.*, 2011). These studies demonstrated the strong influence of mutations in flower traits on the preferences of pollinators.

The absence of fragrance in *P.exserta* is caused by a polymorphism in *cis*-regulatory elements of *ODO1* (Klahre *et al.*, 2011), which is likely caused by mutation of MBS within the 1.2 kbp *ODO1* promoter (Van Moerkercke *et al.*, 2011). Presence of EEs in the same region can be another factor that control night production and emission of volatiles, as well as volatile composition in *P.axillaris*. Therefore, we compared the 1.2 kbp of promoter sequences of *ODO1* among wild and cultivars of petunia (Fig. 2) and summarized the number of EEs in Table 1. The numbers of EEs varied among fragrant petunia and the differences cannot be correlated with day and night emission of volatiles (Table 1). For instance, *P.axillaris* N and *P.integrifolia inflata* S6 have one EE although the flowers emit volatiles during night and day respectively (Hoballah *et al.*, 2005). *ODO1* transcript levels in both petunias are high at noon (PhD thesis M. Boersma, unpublished) similar to those of *P.hybrida* cv. V26 (Chapter 4). This suggests that *ODO1* transcription is not the primary factor that regulates night emission in *P.axillaris*. EEs can also be found in *EPSPS* and *IGS* promoters (Fenske *et al.*, 2015), and possibly in other promoter of FVBP genes as well. LHY may repress the morning expression of the upstream precursor genes (e.g *EPSPS*) which directly controls the availability of precursor at night in *P.axillaris*. Furthermore, *P.axillaris* and *P.inflata* have 2 and 3 copies of the clock gene,

*gigantea* (*GI*), respectively in their genome (Bombarely *et al.*, 2016). *GI* transcripts and proteins are modulated by circadian rhythm, light input and temperature response (Fowler *et al.*, 1999, David *et al.*, 2006, Gould *et al.*, 2006), suggesting that the extra copy of *GI* in *P.inflata* may have a new biological role. It would be interesting to investigate the role of *GI* in regulating the rhythm of volatile emission of *P.axillaris* and *P.inflata*.

Although *ODO1* seems to be one of the molecular players controlling total scent production of benzenoid/phenylpropanoid by *P.hybrida* cv. Mitchell flowers (Verdonk *et al.*, 2005), it may not directly determine the composition of the FVBP bouquet. In wild petunias, *P.axillaris* and *P.integrifolia*, the different levels of benzenoid compounds are unlikely caused by *ODO1* regulation. In Chapter 5, we showed that the higher levels of benzylbenzoate in early during the day are likely caused by a gene on chromosome II. Although we could not correlate *BPBT1* and *BPBT2* expression levels in *P.axillaris* and the selected RILs with high, early benzylbenzoate production, we cannot exclude that these genes are causally related to the QTL. Interestingly, the overall benzaldehyde levels were higher in RILs, but methylbenzoate levels were lower than in *P.axillaris* (Chapter 5). *P.integrifolia* lacks *BSMT* expression, but has an active *ODO1* and *cinnamate-CoA ligase* (*CNL*) (Amrad *et al.*, 2016), thus loss of active *BSMT* copy on chromosome V in the RILs may lead to the absence of methylbenzoate. The methylbenzoate production can be restored in an F<sub>1</sub> of *P.integrifolia* x *P.exserta*, which has an active copy of *BSMT*, but lacks *ODO1* and *CNL* expression, which also causes a strong reduction of benzaldehyde production (Amrad *et al.*, 2016). This suggests that activation of *BSMT* (or/and *BPBT*) could redirect metabolic flux away from benzaldehyde toward methylbenzoate production (Amrad *et al.*, 2016). If no *BSMT* activity is present the default pathway leads to benzaldehyde production. Interestingly, tight genetic linkage between loci on chromosome II that specify scent production, UV absorbance and reproductive organs morphology have been identified, suggests that clustered floral traits create a “speciation island” able to respond to specific pollinators (Hermann *et al.*, 2013). Many FVBP genes are on chromosome II (*phenylacetaldehyde synthase* (*PAAS*), *CNL*, *BPBT*, *PAL1* and *IGS*), indicating that these genes may have strong selection on pollination syndrome traits to keep separating two closely related species in a sympatric population (Hermann *et al.*, 2013). For example: loss of functional *CNL* is very important for transition from hawkmoth to hummingbird pollination (Amrad *et al.*, 2016), and a QTL on chromosome II (Chapter 5) may have contributed to the pollinator shift from bee to hawkmoth.



```

S7          TTCCAATGTAATTC CGCAG-TTAGGGCGTGG-CCTTGTATATCAATAATTC CAAAAAGT 1141
AxillarisN TTCCAATGTAATTC CGCAA-TTAGGACGTGGTCCTTGTATATCAATAATTC CAAAAAGT 1110
Exserta    TTTCAATGTAATTC CGCAG-TTAGGGCG-GG-CCTTGTATATCAATAATTC CAAAAAGT 1130
Inflata    TTCCAATGTAATTC CGCAA-TTAGGACGTGGTCCTTGTATATCAATAATTC CAAAAAGT 1111
W115      TTTCAATGTAATTC CGCAGTTTAGGGCGTGG-CCTTGTATATCAATAATTC CAAAAAGT 1144
R27        TTCCAATGTAATTC CGCAA-TTAGGGCGTGGTCCTTGTATATCAATAATTC CAAAAAGT 1129
V26        TTCCAATGTAATTC CGCAA-TTAGGACGTGGTCCTTGTATATCAATAATTC CAAAAAGT 1110
** .*****.*****.*** ** *****

S7          AC-ATAAATAGGACATAAACCTAATAAAAATATCTTGATACATAATATACCTCTTACT 1081
AxillarisN ACGAAAAATAAGACATAAACCTAATAAAAACATATCTCCATGCATAATATACTTCTTACT 1049
Exserta    AC-ATAAATAGGAC-TAAACCTAATAAAAATATCTTGATACATAATATACCTCTTACT 1071
Inflata    ACGAAAAATAAGACATAAACCTAATAAAAACATATCTCCATGCATAATATACTTCTTACT 1050
W115      ACGATAAATAGGACATAAACCTAATAAAAATATCTTGATACATAATATACCTCTTACT 1083
R27        ACGAAAAATAAGACATAAACTTAATAAAACATATCTCCATGCATAATATACTTCTTACT 1068
V26        ACGAAAAATAAGACATAAACCTAATAAAAACATATCTCCATGCATAATATACTTCTTACT 1049
** * *****.*** *****.***** *****. ** .*****.*****

S7          CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCC-AAATTAAGTACGAAAA 1021
AxillarisN CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCCaAcATTATGTACGAGAAA 988
Exserta    CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCCAAATTAAGTACGAAAA 1010
Inflata    CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCC-ACATTATGTACGAAAA 1010
W115      CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCCAAATTAAGTACGAAAA 1022
R27        CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCCAAATTAAGTACGAGAAA 1007
V26        CAATGCAGTTAGGGCGTGGTCCTTGTATATTAATTAATTCCAACATTATGTACGAGAAA 988
*****

S7          AATTGGACATAAACCAATAAAAAATATCTC 1009
AxillarisN AATAGGACATAAACCTAATAAAAATATCTC 976
Exserta    AATTGGACATAAACCAATAAAAAATATCTC 998
Inflata    AATAGGACATAAACCTAATAAAAATATCTC 998
W115      AATTGGACATAAACCTAATAAAAATATCTC 1010
R27        AATAGGACATAAAACTAATAAAAAATATCTC 995
V26        AATAGGACATAAACCTAATAAAAATATCTC 976
*** ***** * *****

```

**Figure 2:** Sequence alignment of part of the ODO1 promoter from wild petunias (*P.axillaris parodii* S7, *P.axillaris* N, *P.integrifolia inflata* S6, *P.exserta*) and cultivars (*P.hybrida* cv. Mitchell, *P.hybrida* cv. V26, *P.hybrida* cv. R27). The alignment was generated using ClustalW (<http://www.genome.jp/tools-bin/clustalw>). The sequences upstream of the ATG, approximately between 1204 and 995, are shown. Red letters indicate the perfect sequence of MYB binding site (MBS) and blue boxes indicate the perfect sequence of evening elements (EEs). Abbreviations: *P.axillaris parodii* S7 (S7); *P.axillaris* N (AxillarisN); *P.exserta* (Exserta); *P.integrifolia inflata* S6 (Inflata); *P.hybrida* cv. Mitchell (W115); *P.hybrida* cv. V26 (V26); *P.hybrida* cv. R27 (R27).

**Table 1:** Summary of *ODO1* promoter sequences from 1204 bp to 995 bp upstream of ATG

Species/cultivars	MBS	EE	ODO1 expression	Peak of expression	Fragrance
<i>P.axillaris parodii</i> S7	1	2	++	evening	Fragrant
<i>P.axillaris</i> N	2	1	++	evening	Fragrant
<i>P.exserta</i>	1	2	-	-	Non-fragrant
<i>P.integrifolia inflata</i> S6	2	1	++	midday	Fragrant
<i>P.hybrida</i> cv. W115	2	2	++	evening	Fragrant
<i>P.hybrida</i> cv. R27	1	1	-	-	Non-fragrant
<i>P.hybrida</i> cv. V26	2	1	++	midday	Fragrant

## 2. The metabolic connection between scent and color biosynthesis

Flower color and scent are essential for the attraction of pollinators and hence for high pollination efficiency. In petunia flowers, both floral volatile and anthocyanin biosynthesis have been intensively studied in last few decades, including the characterization of transcription factors and biosynthetic genes (Schuurink *et al.*, 2006, Albert *et al.*, 2014, Muhlemann *et al.*, 2014). Benzenoid/phenylpropanoid and anthocyanin biosynthesis are derived from phenylpropanoid pathway and shared common precursors, Phe and 4-coumaryl CoA.

Anthocyanin is a class of flavonoid, which is synthesized in cytoplasm and stored in vacuoles (Mueller *et al.*, 2000). In petunia flowers, anthocyanins are produced at an early stage of flower development and the synthesis is ceased upon flower opening. The R2R3 MYB transcription factors AN2 and AN4 are exclusively expressed in petals and anthers, respectively, and form a complex with basic-helix-loop-helix, AN1, and WD-repeat protein, AN11 to regulate the anthocyanin pathway in petunia (de Vetten *et al.*, 1997, Quattrocchio *et al.*, 1999, Spelt *et al.*, 2000). Unlike the anthocyanin biosynthesis, FVBP production only begins during anthesis when flower are fully developed and reproductive organs are mature, hence ready for pollination (Verdonk *et al.*, 2003). This temporal separation could be a mechanism to prevent metabolic competition between these two pathways in flowers (Verdonk *et al.*, 2005). More arguments against metabolic competition have been reported between anthocyanin and benzenoid/phenylpropanoid production in petunia flowers. Silencing of EOBII in *P.hybrida* line P720 did not affect the anthocyanin levels, although FVBP production and emission were lower than controls (Spitzer-Rimon *et al.*, 2010).

A crosstalk between benzenoid/phenylpropanoid volatiles and anthocyanin can be anticipated on the basis of their common biochemical pathway. Manipulation of the R2R3 MYB transcription factor PH4 and phenylpropanoid biosynthetic genes indeed redirected the metabolic flux from one branch to another. Final petal color is regulated by PH4 that interacts with AN1 and AN11 to control acidification of the vacuole (Quattrocchio *et al.*, 2006). In contrast to AN2, PH4 plays role in floral pigmentation and volatile emission (Cna'ani *et al.*, 2015b). It was suggested that PH4 is involved in flower pigmentation via vacuolar acidification during bud development and shift towards scent emission following anthesis of flowers (Cna'ani *et al.*, 2015b). Like the other FVBP transcription factors, *PH4* transcript levels is developmentally regulated, the transcript in purple petunia, *P.hybrida* cv. V30 increased

throughout the developmental stages and reached the highest levels after anthesis, preceding the peak volatile emission of petunia (Cna'ani *et al.*, 2015b). PH4 plays no role in activation of structural anthocyanin genes, thus functional *AN2* is a “must” requirement to allow anthocyanin synthesis in petunia petals (Quattrocchio *et al.*, 2006).

In last decade, suppression of *flavanone-3-hydrolase (F3H)* in carnation has already suggested a metabolic connection between anthocyanin and benzenoid biosynthesis (Zuker *et al.*, 2002). Suppression lines of *F3H* showed a loss of anthocyanin pigments in carnation flowers and more methylbenzoate emission than in control lines (Zuker *et al.*, 2002). Silencing of *chalcone synthase (CHS)*, a gene encoding the first committed enzyme in flavonoid biosynthesis, increased the production of (iso)eugenol in cultivated strawberry (Hoffmann *et al.*, 2011). However, pigmentation of petals was severely affected upon suppression of *F3H* and *CHS* in petunia flowers but FVBP levels were not increased (Tsuda *et al.*, 2004, Spitzer *et al.*, 2007), suggesting phenotypic effects may depend on the genes targeted gene. In Chapter 2 we showed that downregulation of one phenylpropane gene, *caffeoyl-CoA O-methyltransferase 1 (CCoAOMT1)* in *P.hybrida* cv. Mitchell can lead to the production of pink flowers and purple leaves and stems. Flowers of these plants produce normal amounts of volatiles, although slightly less eugenol is produced. Although *P.hybrida* cv. Mitchell flowers lack a functional *AN2*, *DFR* might have been activated by PURPLE HAZE (PHZ), a R2R3 MYB transcription factor that regulates anthocyanin pathway in flowers and vegetative tissues (Albert *et al.*, 2011). Most likely the accumulation of a metabolite resulted in PHZ activation. The transcript levels of *CCoAOMT1* can be detected in buds (Chapter 2), thus we predict that the anthocyanin has been synthesized early during flower development and ceased when FVBP production started. Interestingly, introduction of *AN2* into *P.axillaris* had no effect on scent production although the anthocyanin biosynthesis was activated in the flowers (Hoballah *et al.*, 2007). The metabolic connection between flower color and scent in the phenylpropanoid pathway is during development, thus metabolic losses in both pathways can be minimized and on the other hand, plant pollination success can be maximized.

### 3. Outlook

There has been a boost in floral volatile research and petunia has become a perfect model to study the regulation of phenylpropanoid/benzenoid biosynthesis. In recent work, the clock gene *LHY* has been identified as a main regulator in controlling the timing of FVBP regulation in flowers (Fenske *et al.*, 2015, Yon *et al.*, 2016). However, more research efforts are needed to discover how *LHY* and other clock genes interact with other transcription factors and FVBP structural genes and how the changes in the floral clock affect the pollinator choice in field. The role of other clock components in regulation of temporal production of phenylpropanoid/benzenoid remains to be investigated (Bombarely *et al.*, 2016).

Color and scent variations can be part of pollination syndromes that contribute to reproductive isolation between closely related species (Hoballah *et al.*, 2007, Klahre *et al.*, 2011, Byers *et al.*, 2014, Bischoff *et al.*, 2015). Both structural and regulatory genes has been shown to be involved in the natural variation in pollination syndromes (Hoballah *et al.*, 2007, Klahre *et al.*, 2011, Amrad *et al.*, 2016). Mutations in the coding region of *AN2* or suppression of *ODO1* do not have an effect on volatile phenylpropanoids and anthocyanin production, respectively, in spite of the fact that these compounds are derived from a common precursor (Verdonk *et al.*, 2005, Hoballah *et al.*, 2007). Tight regulation, both temporally and developmentally, may contribute to the metabolic separation between these pathways. Although

manipulation of *PH4* and structural genes (*CCoAOMT1*) have shown a metabolic connection between color and scent, more studies on metabolic channeling in the phenylpropanoid pathway should be performed. Moreover, it would be of great interest to investigate how the biochemical and genetic changes affect the pollinator perception to provide more evidence on co-evolution between plant and pollinators.

## REFERENCES

- Albert, N.W., Davies, K.M., Lewis, D.H., Zhang, H., Montefiori, M., Brendolise, C., Boase, M.R., Ngo, H., Jameson, P.E. and Schwinn, K.E. (2014) A Conserved Network of Transcriptional Activators and Repressors Regulates Anthocyanin Pigmentation in Eudicots. *Plant Cell*, **26**, 962-980.
- Albert, N.W., Lewis, D.H., Zhang, H., Schwinn, K.E., Jameson, P.E. and Davies, K.M. (2011) Members of an R2R3-MYB transcription factor family in *Petunia* are developmentally and environmentally regulated to control complex floral and vegetative pigmentation patterning. *Plant J*, **65**, 771-784.
- Amrad, A., Moser, M., Mandel, T., de Vries, M., Schuurink, R.C., Freitas, L. and Kuhlemeier, C. (2016) Gain and Loss of Floral Scent Production through Changes in Structural Genes during Pollinator-Mediated Speciation. *Curr Biol*, **26**, 3303-3312.
- Ben Zvi, M.M., Negre-Zakharov, F., Masci, T., Ovadis, M., Shklarman, E., Ben-Meir, H., Tzfira, T., Dudareva, N. and Vainstein, A. (2008) Interlinking showy traits: co-engineering of scent and colour biosynthesis in flowers. *Plant Biotechnol J*, **6**, 403-415.
- Bischoff, M., Raguso, R.A., Jürgens, A. and Campbell, D.R. (2015) Context-dependent reproductive isolation mediated by floral scent and color. *Evolution*, **69**, 1-13.
- Boatright, J., Negre, F., Chen, X., Kish, C.M., Wood, B., Peel, G., Orlova, I., Gang, D., Rhodes, D. and Dudareva, N. (2004) Understanding in vivo benzenoid metabolism in *petunia* petal tissue. *Plant Physiol*, **135**, 1993-2011.
- Bombarely, A., Moser, M., Amrad, A., Bao, M., Bapaume, L., Barry, C.S., Bliik, M., Boersma, M.R., Borghi, L., Bruggmann, R., Bucher, M., D'Agostino, N., Davies, K., Druege, U., Dudareva, N., Egea-Cortines, M., Delledonne, M., Fernandez-Pozo, N., Franken, P., Grandont, L., Heslop-Harrison, J.S., Hintzsche, J., Johns, M., Koes, R., Lv, X., Lyons, E., Malla, D., Martinioia, E., Mattson, N.S., Morel, P., Mueller, L.A., Muhlemann, J., Nouri, E., Passeri, V., Pezzotti, M., Qi, Q., Reinhardt, D., Rich, M., Richert-Pöggeler, K.R., Robbins, T.P., Schatz, M.C., Schranz, M.E., Schuurink, R.C., Schwarzacher, T., Spelt, K., Tang, H., Urbanus, S.L., Vandenbussche, M., Vijverberg, K., Villarino, G.H., Warner, R.M., Weiss, J., Yue, Z., Zethof, J., Quattrocchio, F., Sims, T.L. and Kuhlemeier, C. (2016) Insight into the evolution of the Solanaceae from the parental genomes of *Petunia hybrida*. *Nature Plants*, **2**, 16074.
- Boo, H.O., Heo, B.G., Gorinstein, S. and Chon, S.U. (2011) Positive effects of temperature and growth conditions on enzymatic and antioxidant status in lettuce plants. *Plant Sci*, **181**, 479-484.
- Byers, K.J.R.P., Vela, J.P., Peng, F., Riffell, J.A. and Bradshaw, H.D. (2014) Floral volatile alleles can contribute to pollinator-mediated reproductive isolation in monkeyflowers (*Mimulus*). *The Plant journal : for cell and molecular biology*, **80**, 1031-1042.
- Cheng, S., Fu, X., Mei, X., Zhou, Y., Du, B., Watanabe, N. and Yang, Z. (2016) Regulation of biosynthesis and emission of volatile phenylpropanoids/benzenoids in *petunia* *hybrida* flowers by multi-factors of circadian clock, light, and temperature. *Plant Physiol Biochem*, **107**, 1-8.
- Cna'ani, A., Muhlemann, J.K., Ravid, J., Masci, T., Klempien, A., Nguyen, T.T., Dudareva, N., Pichersky, E. and Vainstein, A. (2015a) *Petunia* x *hybrida* floral scent production is negatively affected by high-temperature growth conditions. *Plant Cell Environ*, **38**, 1333-1346.
- Cna'ani, A., Spitzer-Rimon, B., Ravid, J., Farhi, M., Masci, T., Aravena-Calvo, J., Ovadis, M. and Vainstein, A. (2015b) Two showy traits, scent emission and pigmentation, are finely coregulated by the MYB transcription factor PH4 in *petunia* flowers. *New Phytol*, **208**, 708-714.
- Colquhoun, T.A., Marciniak, D.M., Wedde, A.E., Kim, J.Y., Schwieterman, M.L., Levin, L.A., Van Moerkercke, A., Schuurink, R.C. and Clark, D.G. (2012) A peroxisomally localized acyl-activating enzyme is required for volatile benzenoid formation in a *Petunia* *hybrida* cv. 'Mitchell Diploid' flower. *J Exp Bot*, **63**, 4821-4833.

- Coquoz, J.-L., Buchala, A. and Métraux, J.-P.** (1998) The Biosynthesis of Salicylic Acid in Potato Plants. *Plant Physiology*, **117**, 1095-1101.
- David, K.M., Armbruster, U., Tama, N. and Putterill, J.** (2006) Arabidopsis GIGANTEA protein is post-transcriptionally regulated by light and dark. *FEBS Lett*, **580**, 1193-1197.
- de Vetten, N., Quattrocchio, F., Mol, J. and Koes, R.** (1997) The an11 locus controlling flower pigmentation in petunia encodes a novel WD-repeat protein conserved in yeast, plants, and animals. *Genes Dev*, **11**, 1422-1434.
- Effmert, U., Große, J., Röse, U.S.R., Ehrig, F., Kägi, R. and Piechulla, B.** (2005) Volatile composition, emission pattern, and localization of floral scent emission in *Mirabilis jalapa* (Nyctaginaceae). *American Journal of Botany*, **92**, 2-12.
- Fenske, M.P., Hewett-Hazelton, K.D., Hempton, A.K., Shim, J.S., Yamamoto, B.M., Riffell, J.A. and Imaizumi, T.** (2015) Circadian clock gene LATE ELONGATED HYPOCOTYL directly regulates the timing of floral scent emission in *Petunia*. *Proc Natl Acad Sci U S A*, **112**, 9775-9780.
- Fenster, C.B., Armbruster, W.S., Wilson, P., Dudash, M.R. and Thomson, J.D.** (2004) Pollination Syndromes and Floral Specialization. *Annual Review of Ecology, Evolution, and Systematics*, **35**, 375-403.
- Fowler, S., Lee, K., Onouchi, H., Samach, A., Richardson, K., Morris, B., Coupland, G. and Putterill, J.** (1999) GIGANTEA: a circadian clock-controlled gene that regulates photoperiodic flowering in *Arabidopsis* and encodes a protein with several possible membrane-spanning domains. *EMBO J*, **18**, 4679-4688.
- Gould, P.D., Locke, J.C., Larue, C., Southern, M.M., Davis, S.J., Hanano, S., Moyle, R., Milich, R., Putterill, J., Millar, A.J. and Hall, A.** (2006) The molecular basis of temperature compensation in the *Arabidopsis* circadian clock. *Plant Cell*, **18**, 1177-1187.
- Hermann, K., Klahre, U., Moser, M., Sheehan, H., Mandel, T. and Kuhlemeier, C.** (2013) Tight genetic linkage of prezygotic barrier loci creates a multifunctional speciation island in *Petunia*. *Curr Biol*, **23**, 873-877.
- Hoballah, M.E., Gubitzi, T., Stuurman, J., Broger, L., Barone, M., Mandel, T., Dell'Olivo, A., Arnold, M. and Kuhlemeier, C.** (2007) Single gene-mediated shift in pollinator attraction in *Petunia*. *Plant Cell*, **19**, 779-790.
- Hoballah, M.E., Stuurman, J., Turlings, T.C., Guerin, P.M., Connetable, S. and Kuhlemeier, C.** (2005) The composition and timing of flower odour emission by wild *Petunia axillaris* coincide with the antennal perception and nocturnal activity of the pollinator *Manduca sexta*. *Planta*, **222**, 141-150.
- Hoffmann, T., Kurtzer, R., Skowranek, K., Kiessling, P., Fridman, E., Pichersky, E. and Schwab, W.** (2011) Metabolic engineering in strawberry fruit uncovers a dormant biosynthetic pathway. *Metab Eng*, **13**, 527-531.
- Kaminaga, Y., Schnepf, J., Peel, G., Kish, C.M., Ben-Nissan, G., Weiss, D., Orlova, I., Lavie, O., Rhodes, D., Wood, K., Porterfield, D.M., Cooper, A.J., Schloss, J.V., Pichersky, E., Vainstein, A. and Dudareva, N.** (2006) Plant phenylacetaldehyde synthase is a bifunctional homotetrameric enzyme that catalyzes phenylalanine decarboxylation and oxidation. *J Biol Chem*, **281**, 23357-23366.
- Kim, S.G., Yon, F., Gaquerel, E., Gulati, J. and Baldwin, I.T.** (2011) Tissue specific diurnal rhythms of metabolites and their regulation during herbivore attack in a native tobacco, *Nicotiana attenuata*. *PLoS One*, **6**, e26214.
- Klahre, U., Gurba, A., Hermann, K., Sachsenhofer, M., Bossolini, E., Guerin, P.M. and Kuhlemeier, C.** (2011) Pollinator choice in *Petunia* depends on two major genetic Loci for floral scent production. *Curr Biol*, **21**, 730-739.
- Koeduka, T., Louie, G.V., Orlova, I., Kish, C.M., Ibdah, M., Wilkerson, C.G., Bowman, M.E., Baiga, T.J., Noel, J.P., Dudareva, N. and Pichersky, E.** (2008) The multiple phenylpropene synthases in both *Clarkia breweri* and *Petunia hybrida* represent two distinct protein lineages. *Plant J*, **54**, 362-374.
- Kolosova, N., Gorenstein, N., Kish, C.M. and Dudareva, N.** (2001a) Regulation of circadian methyl benzoate emission in diurnally and nocturnally emitting plants. *Plant Cell*, **13**, 2333-2347.
- Kolosova, N., Sherman, D., Karlson, D. and Dudareva, N.** (2001b) Cellular and subcellular localization of S-adenosyl-L-methionine:benzoic acid carboxyl methyltransferase, the enzyme responsible for biosynthesis of the volatile ester methylbenzoate in snapdragon flowers. *Plant Physiol*, **126**, 956-964.

- Kong, Y., Sun, M., Pan, H.-t. and Zhang, Q.-x.** (2012) Composition and Emission Rhythm of Floral Scent Volatiles from Eight Lily Cut Flowers. *Journal of the American Society for Horticultural Science*, **137**, 376-382.
- Lo Piero, A.R., Puglisi, I., Rapisarda, P. and Petrone, G.** (2005) Anthocyanins accumulation and related gene expression in red orange fruit induced by low temperature storage. *J Agric Food Chem*, **53**, 9083-9088.
- Long, M.C., Nagegowda, D.A., Kaminaga, Y., Ho, K.K., Kish, C.M., Schnepf, J., Sherman, D., Weiner, H., Rhodes, D. and Dudareva, N.** (2009) Involvement of snapdragon benzaldehyde dehydrogenase in benzoic acid biosynthesis. *Plant J*, **59**, 256-265.
- Maeda, H., Shasany, A.K., Schnepf, J., Orlova, I., Taguchi, G., Cooper, B.R., Rhodes, D., Pichersky, E. and Dudareva, N.** (2010) RNAi suppression of Arogenate Dehydratase1 reveals that phenylalanine is synthesized predominantly via the arogenate pathway in petunia petals. *Plant Cell*, **22**, 832-849.
- McClung, C.R.** (2006) Plant circadian rhythms. *Plant Cell*, **18**, 792-803.
- Mueller, L.A., Goodman, C.D., Silady, R.A. and Walbot, V.** (2000) AN9, a Petunia Glutathione S-Transferase Required for Anthocyanin Sequestration, Is a Flavonoid-Binding Protein. *Plant Physiology*, **123**, 1561-1570.
- Muhlemann, J.K., Klempien, A. and Dudareva, N.** (2014) Floral volatiles: from biosynthesis to function. *Plant, Cell & Environment*, **37**, 1936-1949.
- Negre, F., Kish, C.M., Boatright, J., Underwood, B., Shibuya, K., Wagner, C., Clark, D.G. and Dudareva, N.** (2003) Regulation of methylbenzoate emission after pollination in snapdragon and petunia flowers. *Plant Cell*, **15**, 2992-3006.
- Orlova, I., Marshall-Colon, A., Schnepf, J., Wood, B., Varbanova, M., Fridman, E., Blakeslee, J.J., Peer, W.A., Murphy, A.S., Rhodes, D., Pichersky, E. and Dudareva, N.** (2006) Reduction of benzenoid synthesis in petunia flowers reveals multiple pathways to benzoic acid and enhancement in auxin transport. *Plant Cell*, **18**, 3458-3475.
- Oyama-Okubo, N., Ando, T., Watanabe, N., Marchesi, E., Uchida, K. and Nakayama, M.** (2005) Emission mechanism of floral scent in Petunia axillaris. *Biosci Biotechnol Biochem*, **69**, 773-777.
- Qualley, A.V., Widhalm, J.R., Adebessin, F., Kish, C.M. and Dudareva, N.** (2012) Completion of the core beta-oxidative pathway of benzoic acid biosynthesis in plants. *Proc Natl Acad Sci U S A*, **109**, 16383-16388.
- Quattrocchio, F., Verweij, W., Kroon, A., Spelt, C., Mol, J. and Koes, R.** (2006) PH4 of Petunia is an R2R3 MYB protein that activates vacuolar acidification through interactions with basic-helix-loop-helix transcription factors of the anthocyanin pathway. *Plant Cell*, **18**, 1274-1291.
- Quattrocchio, F., Wing, J., van der Woude, K., Souer, E., de Vetten, N., Mol, J. and Koes, R.** (1999) Molecular analysis of the anthocyanin2 gene of petunia and its role in the evolution of flower color. *Plant Cell*, **11**, 1433-1444.
- Ritter, H. and Schulz, G.E.** (2004) Structural basis for the entrance into the phenylpropanoid metabolism catalyzed by phenylalanine ammonia-lyase. *Plant Cell*, **16**, 3426-3436.
- Sagae, M., Oyama-Okubo, N., Ando, T., Marchesi, E. and Nakayama, M.** (2008) Effect of temperature on the floral scent emission and endogenous volatile profile of Petunia axillaris. *Biosci Biotechnol Biochem*, **72**, 110-115.
- Schuurink, R.C., Haring, M.A. and Clark, D.G.** (2006) Regulation of volatile benzenoid biosynthesis in petunia flowers. *Trends Plant Sci*, **11**, 20-25.
- Shaked-Sachray, L., Weiss, D., Reuveni, M., Nissim-Levi, A. and Oren-Shamir, M.** (2002) Increased anthocyanin accumulation in aster flowers at elevated temperatures due to magnesium treatment. *Physiol Plant*, **114**, 559-565.
- Sheehan, H., Moser, M., Klahre, U., Esfeld, K., Dell'Olivo, A., Mandel, T., Metzger, S., Vandenbussche, M., Freitas, L. and Kuhlemeier, C.** (2016) MYB-FL controls gain and loss of floral UV absorbance, a key trait affecting pollinator preference and reproductive isolation. *Nat Genet*, **48**, 159-166.
- Spelt, C., Quattrocchio, F., Mol, J.N. and Koes, R.** (2000) anthocyanin1 of petunia encodes a basic helix-loop-helix protein that directly activates transcription of structural anthocyanin genes. *Plant Cell*, **12**, 1619-1632.
- Spitzer-Rimon, B., Farhi, M., Albo, B., Cna'ani, A., Ben Zvi, M.M., Masci, T., Edelbaum, O., Yu, Y., Shklarman, E., Ovadis, M. and Vainstein, A.** (2012) The R2R3-MYB-like regulatory factor EOBI, acting downstream of EOBI, regulates scent production by activating ODO1 and structural scent-related genes in petunia. *Plant Cell*, **24**, 5089-5105.

- Spitzer-Rimon, B., Marhevka, E., Barkai, O., Marton, I., Edelbaum, O., Masci, T., Prathapani, N.K., Shklarman, E., Ovadis, M. and Vainstein, A.** (2010) EOBII, a gene encoding a flower-specific regulator of phenylpropanoid volatiles' biosynthesis in petunia. *Plant Cell*, **22**, 1961-1976.
- Spitzer, B., Zvi, M.M.B., Ovadis, M., Marhevka, E., Barkai, O., Edelbaum, O., Marton, I., Masci, T., Alon, M., Morin, S., Rogachev, I., Aharoni, A. and Vainstein, A.** (2007) Reverse Genetics of Floral Scent: Application of Tobacco Rattle Virus-Based Gene Silencing in Petunia. *Plant Physiology*, **145**, 1241-1250.
- Tsuda, S., Fukui, Y., Nakamura, N., Katsumoto, Y., Yonekura-Sakakibara, K., Fukuchi-Mizutani, M., Ohira, K., Ueyama, Y., Ohkawa, H., Holton, T., Kusumi, T. and Tanaka, Y.** (2004) *Flower color modification of Petunia hybrida commercial varieties by metabolic engineering.*
- Van Moerkercke, A., Haring, M.A. and Schuurink, R.C.** (2011) The transcription factor EMISSION OF BENZENOIDS II activates the MYB ODORANT1 promoter at a MYB binding site specific for fragrant petunias. *Plant J*, **67**, 917-928.
- Van Moerkercke, A., Schauvinhold, I., Pichersky, E., Haring, M.A. and Schuurink, R.C.** (2009) A plant thiolase involved in benzoic acid biosynthesis and volatile benzenoid production. *Plant J*, **60**, 292-302.
- Verdonk, J.C., Haring, M.A., van Tunen, A.J. and Schuurink, R.C.** (2005) ODORANT1 regulates fragrance biosynthesis in petunia flowers. *Plant Cell*, **17**, 1612-1624.
- Verdonk, J.C., Ric de Vos, C.H., Verhoeven, H.A., Haring, M.A., van Tunen, A.J. and Schuurink, R.C.** (2003) Regulation of floral scent production in petunia revealed by targeted metabolomics. *Phytochemistry*, **62**, 997-1008.
- Whitehead, M.R. and Peakall, R.** (2014) POLLINATOR SPECIFICITY DRIVES STRONG PREPOLLINATION REPRODUCTIVE ISOLATION IN SYMPATRIC SEXUALLY DECEPTIVE ORCHIDS. *Evolution*, **68**, 1561-1575.
- Wright, G.A. and Schiestl, F.P.** (2009) The Evolution of Floral Scent: The Influence of Olfactory Learning by Insect Pollinators on the Honest Signalling of Floral Rewards. *Functional Ecology*, **23**, 841-851.
- Yon, F., Joo, Y., Cortes Llorca, L., Rothe, E., Baldwin, I.T. and Kim, S.G.** (2016) Silencing Nicotiana attenuata LHY and ZTL alters circadian rhythms in flowers. *New Phytol*, **209**, 1058-1066.
- Zuker, A., Tzfira, T., Ben-Meir, H., Ovadis, M., Shklarman, E., Itzhaki, H., Forkmann, G., Martens, S., Neta-Sharir, I., Weiss, D. and Vainstein, A.** (2002) Modification of flower color and fragrance by antisense suppression of the flavanone 3-hydroxylase gene. *Molecular Breeding*, **9**, 33-41.