



UvA-DARE (Digital Academic Repository)

Test of QED in $e^+ e^- \rightarrow \gamma \gamma$ at LEP

Adeva, B.; Adriani, O.; Aguilar-Benitez, M.; Akbari, H.; Alcaraz, J.; Aloisio, A.; Alverson, G.; Alviggi, M.G.; Linde, F.L.

Published in:
Physics Letters B

DOI:
[10.1016/0370-2693\(90\)91177-D](https://doi.org/10.1016/0370-2693(90)91177-D)

[Link to publication](#)

Citation for published version (APA):

Adeva, B., Adriani, O., Aguilar-Benitez, M., Akbari, H., Alcaraz, J., Aloisio, A., ... Linde, F. L. (1990). Test of QED in $e^+ e^- \rightarrow \gamma \gamma$ at LEP. *Physics Letters B*, 250, 199-204. DOI: 10.1016/0370-2693(90)91177-D

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Test of QED in $e^+e^- \rightarrow \gamma\gamma$ at LEP

L3 Collaboration

B. Adeva^a, O. Adriani^b, M. Aguilar-Benitez^c, H. Akbari^d, J. Alcaraz^c, A. Aloisio^e, G. Alverson^f, M.G. Alviggi^e, Q. An^g, H. Anderhub^h, A.L. Andersonⁱ, V.P. Andreev^j, T. Angelovⁱ, L. Antonov^k, D. Antreasyan^l, P. Arce^c, A. Arefiev^m, T. Azemoonⁿ, T. Aziz^o, P.V.K.S. Baba^g, P. Bagnaia^p, J.A. Bakken^q, L. Baksay^r, R.C. Ballⁿ, S. Banerjee^{o,g}, J. Bao^d, L. Barone^p, A. Bay^s, U. Beckerⁱ, J. Behrens^h, S. Beingessner^t, Gy.L. Bencze^{u,r}, J. Berdugo^c, P. Bergesⁱ, B. Bertucci^p, B.L. Betev^k, A. Biland^h, R. Bizzarri^p, J.J. Blaising^t, P. Blömeke^v, B. Blumenfeld^d, G.J. Bobbink^w, M. Bocciaolini^b, W. Böhlen^x, A. Böhm^v, T. Böhlinger^y, B. Borgia^p, D. Bourilkov^k, M. Bourquin^s, D. Boutigny^t, J.G. Branson^z, I.C. Brock^{aa}, F. Bruyant^a, C. Buisson^{ab}, A. Bujak^{ac}, J.D. Burgerⁱ, J.P. Burq^{ab}, J. Busenitz^{ad}, X.D. Cai^g, C. Camps^v, M. Capellⁿ, F. Carbonara^e, F. Carminati^b, A.M. Cartacci^b, M. Cerrada^c, F. Cesaroni^p, Y.H. Changⁱ, U.K. Chaturvedi^g, M. Chemarin^{ab}, A. Chen^{ae}, C. Chen^{af}, G.M. Chen^{af}, H.F. Chen^{ag}, H.S. Chen^{af}, M. Chenⁱ, M.L. Chenⁿ, G. Chiefari^e, C.Y. Chien^d, C. Civinini^b, I. Clareⁱ, R. Clareⁱ, G. Coignet^t, N. Colino^a, V. Commichau^v, G. Conforto^b, A. Contin^a, F. Crijns^w, X.Y. Cui^g, T.S. Daiⁱ, R. D'Alessandro^b, R. de Asmundis^e, A. Degré^{a,t}, K. Deiters^{a,ah}, E. Dénes^u, P. Denes^q, F. De Notaristefani^p, M. Dhina^h, D. DiBitonto^{ad}, M. Diemoz^p, F. Diez-Hedo^a, H.R. Dimitrov^k, C. Dionisi^p, F. Dittus^{ai}, R. Dolinⁱ, E. Drago^c, T. Driever^w, D. Duchesneau^s, P. Duinker^{w,a}, I. Duran^{a,c}, H. El Mamouni^{ab}, A. Engler^{aa}, F.J. Epplingⁱ, F.C. Erné^w, P. Extermann^s, R. Fabbretti^h, G. Faberⁱ, S. Falciano^p, Q. Fan^{g,af}, S.J. Fan^{aj}, M. Fabre^h, J. Fay^{ab}, J. Fehlmann^h, H. Fenker^f, T. Ferguson^{aa}, G. Fernandez^c, F. Ferroni^{p,a}, H. Fesefeldt^v, J. Field^s, G. Finocchiaro^p, P.H. Fisher^d, G. Forconi^s, T. Foreman^w, K. Freudenreich^h, W. Friebel^{ah}, M. Fukushimaⁱ, M. Gailloud^y, Yu. Galaktionov^m, E. Gallo^b, S.N. Ganguli^o, P. Garcia-Abia^c, S.S. Gau^{ae}, S. Gentile^p, M. Glaubman^f, S. Goldfarbⁿ, Z.F. Gong^{g,ag}, E. Gonzalez^c, A. Gordeev^m, P. Göttlicher^v, D. Goujon^s, G. Gratta^{ai}, C. Grinnellⁱ, M. Gruenewald^{ai}, M. Guanziroli^g, A. Gurtu^o, H.R. Gustafsonⁿ, L.J. Gutay^{ac}, H. Haan^v, S. Hancke^v, K. Hangarter^v, M. Harris^a, A. Hasan^g, C.F. He^{aj}, T. Hebbeker^v, M. Hebert^z, G. Hertenⁱ, U. Herten^v, A. Hervé^a, K. Hilgers^v, H. Hofer^h, H. Hoorani^g, L.S. Hsu^{ae}, G. Hu^g, G.Q. Hu^{aj}, B. Ille^{ab}, M.M. Ilyas^g, V. Innocente^{c,a}, E. Isiksal^h, E. Jagel^g, B.N. Jin^{af}, L.W. Jonesⁿ, R.A. Khan^g, Yu. Kamyshkov^m, Y. Karyotakis^{t,a}, M. Kaur^g, S. Khokhar^g, V. Khoze^j, D. Kirkby^{ai}, W. Kittel^w, A. Klimentov^m, A.C. König^w, O. Kornadt^v, V. Koutsenko^m, R.W. Kraemer^{aa}, T. Kramerⁱ, V.R. Krastev^k, W. Krenz^v, J. Krizmanic^d, A. Kuhn^x, K.S. Kumar^{ak}, V. Kumar^g, A. Kunin^m, A. van Laak^v, V. Lalieu^s, G. Landi^b, K. Lanius^{a,ah}, W. Lange^{ah}, D. Lanske^v, S. Lanzano^e, P. Lebrun^{ab}, P. Lecomte^h, P. Lecoq^a, P. Le Coultre^h, I. Leedom^f, J.M. Le Goff^a, A. Leike^{ah}, L. Leistam^a, R. Leiste^{ah}, M. Lenti^b, J. Lettry^h, P.M. Levchenko^j, X. Leytens^w, C. Li^{ag}, H.T. Li^{af}, J.F. Li^g, L. Li^h, P.J. Li^{aj}, Q. Li^g, X.G. Li^{af}, J.Y. Liao^{aj}, Z.Y. Lin^{ag}, F.L. Linde^{aa}, D. Linnhofer^a, R. Liu^g, Y. Liu^g, W. Lohmann^{ah}, S. Lökös^r, E. Longo^p, Y.S. Lu^{af}, J.M. Lubbers^w, K. Lübelmeyer^v, C. Luci^a, D. Luckey^{l,i}, L. Ludovici^p, X. Lue^h, L. Luminari^p, W.G. Ma^{ag}, M. MacDermott^h, R. Magahiz^r, M. Maire^t, P.K. Malhotra^o, R. Malik^g, A. Malinin^m, C. Mañá^c, D.N. Maoⁿ, Y.F. Mao^{af}, M. Maolinbay^h, P. Marchesini^g, A. Marchionni^b, J.P. Martin^{ab}, L. Martinez^a, F. Marzano^p,

G.G.G. Massaro ^w, T. Matsuda ⁱ, K. Mazumdar ^o, P. McBride ^{ak}, T. McMahon ^{ac}, D. McNally ^h, Th. Meinholz ^v, M. Merk ^w, L. Merola ^c, M. Meschini ^b, W.J. Metzger ^w, Y. Mi ^g, M. Micke ^v, U. Micke ^v, G.B. Mills ⁿ, Y. Mir ^g, G. Mirabelli ^p, J. Mnich ^v, M. Möller ^v, B. Monteleoni ^b, G. Morand ^s, R. Morand ^t, S. Morganti ^p, V. Morgunov ^m, R. Mount ^{ai}, E. Nagy ^u, M. Napolitano ^e, H. Newman ^{ai}, M.A. Niaz ^g, L. Niessen ^v, W.D. Nowak ^{ah}, H. Nowak ^{ah}, S. Nowak ^{ah}, D. Pandoulas ^v, G. Passaleva ^b, G. Paternoster ^c, S. Patricelli ^e, Y.J. Pei ^v, D. Perret-Gallix ^t, J. Perrier ^s, A. Pevsner ^d, M. Pieri ^b, P.A. Piroué ^q, V. Plyaskin ^m, M. Pohl ^h, V. Pojidaev ^m, N. Produit ^s, J.M. Qian ^{i,g}, K.N. Qureshi ^g, R. Raghavan ^o, G. Rahal-Callot ^h, P. Razis ^h, K. Read ^q, D. Ren ^h, Z. Ren ^g, S. Reucroft ^f, T. Riemann ^{ah}, O. Rind ⁿ, C. Rippich ^{aa}, H.A. Rizvi ^g, B.P. Roe ⁿ, M. Röhner ^v, S. Röhner ^v, Th. Rombach ^v, L. Romero ^c, J. Rose ^v, S. Rosier-Lees ^t, R. Rosmalen ^w, Ph. Rosselet ^y, J.A. Rubio ^{a,c}, W. Ruckstuhl ^s, H. Rykaczewski ^h, M. Sachwitz ^{ah}, J. Salicio ^{a,c}, J.M. Salicio ^c, G. Sartorelli ^g, G. Sauvage ^t, A. Savin ^m, V. Schegelsky ^j, D. Schmitz ^v, P. Schmitz ^v, M. Schneegans ^t, M. Schöntag ^v, H. Schopper ^{ak}, D.J. Schotanus ^w, H.J. Schreiber ^{ah}, R. Schulte ^v, S. Schulte ^v, K. Schultze ^v, J. Schütte ^{ak}, J. Schwenke ^v, G. Schwering ^v, C. Sciacca ^e, R. Sehgal ^g, P.G. Seiler ^h, J.C. Sens ^w, I. Sheer ^z, V. Shevchenko ^m, S. Shevchenko ^m, X.R. Shi ^{aa}, K. Shmakov ^m, V. Shoutko ^m, E. Shumilov ^m, N. Smirnov ^j, A. Sopczak ^{ai,z}, C. Souyri ^t, C. Spartiotis ^d, T. Spickermann ^v, B. Spiess ^x, P. Spillantini ^b, R. Starosta ^v, M. Steuer ^{l,i}, D.P. Stickland ^q, B. Stöhr ^h, H. Stone ^s, K. Strauch ^{ak}, B.C. Stringfellow ^{ac}, K. Sudhakar ^{o,v}, G. Sultanov ^a, R.L. Sumner ^q, L.Z. Sun ^{ag}, H. Suter ^h, R.B. Sutton ^{aa}, J.D. Swain ^g, A.A. Syed ^g, X.W. Tang ^{af}, E. Tarkovsky ^m, L. Taylor ^f, J.M. Thenard ^t, E. Thomas ^g, C. Timmermans ^w, Samuel C.C. Ting ⁱ, S.M. Ting ⁱ, Y.P. Tong ^{ac}, F. Tonisch ^{ah}, M. Tonutti ^v, S.C. Tonwar ^o, J. Tòth ^u, G. Trowitzsch ^{ah}, K.L. Tung ^{af}, J. Ulbricht ^x, L. Urbàn ^u, U. Uwer ^v, E. Valente ^p, R.T. Van de Walle ^w, H. van der Graaf ^w, I. Vetlitsky ^m, G. Viertel ^h, P. Vikas ^g, U. Vikas ^g, M. Vivargent ^{l,i}, H. Vogel ^{aa}, H. Vogt ^{ah}, M. Vollmar ^v, G. Von Dardel ^a, I. Vorobiev ^m, A.A. Vorobyov ^j, An.A. Vorobyov ^j, L. Vuilleumier ^y, M. Wadhwa ^g, W. Wallraff ^v, C.R. Wang ^{ag}, G.H. Wang ^{aa}, J.H. Wang ^{af}, Q.F. Wang ^{ak}, X.L. Wang ^{ag}, Y.F. Wang ^b, Z. Wang ^g, Z.M. Wang ^{g,ag}, J. Weber ^h, R. Weill ^y, T.J. Wenaus ⁱ, J. Wenninger ^s, M. White ⁱ, R. Wilhelm ^w, C. Willmott ^c, F. Wittgenstein ^a, D. Wright ^q, R.J. Wu ^{af}, S.L. Wu ^g, S.X. Wu ^g, Y.G. Wu ^{af}, B. Wystouch ⁱ, Y.D. Xu ^{af}, Z.Z. Xu ^{ag}, Z.L. Xue ^{aj}, D.S. Yan ^{aj}, B.Z. Yang ^{ag}, C.G. Yang ^{af}, G. Yang ^g, K.S. Yang ^{af}, Q.Y. Yang ^{af}, Z.Q. Yang ^{aj}, C.H. Ye ^g, J.B. Ye ^h, Q. Ye ^g, S.C. Yeh ^{ae}, Z.W. Yin ^{aj}, J.M. You ^g, C. Zaccardelli ^{ai}, L. Zehnder ^h, M. Zeng ^g, Y. Zeng ^v, D. Zhang ^z, D.H. Zhang ^w, Z.P. Zhang ^{ag}, J.F. Zhou ^v, R.Y. Zhu ^{ai}, H.L. Zhuang ^{af} and A. Zichichi ^{a,g}

^a European Laboratory for Particle Physics, CERN, CH-1211 Geneva 23, Switzerland

^b INFN – Sezione di Firenze and University of Firenze, I-50125 Florence, Italy

^c Centro de Investigaciones Energeticas, Medioambientales y Tecnologicas, CIEMAT, E-28040 Madrid, Spain

^d Johns Hopkins University, Baltimore, MD 21218, USA

^e INFN – Sezione di Napoli and University of Naples, I-80125 Naples, Italy

^f Northeastern University, Boston, MA 02115, USA

^g World Laboratory, FBLJA Project, CH-1211 Geneva, Switzerland

^h Eidgenössische Technische Hochschule, ETH Zürich, CH-8093 Zurich, Switzerland

ⁱ Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^j Leningrad Nuclear Physics Institute, SU-188 350 Gatchina, USSR

^k Central Laboratory of Automation and Instrumentation, CLANP, Sofia, Bulgaria

^l INFN – Sezione di Bologna, I-40126 Bologna, Italy

^m Institute of Theoretical and Experimental Physics, ITEP, SU-117 259 Moscow, USSR

ⁿ University of Michigan, Ann Arbor, MI 48109, USA

^o Tata Institute of Fundamental Research, Bombay 400 005, India

^p INFN – Sezione di Roma and University of Rome “La Sapienza”, I-00185 Rome, Italy

^q Princeton University, Princeton, NJ 08544, USA

- ^r Union College, Schenectady, NY 12308, USA
^s University of Geneva, CH-1211 Geneva 4, Switzerland
^t Laboratoire de Physique des Particules, LAPP, F-74519 Annecy-le-Vieux, France
^u Central Research Institute for Physics of the Hungarian Academy of Sciences, H-1525 Budapest 114, Hungary
^v I. Physikalisches Institut, RWTH, D-5100 Aachen, FRG¹
and III. Physikalisches Institut, RWTH, D-5100 Aachen, FRG¹
^w National Institute for High Energy Physics, NIKHEF, NL-1009 DB Amsterdam, The Netherlands
and NIKHEF-H and University of Nijmegen, NL-6525 ED Nijmegen, The Netherlands
^x Paul Scherrer Institut (PSI), Würenlingen, Switzerland
^y University of Lausanne, CH-1015 Lausanne, Switzerland
^z University of California, San Diego, CA 92182, USA
^{aa} Carnegie Mellon University, Pittsburgh, PA 15213, USA
^{ab} Institut de Physique Nucléaire de Lyon, IN2P3-CNRS/Université Claude Bernard, F-69622 Villeurbanne Cedex, France
^{ac} Purdue University, West Lafayette, IN 47907, USA
^{ad} University of Alabama, Tuscaloosa, AL 35486, USA
^{ae} High Energy Physics Group, Taiwan, ROC
^{af} Institute of High Energy Physics, IHEP, Beijing, P.R. China
^{ag} Chinese University of Science and Technology, USTC, Hefei, Anhui 230 029, P.R. China
^{ah} High Energy Physics Institute, DDR-1615 Zeuthen-Berlin, GDR
^{ai} California Institute of Technology, Pasadena, CA 91125, USA
^{aj} Shanghai Institute of Ceramics, SIC, Shanghai, P.R. China
^{ak} Harvard University, Cambridge, MA 02139, USA
^{al} University of Hamburg, D-2000 Hamburg, FRG

Received 3 August 1990

We have measured the cross-section of the reaction $e^+e^- \rightarrow \gamma\gamma$ at center of mass energies around the Z^0 mass. The results are in good agreement with QED predictions. For the QED cutoff parameters the limit of $A_+ > 103$ GeV and $A_- > 118$ GeV are found. For the decays $Z^0 \rightarrow \gamma\gamma$, $Z^0 \rightarrow \pi^0\gamma$, $Z^0 \rightarrow \eta\gamma$ and $Z^0 \rightarrow \gamma\gamma\gamma$ we find upper limits of 2.9×10^{-4} , 2.9×10^{-4} , 4.1×10^{-4} and 1.2×10^{-4} , respectively. All limits are at 95% CL.

1. Introduction

At LEP energies, the reaction $e^+e^- \rightarrow \gamma\gamma$ provides a clean test of QED. In contrast to lepton pair production it is in lowest order not affected by weak interaction effects and hadronic vacuum polarisation. In the absence of rare or theoretically forbidden decays, such as $Z^0 \rightarrow \gamma\gamma$, this reaction can only proceed via the exchange of a virtual electron. On the other hand, deviations from QED in the Z^0 region, could yield information on non-standard-model properties of this boson.

The first order differential cross section for $e^+e^- \rightarrow \gamma\gamma$ is predicted by the Born term of QED. Deviations from QED are generally parametrized by introducing cutoff parameters A_{\pm} , i.e. by generalizing

the lowest order QED differential cross section to [1–3]

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{s} \frac{1 + \cos^2\theta}{1 - \cos^2\theta} \left(1 \pm \frac{s^2}{2A_{\pm}^4} (1 - \cos^2\theta) \right). \quad (1)$$

At LEP energies higher order terms are important, however, and radiative corrections [4] are necessary before comparing the above expression with data. We report here on a comparison between our data on $e^+e^- \rightarrow \gamma\gamma$ obtained in the Z^0 region and the QED prediction for this reaction calculated up to order α^3 .

The same reaction $e^+e^- \rightarrow \gamma\gamma$ can also be used to set limits for the mass of an excited virtual electron (e^*) [3] and various forbidden or rare Z^0 decay modes.

2. The L3 detector

The L3 detector covers 99% of 4π . The detector in-

¹ Supported by the German Bundesministerium für Forschung und Technologie.

cludes a central vertex chamber, a precise electromagnetic calorimeter composed of BGO crystals, a uranium and brass hadron calorimeter with proportional wire chamber readout, a high accuracy muon chamber system, and a ring of scintillation counters. These detectors are installed in a 12 m diameter magnet which provides a uniform field of 0.5 T along the beam direction. The luminosity is determined by measuring small angle Bhabha events in two forward calorimeters consisting of BGO crystals. A detailed description of each detector subsystem, and its performance, is given in ref. [5].

The fine segmentation of the electromagnetic calorimeter allows the measurement of photon and electron showers with an angular resolution of 2 mrad. For the present analysis we only use data collected in the polar angle θ region:

$$\frac{1}{4}\pi < \theta < \frac{3}{4}\pi.$$

3. Event selection

Events collected from the 1990 LEP running period (March–June) corresponding to 2.24 pb^{-1} are used for the present analysis.

The selection of the $e^+e^- \rightarrow \gamma\gamma$ candidates is primarily based on the number of shower peaks in the BGO calorimeter and on their measured energies. We require

(i) at least two but no more than twelve shower peaks;

(ii) two of them should have an energy between 35 GeV and 55 GeV.

The first cut eliminates a major fraction of the hadronic events. The second one eliminates cosmic ray interactions, hadronic and $\tau\tau$ events.

(iii) Next, we eliminate all events in which both major shower peaks have a matching track in the vertex chamber within 50 mrad in transverse projection.

The result is a sample of 35 candidates with no matching vertex chamber tracks and a sample of 4 candidates with one matching vertex chamber track. Further examination of the latter sample indicates that only one of them is a recognizable γ conversion in the beam pipe with both decay tracks clearly separated. It should be noted that from the material in front of the TEC we expect 0.7 γ conversions in our




Fig. 1. The $\gamma\gamma$ energy sum versus $|\theta_1 + \theta_2 - \pi|$ with the solid line showing the energy cut $E_1 + E_2 > 0.9\sqrt{s}$.

sample. The remaining 3 events are classified as $e^+e^- \rightarrow e^+e^-$ events with one track detected and are eliminated from the sample. The contamination in the remaining sample where $e^+e^- \rightarrow e^+e^-$ with both tracks undetected is less than 0.005 events.

Fig. 1 shows a plot of $(E_1 + E_2)/\sqrt{s}$ versus $|\theta_1 + \theta_2 - \pi|$ for the centers of the shower peaks of the remaining 36 events. The figure shows the presence of 4 events with $E_1 + E_2 < 0.9\sqrt{s}$, which combine a relatively large acollinearity with a low total energy. Initial state radiation produces events of this type. It affects the angular distribution in eq. (1), therefore a cut is imposed requiring $E_1 + E_2 > 0.9\sqrt{s}$. Applying the same cut to a Monte Carlo sample [of about 1000 $e^+e^- \rightarrow \gamma\gamma(\gamma)$ events] indicates that it has an acceptance of 89% in the polar angle region under consideration; the number of events removed by the cut agrees with this acceptance.

After the above cut we are left with a sample of 32 events.

4. Analysis $e^+e^- \rightarrow \gamma\gamma$

Table 1 presents the cross sections observed for $e^+e^- \rightarrow \gamma\gamma$ for three different center of mass energy regions and for the polar region under consideration.

Table 1

Region	Energy (GeV)	Luminosity (pb^{-1})	Events	Visible cross section
below Z	89.50	0.567	8	16.2 ± 5.4
on Z	91.28	1.202	18	17.2 ± 3.8
above Z	93.17	0.471	6	14.7 ± 5.6
all	91.22	2.240	32	16.4 ± 2.7

Within this angular range the efficiency of the electromagnetic calorimeter is $(98 \pm 1)\%$. Also given are the corresponding integrated luminosities and the number of events.

Integration of the lowest order QED differential cross section between $\frac{1}{4}\pi$ and $\frac{3}{4}\pi$ yields a prediction of 16.4 pb at 91.2 GeV. Applying the required radiative corrections, up to order α^3 , using a program written by Berends and Kleiss [4] changes the QED-prediction to 16.8 pb, in agreement with the results presented in table 1.

Fig. 2 shows a comparison between the radiatively corrected QED angular dependence and our data points. The $\cos \theta$ dependence of the data is consistent with QED. To test the agreement with QED we used the binning-free Smirnov-Cramér-von Mises test




Fig. 2. The measured number of events versus $\frac{1}{2}(|\cos \theta_1| + |\cos \theta_2|)$. The solid curves show the QED prediction and the expectations with the cutoff parameters $\Lambda_+ = 103$ GeV and $\Lambda_- = 118$ GeV (95% confidence level limit).

Table 2

Experiment	Λ_+ (GeV)	Λ_- (GeV)
L3 (this experiment)	103	118
OPAL	82	89
CELLO	59	44
JADE	61	57
MARK J	72	65
PLUTO	46	-
TASSO	61	56
HRS	59	59
MAC	66	67
AMY	65	-
TOPAZ	94	59

[6]. Parametrizing a possible QED deviation as indicated in the introduction, and varying the cutoff Λ parameters, one obtains the following limits at the 95% confidence level:

$$\Lambda_+ > 103 \text{ GeV}, \quad \Lambda_- > 118 \text{ GeV}.$$

Table 2 shows a comparison between these limits and previously obtained e^+e^- results [7].

It is possible to parametrize a modification of the electron propagator in the reaction $e^+e^- \rightarrow \gamma\gamma$ in terms of the exchange of a virtual excited electron e^* [3]. The two parameters that enter into consideration then are λ , the ratio of the e^* to e coupling, and M_{e^*} , the mass of the e^* . Assuming $\lambda = 1$ we find at 95% CL

$$M_{e^*} > 83 \text{ GeV}.$$

This result is in agreement with limits recently obtained [8-10].

5. Rare Z^0 decays

Any deviation from the QED prediction for the

$e^+e^- \rightarrow \gamma\gamma$ cross section could also be due to rare Z^0 decays such as the theoretically forbidden $Z^0 \rightarrow \gamma\gamma$ [11] or the decays $Z^0 \rightarrow \pi^0\gamma$, $Z^0 \rightarrow \eta\gamma$, in which the π^0 or η would be indistinguishable from a γ . A difference between such Z^0 decays and QED events is the angular distribution. The QED reaction is strongly peaked forward whereas the Z^0 decays show essentially a $1 + \cos^2\theta$ dependence. Upper limits for the Z^0 decays can thus be obtained by determining the amount of extra contribution of the type $1 + \cos^2\theta$ our $\gamma\gamma$ sample is able to accommodate in the θ region under consideration. Taking into account the geometrical acceptances and the η -decay modes one arrives at detection efficiencies of 62%, 62% and 43% for $Z^0 \rightarrow \gamma\gamma$, $Z^0 \rightarrow \pi^0\gamma$ and $Z^0 \rightarrow \eta\gamma$ respectively. Using these numbers one derives [6] at 95% confidence level the following upper limits:

$$\text{BR}(Z^0 \rightarrow \gamma\gamma) < 2.9 \times 10^{-4},$$

$$\text{BR}(Z^0 \rightarrow \pi^0\gamma) < 2.9 \times 10^{-4},$$

$$\text{BR}(Z^0 \rightarrow \eta\gamma) < 4.1 \times 10^{-4}.$$

As an extension of the $e^+e^- \rightarrow \gamma\gamma$ study, a search was made for the higher order QED process $e^+e^- \rightarrow \gamma\gamma\gamma$. The selection criteria used were the following:

- (i) at least three shower peaks, together having an energy greater than $0.5\sqrt{s}$;
- (ii) an acoplanarity between these clusters of less than 9 degrees;
- (iii) no matching tracks in the vertex chamber.

No candidate events are found. The acceptance of the above cuts, for the process $Z^0 \rightarrow \gamma\gamma\gamma$, roughly corresponds to the 3γ acceptance of the BGO barrel i.e. 35%. Using Poisson statistics we derive

$$\text{BR}(Z^0 \rightarrow \gamma\gamma\gamma) < 1.2 \times 10^{-4}$$

at the 95% confidence level.

The standard model predicts a $\text{BR}(Z^0 \rightarrow \gamma\gamma\gamma)$ of 8×10^{-10} [12]. However, in some composite models branching ratios as high as 2×10^{-4} have been predicted [13].

Acknowledgement

We wish to thank CERN for its hospitality and help. We want particularly to express our gratitude to the LEP division: it is their excellent achievements which made this experiment possible. We acknowledge the support of all the funding agencies which contributed to this experiment.

References

- [1] S. Drell, Ann. Phys. (NY) 4 (1958) 75.
- [2] F.E. Low, Phys. Rev. Lett. 14 (1965) 238.
- [3] A. Litke, Harvard University, Ph.D. Thesis (1970), unpublished.
- [4] F.A. Berends and R. Kleiss, Nucl. Phys. B 186 (1981) 22.
- [5] L3 Collab., B. Adeva et al., Nucl. Instrum. Methods A 289 (1990) 35.
- [6] W.T. Eadie et al., Statistical methods in experimental physics (North-Holland, Amsterdam, 1971, p. 268).
- [7] OPAL Collab., M.Z. Akrawy et al., Phys. Lett. B 241 (1990) 133;
CELLO Collab., H.J. Behrend et al., Phys. Lett. B 123 (1983) 127;
JADE Collab., W. Bartel et al., Z. Phys. C 19 (1983) 197;
MARK J. Collab., B. Adeva et al., Phys. Lett. B 152 (1985) 439;
H.S. Chen, Riv. Nuovo Cimento 1 (1988) 11;
PLUTO Collab., Ch. Berger et al., Phys. Lett. B 94 (1980) 87;
TASSO Collab., M. Althoff et al., Z. Phys. C 26 (1984) 337;
HRS Collab., M. Derrick et al., Phys. Lett. B 166 (1986) 468; Phys. Rev. D 34 (1986) 3286;
MAC Collab., E. Fernandez et al., Phys. Rev. D 35 (1987) 1;
AMY Collab., S.K. Kim et al., Phys. Lett. B 223 (1989) 476;
TOPAZ Collab., I. Adachi et al., Phys. Lett. B 200 (1988) 391;
- [8] ALEPH Collab., D. Decamp et al., Phys. Lett. B 236 (1990) 501.
- [9] OPAL Collab., M.Z. Akrawy et al., Phys. Lett. B 244 (1990) 135.
- [10] L3 Collab., B. Adeva et al., Phys. Lett. B 247 (1990) 177.
- [11] C.N. Yang, Phys. Rev. 77 (1950) 242.
- [12] M. Laursen et al., Phys. Rev. D 23 (1981) 2795.
- [13] Compositeness Working Group, F.M. Renard et al., CERN report 89-08, Vol. 2, p. 185.