



UvA-DARE (Digital Academic Repository)

Differential effects of elevated extracellular calcium on field potentials in dentate gyrus and CA1 of the rat hippocampal slice preparation

Melchers, B.P.C.; Pennartz, C.M.A.; Lopes da Silva, F.H.

Published in:
Neuroscience Letters

DOI:
[10.1016/0304-3940\(87\)90603-3](https://doi.org/10.1016/0304-3940(87)90603-3)

[Link to publication](#)

Citation for published version (APA):

Melchers, B. P. C., Pennartz, C. M. A., & Lopes da Silva, F. H. (1987). Differential effects of elevated extracellular calcium on field potentials in dentate gyrus and CA1 of the rat hippocampal slice preparation. *Neuroscience Letters*, 77, 37-42. DOI: 10.1016/0304-3940(87)90603-3

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

NSL 04602

Differential effects of elevated extracellular calcium concentrations on field potentials in dentate gyrus and CA₁ of the rat hippocampal slice preparation

B.P.C. Melchers, C.M.A. Pennartz and F.H. Lopes da Silva

University of Amsterdam, Department of General Zoology, Amsterdam (The Netherlands)

(Received 2 November 1986; Revised version received 24 December 1986; Accepted 11 February 1987)

Key words: Hippocampal slice; Dentate gyrus; CA₁; Calcium; Long-term potentiation; Field potential; Rat

The effect of a short-lasting elevation of the extracellular Ca²⁺ concentration (from 2 to 6 mM) on the field potentials in CA₁ of the rat hippocampal slice was an increase of both the CA₁ population spike (PS) and the excitatory postsynaptic potential (EPSP); this effect persisted after returning to 2 mM Ca²⁺ Ringer, and thus can be considered as a Ca²⁺-induced long-term potentiation (LTP). In the dentate gyrus (DG) a quite different effect was encountered; here the PS decreased and the EPSP increased only slightly during the perfusion with 6 mM Ca²⁺, and no reproducible long-term effect was induced. The results indicate that substantial differences exist in the balance between inhibitory and excitatory processes in the neural networks of the two hippocampal subregions; these differences are enhanced during perfusion with high Ca²⁺, which induces LTP in CA₁ but not in the DG.

The crucial role of Ca²⁺ in neurotransmitter release and in paired-pulse facilitation, especially at the neuromuscular junction [6, 8] but also in the squid giant synapse [10], is well recognized. The importance of Ca²⁺ has been shown also in long-term potentiation (LTP), a relatively long-lasting form of synaptic plasticity (for a review see ref. 4). LTP is typically elicited by tetanic stimulation of afferents and is characterized by a facilitation of synaptic transmission that may last for several hours in e.g. hippocampal slices. The importance of Ca²⁺ in the induction of LTP has been shown in a number of studies: (i) LTP cannot be elicited at low extracellular Ca²⁺-concentrations [3]; (ii) the injection of a Ca²⁺-chelating agent prevents LTP induction in the postsynaptic neuron [7]; (iii) an increased uptake and retention of Ca²⁺ in hippocampal slices has been reported following LTP induction [2]; and (iv) Agoston and Kuhnt [1] have recently demonstrated an increase in Ca²⁺ accumulation in synaptosomes prepared from slices in which LTP was elicited.

Correspondence: B.P.C. Melchers, University of Amsterdam, Department of General Zoology, Kruislaan 320, 1098 SM Amsterdam, The Netherlands.

In fact, LTP can be induced in the pyramidal cells of area CA₁ in the hippocampal slice simply by an elevation of the extracellular Ca²⁺-concentration for some minutes [13]. It was to be expected that the effects of an elevated Ca²⁺-concentration on neurons are general phenomena. However, pilot studies suggested that the effects of elevated Ca²⁺ on the granule cells of the dentate gyrus (DG) were quite different from those reported for the CA₁ area [13]. To investigate this in more detail we have tested simultaneously, and in the same hippocampal slice, the effects of a rise in extracellular Ca²⁺ on two subsystems, the pyramidal cells of area CA₁ and the granule cells of the DG. In this way we could show an additional physiological difference between the CA₁ area and the DG to add to previously reported ones [9, 11, 12]. Furthermore, it is demonstrated that the effects of an elevated Ca²⁺-concentration on synaptic transmission have to be interpreted taking into consideration the particular network involved.

Hippocampal slices were made following standard procedures: rats (160–200 g) were decapitated after ether anesthesia, the brain was removed, rinsed immediately with ice-cold (0–4°C) Ringer solution (in mM: NaCl 124, KCl 5, CaCl₂ 2, MgSO₄ 2, NaH₂PO₄ 1.25, NaHCO₃ 26, glucose 10) and the hippocampi were prepared free. Transverse slices of 500 μm were cut manually with razor blades and stored at room temperature until used. One hour before the start of an experiment one slice was transferred to the recording chamber which was continuously perfused with Ringer, saturated with 95% O₂–5% CO₂ (2–3 ml/min) at 34–35°C. Stimulation electrodes (two trimel-insulated stainless-steel wires of 60 μm diameter) were placed under visual guidance both in the stratum moleculare of the DG, in order to stimulate the perforant path, and in the stratum radiatum of CA₁, in order to stimulate the Schaffer collaterals. For recording evoked field potentials in DG and CA₁, glass microelectrodes, filled with 3 M NaCl (resistance 3–5 MΩ) were positioned in the stratum granulosum and stratum pyramidale. Every 15 s paired test stimuli (interval 20 ms) consisting of bipolar, biphasic current pulses of 0.2 ms duration were delivered via one of the stimulation electrodes. The current intensity was adjusted to 50% of the intensity that elicited a saturated response, and ranged between 50 and 150 μA. The field potentials were digitized at 4 kHz using a Motorola Exorset microcomputer. Series of 4 test-stimulus pairs were applied alternatively to the DG and CA₁; the corresponding field potentials were recorded, averaged and stored on a diskette. The 6 mM Ca²⁺ Ringer was made by adding extra CaCl₂ to the perfusion Ringer solution and was perfused for a period of 20 min.

The observed changes in CA₁ field potentials *during* the perfusion with 6 mM Ca²⁺ Ringer are in good agreement with the report of Turner et al. [13]. The amplitude of both the population spike (PS) and the slope of the field excitatory postsynaptic potential (EPSP) in response to the first stimulus of each pair (R₁) increased gradually in 10 out of 11 slices. On average, the maximal rising slope of the EPSP increased up to 175 ± 17% (S.E.M.) of control level ($P < 0.01$, Wilcoxon's matched-pairs signed-rank test, two-tailed) and the PS up to 198 ± 31% of control level ($P < 0.01$). The perfusion with 6 mM Ca²⁺ Ringer induced a clear LTP-like effect in CA₁: the average response amplitudes decreased only slightly after returning to




Fig. 1. Field potentials obtained in the stratum radiatum of CA₁ (A) and the stratum granulosum of the DG (B), during the control period. The arrows indicate the time points at which the paired stimuli were given to either the Schaffer collaterals or the perforant path. The PPF of the PS is clearly demonstrated. C-F: average of all slices ($n=11$) of the relative amplitude of the PS (C) and rising slope of the EPSP (E) of the R₁ in CA₁ and in the DG, respectively D and F. The black bar indicates the time during which 6 mM Ca²⁺ Ringer was perfused. The white bar indicates the time range in which significant differences (Wilcoxon's matched-pairs signed-rank test, $P<0.05$) with respect to the control value (set at 100%) were found. The number of slices per point varied, due to the fact that not all slices were monitored during the whole period, but it was always at least 6. Note the increase in the amplitude of both PS and rising slope of the EPSP in CA₁ during and for a long period after the perfusion with 6 mM Ca²⁺, whereas in the DG a decrease in PS was found and just a slight increase in EPSP.

2 mM Ca²⁺, staying well above control level for at least 45 min (Fig. 1C, E). In 8 slices (two were only monitored before and during perfusion with 6 mM Ca²⁺) the responses remained elevated during the whole recording period, which ranged from 15 to 180 min after returning to 2 mM Ca²⁺. In 4 *additional* slices CA₁ field potentials were elicited only before and 25 min after the perfusion with 6 mM Ca²⁺ Ringer, in order to check whether the series of test stimuli delivered throughout the experiment (as described above) had any effect on the Ca²⁺-induced LTP. In all 4 slices a clear potentiation of both the EPSP and the PS was encountered, indicating that repeated afferent stimulation is not necessary for eliciting Ca²⁺-induced LTP.

In the DG a clearly different effect was found. During the perfusion with 6 mM Ca^{2+} Ringer a gradual decrease was found in the amplitude of the R_1 PS in all 11 slices reaching an average value as low as $33 \pm 10\%$ of control level ($P < 0.01$; Fig. 1D). After returning to 2 mM Ca^{2+} the R_1 PS returned to a level which was not significantly different from control. The rising slope of the EPSP, in contrast, showed an increase – although less pronounced than in CA_1 – in 7 out of 11 slices (on average up to $127 \pm 10\%$ of control value ($P < 0.05$; Fig. 1F)). After returning to 2 mM Ca^{2+} Ringer, the rising slope of the EPSP in R_1 remained elevated in 5 slices; in the remaining two slices the response returned to the control level or even lower. From 30 min after returning to 2 mM Ca^{2+} the overall increase was no longer significant. In DG the long-term effects of the elevation of the extracellular Ca^{2+} concentration were variable among different slices. In 4 slices both the PS and the EPSP showed a potentiation after returning to 2 mM Ca^{2+} , but in the remaining slices either no long-term effect or even a long-lasting decrease in the PS was found.

It should be noted that an increase in extracellular calcium can exert effects on both excitatory and inhibitory synapses. We therefore tested whether paired-pulse facilitation (PPF) of the PS, which depends at least partly on the relative strength of inhibitory processes, was altered during and/or after the perfusion with 6 mM Ca^{2+} . In the neuromuscular junction a decrease in PPF of the EPSP during elevated extracellular Ca^{2+} levels has been reported [8]. In most slices during the control period, both in DG and CA_1 , a PPF of the PS and of the EPSP in the response to the second stimulus of a pair (R_2) was encountered at the intensity and inter-stimulus interval (20 ms) used. During perfusion with 6 mM Ca^{2+} Ringer the amount of PPF of the PS decreased in all 11 slices, both in the CA_1 and DG. This decrease was larger in DG than in CA_1 (the difference was significant at $P < 0.01$, Wilcoxon). After returning to 2 mM Ca^{2+} , the amount of PS PPF in CA_1 remained below control level in most slices; in the DG the PPF of the PS recovered to approximately control levels. The PPF of the EPSP also showed a decrease during perfusion with 6 mM Ca^{2+} Ringer, both in the CA_1 and DG, and returned to the control level after switching to 2 mM Ca^{2+} Ringer.

In discussing these findings the short- and the long-term (LTP) effects will be considered separately. The *short term* effects (i.e. during elevated Ca^{2+}) in CA_1 can be explained by taking into consideration that an increased extracellular Ca^{2+} concentration is known to lead to an increase in transmitter release both at the neuromuscular junction [8] and in the squid giant synapse [10]. Our results would be explained if we assume that the same process takes place at the excitatory synapses of the Schaffer collaterals with the CA_1 pyramidal cells. The increase in R_1 PS indicates an enhancement of synchronous firing of these cells. In this way more interneurons, responsible for recurrent inhibition, may be recruited, which would account for the relative decrease of the R_2 PS (decrease in PPF) observed in CA_1 . In the DG, however, different processes are apparently at work, since here the EPSP slope increases while the PS decreases. The former effect is of the same type as seen in CA_1 and may be explained in a similar way. The decrease in PS indicates a decrease in firing probability of the granule cells in contrast to that of the pyramidal cells. Two explanations

may be advanced for this effect of the elevated extracellular Ca^{2+} concentration: (i) it could exert a stabilizing effect on cell membranes [5] and thus elevate the firing threshold, but it is unlikely that this effect would appear only in the DG and not in CA_1 ; (ii) it could elicit an increase of hyperpolarizing potentials to a larger degree in the DG than in CA_1 . The latter possibility is more plausible since a number of experimental findings suggest that CA_1 and DG differ in excitatory/inhibitory circuits. Tielen et al. [12] have shown that Met-enkephalin can cause an increase in PS amplitude in CA_1 and at the same time, a decrease in PS in the DG. Furthermore, the DG may possess a larger amount of inhibitory control than CA_1 , as suggested by GABA-receptor binding studies [9]. In addition LTP appears to be more readily elicited (after a tetanus) in CA_1 than in the DG; this difference can be neutralized by a blockade of the Cl^- -mediated inhibition [14]. An increase in inhibitory processes in the DG by an elevated Ca^{2+} concentration would also explain the decrease in PPF in the DG during elevated Ca^{2+} , in spite of the decrease in the R_1 PS.

Therefore, the *short-term* effects can be accounted for by postulating that the DG has a larger inhibitory 'reserve' than CA_1 . We assume that elevated extracellular Ca^{2+} would lead to an enhancement of both excitatory and inhibitory synaptic activity. The net effect would depend on the dynamic equilibrium between inhibitory and excitatory processes within the subregion. The *long-term* effect seems to be in agreement with these considerations. The difference between the CA_1 and DG in this respect can be explained by a disproportionate potentiation of inhibitory processes versus excitatory processes within the DG. This leads even to an apparent decrease in the DG field potentials in some slices. In CA_1 , however, a reproducible LTP was induced, even if the slices were not stimulated during the period of elevated Ca^{2+} .

This work was subsidized partially by MEDIGON grant 900-548-72 of the Dutch organization for the advancement of pure research (Z.W.O.). The authors thank Drs. W.E.J.M. Ghysen, W. Kamphuis and W.J. Wadman for their comments.

- 1 Agoston, D.V. and Kuhnt, U., Increased Ca-uptake of presynaptic terminals during long-term potentiation in hippocampal slices, *Exp. Brain Res.*, 62 (1986) 663-668.
- 2 Baimbridge, K.G. and Miller, J.J., Calcium uptake and retention during long-term potentiation of neuronal activity in the rat hippocampal slice preparation, *Brain Res.*, 221 (1981) 299-305.
- 3 Dunwiddie, T.V. and Lynch, G., The relationship between extracellular calcium concentrations and the induction of hippocampal long-term potentiation, *Brain Res.*, 169 (1979) 103-110.
- 4 Eccles, J.C., Calcium in long-term potentiation as a model for memory, *Neuroscience*, 10 (1983) 1071-1081.
- 5 Frankenhaeuser, B. and Hodgkin, A.L., The action of calcium on the electrical properties of squid axons, *J. Physiol. (London)*, 137 (1957) 217-245.
- 6 Katz, B. and Miledi, T., The role of calcium in neuromuscular facilitation, *J. Physiol. (London)*, 195 (1968) 481-492.
- 7 Lynch, G., Larson, J., Kelso, S., Barrionuevo, G. and Schottler, F., Intracellular injections of EGTA block induction of hippocampal long-term potentiation, *Nature (London)*, 305 (1983) 719-721.
- 8 Rahamimoff, R., A dual effect of calcium ions on neuromuscular facilitation, *J. Physiol. (London)*, 195 (1968) 471-480.
- 9 Shin, C., Pedersen, H.B. and McNamara, J.O., γ -Aminobutyric acid and benzodiazepine receptors in the kindling model of epilepsy: a quantitative radiohistochemical study, *J. Neurosci.*, 5 (1985) 2696-2701.

- 10 Stanley, E.F., Decline in calcium cooperativity as the basis of facilitation at the squid giant synapse, *J. Neurosci.*, 6 (1986) 782-789.
- 11 Stanton, P.C. and Sarvey, J.M., Depletion of norepinephrine but not serotonin reduces long-term potentiation in the dentate gyrus of the rat hippocampal slice, *J. Neurosci.*, 5 (1985) 2169-2177.
- 12 Tielen, A.M., Lopes da Silva, F.H., Mollevanger, W.J. and de Jonge, F.H., Differential effects of enkephalin within hippocampal areas, *Exp. Brain Res.*, 44 (1982) 343-346.
- 13 Turner, R.W., Bainbridge, K.G. and Miller, J.J., Calcium induced long-term potentiation in the hippocampus, *Neuroscience*, 7 (1982) 1411-1416.
- 14 Wigström, H. and Gustafsson, B., Large long-lasting potentiation in the dentate gyrus in vitro during blockade of inhibition, *Brain Res.*, 275 (1983) 153-158.