UvA-DARE (Digital Academic Repository)

Planning skills of adolescents with ADHD

Boyer, B.E.; Geurts, H.M.; Van der Oord, S.

DOI

[10.1177/1087054714538658](https://doi.org/10.1177/1087054714538658)

Publication date

2018

Document Version

Final published version

Published in

Journal of Attention Disorders

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

Boyer, B. E., Geurts, H. M., & Van der Oord, S. (2018). Planning skills of adolescents with ADHD. *Journal of Attention Disorders*, 22(1), 46-57.
<https://doi.org/10.1177/1087054714538658>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Planning Skills of Adolescents With ADHD

Journal of Attention Disorders
2018, Vol. 22(1) 46–57
© The Author(s) 2014
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1087054714538658
journals.sagepub.com/home/jad

Bianca E. Boyer^{1,3}, Hilde M. Geurts^{1,3,4}, and Saskia Van der Oord^{1,2,4} SAGE

Abstract

Objective: This study assesses planning skills of adolescents with ADHD using a multi-method assessment and explores subtype differences between inattentive and combined subtypes. **Method:** One hundred fifteen adolescents with ADHD and 34 typically developing controls completed subtests of the Delis–Kaplan Executive Function System (D-KEFS), Behavioral Assessment of the Dysexecutive Syndrome (BADS), and Behavior Rating Inventory of Executive Function (BRIEF). **Results:** No group differences were found on the D-KEFS. The BADS Zoo Map test showed longer completion times in the inattentive than in the combined subtype and controls. The BRIEF showed more planning problems in adolescents with ADHD (both subtypes) than in controls, with the combined subtype showing most problems on overall executive functioning. **Conclusion:** Only a proportion of adolescents with ADHD shows planning deficits. To capture these planning problems in adolescents with ADHD, it seems important to use multiple measures of planning, both ratings and “less structured” neurocognitive measures of planning. (*J. of Att. Dis.* 2018; 22(1) 46–57)

Keywords

ADHD, subtype differences, executive function, assessment, adolescent ADHD

Introduction

Given that in adolescence, parental control diminishes and a need for independence arises (Spear & Kulbok, 2004), planning is an important skill to secure an independent life-style (Nurmi, 1991). Ample research shows that planning skills of typically developing adolescents (TDA) increase during adolescence (Anderson, Anderson, Northam, Jacobs, & Catroppa, 2001), but no studies have investigated the planning skills of adolescents with ADHD in detail using multiple measures of planning, and few studies have focused on subtype differences (*but* Qian, Shuai, Chan, Qian, & Wang, 2013). To investigate planning skills of adolescents with ADHD in more detail, this study assesses planning skills of adolescents with different subtypes of ADHD using a multi-method assessment with both cognitive tasks and parental ratings of planning.

Executive Functions (EFs) play a pivotal role in most theories of ADHD (Barkley, 1997; Sonuga-Barke, 2003; Sonuga-Barke, Bitsakou, & Thompson, 2010). EFs are described as neurocognitive processes that enable self-control (Lezak, 1995). EFs are necessary to maintain an appropriate problem solving set to attain a future goal (Welsh & Pennington, 1988). Individuals with ADHD (mainly based on results of studies with children younger than 12 years) show most problems with the EFs inhibition and working memory (Willcutt, Doyle, Nigg, Faraone, & Pennington, 2005). In adolescence, the symptomatology of ADHD changes, as the symptoms of hyperactivity and impulsivity decrease in comparison with

childhood (Fischer, Barkley, Smallish, & Fletcher, 2005), and in daily life, problems with the EF planning are described to be more pronounced (Barkley, 2004; Wolraich et al., 2005). Planning is generally defined as the process of formulating a sequence of operations intended to achieve a goal in an organized, strategic, and efficient manner (e.g., Hayes-Roth & Hayes-Roth, 1979).

In adolescence, the control and help of parents and teachers diminishes as compared with childhood, while at the same time, the transition to secondary school increases the need for EFs, particularly for planning skills. For example, in secondary school, adolescents have to change classes and teachers every hour, are expected to be where they are supposed to be on time, and are expected to bring necessary materials for all their classes. Adolescents are expected to independently manage planning for long-term projects, study for tests, and complete and turn-in assignments for multiple classes and teachers on time (Evans et al., 2009). Therefore, it is not surprising that planning aspects of EF

¹University of Amsterdam, The Netherlands

²KU Leuven, Belgium

³Autism & ADHD Research Center, The Netherlands

⁴Center for Cognitive Science, University of Amsterdam, The Netherlands

Corresponding Author:

Bianca E. Boyer, Department of Developmental Psychology, University of Amsterdam, Weesperplein 4, 1018 XA Amsterdam, The Netherlands.
Email: b.e.boyer@uva.nl

are highly predictive of academic functioning in adolescents with ADHD (Langberg, Dvorsky, & Evans, 2013).

Even though planning problems appear to have important consequences for adolescents in particular, studies assessing planning skills in ADHD focusing on the adolescent age are sparse (Gau, Chiu, Shang, Cheng, & Soong, 2009; Gau & Shang, 2010; Qian et al., 2013; Toplak, Bucciarelli, Jain, & Tannock, 2009). Planning problems in adolescents with ADHD have been described on a behavioral level (Barkley, 2004; Wolraich et al., 2005) and by using a parent-rated questionnaire (using the Behavior Rating Inventory of EF; Langberg et al., 2013; Toplak et al., 2009). However, the few studies using neuropsychological measures of planning skills in adolescents with ADHD show mixed results. In three studies, adolescents with ADHD show significantly impaired performance on a neurocognitive measure of planning, the computerized Stockings of Cambridge task of the Cambridge Neuropsychological Test Automated Battery (CANTAB; Gau et al., 2009; Gau & Shang, 2010; Toplak et al., 2009) as compared with TDA. However, in contrast, a recent large study investigating the development of planning skills of children and adolescents using another neurocognitive measure of planning (Tower of Hanoi) showed no significant differences in planning skills between adolescents with ADHD and TDA (Qian et al., 2013). Studies focusing on children younger than 12 years of age show the same contradiction: even though studies predominantly found children with ADHD have impaired planning skills on neurocognitive measures as compared with controls (e.g., Gioia, Isquith, Kenworthy, & Barton, 2002; McCandless & O'Laughlin, 2007; Nigg, Blaskey, Huang-Pollock, & Rapaport, 2002; O'Brien, Dowell, Mostofsky, Cenckla, & Mahone, 2010; Shimon, Engel-Yeger, & Tirosh, 2012; Wodka, Loftis, et al., 2008), some studies did *not* find these differences (Qian et al., 2013; Riccio, Homack, Jarratt, & Wolfe, 2006; Wodka, Loftis, et al., 2008; Wodka, Mostofsky, et al., 2008).

In addition, there are few studies on subtype differences in planning skills. The only study conducted in an adolescent sample shows no subtype differences in planning skills as assessed by a neurocognitive measure (Qian et al., 2013), whereas studies focusing on subtype differences in planning skills of children younger than 12-year old show inconsistent results. Two studies show that children with the ADHD combined subtype (ADHD-C) showed significantly more planning problems in comparison with the inattentive subtype (ADHD-I; Klorman et al., 1999; Lockwood, Marcotte, & Stern, 2001); other studies did not find significant differences between subtypes (Nigg et al., 2002; Riccio et al., 2006; Salcedo-Marin, Moreno-Granados, Ruiz-Veguilla, & Ferrin, 2013; Wodka, Loftis, et al., 2008; Wodka, Mostofsky, et al., 2008). Different factors may account for these inconsistencies in the literature.

First of all, planning skills go through major developments during late childhood and adolescence and some studies draw conclusions regarding planning skills based on samples with broad age ranges from childhood to adolescence. These studies thereby possibly do not detect differences in planning skills or finding differences that may not be there when focusing on a specific developmental stage (Gioia et al., 2002; Riccio et al., 2006; Salcedo-Marin et al., 2013; Wodka, Loftis, et al., 2008; Wodka, Mostofsky, et al., 2008). Second, studies often use only one task to measure planning. Between these studies, there is a large variability of the measures used, while research shows only small and non-significant correlations between these different measures of planning (Welsh, Satterlee-Cartmell, & Stine, 1999). In addition, these different measures of planning differ greatly in level of ecologic validity (Gau et al., 2009; Gau & Shang, 2010; Gioia et al., 2002; Harrier & DeOrnellas, 2005; Klorman et al., 1999; Kofman, Larson, & Mostofsky, 2008; Lockwood et al., 2001; McCandless & O'Laughlin, 2007; Qian et al., 2013; Salcedo-Marin et al., 2013; Toplak et al., 2009). Ecological validity refers to the extent to which the results of controlled tests generalize to performance in naturalistic settings (Wilson, Alderman, Burgess, Emslie, & Evans, 2003). Whether a neuropsychological task to assess planning skills (low ecologic validity) measures the same construct as a parent-rated questionnaire (high ecologic validity) is debatable (Toplak, West, & Stanovich, 2013). Therefore, to get a clear picture of planning skills of adolescents with ADHD in this specific developmental stage, it may be important to use *multiple* neuropsychological tests and complement them with questionnaires to get a broad assessment of planning skills.

Given the importance of planning skills for the daily life of adolescents and the planning impairments that are reported within samples of adolescents with ADHD, a broad assessment of planning skills of adolescents with ADHD as compared with adolescents without ADHD is needed. Therefore, the first goal of this study is to assess planning skills of adolescents with ADHD as compared with planning skills of TDAs. To do this, multiple tasks are used that have varying levels of ecological validity. First, neuropsychological tasks are utilized that are proposed to be pure measures of planning but often have relatively low ecologic validity. Second, neuropsychological tasks are used with higher ecologic validity (i.e., are more comparable with daily life situations; Wilson et al., 2003). Finally, in this study, the most ecologically valid assessment of planning skills is the parental rating of daily life planning problems (Langberg et al., 2013; Toplak et al., 2013). Group differences between adolescents with ADHD and TDA are assessed and, in addition, the clinical relevance of results will be assessed by comparing the percentage of adolescents scoring within the clinical range in both groups of adolescents. Based on behavioral descriptions of ADHD in

adolescence (Barkley, 2004; Wolraich et al., 2005) and in line with previous research (Gau et al., 2009; Gau & Shang, 2010; Toplak et al., 2009 but Qian et al., 2013), it is hypothesized that compared with TDAs, adolescents with ADHD will show clinically impaired planning skills on all planning measures and thus on all three levels of ecologic validity.

The second goal is investigating subtype differences in planning skills, with focus on differences between ADHD-I and ADHD-C, because the hyperactive/impulsive subtype (ADHD-H) is uncommon during adolescence (Farone, Biederman, & Mick, 2006). Given that the only study assessing subtype differences within adolescence (with only one neurocognitive measure) showed no significant differences between subtypes (Qian et al., 2013), no differences are expected between the ADHD-C and ADHD-I subtypes (Qian et al., 2013).

Method

Participants

Thirty-four TDAs, 77 adolescents with ADHD-I, and 38 adolescents with ADHD-C participated. The participants met the following inclusion criteria: (a) age range between 12 and 17 years, (b) attending a regular middle or high school, (c) no active non-pharmacological treatment for ADHD or associated symptoms, and (d) IQ above 80, measured by two subtests (Vocabulary and Block Design) of the Dutch Wechsler Intelligence Scale for Children–III (WISC-III; Kort et al., 2005); these two subtests correlate highly with Full Scale IQ (FSIQ; Sattler, 2001). The study was approved by the Ethics Committee of the University of Amsterdam.

Selection of the groups

ADHD group. All adolescents applied for psychosocial treatment at 1 of 16 participating outpatient mental health institutes in the Netherlands. Before receiving treatment, they participated in the current study. All of these adolescents ($N = 201$) had a prior clinical *Diagnostic and Statistical Manual of Mental Disorders* (4th ed., text rev.; *DSM-IV-TR*; American Psychiatric Association [APA], 2000) diagnosis of ADHD. Diagnosis was confirmed by parent versions of the Dutch Disruptive Behavior Disorder (DBD) rating scale (Oosterlaan, Scheres, Antrop, Roeyers, & Sergeant, 2000; Pelham, Gnagy, Greenslade, & Milich, 1992) and the Diagnostic Interview Schedule for Children for *DSM-IV* (DISC-IV; Shaffer, Fisher, Lucas, Dulcan, & Schwab-Stone, 2000). The DBD contains four scales composed of the *DSM-IV* (APA, 1994) criteria for ADHD Inattention, ADHD Hyperactivity/Impulsivity, Oppositional Defiant Disorder, and Conduct Disorder. Adequate psychometric properties have been reported (Oosterlaan et al., 2000). Only adolescents with a clinical score on any of the two ADHD scales of the DBD were included in this study;

47 adolescents did not meet this criterion and were excluded. Also, 15 adolescents were excluded because their parents failed to complete the questionnaires due to technical problems with the questionnaire software.

The DISC-IV, a structured diagnostic interview, establishes ADHD group membership and subtype status based on a diagnostic algorithm, including a check of cross-situational impairment (Shaffer et al., 2000). Only adolescents with an ADHD-I or ADHD-C diagnosis on the DISC-IV were included; 15 adolescents did not reach ADHD criteria and 6 adolescents met criteria for the ADHD-H subtype and were excluded. Another 3 adolescents had an FSIQ below 80 and were excluded from the study. This resulted in an ADHD-I group of 77 and an ADHD-C group of 38 participants. The adolescents with ADHD taking methylphenidate discontinued medication at least 24 hr before testing, allowing for a complete wash out (Paykina & Greenhill, 2007).

TDA group. TDAs were recruited from a community school in the Netherlands ($n = 39$). All typically developing participants scored below the 90th percentile on the DBD. Parents of one adolescent failed to complete the questionnaires, one adolescent had a prior ADHD diagnosis, and three adolescents had an FSIQ under 80: They were excluded from the study, resulting in 34 TDAs.

Instruments

Neuropsychological measures. Two subtests of the Delis–Kaplan Executive Function System (D-KEFS; Delis, Kaplan, & Kramer, 2001) were used: the Tower Test and the Trail Making Test (TMT). The D-KEFS has good reliability and validity (Homack, Lee, & Riccio, 2005).

Tower Test. The Tower Test assesses several key EFs, including planning. The objective of this subtest is to move disks, varying in size from small to large, across three pegs to build a designated tower in the fewest number of moves possible. In constructing the target towers, the examinee must follow two rules: (a) move only one disk at a time and (b) never place a larger disk over a smaller disk.

Two scores from the Tower Test were used: the “total achievement score” and the “move accuracy ratio.” The total achievement score gives an overall planning score and is based on the number of moves to build the tower, rule violations, completion time, and final tower correctness. Scores range from 0 to 30, with higher scores indicating better overall planning skills. The move accuracy ratio is calculated by dividing the total number of moves set by the minimum number of moves necessary to build the designated tower. Scores range from 0 to 3.1: The lower this score, the less moves were necessary to obtain the correct

tower; therefore, a lower score indicates good planning skills.

TMT. We used three conditions of the TMT to calculate mental flexibility (that is considered necessary for planning). In the “switching” condition, adolescents were instructed to connect items on a page, switching between numbers and letters beginning with 1 and A, and connecting them in numerical and alphabetical order, respectively. To control for possible differences in motor speed and familiarity with numbers and letters, the “number” condition (connecting numbers in numerical order) and the “letter” condition (connecting letters in alphabetical order) were also administered. In this study, mental flexibility is measured, using the time to complete the switching condition minus the average time to complete the number and letter conditions. Scores can range from –150 to 240 s, with a higher score indicating more difficulty to switch between two sets.

Ecologically valid neuropsychological measures. Two subtests of the Behavioral Assessment of the Dysexecutive Syndrome (BADS; Wilson, Alderman, Burgess, Emslie, & Evans, 1996) were used: Key Search and Zoo Map. These subtests improve the ability to predict daily planning difficulties by assessing more complex, real-life situations than do neuropsychological measures (Wilson et al., 2003). Inter-rater reliability was found to be high ($r = .88$) and discriminative validity was supported (Baron, 2007). Usually the BADS is used for ages 16 and older and, for children, the Behavioral Assessment of the Dysexecutive Syndrome in Children (BADS-C) is used (norms up to 16 years). As ceiling effects are reported for subtests of the BADS-C (Henry & Bettenay, 2010; Shimoni et al., 2012) and our sample is on the older end of the norm group of the BADS-C, we decided to use the BADS.

Key Search. To examine the ability to plan an efficient, systematic plan of action, adolescents are presented with a drawing of a 100-mm square that represents a field in which they have lost their key. The adolescents are asked to draw a pathway through which it would be reasonable to search for the lost key. Two measures were used: (a) the strategy score (ranging from 0 to 16, with a higher score indicating a better performance), based on whether the rater believes the strategy to be systematic, efficient, and likely to be effective; and (b) completion time.

Zoo Map. This subtest consists of two conditions. In Condition 1, adolescents are asked to plan a route on a given zoo map, passing through a list of animals (the ability to formulate and implement a plan). In Condition 2, they have to pass the list of animals in a given order (follow a pre-formulated plan), thereby making less of an appeal to

planning skills than Condition 1. For each condition, two measures were used: (a) the order score, based on successful implementation of the plan (ranging from minus the number of errors to 8, with a higher score indicating a better performance); and (b) completion time.

Parental rating of daily life planning. The Dutch Behavior Rating Inventory of Executive Function (BRIEF; Gioia, Isquith, Guy, & Kenworthy, 2000; Smidts & Huizinga, 2009) is a normative behavioral rating scale for children 5 to 18 years old, designed to elicit everyday EF as observed by the parents in natural everyday environments. The internal consistency, the test–retest stability, and convergent and discriminant validity of the Dutch version of the BRIEF are considered good (Huizinga & Smidts, 2010).

The BRIEF consists of 75 items of which each item pertains to specific everyday behavior, relevant to EF. Parents were asked to indicate how often their child displayed a given behavior in the past 2 weeks by giving one of three responses, “Never,” “Sometimes,” or “Often.” The items of the BRIEF are categorized into eight clinical scales: Inhibit, Shift, Emotional Control, Initiate, Working Memory, Plan/Organize, Organization of Materials, and Monitor.

In this study, three scales were used: (a) Plan/Organize, (b) Metacognition Index (MI), and (c) Global Executive Composite (GEC). The Plan/Organize scale measures the child’s capacity to anticipate future events, set goals, develop appropriate steps to carry out associated tasks or actions, and manage current and future-oriented task demands (12 items, scores range from 12 to 36). The MI is a composite score of the scales Initiate, Working Memory, Plan/Organize, Organization of Materials, and Monitor Scales, and represents the ability to plan, organize, initiate, and hold information in mind for future-oriented problem solving (44 items, scores range from 44 to 132). The GEC is a summary score that includes all eight clinical scales (72 items, scores range from 72 to 216). Higher scores on the subscales of the BRIEF indicate more EF problems.

Procedure. Prior to face-to-face testing, a telephonic interview was conducted with parents to obtain demographic information and check inclusion criteria. After completion of informed consent by parent and adolescent, the parent was interviewed using the DISC-IV and questionnaires were filled out. At the same time, the adolescent was tested. As the tasks administered were part of a multi-center study on treatment of adolescents with ADHD, in addition to the tasks described above, other tasks were also administered. Duration of assessment was 2.5 hr including breaks. All assessments started with the WISC-III followed by the neuropsychological batteries (BADS/D-KEFS), which were presented in counterbalanced order across participants, to account for order effects. After assessment, participants received a small gift (worth approximately 2 euro).

Missing data and outliers. Either examiner error or adolescent non-compliance led to missing data. The number of missing cases ranged from 0 to 5 (D-KEFS, $n = 5$; BADS, $n = 0$; BRIEF, $n = 0$). Of all tests, only the subtest Key search showed outliers ($n = 2$). As analyses with and without outlier truncation showed similar results, original analyses without outlier truncation are reported.

Statistical analysis. Correlations between dependent measures were calculated (Table 1). Next, differences in planning skills between diagnostic groups (TD, ADHD-I, ADHD-C) on three levels of measurement (D-KEFS, BADS, BRIEF) were tested using MANOVAs on the raw scores. As assumptions of normality and homogeneity of variances were not retained for all dependent variables, Pillai's trace was chosen as a test statistic. Pillai's trace has been shown to be most robust to violations of unequal sample sizes (Stevens, 1996). In case of main group effects, Games-Howell post hoc tests (accurate with unequal sample sizes; Field, 2009) were performed. Analyses were conducted with and without covarying for medication use (yes/no); when medication use affected results, this will be reported.

Finally, based on age-based norms, adolescents' scores were classified as "clinical" or "non-clinical." The D-KEFS has been standardized with large normative and patient samples of different age ranges (Homack et al., 2005): On the TMT and the Tower Test (Delis et al., 2001) scaled scores ($M = 10$, $SD = 3$) of 6 or below (comparable with the 10th percentile) are considered to represent clinical planning problems. The norms of the BADS are based on 216 healthy individuals (M age = 46.6, $SD = 19.8$), but age-based data on the individual tasks are not provided. As EF develops dramatically during adolescence, age-based norms are important. Therefore, we used the best available norm group: the scores of our TDA. On all BADS measures, the average score of TDA was calculated, and based on their standard deviation, the 10th or 90th percentile was calculated (depending on the measure used) to represent the cut-off score for clinical planning problems. On the BRIEF, age-based norms lead to T -scores: T -scores greater than 65 (comparable with the 90th percentile) are considered to represent clinical planning problems. Differences between groups in clinical status are assessed with chi-square analyses.

Results

Sample description. Groups did not differ from each other with respect to age, gender, and FSIQ (Table 2). As expected, ADHD groups showed more symptoms than the typically developing (TD) group and the ADHD-C group showed more hyperactive/impulsive symptoms than did the ADHD-I group. Also, the ADHD-C group showed more

inattentive symptoms than the ADHD-I group on the DISC-IV. The ADHD groups showed no differences in the number of methylphenidate (MPH) users or average school week dosage of immediate release MPH.

MANOVA. Main group effects of the MANOVAs for both the D-KEFS tasks (TMT and Tower Test) and the BADS tasks (Key Search and Zoo Map Test) were non-significant (respectively $p = .40$, $\eta_p^2 = .02$ and $p = .23$, $\eta_p^2 = .05$; Table 3). However, univariate tests revealed a group difference on the BADS Zoo Map completion time in Condition 1, $F(2, 146) = 3.05$, $p = .05$, $\eta_p^2 = .04$, showing a longer completion time in the ADHD-I group in comparison with the two other groups. When covarying for medication use, completion time was no longer significant, but the order score in Condition 1 of the BADS Zoo Map was.

There was a significant main group effect for the BRIEF, $V = .92$, $F(6, 290) = 43.26$, $p < .0001$, $\eta_p^2 = .46$. Univariate group comparisons showed that groups differed significantly on the Plan/Organize Scale, $F(2, 146) = 106.64$, $p < .0001$, $\eta_p^2 = .59$; MI, $F(2, 146) = 133.54$, $p < .0001$, $\eta_p^2 = .65$; and GEC, $F(2, 146) = 143.67$, $p < .0001$, $\eta_p^2 = .66$. Games-Howell post hoc tests showed more impaired EFs in the ADHD groups than in the TD group on all BRIEF scales. Also, on the GEC, the ADHD-C group showed a significantly higher score than the ADHD-I group.

Clinical significance. Chi-square tests were conducted, examining group differences in the number of adolescents scoring in the clinical range of all three measures (Table 4). On the D-KEFS and BADS, most group effects were non-significant (Tower Test total achievement score, move accuracy, and TMT, $p = 1.00$, $p = .25$, $p = .46$, respectively; BADS Key Search strategy score, $p = .76$; time, $p = .63$; Zoo Map 1 order score, $p = .67$; Zoo Map 2 order score, $p = .46$; time, $p = .83$), except for completion time of Zoo Map Condition 1, $\chi^2(2) = 6.11$, $p = .05$, in which more adolescents with ADHD-I showed a longer completion time than in the other two groups.

On the BRIEF, there were significant differences between groups on the Plan/Organize scale, $\chi^2(2) = 28.42$, $p < .0001$; MI scale, $\chi^2(2) = 22.06$, $p < .0001$; and GEC, $\chi^2(2) = 34.19$, $p < .0001$. On all scales of the BRIEF, significantly more ADHD adolescents (both subtypes) than TDA scored within the clinical range. Furthermore, on the GEC, significantly more adolescents with ADHD-C than with ADHD-I scored within the clinical range.

Discussion

This study investigated planning skills of adolescents with ADHD and subtype differences using a multi-method assessment on different levels of ecological validity. In general, on our neuropsychological measures, adolescents with

Table 1. Correlations Between IQ, Symptoms, and Planning Measures.

	FSIQ	Inattention	Hyperactivity/ impulsivity	Tower Test total achievement	Tower Test move accuracy	TMT	Key Search strategy score	Key Search time	Zoo Map 1 order score	Zoo Map 1 time	Zoo Map 2 order score	Zoo Map 2 time	Plan/ organize	MI	GEC
FSIQ	1														
Inattention	-.196*	1													
Hyperactivity/ impulsivity	-.111	.489**	1												
Tower Test total achievement	-.041	.03	.01	1											
Tower Test move accuracy	.092	-.074	.095	-.496**	1										
TMT	-.149	.164*	-.007	-.103	.051	1									
Key Search strategy score	.169*	-.081	-.082	.193*	-.078	-.026	1								
Key Search time	.01	-.017	.107	-.166*	.104	.003	.001	1							
Zoo Map 1 order score	.074	-.061	-.103	.023	.056	.142	.076	.039	1						
Zoo Map 1 time	-.105	.101	.053	.027	-.156	-.042	.097	.151	-.292**	1					
Zoo Map 2 order score	-.099	-.075	-.148	-.132	-.143	-.02	.144	-.081	.01	.141	1				
Zoo Map 2 time	-.084	.136	.08	.089	-.104	.131	.006	.048	-.310**	.353**	.01	1			
Plan/organize	-.168*	.834**	.449**	-.01	.023	.099	-.068	-.021	-.068	.113	-.192*	.042	1		
MI	-.16	.855**	.499**	-.023	.019	.089	-.053	-.057	-.057	.106	-.16	.034	.948**	1	
GEC	-.190*	.810**	.622**	.011	.027	.057	-.043	-.031	-.082	.105	-.143	.046	.894**	.954**	1

Note. FSIQ = Full Scale IQ; TMT = Trail Making Test; MI = Metacognition Index; GEC = Global Executive Composite.

* $p < .05$. ** $p < .01$.

Table 2. Demographic and Clinical Description of the Sample.

Measure	TDA (<i>n</i> = 34)		ADHD-I (<i>n</i> = 77)		ADHD-C (<i>n</i> = 38)		Group comparisons
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	
Age (years)	14.66	1.20	14.40	1.3	14.11	1.21	<i>ns</i>
<i>n</i> Boys (%)	21	(61.8)	60	(77.9)	23	(60.5)	<i>ns</i>
FSIQ	106.85	13.95	101.99	11.34	103.74	13.25	<i>ns</i>
DBD							
Inattention	3.28	3.17	18.71	4.03	18.42	4.53	TDA < ADHD-I = ADHD-C**
Hyperactivity/impulsivity	1.33	1.55	6.35	4.20	14.48	4.62	TDA < ADHD-I < ADHD-C**
DISC-IV							
Inattentive	—	—	15.65	1.95	16.32	2.01	ADHD-I < ADHD-C*
Hyperactive	—	—	3.99	3.13	13.89	2.38	ADHD-I < ADHD-C**
Medication							
<i>n</i> MPH users (%)	—	—	51	(66.2)	31	(81.6)	<i>ns</i>
MPH IR school week dose	—	—	36.28	18.48	33.91	14.52	<i>ns</i>

Note. TDA = typically developing adolescents; I = inattentive subtype; C = combined subtype; *ns* = not significant; FSIQ = Full Scale IQ; DBD = Disruptive Behavior Disorder Rating Scale; DISC-IV = Diagnostic Interview Schedule for Children; MPH = methylphenidate; IR = all medication was converted into immediate release dose.

*p < .05. **p < .01.

ADHD (both subtypes) showed few planning deficits, which is in contrast to our predictions and other studies (Gau et al., 2009; Gau & Shang, 2010; Toplak et al., 2009; *but see* Qian et al., 2013). Only on one outcome, of the more ecologically valid task (BADs), there was a group difference, showing that adolescents with ADHD-I need more time for adequate planning than TDAs and ADHD-C. Finally, parents of adolescents with both subtypes of ADHD reported more daily life planning problems than parents of TDAs. In addition, parents of adolescents with ADHD-C reported more overall EF problems than those with ADHD-I. Our findings are comparable with the mixed findings of previous studies investigating planning skills of adolescents with ADHD (Gau et al., 2009; Gau & Shang, 2010; Qian et al., 2013; Toplak et al., 2009).

These findings lead to the question what is the best way to measure planning skills in adolescents with ADHD, by questionnaires or by neurocognitive measures. Unfortunately, there is no “golden standard” on how to adequately measure daily life planning problems. Even though observations of daily life planning skills by independent raters would probably be the most ecologically valid measure of planning, in general, parent-rated questionnaires are considered to capture these real-life planning problems (Siu & Zhou, 2014; Toplak et al., 2013). On parent-rated questionnaires, children (Gioia et al., 2002; Shimoni et al., 2012; Siu & Zhou, 2014) as well as adolescents with ADHD (this study; Toplak et al., 2009) consistently show more planning problems than controls. With regard to neurocognitive tasks of planning skills, a discrepancy is visible: On “paper-and-pencil,” tasks results are mixed in both child and adolescent samples (this study; Galasso et al., 2014; Klorman et al.,

1999; Nigg et al., 2002; O’Brien et al., 2010; Qian et al., 2013; Shimoni et al., 2012; Siu & Zhou, 2014; Wodka, Loftis, et al., 2008; Wodka, Mostofsky, et al., 2008), whereas on computerized tasks, studies consistently show planning deficits in both children and adolescents with ADHD (Gau et al., 2009; Gau & Shang, 2010; Song & Hakoda, 2014; Toplak et al., 2009). In line with the literature, also in our study on parent-rated measures, planning deficits were found, and in our paper-and-pencil neuropsychological measures, we found a significant deficit on one task (BADs) but not on another (D-KEFS). In summary, in the literature, the two measures consistently showing planning problems in children and adolescents with ADHD are on both ends of the continuum of ecological validity: parent-rated measures on one end and computerized neurocognitive tests on the other. Thus, the degree of ecological validity does not seem to be the best explanation for the mixed findings of studies assessing planning skills of adolescents with ADHD.

A better explanation may be that parent ratings of planning skills, neurocognitive paper-and-pencil and computerized planning measures, assess different components of planning (Toplak et al., 2013). However, another plausible influence to performance on planning measures, or to planning skills overall, are motivational incentives. Studies using a computerized tower test and parent report showed impaired planning skills of adolescents with ADHD (Gau et al., 2009; Gau & Shang, 2010; Toplak et al., 2009), whereas studies using paper-and-pencil tests did not (this study; Qian et al., 2013). Paper-and-pencil tests inevitably require more involvement of an examiner, create a more structured environment, and are more “hands-on,” therefore possibly

Table 3. Group Comparison of Subtypes of ADHD and TDA on Planning Measures.

	TDA		ADHD-I		ADHD-C		
Measure	M	SD	M	SD	M	SD	Group comparisons
D-KEFS							
Tower Test total achievement	17.47	2.76	17.40	2.51	18.11	2.66	ns
Tower Test move accuracy	1.92	0.56	1.85	0.5	1.92	0.59	ns
TMT	38.89	21.14	43.18	19.61	42.82	19.71	ns
BADS							
Key Search strategy score	12.82	3.47	12.38	3.54	12.53	3.44	ns
Key Search time	54.09	36.77	54.40	38.27	62.47	59.10	ns
Zoo Map 1 order score	4.44	3.65	4.27	3.81	3.24	4.23	ns ^a
Zoo Map 1 time	117.47	42.12	150.04	73.18	135.47	62.97	TDA = ADHD-C < ADHD-I ^{*b}
Zoo Map 2 order score	7.97	0.39	7.82	0.81	7.74	0.80	ns
Zoo Map 2 time	56.68	17.15	63.48	18.27	62.13	17.18	ns
BRIEF							
Plan/organize	17.38	4.99	29.14	3.77	29.45	4.11	TDA < ADHD-I = ADHD-C**
MI	68.97	16.19	107.52	10.68	109.29	11.27	TDA < ADHD-I = ADHD-C**
GEC	104.68	20.22	155.13	15.68	168.45	16.63	TDA < ADHD-I < ADHD-C**

Note. The number of participants differs for some dependent variables due to missing data (see text). TDA = typically developing adolescents; I = inattentive subtype; C = combined subtype; D-KEFS = Delis-Kaplan Executive Function System; ns = not significant; TMT = Trail Making Test; BADS = Behavioral Assessment of the Dysexecutive Syndrome; BRIEF = Behavior Rating Inventory of Executive Function; MI = Metacognition Index; GEC = Global Executive Composite.

^aWhen covarying for medication use, this comparison becomes significant.

^bWhen covarying for medication use, this comparison is no longer significant.

* $p < .05$. ** $p < .01$.

causing adolescents to be more actively involved and more engaged in the task (Power, 1992; Toplak et al., 2013). As a result, these motivational incentives could improve their performance as compared with computerized tests or daily life tasks without such incentives, for example, homework. If this is the case, one might argue that adolescents with ADHD know how to plan, as long as they are motivated enough.

Another explanation for mixed findings between planning measures could be that tasks, used to assess planning skills, differ in the extent to which they appeal to other EFs. As finding your way in a zoo in the BADS Zoo Map Test could be considered less structured and possibly posits more distracters than building a tower in the Tower Test, it could make a stronger appeal to, for instance, working memory (remembering the assignment) or inhibition (not being distracted by other animals in the zoo). Therefore, our results could imply that planning problems observed in daily lives of adolescents with ADHD are not caused by deficits in underlying planning skills but by other underlying EF deficits (Phillips, Wynn, Gilhooly, Della Sala, & Logie, 1999; Salcedo-Marin et al., 2013; Sonuga-Barke, Dalen, Daley, & Remington, 2002). This is consistent with the idea that planning skills are part of the later developing “higher-order” EFs such as set shifting, whereas, for example, working memory and inhibition are “lower order” earlier developing EFs (Best, Miller, & Jones, 2009; Lin, Hsiao,

& Chen, 1999; O’Brien et al., 2010). It may well be that planning problems in daily life derive from problems in these lower order EFs or from other basic cognitive processes such as sustained attention. Unfortunately, assessment of any EF also implicates other cognitive processes that are not directly relevant to the target EF (Miyake, Friedman, Emerson, Witzki, & Howerter, 2000). Therefore, it cannot be determined to what extent the different EFs have contributed to performance on our neurocognitive measures. More work in this field is needed to investigate to what extent daily life planning problems in individuals with ADHD are caused by deficits in other cognitive processes or “lower order” EFs.

In line with Barkley’s behavioral description of planning problems of adolescents with ADHD (Barkley, 2004), prior research with children (Gioia et al., 2002; McCandless & O’Laughlin, 2007; Riccio et al., 2006; Shimon et al., 2012) and adolescents (Toplak et al., 2009), in our study adolescents with ADHD did show impairments on daily life planning skills as rated by their parents. However, only half to two third of the adolescents with ADHD scored within the clinical range of planning or EF problems as rated by parents. This is in line with studies showing that only a proportion of children with ADHD have EF deficits (Willcutt et al., 2005). In addition, this study also shows that there are differences between the subtypes of ADHD in planning skills. Both subtypes of ADHD show planning problems as

Table 4. Number of Adolescents Scoring Within the Clinical Range.

	TDA	ADHD-I	ADHD-C	
Measure	n in clinical range (%)			Group comparisons
D-KEFS				
Tower Test total achievement	0 (0.0)	0 (0.0)	0 (0.0)	ns
Tower Test move accuracy	9 (26.5)	12 (15.6)	10 (26.3)	ns
TMT	3 (8.8)	13 (16.9)	7 (18.4)	ns
BADS				
Key Search strategy score	5 (14.7)	14 (18.2)	5 (13.2)	ns
Key Search time	0 (0.0)	1 (1.3)	1 (2.6)	ns
Zoo Map 1 order score	3 (8.8)	10 (13.0)	6 (15.8)	ns
Zoo Map 1 time	2 (5.9)	20 (26.0)	7 (18.4)	TDA = ADHD-C < ADHD-I*
Zoo Map 2 order score	1 (2.9)	6 (7.8)	4 (10.5)	ns
Zoo Map 2 time	5 (14.7)	7 (9.1)	15 (39.5)	ns
BRIEF				
Plan/organize	1 (2.9)	38 (49.4)	23 (60.5)	TDA < ADHD-I = ADHD-C**
MI	3 (8.8)	40 (51.9)	22 (57.9)	TDA < ADHD-I = ADHD-C**
GEC	0 (0.0)	35 (45.5)	25 (65.8)	TDA < ADHD-I < ADHD-C**

Note. The number of participants differs for some dependent variables due to missing data (see text). TDA = typically developing adolescents; I = inattentive subtype; C = combined subtype; D-KEFS = Delis-Kaplan Executive Function System; ns = not significant; TMT = Trail Making Test; BADS = Behavioral Assessment of the Dysexecutive Syndrome; BRIEF = Behavior Rating Inventory of Executive Function; MI = Metacognition Index; GEC = Global Executive Composite.

* $p < .05$. ** $p < .01$.

perceived by parents but when assessing performance on neurocognitive tasks that resemble daily life, the ADHD-I subtype needed more time to adequately plan, whereas the ADHD-C subtype showed no deficits in planning. This could imply that within both subtypes, planning problems are apparent in daily life, but these planning problems may have different causes, with the ADHD-I subtype needing more time to plan. This is in accordance with the sluggish cognitive tempo that especially the ADHD-I subtype has been associated with (Langberg, Becker, & Dvorsky, 2014).

This study has several limitations. First, although we already had a multi-method assessment, having additional raters of planning problems (self-report, teacher report) would have given an even more all-round picture of EF. Furthermore, one could argue about the accuracy and/or validity of parental report in adolescence, as parents are likely to spend less time with their adolescent child in comparison with childhood (Achenbach, McConaughy, & Howell, 1987). Second, the TMT could be questioned as a measure of planning. Even though the TMT is designed to measure mental flexibility or set shifting, one could argue that the switching condition does make an appeal to planning skills, as planning is defined as the ability to formulate actions in advance and to approach a task in an organized, strategic, and efficient manner (Hayes-Roth & Hayes-Roth, 1979). Third, given that on most measures, we show no differences between ADHD and control groups, one may argue that our sample was selective or relatively unimpaired.

However, one could argue that the sample may be biased (more severe), given that all adolescents applied for psychosocial treatment. However, cross-situational impairment was part of the diagnostic procedure; 71.3% of the adolescents were taking medication for their ADHD and still applied for additional treatment at an outpatient mental health institute, implying a non-selective, impaired, and clinical representative sample of adolescents with ADHD. Finally, it has been proposed that some of the BRIEF items overlap with symptoms of ADHD (Gioia et al., 2000), and our study also showed strong correlations between ADHD scores on the ADHD rating scale and the three BRIEF scales used. However, after removing items from the BRIEF subscales that based on face validity, reflected *DSM-IV-TR* ADHD symptoms (APA, 2000), analyses still showed significant differences between adolescents with ADHD and TDAs (data available from first author).

This study brings forward some clinical implications. Only a proportion (about half to one third) of adolescents with ADHD appears to have planning deficits according to parents. Although both the ADHD-I and ADHD-C subtypes can experience planning problems, these problems seem to arrive from different underlying deficits: In ADHD-I, this may be related to a more sluggish cognitive tempo (Langberg et al., 2014). In the literature, planning deficits in ADHD have most consistently been shown using parent-rated measures and computerized neurocognitive tests: Given that questionnaires reporting daily life behavior

might assess different components of planning skills than neurocognitive measures, it seems important to use them complementary to get an overall picture of planning skills (Barkley & Edwards, 2006; Toplak et al., 2013). As the mixed findings on more “hands-on” neurocognitive tests suggest that motivation positively affects planning skills, enhancing motivation by adding structure and dividing large tasks in smaller steps to keep the adolescent actively involved may help resolve daily life planning problems in adolescents with ADHD.

All in all, in this study, a multi-method assessment was used to investigate planning skills in a large sample of adolescents with ADHD, while exploring differences between the two most prevalent subtypes in adolescence. More than half of adolescents with ADHD showed planning deficits as reported by parents, with the ADHD-C subtype showing additional overall EF deficits. However, “hands-on” neurocognitive measures showed mixed results. Future research is needed to investigate the causal processes that give rise to planning deficits in ADHD and to determine the best way to assess these deficits. For now, to capture planning deficits in adolescents with ADHD, the combined use of rating scales and “less structured” neurocognitive measures is recommended.

Acknowledgments

We thank ZonMW for providing this research grant. We are grateful to the participating Mental Health providers, families, and research assistants for their participation and collaboration.

Declaration of Conflicting Interests

Saskia van der Oord has been a paid consultant for Janssen pharmaceuticals in the development and evaluation of a serious game “Heelseeker” aimed at training cognitive functions. The other author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study is funded by ZonMW; project number 15700.3006.

References

- Achenbach, T. M., McConaughy, S. H., & Howell, C. T. (1987). Child/adolescents behavioral and emotional problems: Implications of cross-informant correlations for situational specificity. *Psychological Bulletin*, 101, 213-232.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., text rev.). Washington, DC: Author.
- Anderson, V. A., Anderson, P., Northam, E., Jacobs, R., & Catroppa, C. (2001). Development of executive functions through late childhood and adolescence in an Australian sample. *Developmental Neuropsychology*, 20, 385-406.
- Barkley, R. A. (1997). Behavioural inhibition, sustained attention, and executive functions: Constructing a unifying theory of AD/HD. *Psychological Bulletin*, 121, 65-94.
- Barkley, R. A. (2004). Adolescents with attention deficit/hyperactivity disorder: An overview of empirically based treatments. *Journal of Psychiatric Practice*, 10, 39-56.
- Barkley, R. A., & Edwards, G. (2006). Diagnostic interview, behavior rating scales, and the medical examination. In R. A. Barkley & G. Edwards (Eds.), *Attention-deficit hyperactivity disorder: A clinical workbook* (Vol. 2, pp. 337-368). New York, NY: Guilford Press.
- Baron, I. S. (2007). Behavioural assessment of the dysexecutive syndrome for children (BADS-C) by Emslie, H., Wilson, F. C., Burden, V., Nimmo-Smith, I., & Wilson, B. A. (2003). *Child Neuropsychology*, 13, 539-542.
- Best, J. R., Miller, P. H., & Jones, L. L. (2009). Executive functions after age 5: Changes and correlates. *Developmental Review*, 29, 180-200.
- Delis, D., Kaplan, E., & Kramer, J. (2001). *Delis-Kaplan Executive Function Scale*. San Antonio, TX: The Psychological Corporation.
- Evans, S. W., Schultz, B. K., White, L. C., Brady, C., Sibley, M. H., & Van Eck, K. (2009). A school-based organization intervention for young adolescents with attention-deficit/hyperactivity disorder. *School Mental Health*, 1, 78-88.
- Farone, S., Biederman, J., & Mick, E. (2006). The age dependent decline of attention-deficit/hyperactivity disorder: A meta-analysis of follow-up studies. *Psychological Medicine*, 36, 159-165.
- Field, A. (2009). *Discovering statistics using SPSS* (3rd ed.). London, England: SAGE.
- Fischer, M., Barkley, R. A., Smallish, I., & Fletcher, K. (2005). Executive functioning in hyperactive children as young adults: Attention, inhibition, and response perseveration and the impact of comorbidity. *Developmental Neuropsychology*, 27, 107-133.
- Galasso, C., Lo-Castro, A., Di Carlo, L., Pitzianti, M. B., D'Agati, E., Curatolo, P., & Pasini, A. (2014). Planning deficit in children with neurofibromatosis Type 1: A neurocognitive trait independent from attention-deficit hyperactivity disorder (ADHD)? *Journal of Child Neurology*. Advance online publication. doi:10.1177/0883073813517001
- Gau, S. S., Chiu, C., Shang, C., Cheng, A. T., & Soong, W. (2009). Executive function in adolescence among children with attention-deficit/hyperactivity disorder in Taiwan. *Journal of Developmental & Behavioral Pediatrics*, 30, 525-534.
- Gau, S. S., & Shang, C. (2010). Executive functions as endophenotypes in ADHD: Evidence from the Cambridge Neuropsychological Test Battery (CANTAB). *Journal of Child Psychology and Psychiatry*, 51, 838-849.
- Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). *Behavior Rating Inventory of Executive Function*. Odessa, FL: Psychological Assessment Resources.
- Gioia, G. A., Isquith, P. K., Kenworthy, L., & Barton, R. M. (2002). Profiles of everyday executive function in acquired

- and developmental disorders. *Child Neuropsychology*, 8, 121-137.
- Harrier, L. K., & DeOrnellas, K. (2005). Performance of children diagnosed with ADHD on selected planning and reconstitution tests. *Applied Neuropsychology: Adult*, 12, 106-119.
- Hayes-Roth, B., & Hayes-Roth, F. (1979). A cognitive model of planning. *Cognitive Science*, 3, 245-262.
- Henry, L. A., & Bettenay, C. (2010). The assessment of executive functioning in children. *Child and Adolescent Mental Health*, 15, 110-119.
- Homack, S., Lee, D., & Riccio, C. A. (2005). Test review: Delis-Kaplan executive function system. *Journal of Clinical and Experimental Neuropsychology*, 27, 599-609.
- Huizinga, M., & Smidts, D. P. (2010). Age-related changes in executive function: A normative study with the Dutch version of the Behavior Rating Inventory of Executive Function (BRIEF). *Child Neuropsychology*, 17, 51-66.
- Klorman, R., Hazel-Fernandez, L. A., Shaywitz, S. E., Fletcher, J. M., Marchione, K. E., Holahan, J. M., . . . Shaywitz, B. A. (1999). Executive functioning deficits in attention-deficit/hyperactivity disorder are independent of oppositional defiant or reading disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 38, 1148-1155.
- Kofman, O., Larson, J. G., & Mostofsky, S. H. (2008). A novel task for examining strategic planning: Evidence for impairment in children with ADHD. *Journal of Clinical and Experimental Neuropsychology*, 30, 261-271.
- Kort, W., Schittekatte, M., Bosmans, M., Compaa, E. L., Dekker, P. H., Vermeir, G., & Verhaeghe, P. (2005). *Wechsler Intelligence Scale for Children-III: Handleiding*. Amsterdam, The Netherlands: Pearson.
- Langberg, J. M., Becker, S. P., & Dvorsky, M. R. (2014). The association between sluggish cognitive tempo and academic functioning in youth with attention-deficit/hyperactivity disorder (ADHD). *Journal of Abnormal Child Psychology*, 42, 91-103. doi:10.1007/s10802-013-9722-3
- Langberg, J. M., Dvorsky, M. R., & Evans, S. W. (2013). What specific facets of executive function are associated with academic functioning in youth with attention-deficit/hyperactivity disorder? *Journal of Abnormal Child Psychology*, 41, 1145-1159.
- Lezak, M. (1995). *Neuropsychological assessment* (3rd ed.). New York, NY: Oxford University Press.
- Lin, C. C. H., Hsiao, C. K., & Chen, W. J. (1999). Development of sustained attention assessed using the Continuous Performance Test among children 6-15 years of age. *Journal of Abnormal Child Psychology*, 27, 403-412.
- Lockwood, K. A., Marcotte, A. C., & Stern, C. (2001). Differentiation of attention-deficit/hyperactivity disorder subtypes: Application of a neuropsychological model of attention. *Journal of Clinical and Experimental Neuropsychology*, 23, 317-330.
- McCandless, S., & O'Laughlin, L. (2007). The clinical utility of the Behavior Rating Inventory of Executive Function (BRIEF) in the diagnosis of ADHD. *Journal of Attention Disorders*, 10, 381-389.
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., & Howerter, A. (2000). The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: A latent variable analysis. *Cognitive Psychology*, 41, 49-100.
- Nigg, J. T., Blaskey, L., Huang-Pollock, C., & Rappley, M. D. (2002). Neuropsychological executive functions and ADHD DSM-IV subtypes. *Journal of the American Academy of Child & Adolescent Psychiatry*, 41, 59-66.
- Nurmi, J. E. (1991). How do adolescents see their future? A review of the development of future orientation and planning. *Developmental Review*, 11, 1-59.
- O'Brien, J. W., Dowell, L. R., Mostofsky, S. H., Cenckla, M. B., & Mahone, E. M. (2010). Neuropsychological profile of executive function in girls with attention deficit/hyperactivity disorder. *Archives of Clinical Neuropsychology*, 25, 656-670.
- Oosterlaan, J., Scheres, A., Antrop, I., Roeyers, H., & Sergeant, J. A. (2000). *Handleiding bij de Vragenlijst voor Gedragsproblemen bij Kinderen VvGK* [User's manual for the Disruptive Behavior Disorder Rating Scale in 6- to 16-year-old children]. Lisse, The Netherlands: Swets Test Publishers.
- Paykina, N. L., & Greenhill, L. L. (2007). Pharmacological treatments for attention deficit hyperactivity disorder. In P. E. Nathan & J. Gorman (Eds.), *A guide to treatments that work* (pp. 29-70). New York, NY: Oxford University Press.
- Pelham, W., Gnagy, E. M., Greenslade, K. E., & Milich, R. (1992). Teacher ratings of DSM-III-R symptoms for the disruptive behavior disorders. *Journal of the American Academy of Child & Adolescent Psychiatry*, 31, 210-218.
- Phillips, L. H., Wynn, V., Gilhooly, K. J., Della Sala, S., & Logie, R. H. (1999). The role of memory in the Tower of London task. *Memory*, 7, 209-231.
- Power, T. J. (1992). Contextual factors in vigilance testing of children with ADHD. *Journal of Abnormal Child Psychology*, 20, 579-593.
- Qian, Y., Shuai, L., Chan, R. C. K., Qian, Q. J., & Wang, Y. (2013). The developmental trajectories of executive function of children and adolescents with attention deficit hyperactivity disorder. *Research in Developmental Disabilities*, 34, 1434-1445.
- Riccio, C. A., Homack, S., Jarratt, K. P., & Wolfe, M. E. (2006). Differences in academic and executive function domains among children with ADHD predominantly inattentive and combined types. *Archives of Clinical Neuropsychology*, 21, 657-667.
- Salcedo-Marin, M. D., Moreno-Granados, J. M., Ruiz-Veguilla, M., & Ferrin, M. (2013). Evaluation of planning dysfunction in attention deficit hyperactivity disorder and autistic spectrum disorders using the Zoo Map Task. *Child Psychiatry & Human Development*, 44, 166-185. doi:10.1007/s10578-012-0317-y
- Sattler, J. M. (2001). *Assessment of children: Cognitive applications* (4th ed.). San Diego, CA: Author.
- Shaffer, D., Fisher, P., Lucas, C. P., Dulcan, M. K., & Schwab-Stone, M. E. (2000). NIMH Diagnostic Interview Schedule for Children Version IV (NIMH DISC-IV): Description, differences from previous versions, and reliability of some common diagnoses. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39, 28-38.
- Shimoni, M., Engel-Yeger, B., & Tirosh, E. (2012). Executive dysfunctions among boys with attention deficit hyperactivity disorder (ADHD): Performance-based test and parents report. *Research in Developmental Disabilities*, 33, 858-865.

- Siu, A. F. Y., & Zhou, Y. (2014). Behavioral assessment of the dysexecutive syndrome for children: An examination of clinical utility for children with attention-deficit hyperactivity disorder (ADHD). *Journal of Child Neurology*, 29, 608-616. doi:10.1177/0883073813516191
- Smidts, D. P., & Huizinga, M. (2009). *Handleiding Executieve Functies Gedragsvragenlijst* [Dutch user's manual of the Behavior Rating Inventory of Executive Function]. Amsterdam, The Netherlands: Hogrefe.
- Song, Y., & Hakoda, Y. (2014). Executive and non-executive functions in attention deficit hyperactivity disorder of the inattentive type (ADHD-I): A cognitive profile. *Journal of Behavioral and Brain Science*, 4, 1-10.
- Sonuga-Barke, E. J. S. (2003). The dual pathway model of AD/HD: An elaboration of neuro-developmental characteristics. *Neuroscience & Biobehavioral Reviews*, 27, 593-604.
- Sonuga-Barke, E. J. S., Bitsakou, P., & Thompson, M. (2010). Beyond the dual pathway model: Evidence for the dissociation of timing, inhibitory, and delay-related impairments in attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 49, 345-355.
- Sonuga-Barke, E. J. S., Dalen, L., Daley, D., & Remington, B. (2002). Are planning, working memory, and inhibition associated with individual differences in preschool ADHD symptoms? *Developmental Neuropsychology*, 21, 255-272.
- Spear, H. J., & Kulbok, P. (2004). Autonomy and adolescence: A concept analysis. *Public Health Nursing*, 21, 144-152.
- Stevens, J. (1996). *Applied multivariate statistics for the social sciences*. Hillsdale, NJ: Lawrence Erlbaum.
- Toplak, M. E., Bucciarelli, S. M., Jain, U., & Tannock, R. (2009). Executive functions: Performance-based measures and the Behavior Rating Inventory of Executive Function (BRIEF) in adolescents with attention deficit/hyperactivity disorder (ADHD). *Child Neuropsychology*, 15, 53-72.
- Toplak, M. E., West, R. F., & Stanovich, K. E. (2013). Do performance-based measures and ratings of executive function assess the same construct? *Journal of Child Psychology and Psychiatry*, 54, 131-143.
- Welsh, M. C., & Pennington, B. F. (1988). Assessing frontal lobe functioning in children: Views from developmental psychology. *Developmental Neuropsychology*, 4, 199-230.
- Welsh, M. C., Satterlee-Cartmell, T., & Stine, M. (1999). Towers of Hanoi and London: Contribution of working memory and inhibition to performance. *Brain and Cognition*, 41, 231-242.
- Willcutt, E. G., Doyle, A. E., Nigg, J. T., Faraone, S. V., & Pennington, B. F. (2005). Validity of the executive function theory of attention deficit/hyperactivity disorder: A meta-analytic review. *Biological Psychiatry*, 57, 1336-1346.
- Wilson, B. A., Alderman, N., Burgess, P. W., Emslie, H., & Evans, J. J. (1996). *Behavioral Assessment of Dysexecutive Syndrome*. St. Edmunds, UK: Thames Valley Test Company.
- Wilson, B. A., Alderman, N., Burgess, P. W., Emslie, H., & Evans, J. J. (2003). Behavioral Assessment of the Dysexecutive Syndrome (BADS). *Journal of Occupational Psychology, Employment and Disability*, 5(2), 33-37.
- Wodka, E. L., Loftis, C., Mostofsky, S. H., Prahme, C., Gidley Larson, J. C., Denckla, M. B., & Mahone, E. M. (2008). Prediction of ADHD in boys and girls using the D-KEFS. *Archives of Clinical Neuropsychology*, 23, 283-293.
- Wodka, E. L., Mostofsky, S. H., Prahme, C., Gidley Larson, J. C., Loftis, C., Denckla, M. B., & Mahone, E. M. (2008). Process examination of executive function in ADHD: Sex and subtype effects. *The Clinical Neuropsychologist*, 22, 826-841.
- Wolraich, M. L., Wibbelsman, C. J., Brown, T. E., Evans, S. W., Gotlieb, E. M., Knight, J. R., . . . Wilens, T. (2005). Attention-deficit/hyperactivity disorder among adolescents: A review of the diagnosis, treatment, and clinical implications. *Pediatrics*, 115, 1734-1746.

Author Biographies

Bianca E. Boyer is a neuropsychologist, working at the Developmental Psychology department of the University of Amsterdam. In 2009, she started her PhD project regarding the effectiveness of cognitive behavioral treatment for adolescents with ADHD 12 to 17 years old. Before she started this project, she worked as a clinician at several outpatient clinics, diagnosing and treating children and adolescents with ADHD and Autism Spectrum Disorders (ASD).

Hilde M. Geurts is currently a professor (focus Autism: Cognition across the life span) at the Department of Brain and Cognition of the University of Amsterdam. Also, she works as a senior researcher and psychologist in the autism expert team of the "Dr. Leo Kannerhuis." The main focus of her work is on ASD and ADHD, two of the major neurodevelopmental disorders. The main question of her work is whether there are dissociations between these neurodevelopmental disorders and how we can account for the overlap. She focuses on cognitive control, intra-individual variability, and reinforcement learning across the life span.

Saskia van der Oord works as an assistant professor at the Department of Clinical Psychology of the University of Leuven. Furthermore, she is a part-time assistant professor at the Department of Developmental Psychology of the University of Amsterdam. Her research is focused on childhood psychopathology. Areas of interest are diagnosis and treatment of disruptive behavior disorders, such as ADHD. Another research field is the treatment of posttraumatic stress disorder (PTSD) in children. She is a licensed behavior therapist and has clinical experience in (cognitive) behavioral treatment for children and adolescents with psychopathology such as ADHD.