UvA-DARE (Digital Academic Repository)

The Indonesian sponge fauna: a status report

van Soest, R.W.M.

DOI

[10.1016/0077-7579\(89\)90016-1](https://doi.org/10.1016/0077-7579(89)90016-1)

Publication date

1989

Published in

Netherlands Journal of Sea Research

[Link to publication](#)

Citation for published version (APA):

van Soest, R. W. M. (1989). The Indonesian sponge fauna: a status report. *Netherlands Journal of Sea Research*, 23(2), 223-230. [https://doi.org/10.1016/0077-7579\(89\)90016-1](https://doi.org/10.1016/0077-7579(89)90016-1)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

THE INDONESIAN SPONGE FAUNA: A STATUS REPORT

R.W.M. VAN SOEST

Institute of Taxonomic Zoology, University of Amsterdam, P.O. Box 4766, 1009 AT Amsterdam, The Netherlands

ABSTRACT

Snellius-II sponge collecting yielded 355 species (together about 1300 specimens). An illustrated simplified key to the more common reef forms is presented using only field characters. A preliminary taxonomic review of the Indonesian sponge fauna is given based on an analysis of material collected during the Snellius-II Expedition, of older material of the Siboga Expedition (so far largely unpublished), and of data in the literature (most recent dating from before 1935).

The Indonesian sponge fauna, which is provisionally estimated to comprise 830 nominal species (obvious synonyms not counted) is compared to the fauna of relatively well-studied areas elsewhere in the Indo-Pacific region and to the West Indian sponge fauna, using numbers of species of each of 188 genera occurring in more than one of the areas concerned. Based on published reports the Australian tropical sponge fauna appears to be dissimilar to that of Indonesia.

1. INTRODUCTION

In a previous, preliminary report on the Snellius-II sponge project (VAN SOEST, in press) attention was focussed on the common reef sponges of eastern Indonesia. A total of 56 species was indicated as common in the area and their depth distribution was presented. Since then all collected sponge specimens were examined and this has resulted in a revision of this list. It is extended to comprise 59 species and some nomenclatorial corrections appeared necessary. In this final report the definitive list of common reef sponges is presented in the form of an illustrated key, which will enable provisional identification by non-specialists. Besides, the Snellius-II results were added to the information on Indonesian sponges already published or otherwise available to me (viz. identified collections of the Siboga Expedition in the 'Zoological Museum' Amsterdam). The combined in-

formation permitted a review of the Indonesian Demospongiae fauna (90% of the sponges belong in this group), which is also presented in this paper.

Despite the preliminary status of our knowledge of Indonesian demosponges, this fauna is compared to that of other tropical regions, in order to assess similarities (indicating shared histories) and differences (separate histories) between the Indonesian region and those other regions. Some incidental results of the Snellius-II sponge project were already published elsewhere (VAN SOEST, in press; VAN SOEST & VERSEVELDT, 1987).

Acknowledgments.—Dr J. van der Land enabled me to join the first cruise of theme 4. Drs Mario de Kluijver executed the statistic comparison of tropical sponge faunas using the University of Amsterdam Cyber computer.

2. COMMON REEF SPONGES OF EASTERN INDONESIA

Below, an illustrated key is presented of sponges found to occur commonly on eastern Indonesian reefs. As was explained elsewhere (VAN SOEST, in press), the list is based on the first exploratory cruise of theme 4, in which 8 localities were visited. The nomenclature of a previously published list of these sponges was corrected in several instances: *Hymeniacidon conulosa* is now *H. massa*, *Psammaplysilla purpurea* is now *Druinella p.*, *Haliclona cf. nematifera* is merged with *Haliclona violacea* (*Nara nematifera* is different), *Axinella carteri* is now *Acanthella c.*, *Haliclona cf. turquoisia* is *H. fascigera*, *Callyspongia confederata* is *C. schulzei*. Three species, *Spirastrella decumbens*, *Cliona spec.* and *Ircinia irregularis* turned out to be not as common as was previously assumed, and consequently are removed from the list. Seven others appeared to fall within the criterion of commonness (common in three or more of the localities visited) and are thus added to the list: *Dysidea arenaria* (depth range 4 - > 15 m), *Biemna fortis*

(0 - 4 m), *Paratetilla bacca* (4 - > 15 m), *Iotrochota baculifera* (0 - > 15 m), *Clathria fasciculata* (4 - 15 m), *Haliclona pigmentifera* (0 - 4 m), *Chondrilla australiensis* (0 - 4 m). As a result the list is now extended to 59 species.

Nomenclature still remains problematical because

not all species have been checked against type specimens yet. The revision of Indo-Pacific type specimens of common sponges is difficult since these are distributed over a large number of institutions and museums, all over the world.

Figs 1 and 2. Sketch drawings of characteristic habits of 59 species of common sponges, keyed out in the text.

Simplified illustrated key to the common reef sponges of eastern Indonesia:

The key is intended as a quick field guide for non-specialists, omitting all scientific terms and descriptions of internal anatomy (which is indispensable for a definite identification). The chosen species (choice based upon their relatively common occurrence in three or more of the eight areas visited) are subdivided on their characteristic growth form into nine groups. Within these groups some further characteristics are provided which permit provisional identification. All species are represented by sketch drawings in Figs 1-2.

A. Tubes or pipes:

1. blue-green-grey, with sharp conules, resilient *Callyspongia schulzei* (Kieschnick, 1900)
2. purple, quite smooth, fragile, easily torn *Haliclona fascigera* (Hentschel, 1912)
3. grey, muddy appearance, undulating surface with network of fine grooves, spongy consistency
Liosina paradoxa Thiele, 1899
4. purple, indented-grooved surface, stony hard *Theonella conica* (Kieschnick, 1896)
5. brown-red (white interior), smooth in small, grooved-indented in larger specimens, compressible but fragile *Theonella swinhoei* Sollas, 1888
6. red-orange, indented or holed, spongy-compressible but tough *Clathria basilana* Lévi, 1951
7. grey-blue, surface pilose- conulose, soft *Niphates olemda* De Laubenfels, 1954

B. Cups and tubs:

8. often giant sized, red-grey, with heavily ridged outer walls, stony hard
Petrosia (Xestospongia) testudinaria (Lamarck, 1815)
9. cups or fans, thin-walled, mauve-grey, with characteristic grooved pattern on inside surface, tough *Phyllospongia foliascens* (Pallas, 1766)
10. cups or fans, thin-walled, grey, regularly spaced star-shaped oscules on inside surface, tough . *Phyllospongia papyracea* (Esper, 1794)
11. globular, with deep vent, red-brown, with widely spaced low rounded conules, hard-incompressible ...
Stelletta globostellata (Carter, 1883)
12. irregular tubs or laminated masses, smooth surface, dark brown, stony hard
Petrosia nigricans Lindgren, 1897
13. stalked cup with narrow deep vent, reddish light brown, with rounded low conules, tough-compressible
Asteropus sarassinorum (Thiele, 1899)

C. Fan-shaped:

14. irregular, with sharp ridges and conules, orange-red with yellow-orange sheen, rubbery
Acanthella carteri (Dendy, 1887)
15. irregular, often sending out thin branches, smooth, bright bluish purple, firm but easily damaged
Haliclona amboinensis Lévi, 1961
16. sharply conulose, laminated, red-orange, fragile, easily torn *Dysidea cinerea* (Keller, 1881)
17. thin-walled, finely conulose, yellow-grey-purple, very soft *Dysidea herbacea* (Keller, 1881)
18. irregular, roughly and heavily conulose, slate-grey, sandy-fragile ... *Dysidea arenaria* Bergquist, 1965
19. thickly, even massively laminated, with holes and pits, bright orange, spongy
Agelas mauritiana Carter, 1882

D. Massive structures:

20. thick masses with deeply grooved surface, yellow-orange, tough *Epipolasis suluensis* (Wilson, 1925)
21. wall-shaped, smooth, bright yellow-orange, soft, easily torn *Hymeniacidon massa* (Carter, 1886)
22. tower-shaped, with side-branches, part-buried in the sandy substrate, oscules concentrated topside, where colour is darker than the pale-beige shaft, tough-hard *Spirastrella vagabunda* Ridley, 1884
23. tower-shaped masses similar to previous species, but with corrugated upper parts, conulose sides, compressible, fragile *Biemna fortis* (Topsent, 1897)
24. large masses with reticulate surface, dark brown to black, hard but fibrous
Fascaplysinopsis reticulata (Hentschel, 1912)

E. Globular or ball-shaped:

25. rounded clathrate-cavernous masses consisting of intertwined trabeculae surrounding rounded cavities, bright orange *Acanthella cavernosa* Dendy, 1922

26. ball-shaped with regularly spaced rounded crater-like pits between which the surface is hispid, bright yellow *Paratetilla bacca* (Selenka, 1864)
27. ball-shaped, similar to previous species, but more squat and yellow colour obscured by sediment; crater-like pits smaller and more numerous *Cinachyra australiensis* (Carter, 1886)
28. small, berry-shaped, olive-brown *Stelletta clavosa* (Ridley, 1884)
29. masses of globular osculiferous lobes, orange-brown, rough to the touch but compressible; turns black in alcohol *Aaptos cf. suberitoides* Brondsted, 1934
30. ball-shaped, covered with rounded tubercles, red-green-orange ... *Tethya robusta* (Bowerbank, 1875)

F. Fistulose:

31. main body buried in the sandy substrate, golden brown parchment-like fistules
Coelocarteria singaporense (Carter, 1883)
32. turnip-shaped main body, long smooth fistules with rounded closed ends, pale wine-red, tough but fragile
Oceanapia amboinensis (Topsent, 1897)

G. Ramose:

33. very spinous, fibrous branches, erect or creeping, longer spines are themselves spined, light purple *Gelliodes fibulatus* (Ridley, 1884)
34. irregular undulating-conulose surface, often covered by sediment, salmon-coloured, compressible
Clathria reinwardti Vosmaer, 1882
35. flattened branches, irregularly conulose, conules blunt, erect-creeping, light brown, very toughly spongy
Dactylospongia elegans (Thiele, 1899)
36. flattened branches, regularly conulose, dark olive-brown, toughly spongy *Ircinia ramosa* (Keller, 1881)
37. flattened branches, regularly conulose, black, toughly spongy *Hyrtios erectus* (Keller, 1881)
38. thin, rounded branches, finely conulose, purple, spongy *Dysidea granulosa* Bergquist, 1965
39. limp, soft branches rising from a broadly incrusting base, star-shaped oscules, furry black, soft
Acervochalina confusa Dendy, 1905
40. intertwined spinous branches, purple-grey, toughly spongy *Callyspongia joubini* (Topsent, 1897)
41. smooth-lumpy branches, mottled wine-red-white, spongy, compressible
Agelas "ceylonica" (Dendy, 1905)
42. erect slender-branched bushes, brownish green-red, cartilaginous *Haliclona cymaeiformis* (Esper, 1794)

H. Massive-creeping with upright branches:

43. regularly conulose, conules widely spaced and blunt, surface slippery, green (turns dark purple in alcohol), compressible *Druinella purpurea* (Carter, 1880)
44. surface ornamented with system of angular grooves, brown-orange-white, stony hard
Placospongia melobesioides Gray, 1867
45. arm-thick creeping branches with thick-walled oscular chimneys, red-brown, stony hard
Petrosia contignata Thiele, 1899
46. mats covering the substrate with irregular thin upright branches, bright green, cartilaginous
Halichondria cartilaginea (Esper, 1794)
47. very spinous creeping masses-branches, large oscules, orange-red, toughly spongy
Clathria fasciculata (Wilson, 1925)
48. spinous masses issuing thick branches, black with dark purple exudate when squeezed, toughly spongy
Iotrochota baculifera Ridley, 1884
49. creeping masses with variable presence of low conules, large oscules, purple, spongy-resilient
Gelliodes pumila (Von Lendenfeld, 1887)
50. massively incrusting with stiff erect branches or laminations, smooth, dark brown, tough-crumblly
Xestospongia exigua (Kirkpatrick, 1900)

I. Massively incrusting:

51. irregular, smooth masses, with flaky surface, black, fragile-brittle
Xestospongia cf. carbonaria (Lamarck, 1814)

[illegible]

ably representative collection of Indonesian sponges, collected from shallow reef flats, mangroves and sea grass meadows, coral reefs, continental slopes and abyssal localities. The data of the two large collections were combined with all data published on Indonesian sponges (HARTING, 1870; KIESCHNICK, 1896, 1898, 1900; TOPSENT, 1897 (recently redescribed by DESQUEYROUX, 1981); LINDGREN, 1897; SCHULZ, 1898; THIELE, 1899, 1903a, 1903b; HENTSCHEL, 1912; BRØNDSTED, 1934; many incidental records in different expedition reports, monographs, and notes). After elimination of all obvious synonyms, a preliminary checklist of Indonesian Demospongiae was constructed, omitting Hexactinellids (nothing substantial has been described since Ijima's excellent monograph) and Calcarea (at present very difficult to deal with pending generic and familial revisions). The checklist comprises about 780 nominal species; another 50 are assumed to be undescribed; in all, 43% of the listed sponges are endemic to Indonesia. The checklist is available on request.

The revision of this formidable fauna of over 800 species represents a major challenge to sponge taxonomy, with considerable bearing upon and importance for the taxonomy of sponges in general. Such a challenge cannot be met with by single museum taxonomists working in isolation; it is imperative that teams of scientists are formed consisting of experienced senior taxonomists working in close cooperation with local scientists having access to the field.

6. COMPARISON OF THE INDONESIAN DEMOSPONGIAE FAUNA WITH OTHER TROPICAL REGIONS

Although the checklist of Indonesian demosponges is preliminary and many future emendations are to be expected, it is now possible to compare this list with (often equally preliminary) faunal lists from other regions of the tropical oceans (more or less comprehensively extracted from a large amount of literature), in order to find out to which areas Indonesia is most (dis)similar. The general biogeographic trend (BRIGGS, 1974) for major benthic taxa is that Indonesia is the central and richest part of a larger Indo-West Pacific region, which includes the north and central Pacific, the north and east coast of Australia, the central and western Indian Ocean, and the Red Sea. All these areas are reported to have some endemic sponge species (varying between 25 and 45% of the species reported from them in the literature); in the complement of their common sponges they are all dissimilar pointing at large local differences (VAN SOEST, in press). Endemism at the generic level, however, is quite low.

Unfortunately the description of the species,

Fig. 3. Dendrogram of sponge faunas of eight tropical areas. Values indicate dissimilarity (0 = completely similar, 1 = completely dissimilar) as determined by Bray-Curtis dissimilarity index. (INDO = Indonesia, CIND = Central Indian Ocean, WIND = Western Indian Ocean, WI = West Indian region, NPAC = North Pacific, REDS = Red Sea, CAPC = Central Pacific, GBR = North and East Australia).

especially in the older literature, is often too vague and type specimens have been so seldom re-examined that a comparison of lists of actual species seems a meaningless and unreliable effort. However, it is possible at this stage of our knowledge, and with some editing of the generic assignments made by the different authors, to compare the numbers of species of each genus (regardless of their identities) reported from more than one of the areas concerned. This method also makes it possible to draw the West Indian region into the comparison. A matrix table was constructed of 188 genera (the total of all tropical sponge genera diminished for practical reasons by the number of genera occurring in only a single of the eight areas considered) and their numbers of species in the eight areas, expressed as percentage of the total number of species recorded for each area. Similarities of the areas considered were compared using the Bray-Curtis dissimilarity index in combination with Average Linkage clustering procedure (see KAANDORP, 1986). The processing of the data yielded the dendrogram of Fig. 3, showing that the Indonesian fauna is most similar to Indian Ocean faunas and also to that of the West Indies (although endemism in the latter area is close to 100%), and the neighbouring north Pacific. Such similarities are easily explained by common history of the tropical seas dating back to the Tethys. More peripheral

areas such as the central Pacific and the Red Sea are increasingly dissimilar, which is in accordance with the general biogeographic trend. Remarkably, however, the Australian area (Great Barrier Reef) is highly dissimilar to Indonesia. It is possible that we are again dealing with a sampling bias, because the published accounts of the Great Barrier Reef sponge fauna probably do not nearly cover what really occurs there.

On the other hand, there is room for the hypothesis that the Australian region has a distinct element of its own, no doubt dating back from the time that Australia was not in contact with the tropical Asian fauna (KNOX, 1980). The south Australian subtropical/temperate fauna shows a high level of generic endemism with many peculiar forms, e.g. the 'sand sponges'. Possibly, invasions of tropical (reef-) sponges from the Asian tropical region into the NE Australian region has not so completely replaced local Australian elements as was apparently the case in other animal groups.

7. REFERENCES

- BRIGGS, J.C., 1974. Marine Zoogeography. McGraw-Hill, New York: 1-475.
- BRØNDSTED, H.V., 1934. Sponges.—Résult. scient. Voyage Indes orient. néerl. **2** (15): 1-27.
- BURTON, M., 1930. The Porifera of the Siboga Expedition III. Calcareous.—Monogr. Siboga Exp. **6** (a 2): 1-18.
- DESQUEYROUX, R., 1981. Révision de la collection d'éponges d'Amboine (Moluques, Indonésie) constituée par Bedot et Pictet et conservée au Muséum d'Histoire Naturelle de Genève.—Rev. suisse Zool. **88**: 723-764.
- HARTING, P., 1870. Mémoire sur le genre Poterion.—Natuurk. Verhandl. Prov. Utrechts Genootsch. Kunst. Wetensch. N.S. **2** (2): 1-40.
- HENTSCHEL, E., 1912. Kiesel- und Hornschwämme der Aru- und Kei-Inseln.—Abh. Senckenb. naturforsch. Ges. **34**: 293-448.
- IJIMA, I., 1927. The Hexactinellida of the Siboga Expedition.—Monogr. Siboga Exp. **6** (106): 1-383.
- KAANDORP, J.A., 1986. Rocky substrate communities of the infralittoral fringe of the Boulonnais coast, N.W. France: a quantitative survey.—Mar. Biol. **92**: 255-265.
- KIESCHNICK, O., 1896. Silicispongiae von Ternate.—Zool. Anz. **19**: 526-534.
- KIESCHNICK, O., 1898. Die Kiesel Schwämme von Amboina. Inaug. Diss. Jena: 1-66.
- , 1900. Die Kiesel Schwämme von Amboina.—Denkschr. mediz. naturwiss. Ges. Jena **8**: 545-582.
- KNOX, G.A., 1980. Plate tectonics and the evolution of intertidal and shallow-water benthic biotic distribution patterns of the South West Pacific.—Palaeogeogr. Palaeoclim. Palaeoecol. **31**: 267-297.
- LINDGREN, N.G., 1897. Beitrag zur Kenntniss der Spongien-Fauna des Malaiischen Archipels und des Chinesischen Meeres.—Zool. Anz. **20**: 480-487.
- SCHULZ, E., 1898. Hornschwämme von Ternate.—Abh. Senckenb. naturforsch. Ges. **24**: 185-188.
- SOEST, R.W.M. VAN, in press. Shallow-water reef sponges of Eastern Indonesia.—Proc. 3rd Int. Conf. Biol. Sponges, Woods Hole.
- SOEST, R.W.M. VAN & J. VERSEVELDT, 1987. Unique symbiotic octocoral- sponge association from Komodo.—Indo-Malayan Zool. **4**: 27-32.
- THIELE, M.E., 1899. Studien über Pazifische Spongien, II. Über einige Spongien von Celebes.—Zoologica, Stuttg. **24**: 1-33.
- , 1903a. Kiesel Schwämme von Ternate, I.—Abh. Senckenb. naturforsch. Ges. **25**: 19-80.
- , 1903b. Kiesel Schwämme von Ternate, II.—Abh. Senckenb. naturforsch. Ges. **25**: 933-968.
- TOPSENT, E., 1897. Spongiaires de la baie d'Amboine.—Rev. suisse Zool. **4**: 421-487.
- VOSMAER, G.C.J., 1911. The Porifera of the Siboga Expedition II. The genus *Spirastrella*.—Monogr. Siboga Exp. **6** (a1): 1-69.
- VOSMAER, G.C.J. & J.H. VERNHOUT, 1902. The Porifera of the Siboga Expedition I. The genus *Placospongia*.—Monogr. Siboga Exp. **6** (a): 1-17.