



**UvA-DARE (Digital Academic Repository)**

**Charged-hadron inclusive cross sections and fractions in  $e^+ e^-$  annihilation at  $\sqrt{s} = 29\text{-GeV}$**

Aihara, H.; Alston-Garnjost, M.; Avery, R.E.; Barbaro-Galtieri, A.; Barker, A.R.; Barnett, B.A.; Bauer, D.A.; Bay, A.; Linde, F.L.

*Published in:*  
Physical Review Letters

*DOI:*  
[10.1103/PhysRevLett.61.1263](https://doi.org/10.1103/PhysRevLett.61.1263)

[Link to publication](#)

*Citation for published version (APA):*

Aihara, H., Alston-Garnjost, M., Avery, R. E., Barbaro-Galtieri, A., Barker, A. R., Barnett, B. A., ... Linde, F. L. (1988). Charged-hadron inclusive cross sections and fractions in  $e^+ e^-$  annihilation at  $\sqrt{s} = 29\text{-GeV}$ . *Physical Review Letters*, 61(11), 1263-1266. DOI: 10.1103/PhysRevLett.61.1263

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

## Charged-Hadron Inclusive Cross Sections and Fractions in $e^+e^-$ Annihilation at $\sqrt{s}=29$ GeV

H. Aihara, M. Alston-Garnjost, R. E. Avery, A. Barbaro-Galtieri, A. R. Barker, B. A. Barnett, D. A. Bauer, A. Bay, G. J. Bobbink, C. D. Buchanan, A. Buijs, D. O. Caldwell, H.-Y. Chao, S.-B. Chun, A. R. Clark, G. D. Cowan, D. A. Crane, O. I. Dahl, M. Daoudi, K. A. Derby, J. J. Eastman, P. H. Eberhard, T. K. Edberg, A. M. Eisner, R. Enomoto, F. C. Ern , K. H. Fairfield, J. M. Hauptman, W. Hofmann, J. Hylen, T. Kamae, H. S. Kaye, R. W. Kenney, S. Khacheryan, R. R. Kofler, W. G. J. Langeveld, J. G. Layter, W. T. Lin, F. L. Linde, S. C. Loken, A. Lu, G. R. Lynch, R. J. Madaras, B. D. Magnuson, G. E. Masek, L. G. Mathis, J. A. J. Matthews, S. J. Maxfield, E. S. Miller, W. Moses, D. R. Nygren, P. J. Oddone, H. P. Paar, S. K. Park, D. E. Pellett, M. Pripstein, M. T. Ronan, R. R. Ross, F. R. Rouse, K. A. Schwitkis, J. C. Sens, G. Shapiro, B. C. Shen, J. R. Smith, J. S. Steinman, R. W. Stephens, M. L. Stevenson, D. H. Stork, M. G. Strauss, M. K. Sullivan, T. Takahashi, S. Toutouchi, R. van Tyen, W. Vernon, W. Wagner, E. M. Wang, Y.-X. Wang, W. A. Wenzel, Z. R. Wolf, H. Yamamoto, S. J. Yellin, and C. Zeitlin

(TPC/Two-Gamma Collaboration)

*Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720*

*University of California at Davis, Davis, California 95616*

*University of California Institute for Research at Particle Accelerators, Stanford, California 94305*

*University of California at Los Angeles, Los Angeles, California 90024*

*University of California at Riverside, Riverside, California 92521*

*University of California at San Diego, San Diego, California 92093*

*University of California at Santa Barbara, Santa Barbara, California 93106*

*Carnegie-Mellon University, Pittsburgh, Pennsylvania 15213*

*Ames Laboratory, Iowa State University, Ames, Iowa 50011*

*Johns Hopkins University, Baltimore, Maryland 21218*

*University of Massachusetts, Amherst, Massachusetts 01003*

*New York University, New York, New York 10003*

*National Institute for Nuclear and High Energy Physics, Amsterdam, The Netherlands*

*University of Tokyo, Tokyo, Japan*

(Received 9 May 1988)

We report measurements of  $\pi^\pm$ ,  $K^\pm$ , and  $p,\bar{p}$  inclusive cross sections and fractions in  $e^+e^-$  annihilation at  $\sqrt{s}=29$  GeV, for the momentum interval  $0.01 < z = p/p_{\text{beam}} < 0.90$ . The analysis is based on approximately  $70 \text{ pb}^{-1}$  of data collected with the TPC/ $2\gamma$  detector facility at the SLAC storage ring PEP. Detector upgrades result in significantly improved momentum coverage and precision of the data, compared to previous measurements.

PACS numbers: 13.65.+i, 13.87.Fh

Inclusive hadron cross sections and charged-hadron fractions in  $e^+e^-$  annihilation are fundamental quantities providing information about quark fragmentation. In this paper we present measurements of these quantities with substantially improved accuracy and extending to higher  $z$ , where  $z \equiv p_{\text{hadron}}/p_{\text{beam}}$ , than previously reported.<sup>1-4</sup> The particle spectra in the high- $z$  region provide a sensitive probe of the hadron production mechanism, especially for the case of baryons. For example, dimensional-counting arguments<sup>5</sup> predict that the baryon to meson ratio should fall as  $(1-z)^1$  or  $(1-z)^2$  for  $z \rightarrow 1$ , whereas the Lund string model,<sup>6</sup> e.g., predicts a rising baryon fraction. The potential to discriminate between competing hadron production models is enhanced at high  $z$  by the fact that a comparatively large fraction of the particles in this region are produced

directly in the hadronization process, rather than as the result of resonance decays.

We have measured the  $\pi^\pm$ ,  $K^\pm$ , and  $p,\bar{p}$  inclusive cross sections and fractions in the interval  $0.01 < z < 0.90$  with the TPC/ $2\gamma$  detector at the PEP  $e^+e^-$  storage ring at SLAC, operating at  $\sqrt{s}=29$  GeV. The TPC/ $2\gamma$  facility has been described in detail in Ref. 7. The present analysis is based on approximately  $70 \text{ pb}^{-1}$  of data collected between 1984 and 1986, and uses only charged-particle information from the time projection chamber (TPC) itself. The detector configuration during this period included a 13.25-kG superconducting magnet and a gating-grid system to reduce track distortions, resulting in a momentum resolution of typically  $(\Delta p/p)^2 = (1.5\%)^2 + (0.65\% p)^2$  ( $p$  in GeV/ $c$ ).

Charged particles are identified by a simultaneous

measurement of momentum and ionization energy loss ( $dE/dx$ ). A charged particle passing through the TPC's gas volume (80% Ar and 20% CH<sub>4</sub> at a pressure of 8.5 atm) produces several hundred ionization electrons per cm. These electrons are detected by multiwire proportional chambers (sectors). Up to 183  $dE/dx$  measurements are obtained for each track. These  $dE/dx$  values are corrected for electron capture and diffusion along the drift path, variations in wire gain over the surface of a sector, variations between sectors, polar-angle dependence (resulting from dependence on the length of track sampled by each sense wire), and gain variations due to changes in gas density and composition. The most probable  $dE/dx$  is estimated by forming the mean of the lowest 65% of the usable measurements. In the following, this quantity is referred to as the truncated mean  $dE/dx$ , or simply the " $dE/dx$ " of a particle. For typical tracks with 120  $dE/dx$  measurements a resolution for the truncated mean  $dE/dx$  of approximately 3.5% is obtained.

Much of the momentum range studied in this analysis corresponds to the relativistic-rise region of the  $dE/dx$  versus velocity curve. In this region the separation between particle species is at most 3.5 standard deviations. Hence, the particle identity cannot be established on a track-by-track basis and the relative abundances must be determined by a statistical analysis. This procedure requires an accurate knowledge of both the expected  $dE/dx$  as a function of the particle's velocity and the  $dE/dx$  resolution.

An atomic physics model for ionization energy loss was used to predict the expected  $dE/dx$  as a function of the particle's speed.<sup>7-9</sup> The model prediction was scaled to fit the average  $dE/dx$  measured for samples of well-identified particles, such as protons from the  $1/\beta^2$  region, pions from the minimum-ionizing and relativistic-rise regions, cosmic-ray muons, conversion electrons, and Bhabha electrons. The protons, pions, and conversion electrons were taken from multihadron events. The data points and resulting fit are shown in Fig. 1(a). The systematic uncertainty in the  $dE/dx$  versus velocity relation is estimated to 0.2% [determined from systematic differences between different data sets, such as pions in hadronic events and cosmic-ray muons; see inset in Fig. 1(a)].

The resolution for the truncated mean,  $\sigma_{dE/dx}$ , was determined from a sample of approximately 47000 minimum-ionizing pions in multihadron events. The resolution is mainly a function of the number  $m$  of usable  $dE/dx$  measurements and decreases as  $1/\sqrt{m}$ , with a floor of about 1.8%. The  $dE/dx$  resolution function is Gaussian out to 3 standard deviations [Fig. 1(b)]. Studies with cosmic-ray muons indicate that the relative resolution  $\sigma_{dE/dx}/(dE/dx)$  remains approximately independent of the mean  $dE/dx$  of the track sample for the relativistic-rise region. For the  $1/\beta^2$  region, the relative resolution was observed to improve somewhat with in-




FIG. 1. (a) Average truncated mean  $dE/dx$  as a function of  $\beta\gamma = p/m$  of a particle, as determined from protons, pions, and conversion electrons in hadronic events, from cosmic-ray muons, and from Bhabha electrons. Line: (adjusted) atomic physics model. Inset: ratio  $R$  of data points and model fit in the relativistic-rise region. The dashed lines indicate the 0.2% systematic uncertainty. (b) Distribution in difference between measured and predicted  $dE/dx$  for minimum-ionizing pions, normalized to the  $dE/dx$  resolution for each track. The curve represents a normal distribution ( $\sigma = 1$ ).

creasing mean  $dE/dx$ . [Empirically,  $\sigma_{dE/dx}/(dE/dx) \propto (dE/dx)^{-0.35}$ .] The fractional systematic uncertainty in the  $dE/dx$  resolution is estimated to be 8%. A more detailed discussion of the  $dE/dx$  analysis can be found in Refs. 7 and 9.

The selection of multihadron events is described in Ref. 3. For this analysis, it was further required that the angle between the sphericity axis and the beam line be larger than  $45^\circ$ , resulting in a final sample of 20126 events. Tracks in these events were required to have at least forty  $dE/dx$  measurements for determination of the truncated mean, an error in the measurement of the momentum component transverse to the beam of  $\Delta p_T/p_T < 0.15$  or  $\Delta p_T/p_T^2 < 0.15 \text{ GeV}^{-1}$ , and a polar angle

larger than  $30^\circ$ , and to extrapolate back to within 3 cm of the event vertex in the bend plane and to within 5 cm along the direction of the beam. At low momenta, a substantial fraction of the observed protons come from nuclear interactions in the material between the interaction point and the TPC. Therefore, only negative particles are used for  $z < 0.25$ .

For determination of the hadron fractions and cross sections, the track sample is divided into narrow momentum intervals. For each interval, the number of particles  $\phi_i$  of each type ( $i = e, \pi, K, \text{ and } p$ ) is determined by an extended maximum-likelihood fit<sup>10</sup> to the  $dE/dx$  distribution (Fig. 2). The fit does not require binning in  $dE/dx$  and makes maximal use of the information available for each track. The likelihood function is given by

$$L = \exp\left(-\sum_i \phi_i\right) \prod_j \left[ \sum_i \frac{\phi_i}{\sqrt{2\pi}\sigma_{ij}} \exp\left(-\frac{[R_j - \mu_i(p_j)]^2}{2\sigma_{ij}^2}\right) \right],$$

where the index  $j$  runs over all tracks in the momentum interval, and  $i$  runs over the four particle species: electrons, pions, kaons, and protons.  $R_j$  is the measured  $dE/dx$  for track  $j$  divided by the predicted value for a pion at the measured momentum of the track;  $\sigma_{ij}$  is the resolution in  $R_j$  for particle hypothesis  $i$ , which is estimated from the number of  $dE/dx$  measurements, the polar angle, and the curvature error of the track;  $\mu_i(p_j)$  is the expected  $R$  value for a track of momentum  $p_j$  and particle species  $i$ . We do not attempt to separate pions from muons, but instead subtract the muon rate obtained in an independent measurement.<sup>11</sup>

The number of hadrons in each momentum interval was then corrected for effects of geometrical acceptance, nuclear interactions in the material between the interaction point and the TPC, decays in flight, event and track selection cuts, muon contamination of the pion sample, momentum smearing, initial-state radiation, and back-




FIG. 2. Distribution in  $dE/dx$  vs momentum for particles in multihadron events. Lines indicate the predicted average  $dE/dx$  as a function of momentum for different species.

ground from the reactions  $e^+e^- \rightarrow \tau^+\tau^-$  and  $e^+e^- \rightarrow e^+e^- + \text{hadrons}$ . The background corrections are small compared to the estimated errors in the fractions and cross sections. Protons are defined to include decay protons of weakly decaying baryons; pions include decay products of  $K_S^0$  but not those of  $K_L^0$ .

The inclusive cross sections  $(1/\sigma\beta)(d\sigma/dx)$ , with  $x = 2E/\sqrt{s}$  and  $\beta = p/E$ , are presented in Fig. 3, which also includes previously reported data by Aihara *et al.* (TPC Collaboration)<sup>3</sup> and Althoff *et al.* (TASSO Collaboration).<sup>4</sup> The cross sections are normalized to the total annihilation cross section into hadrons. Figure 4 shows the corresponding hadron fractions as a function of  $z \equiv p_{\text{hadron}}/p_{\text{beam}}$ . The errors shown in Figs. 3 and 4 correspond to the quadratic sum of systematic and statistical uncertainties. For the cross sections, statistical errors dominate above  $z \sim 0.5$ . The dominant sources of systematic error for the proton and kaon cross sections come from uncertainty in the  $dE/dx$  versus velocity curve and in the relative  $dE/dx$  resolution. The systematic error in the pion cross section is dominated by a 2% uncertainty in the pattern-recognition efficiency. For the particle fractions, many of the systematic uncertainties cancel to first order, and errors in the fractions are predominantly statistical.

In confirmation of earlier results,<sup>3,4</sup> the particle fractions at low  $z$  show a steep momentum dependence of the  $K^\pm$  and  $p, \bar{p}$  content of jets (Fig. 4). Above  $z \approx 0.3$ , the fractions level off. In particular, the  $p, \bar{p}$  fraction remains constant within errors for  $z > 0.2$ , in contrast with the decreasing baryon fraction predicted by counting rules for  $z \rightarrow 1$ ,<sup>5</sup> and the rising baryon fraction pre-




FIG. 3. Normalized cross sections  $(1/\sigma\beta)(d\sigma/dx)$  for  $\pi^\pm$ ,  $K^\pm$ , and  $p, \bar{p}$ , as a function of  $x = 2E/\sqrt{s}$ , including earlier TPC Collaboration (Ref. 3) and TASSO Collaboration (Ref. 4) data.




FIG. 4. Fractions of pions, kaons, and protons among charged hadrons in multihadron events, as a function of  $z = p/p_{\text{beam}}$ . Included are earlier TPC Collaboration (Ref. 3) and TASSO Collaboration (Ref. 4) data.

dicted by typical fragmentation models with a diquark mechanism for baryon production.<sup>6,12</sup> With use of their default parameters, none of the most commonly used fragmentation models<sup>6,12,13</sup> is consistent with our data. More details and comparisons with model predictions are given in Ref. 9.

The efforts of the SLAC PEP staff, and the engineers, programmers, and technicians who made this work possible are gratefully acknowledged. This work was supported in part by the U.S. Department of Energy, the Na-

tional Science Foundation, the Joint Japan-U.S. Collaboration in High Energy Physics, and the Foundation for Fundamental Research on Matter in The Netherlands.

<sup>1</sup>M. Derrick *et al.* (HRS Collaboration), Phys. Rev. D **35**, 2639 (1987).

<sup>2</sup>H. Schellman *et al.* (Mark II Collaboration), Phys. Rev. D **31**, 3013 (1985).

<sup>3</sup>H. Aihara *et al.* (TPC Collaboration), Phys. Rev. Lett. **52**, 577 (1984).

<sup>4</sup>M. Althoff *et al.* (TASSO Collaboration), Z. Phys. C **17**, 5 (1983), and contribution to the International Symposium on Lepton and Photon Interactions at High Energies, Kyoto, Japan, August 1985 (unpublished), paper No. 399.

<sup>5</sup>S. Brodsky and G. Farrar, Phys. Rev. Lett. **31**, 1153 (1973); V. A. Matveev *et al.*, Nuovo Cimento Lett. **1**, 719 (1973); S. Brodsky and J. Gunion, Phys. Rev. D **17**, 848 (1978).

<sup>6</sup>B. Andersson, G. Gustafson, G. Ingelman, and T. Sjöstrand, Phys. Rep. **97**, 31 (1983).

<sup>7</sup>H. Aihara *et al.* (TPC/ $2\gamma$  Collaboration), LBL Report No. LBL-23737, 1988 (to be published).

<sup>8</sup>W. Allison and J. Cobb, Ann. Rev. Nucl. Part. Sci. **30**, 253 (1980).

<sup>9</sup>G. D. Cowan, Ph.D. thesis, University of California, Berkeley, LBL Report No. LBL-24715, 1988 (unpublished).

<sup>10</sup>L. Lyons, *Statistics for Nuclear and Particle Physicists* (Cambridge Univ. Press, Cambridge, 1986), p. 100.

<sup>11</sup>H. Aihara *et al.* (TPC Collaboration), Phys. Rev. D **31**, 2719 (1985), and Z. Phys. C **27**, 39 (1985).

<sup>12</sup>T. D. Gottschalk and D. A. Morris, Nucl. Phys. **B288**, 927 (1987).

<sup>13</sup>B. R. Webber, Nucl. Phys. **B238**, 492 (1984).




FIG. 2. Distribution in  $dE/dx$  vs momentum for particles in multihadron events. Lines indicate the predicted average  $dE/dx$  as a function of momentum for different species.