UvA-DARE (Digital Academic Repository)

Currents and Ward-Takahashi Identities for Nonlocal Quantum-Field Theories

Bos, J.W.; Koch, J.H.; Naus, H.W.L.

DOI

[10.1103/PhysRevC.44.485](https://doi.org/10.1103/PhysRevC.44.485)

Publication date

1991

Published in

Physical Review C

[Link to publication](#)

Citation for published version (APA):

Bos, J. W., Koch, J. H., & Naus, H. W. L. (1991). Currents and Ward-Takahashi Identities for Nonlocal Quantum-Field Theories. *Physical Review C*, 44, 485-490.
<https://doi.org/10.1103/PhysRevC.44.485>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Currents and Ward-Takahashi identities for nonlocal quantum field theories

J. W. Bos

National Institute for Nuclear Physics and High Energy Physics, Section K (NIKHEF-K), P. O. Box 41882, NL-1009 DB Amsterdam, The Netherlands

J. H. Koch

*National Institute for Nuclear Physics and High Energy Physics, Section K (NIKHEF-K)
and Institute for Theoretical Physics, University of Amsterdam, 1018 XE Amsterdam, The Netherlands*

H. W. L. Naus

*Institute for Theoretical Physics, University of Erlangen-Nürnberg, W-8520 Erlangen, Federal Republic of Germany
(Received 26 February 1991)*

We consider a nonlocal quantum field theory, which can be used for the description of a composite particle, such as the nucleon interacting with its virtual pion field. Assuming a global symmetry of the theory, we discuss how to find the corresponding conserved currents. For the case of a U(1) symmetry, we then examine the conditions for second quantization and derive a general Ward-Takahashi identity. The formalism is then applied to the electromagnetic interaction.

I. INTRODUCTION

There has been a lot of interest in the composite nature of hadrons, and, more recently, in their modifications inside a nuclear medium. The most often discussed particle is the nucleon, and there have been several models to describe its internal structure. An important test for these models is to compare their predictions to the charge and magnetic form factors determined in electron scattering from a free proton. For two-step processes on a single nucleon, such as Compton scattering, or, in general, for nucleons not on their mass shell, more form factors are needed to describe the electromagnetic interaction of such an off-shell particle. This has been discussed extensively by Bincer [1] and recently studied in the context of a simple field theoretical model [2,3]. An important constraint on the electromagnetic vertex is the Ward-Takahashi identity, which relates vertex structure to the self-energy of the particle. This constraint, a consequence of gauge invariance, was derived originally for QED to all orders in the electromagnetic coupling constant e [4]. It was extended on the basis of a specific model for composite fields by Nishijima [5].

Rather than dealing explicitly with all the internal degrees of freedom of a composite particle, such as the virtual pion field for the nucleon, one can project them out and obtain an "effective"-field-theoretical description. In such a description, the Lagrangian density is nonlocal, i.e., contains derivatives of the particle field of arbitrary high order [6–8]. This makes it impossible to arrive at the conserved Noether current, corresponding to a symmetry of the system, by using the standard procedure, which assumes that there are at most first-order derivatives of the fields. In this paper we show how to derive the "Noether currents" corresponding to a global U(1) symmetry of the effective nonlocal second-quantized field

theory for a composite particle. From this point of view we then derive a general Ward-Takahashi identity for the quantum operators. Finally, in Sec. III we apply this to the electromagnetic interaction.

II. NOETHER'S THEOREM AND U(1) TRANSFORMATIONS

Consider an elementary particle described by a field operator ϕ which gets dressed, i.e., acquires a composite character by coupling to another field. Upon elimination of these dressing fields, the equation of motion will be determined by a nonlocal action which is a functional of the fields ϕ ,

$$I = \int d^4x d^4y \mathcal{J}(\phi(x), \phi(y)) . \quad (1)$$

This nonlocality of the action kernel $\mathcal{J}$, involving the field operators at two space-time points x and y , can be translated into the dependence on a single variable at the price that arbitrary high derivatives of the field appear. This shows that the Noether currents, associated with a symmetry of the system, cannot be obtained by the usual procedure, since it is there assumed that only first-order derivatives of the field occur. We therefore examine such symmetry currents for a nonlocal-field theory in a more general fashion. This will be done on the level of second-quantized field operators.

Consider a space- and time-dependent infinitesimal transformation of the field operators in the action integral [Eq. (1)]:

$$\phi_i(x) \rightarrow \phi_i(x) + i\xi^s(x) T_{ij}^s \phi_j(x) . \quad (2)$$

The transformation of the different components of the field ϕ_i is characterized by the transformation matrices T^s , which obey the Lie algebra of the group G associated with this transformation,

$$[T^a, T^b] = iC^{abc}T^c, \quad (3)$$

characterized by the structure constants C^{abc} . The effect of the infinitesimal transformation is that I changes by $\Delta I = \hat{I} - I$, with

$$\Delta I = \int d^4y \xi^s(y) \frac{\delta \hat{I}}{\delta \xi^s}(y), \quad (4)$$

which involves the functional derivative of $\hat{I}$ with respect to ξ^s .

If the action remains unchanged under a *global* transformation, $\xi^s(x) = c^s$, then the change ΔI for a *local* transformation can be written as [9]

$$\Delta I = \int d^4y \xi^s(y) \partial_\mu j_s^\mu(y), \quad (5)$$

i.e., in terms of the divergence of a current. From Eqs. (4) and (5), this means

$$\frac{\delta \hat{I}}{\delta \xi^s}(y) = \partial_\mu j_s^\mu(y). \quad (6)$$

This equation defines a current in terms of the action integral, up to contributions with a vanishing four-divergence. We stress that for the derivation of this equation, a local transformation was made in Eq. (2), while only a global symmetry of the system was assumed.

If the fields ϕ_i satisfy the equations of motion, obtained from Hamilton's principle

$$\frac{\delta \hat{I}}{\delta \xi^s}(y) = 0, \quad (7)$$

one sees that the current in Eq. (6) is conserved, i.e., is a "Noether current,"

$$\partial_\mu j_s^\mu(y) = 0. \quad (8)$$

Associated with such a conserved current is a "charge" operator Q^s ,

$$Q^s(t) = \int d\mathbf{y} j_s^0(\mathbf{y}, t), \quad (9)$$

which for a system with a finite spatial extension is time independent.

In this paper we will use the above approach for a U(1) transformation. In local-field theory, the operator Q [Eq. (9)] is then the generator of the unitary transformation U , which transforms the field operators according to

$$U^{-1} \phi_i U = e^{iq_i \alpha} \phi_i, \quad (10)$$

with α and q_i real. The operator U is given in terms of the Hermitian operator Q as

$$U = e^{i\alpha Q}. \quad (11)$$

By looking at an infinitesimal transformation $\alpha \rightarrow 0$ and comparing to Eq. (2), one sees that this corresponds to using only one transformation parameter $\xi^s = \alpha$ and a diagonal transformation matrix, with elements

$$T_{ij} = \delta_{ij} q_i. \quad (12)$$

Expanding Eq. (10), up to terms linear in α , one finds that the field operators and generator Q satisfy the commuta-

tion relation

$$[Q, \phi_i(\mathbf{x}, t)] = -q_i \phi_i(\mathbf{x}, t). \quad (13)$$

In a local-field theory, an explicit expression for the Noether current is obtained. In that case one can use the canonical commutation relations for the second-quantized field operators to verify Eq. (13) explicitly.

For a nonlocal-field theory, lacking canonical commutation relations for the field operators, it is not possible to show that the charge Q defined in Eq. (9) is the generator of a U(1) transformation as given by Eq. (10) or (13). As was postulated by Pauli [10], Eq. (13) is the condition for the second-quantized field operators in the general case of nonlocal-field theory.

We now look at the nonlocal action that describes an extended particle. For simplicity, a complex field with $q_i = 1$ is here assumed; the extension to a more general case is straightforward. We start from an action that is given in terms of the elementary field operators ϕ and $\phi^\dagger$ in the general form

$$I = \int d^4x d^4x' \phi^\dagger(x') F(x' - x) \phi(x). \quad (14)$$

As an illustration, we briefly consider an extended nucleon. In terms of the elementary nucleon field operators ψ and $\bar{\psi}$, the action can be written as [7]

$$I = \int d^4x d^4x' \bar{\psi}(x') F(x' - x) \psi(x), \quad (15)$$

where we have dropped isospin indices. For a spin- $\frac{1}{2}$ field, the function F in the action integral is a Dirac matrix of the general form [8]

$$F(y) = g_1(y) + \not{y} g_2(y). \quad (16)$$

The scalar functions g_1 and g_2 are the so-called "form functions" which describe the nucleon structure. They fall off rapidly for $|y^2| \gg d^2$, where d characterizes the space-time extension of the nucleon. To arrive at the local-field theory appropriate for a structureless point nucleon, it is convenient to first rewrite $F(x' - x)$ as

$$F(x' - x) = i\gamma \cdot \partial_x g(x' - x) - M f(x' - x). \quad (17)$$

Then, with f and g chosen as

$$f(x' - x) = g(x' - x) = \delta^4(x' - x), \quad (18)$$

one finds by integration by parts the well-known local action for a spin- $\frac{1}{2}$ point particle,

$$I^{\text{point}} = \int d^4x \bar{\psi}(x) (i\partial - M) \psi(x). \quad (19)$$

Returning to the nonlocal action in Eq. (14), one sees that the integrand is clearly invariant under global U(1) phase transformations of the fields according to Eq. (10):

$$\begin{aligned} \phi^\dagger(x') F(x' - x) \phi(x) &\rightarrow e^{-i\alpha} \phi^\dagger(x') F(x' - x) e^{i\alpha} \phi(x) \\ &= \phi^\dagger(x') F(x' - x) \phi(x). \end{aligned} \quad (20)$$

We now follow the general procedure given above to derive the current and charge operators corresponding to this symmetry. First, we make an infinitesimal and space-time-dependent transformation, i.e.,

$$\begin{aligned}\phi(y) &\rightarrow \phi(y) + i\xi(y)\phi(y), \\ \phi^\dagger(y) &\rightarrow \phi^\dagger(y) - i\xi(y)\phi^\dagger(y).\end{aligned}\quad (21)$$

The transformed action is then

$$\begin{aligned}\hat{I} &= \int d^4x' d^4x [1 - i\xi(x')] \phi^\dagger(x') F(x' - x) \\ &\quad \times [1 + i\xi(x)] \phi(x).\end{aligned}\quad (22)$$

The functional derivative with respect to ξ is, to lowest order in ξ ,

$$\begin{aligned}\frac{\delta \hat{I}}{\delta \xi}(y) &= i \int d^4x' d^4x \phi^\dagger(x') F(x' - x) \phi(x) \\ &\quad \times [\delta^4(x - y) - \delta^4(x' - y)].\end{aligned}\quad (23)$$

Therefore, according to the definition in Eq. (6), we find that the four-divergence of the current operator is given by

$$\begin{aligned}\partial^\mu j_\mu(y) &= i \int d^4x' d^4x \phi^\dagger(x') F(x' - x) \phi(x) \\ &\quad \times [\delta^4(x - y) - \delta^4(x' - y)].\end{aligned}\quad (24)$$

The general form for the current j_μ is

$$j_\mu(y) = i \int d^4x' d^4x \phi^\dagger(x') s_\mu(x' - y, x - y) \phi(x), \quad (25)$$

which expresses the current of the composite particle in terms of the elementary fields. The compositeness is contained in the "smearing function" s_μ in Eq. (25), which governs the contributions of the fields at x and x' to the composite particle current at y . Causality requires that the fields contribute only if x and x' can be causally connected to y , i.e., only contributions for timelike arguments of s_μ :

$$\begin{aligned}s_\mu(x' - y, x - y) &= 0 \\ \text{if } (x - y)^2 < 0 \text{ or } (x' - y)^2 < 0.\end{aligned}\quad (26)$$

Substituting the above general form for j_μ in Eq. (24) yields the following condition for s_μ :

$$\begin{aligned}\frac{\partial}{\partial y_\mu} s_\mu(x' - y, x - y) \\ = F(x' - x) [\delta^4(x - y) - \delta^4(x' - y)].\end{aligned}\quad (27)$$

Of course, the current is defined through Eq. (6) or (24) only up to contributions with a vanishing four divergence. For the smearing function this means that to a solution s_μ of Eq. (27) one may always add a term Δs_μ which satisfies

$$\frac{\partial}{\partial y_\mu} \Delta s_\mu(x' - y, x - y) = 0. \quad (28)$$

Therefore, there are always possible additional currents Δj_μ which are conserved as a consequence of Eq. (28). This is of course analogous to the situation in quantum mechanics and electrodynamics, where, e.g., a separately conserved magnetic-moment term for the nucleon can be added. As mentioned before, the current j_μ [Eq. (25)], constrained by the condition Eq. (27), is conserved if the fields operators satisfy the equations of motion. To see

this one makes use of the equation of motion for the fields, obtained by requiring the action I [Eq. (14)] to be stationary under variation of ϕ (or $\phi^\dagger$):

$$\begin{aligned}\int d^4x' \phi^\dagger(x') F(x' - x) &= 0, \\ \int d^4x F(x' - x) \phi(x) &= 0.\end{aligned}\quad (29)$$

From the above it is easily seen that the current is conserved, i.e., $\partial^\mu j_\mu(y) = 0$.

The charge operator Q can be found by integrating the component j_0 over y . To do this we first note that

$$\begin{aligned}\int d\mathbf{y} \frac{\partial}{\partial y_\mu} s_\mu(x' - y, x - y) \\ = \frac{\partial}{\partial y_0} \int d^3\mathbf{y} s_0(x' - y, x - y),\end{aligned}\quad (30)$$

by using the divergence theorem and the fact that the system is of finite extension in space, which allows one to drop the surface term. Using Eq. (27) and

$$\frac{\partial}{\partial t} \theta(x_0 - t) = -\delta(x_0 - t), \quad (31)$$

we can rewrite the left-hand side of Eq. (30) as

$$\begin{aligned}\frac{\partial}{\partial y_0} \int d\mathbf{y} s_0(x' - y, x - y) \\ = \frac{\partial}{\partial y_0} F(x' - x) [\theta(x'_0 - y_0) - \theta(x_0 - y_0)].\end{aligned}\quad (32)$$

Integrating both sides of (32) over y_0 from $-\infty$ to t , we obtain

$$\begin{aligned}\int d\mathbf{y} s_0(x' - y, x - y)|_{y_0=t} \\ = F(x' - x) [\theta(x'_0 - t) - \theta(x_0 - t)],\end{aligned}\quad (33)$$

where again the finite extension in space-time was used to drop the boundary contribution. Using Eq. (33) in (25) and the definition of the charge finally yields

$$\begin{aligned}Q(t) &= i \int d^4x' d^4x \phi^\dagger(x') F(x' - x) \phi(x) \\ &\quad \times [\theta(x'_0 - t) - \theta(x_0 - t)].\end{aligned}\quad (34)$$

Note that the undetermined contribution Δs_μ would not have changed this expression for $Q(t)$, since Eq. (30) remains unchanged. This is guaranteed by the divergence condition, Eq. (28) and the fact that the system is finite in space-time.

To derive the Ward-Takahashi identity, it is necessary to know the equal-time commutation relation of j_0 with the fields. The charge Q and field ϕ satisfy the commutation relation Eq. (13). This can be expressed in terms of j_0 as

$$\int d\mathbf{y} [j_0(\mathbf{y}, t), \phi(\mathbf{x}, t)] = -\phi(\mathbf{x}, t). \quad (35)$$

We see at once that a possible solution is

$$[j_0(\mathbf{y}, t), \phi(\mathbf{x}, t)] = -\delta^3(\mathbf{x} - \mathbf{y}) \phi(\mathbf{x}, t). \quad (36)$$

Assume that a different solution of Eq. (35) exists:

$$[j_0(\mathbf{y}, t), \phi(\mathbf{x}, t)] = -\delta^3(\mathbf{x} - \mathbf{y})\phi(\mathbf{x}, t) + a(\mathbf{x}, \mathbf{y}, t). \quad (37)$$

Since as in s_μ , x and y are causally connected, we can have contributions only for timelike separations,

$$[j_0(y), \phi(x)] \neq 0 \quad \text{only if } (x_0 - y_0)^2 - (\mathbf{x} - \mathbf{y})^2 \geq 0. \quad (38)$$

For equal times we can therefore only obtain a nonvanishing contribution if $\mathbf{x} = \mathbf{y}$. Provided that the commutator is no more singular than a δ function, this means that the additional contribution in Eq. (37) must be of the form

$$a(\mathbf{x}, \mathbf{y}, t) = b(\mathbf{x}, t)\delta^3(\mathbf{x} - \mathbf{y}). \quad (39)$$

Furthermore, the condition for the space integral [Eq. (35)] leads to the constraint

$$\int d^3y a(\mathbf{x}, \mathbf{y}, t) = 0. \quad (40)$$

From Eq. (39) one sees that this implies $b(\mathbf{x}, t) = 0$, and

therefore the extra term in Eq. (37) vanishes. The adjoint field operator must satisfy

$$[Q, \phi^\dagger(\mathbf{x}, t)] = \phi^\dagger(\mathbf{x}, t), \quad (41)$$

and one obtains, in a similar fashion,

$$[j_0(y), \phi^\dagger(x)] = \delta^3(\mathbf{x} - \mathbf{y})\phi^\dagger(\mathbf{x}, t). \quad (42)$$

The commutators [Eqs. (36) and (42)] are implicitly assumed in the derivation of the two-body Ward-Takahashi identity of Bentz [11].

We can now use these equal-time commutation relations [Eqs. (36) and (42)] to establish a connection between the vertex function Γ_μ ,

$$\Gamma_\mu(x, x', y) = \langle 0 | T\phi(x)\phi^\dagger(x')j_\mu(y) | 0 \rangle, \quad (43)$$

and the propagator or Green's function $G(x, x')$,

$$iG(x - x') = \langle 0 | T\phi(x)\phi^\dagger(x') | 0 \rangle. \quad (44)$$

Taking the four-divergence of the vertex function [Eq. (43)] with respect to y_μ , one obtains

$$\begin{aligned} \frac{\partial}{\partial y_\mu} \langle 0 | T\phi(x)\phi^\dagger(x')j_\mu(y) | 0 \rangle &= \langle 0 | T\phi(x)\phi^\dagger(x')\partial^\mu j_\mu(y) | 0 \rangle + \langle 0 | T\phi(x)\delta(x'_0 - y_0)[j_0(y), \phi^\dagger(x')] | 0 \rangle \\ &\quad + \langle 0 | T\delta(x_0 - y_0)[j_0(y), \phi(x)]\phi^\dagger(x') | 0 \rangle. \end{aligned} \quad (45)$$

We now use the commutators of j_0 with the fields [Eqs. (36) and (42)] and, furthermore, current conservation, i.e.,

$$\partial^\mu j_\mu(y) = 0. \quad (46)$$

This yields the general Ward-Takahashi identity

$$\frac{\partial}{\partial y_\mu} \langle 0 | T\phi(x)\phi^\dagger(x')j_\mu(y) | 0 \rangle = \delta^4(x' - y) \langle 0 | T\phi(x)\phi^\dagger(x') | 0 \rangle - \delta^4(x - y) \langle 0 | T\phi(x)\phi^\dagger(x') | 0 \rangle \quad (47)$$

or

$$-i \frac{\partial}{\partial y_\mu} \Gamma_\mu(x, x', y) = \delta^4(x' - y)G(x - x') - \delta^4(x - y)G(x - x'). \quad (48)$$

Defining the irreducible vertex function Γ_μ^{irr} as

$$\Gamma_\mu(x, x', y) = \int d^4u d^4u' iG(x - u)\Gamma_\mu^{\text{irr}}(u - y, u' - y)iG(u' - x'), \quad (49)$$

and using the Fourier transforms

$$\begin{aligned} G(x' - x) &= \int \frac{d^4p}{(2\pi)^4} e^{-ip(x' - x)} G(p), \\ \Gamma_\mu^{\text{irr}}(x - y, x' - y) &= \int \frac{d^4p'}{(2\pi)^4} \frac{d^4p}{(2\pi)^4} e^{ip'(x' - y) - ip(x - y)} \Gamma_\mu^{\text{irr}}(p', p), \end{aligned} \quad (50)$$

one obtains in momentum space the familiar form

$$(p' - p)^\mu \Gamma_\mu^{\text{irr}}(p', p) = G^{-1}(p') - G^{-1}(p). \quad (51)$$

III. THE ELECTROMAGNETIC FIELD

We now consider a composite particle and include the electromagnetic interaction into the discussion. The general approach in Sec. II, of course, also applies if one con-

siders the photon field, along with others, as responsible for dressing the nucleon. The action, smearing function, etc. will therefore also contain contributions from the electromagnetic field and the above discussion applies with this extended interpretation. However, the electromagnetic field is introduced into the action in a special way, namely, such that a locally gauge-invariant theory is obtained. We now will use this feature to obtain the Ward-Takahashi identity appropriate for the electromagnetic vertex operator.

The action Eq. (14) can be made locally gauge invariant through a "Bloch factor" $\exp[-ie\chi(x',x)]$, which contains the electromagnetic field A_μ [7,6],

$$\begin{aligned} I &= \int d^4x' d^4x \phi^\dagger(x') \tilde{F}(x',x) \phi(x) \\ &= \int d^4x' d^4x \phi^\dagger(x') \\ &\quad \times \exp[-ie\chi(x',x)] F(x'-x) \phi(x), \end{aligned} \quad (52)$$

in which e is the charge of the field ϕ and the function $\chi(x',x)$ is defined as

$$\chi(x',x) \equiv \int d^4y \xi_\mu(x'-y, x-y) A^\mu(y). \quad (53)$$

The function $\xi_\mu(x'-y, x-y)$ has to satisfy the condition

$$\frac{\partial}{\partial y_\mu} \xi_\mu(x'-y, x-y) = \delta^4(x-y) - \delta^4(x'-y). \quad (54)$$

One can easily verify that this action, with the above condition, is indeed invariant under local gauge transformations of the fields given by

$$\begin{aligned} A_\mu(x) &\rightarrow A_\mu(x) + \partial_\mu \alpha(x), \\ \phi(x) &\rightarrow e^{-ie\alpha(x)} \phi(x), \\ \phi^\dagger(x) &\rightarrow e^{ie\alpha(x)} \phi^\dagger(x). \end{aligned} \quad (55)$$

Note that α is a function of space-time.

We return, as an example, to the case of a point spin- $\frac{1}{2}$ particle; i.e., $F(x'-x)$ is given by Eqs. (17) and (18). If χ in Eq. (52) satisfies

$$\chi(x,x) = 0 \quad (56)$$

and

$$\frac{\partial}{\partial x^\mu} \chi(x',x) \big|_{x'=x} = A_\mu(x), \quad (57)$$

the action [Eq. (52)] becomes

$$I^{\text{point}} = \int d^4x \bar{\psi}(x) (i\partial + eA - M) \psi(x). \quad (58)$$

Consequently, minimal substitution $\partial_\mu \rightarrow \partial_\mu - ieA_\mu$ in Eq. (19) is a special case of the above procedure of obtaining a gauge-invariant action. In fact, it is for the point spin- $\frac{1}{2}$ particle produced by the choice of Bloch [6],

$$\chi(x',x) = \int_x^{x'} A_\mu(y) d^\mu y, \quad (59)$$

where the integration path is taken along a straight space-time line connecting x to x' . It is readily seen that this choice indeed satisfies the conditions in Eqs. (56) and (57). Other choices for χ , in the context of nonrelativistic quantum mechanics, have been discussed by Heller [12]. There it is also shown that the first condition [Eq. (56)] for χ follows from the requirement of time-reversal invariance and Hermiticity.

The way of writing a gauge-invariant action [Eq. (52)] is not unique. First, the function ξ_μ is not uniquely determined through the condition in Eq. (54). Second, one can add separately gauge-invariant terms to the action integral, e.g., such as the anomalous magnetic moment or "Pauli" term in local-field theory. Such terms do not

alter the Ward-Takahashi identity we consider below.

The action in Eq. (52) is of course invariant under a global U(1) transformation, and analogous to Eqs. (25) and (27), we obtain for the current

$$J_\mu(y) = i \int d^4x' d^4x \phi^\dagger(x') S_\mu(x'-y, x-y) \phi(x), \quad (60)$$

with

$$\begin{aligned} \frac{\partial}{\partial y_\mu} S_\mu(x'-y, x-y) \\ = \tilde{F}(x',x) [\delta^4(x-y) - \delta^4(x'-y)]. \end{aligned} \quad (61)$$

As stressed in Sec. II, the condition Eq. (61) for a given action [Eq. (52)] does not uniquely define the smearing function S_μ and, therefore, the total current. To obtain the Ward-Takahashi identity involving the electromagnetic vertex, we first define the electromagnetic current as

$$\begin{aligned} J_\mu^{\text{em}}(y) &\equiv \frac{\delta I}{\delta A_\mu(y)} \\ &= -ie \int d^4x' d^4x \phi^\dagger(x') \tilde{F}(x',x) \\ &\quad \times \xi_\mu(x'-y, x-y) \phi(x), \end{aligned} \quad (62)$$

which defines the electromagnetic smearing function S_μ^{em} in terms of $\tilde{F}$ and ξ_μ as

$$S_\mu^{\text{em}}(x'-y, x-y) \equiv -e \tilde{F}(x',x) \xi_\mu(x'-y, x-y). \quad (63)$$

Because of Eq. (54), this smearing function satisfies the condition

$$\begin{aligned} \frac{\partial}{\partial y_\mu} S_\mu^{\text{em}}(x'-y, x-y) \\ = -e \tilde{F}(x',x) [\delta^4(x-y) - \delta^4(x'-y)]. \end{aligned} \quad (64)$$

Up to a constant factor, both the general smearing function S_μ [Eq. (61)] and the electromagnetic smearing function S_μ^{em} have the same four-divergence. This is a consequence of introducing the electromagnetic field in Eq. (14) to obtain a locally gauge-invariant action through the Bloch function. This implies that also J_μ^{em} is conserved and that the parts of the currents J_μ and J_μ^{em} which are constrained by the divergence condition are identical up to a factor e , the charge of the particle. In nonrelativistic quantum mechanics, for example, this means that to lowest order in e the electromagnetic current density of the charged particle is e times the probability current density. Nontrivial differences between the two currents can, for example, be due to separately conserved current contributions. All relevant arguments concerning the vertex operator and propagator, therefore, go through as in Sec. II, and we obtain, for the electromagnetic operator,

$$\begin{aligned} \frac{\partial}{\partial y_\mu} \langle 0 | T \phi(x) \phi^\dagger(x') J_\mu^{\text{em}}(y) | 0 \rangle \\ = -e [\delta^4(x'-y) \langle 0 | T \phi(x) \phi^\dagger(x') | 0 \rangle \\ - \delta^4(x-y) \langle 0 | T \phi(x) \phi^\dagger(x') | 0 \rangle]. \end{aligned} \quad (65)$$

Again, as in Eq. (49), one obtains for the irreducible elec-

tromagnetic vertex, in momentum space, the condition

$$(p' - p)^\mu \Gamma_\mu^{\text{em,irr}}(p', p) = -e [G^{-1}(p') - G^{-1}(p)] . \quad (66)$$

This is what one commonly refers to as the “Ward-Takahashi” identity, which relates the dressed irreducible electromagnetic vertex to the propagator (or self-energy) of the particle in question. Note that this holds to all orders in e since the fields ϕ and $\phi^\dagger$ in Eq. (65) are assumed to satisfy the full equation of motion, i.e., to all order in e . In the literature this is often quoted only to lowest order in e . To that order our general field-theoretical derivation reproduces the results of Ohta [8] for the case of a composite nucleon, obtained in the context of nonlocal quantum mechanics.

IV. SUMMARY

Recently, there has been considerable interest in the electromagnetic properties of composite systems, such as the nucleon, under kinematical conditions where not only the free form factors are sufficient. To address such problems, one has to consider a field theory for composite systems. We have here studied a second quantized, field-theoretical description for a composite particle, in which the equations of motion for the particle field are derived from a nonlocal action. Such a nonlocal action contains derivatives of the field operators of arbitrary order. This makes it impossible to follow the usual steps leading to the conserved Noether currents, since this derivation assumes a local theory, containing only first-order derivatives of the field operators. We have shown how one can

derive conditions for the current operator in a nonlocal theory, corresponding to a global symmetry of the nonlocal action. We then considered a nonlocal action that describes a single composite particle in terms of the elementary field operators ϕ and $\phi^\dagger$ and is invariant under a global U(1) transformation. We derived the charge operator and conditions for the current operator j_μ , expressed in terms of the “form function” appearing in the action integral and the second-quantized field operators. Based on the requirement of causality and the singularity structure of the fields, we then also gave the commutation relation of the current operator j_0 with the second-quantized field operators. This then was used to arrive at a general connection between the vertex and propagation functions, the Ward-Takahashi identity. We then applied this general formalism to examine the electromagnetic properties of a composite particle. Since the electromagnetic field is included in the action in a way to preserve local gauge invariance, the resulting Ward-Takahashi identity for the electromagnetic vertex operator leads to an analogous connection with the dressed propagator of the particle.

ACKNOWLEDGMENTS

The authors would like to thank Dr. J. Kunz for helpful discussions. This work was supported by the Foundation for Fundamental Research on Matter (FOM), the Netherlands Organization for Scientific Research (NWO), and the Bundesministerium für Forschung und Technologie.

-
- [1] A. M. Bincer, Phys. Rev. **118**, 855 (1960).
 - [2] H. W. L. Naus and J. H. Koch, Phys. Rev. C **36**, 2459 (1987).
 - [3] P. C. Tiemeijer and J. A. Tjon, Phys. Rev. C **42**, 599 (1990).
 - [4] Y. Takahashi, Nuovo Cimento **6**, 371 (1957).
 - [5] K. Nishijima, Phys. Rev. **111**, 995 (1958).
 - [6] C. Bloch, K. Dan. Vidensk. Selsk. Mat. Fys. Medd. **27**, 1 (1952).
 - [7] M. Chrétien and R. E. Peierls, Proc. R. Soc. London A **223**, 468 (1954).
 - [8] K. Ohta, Phys. Rev. C **41**, 1213 (1990).
 - [9] B. de Wit and J. Smith, *Field Theory in Particle Physics* (North-Holland, Amsterdam, 1986), Chap. 1.
 - [10] W. Pauli, Nuovo Cimento **10**, 648 (1953).
 - [11] W. Bentz, Nucl. Phys. **A446**, 678 (1985).
 - [12] L. Heller, in *The Two-Body Force in Nuclei*, edited by S. M. Austin and G. M. Crawley (Plenum, New York, 1972).