



UvA-DARE (Digital Academic Repository)

Models of molecular clouds

Boland, W.H.M.

[Link to publication](#)

Citation for published version (APA):

Boland, W. H. M. (1982). Models of molecular clouds

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Chapter III

A Model for the Formaldehyde Maser near NGC 7538–IRS 1

W. Boland and T. de Jong

Astronomical Institute, Roetersstraat 15, 1018 WB Amsterdam, The Netherlands

Received October 6, accepted November 25, 1980

Summary. It is shown that the population of the $1_{10}-1_{11}$ transition of Formaldehyde can be inverted by the free-free radio continuum radiation of a nearby compact H II region. The H II region must be very compact with emission measures of $10^8-10^{10} \text{ cm}^{-6} \text{ pc}$ so that it is optically thick at cm wavelengths but still optically thin at millimeter wavelengths. A detailed model of the compact H II region NGC 7538–IRS 1 and its surrounding cloud is constructed to explain the observed $1_{10}-1_{11}$ brightness temperatures of about 10^6 K . The masing gas is in front of the H II region so that it amplifies the radio continuum radiation. To allow sufficiently rapid radiative pumping mm line photons must be able to escape from the cloud which requires a large velocity gradient in the gas. To create sufficient amplification molecules must be lined up in velocity over long pathlengths. Consistent with observations of CO, OH, and H110 α lines we assume that NGC 7538–IRS 1 moves towards us through (or expands into) a molecular cloud with a velocity $V \approx 10 \text{ km s}^{-1}$ and that the molecular gas in front of it is pushed sideways creating velocity gradients of $\sim V/R_{II}$ perpendicular to the line of sight and coherent pathlengths of $\sim R_{II}$ along the line of sight. With these assumptions an amplification factor of ~ 100 required to explain the observed brightness temperatures can be achieved if $n(\text{H}_2) \approx 10^6 \text{ cm}^{-3}$, $T_k \approx 20 \text{ K}$, and $x(\text{H}_2\text{CO}) \approx 8 \cdot 10^{-7}$. We show that those Formaldehyde abundances can be obtained in dark clouds. Our model also explains the observed maser spot sizes. The fact that the maser is saturated explains why both observed maser spots have about equal intensities.

Key words: Formaldehyde maser - compact H II region - molecular clouds

1. Introduction

Using the Westerbork Radio Synthesis Telescope at 6 cm Forster et al. (1980) observed two emission lines of the $1_{10}-1_{11}$ transition of Formaldehyde (H_2CO) at radial velocities of -57.6 and -59.8 km s^{-1} near the ultracompact H II region and infrared source NGC 7538–IRS 1. They found that the emission was spatially unresolved (source size $< 4''$) corresponding to a lower limit to the brightness temperature of the emission of 800 K .

More recently using the Very Large Array Rots et al. (1980) have resolved the source into two spots separated by about $0''.1$. The emission spots have angular sizes smaller than $0''.15$ cor-

responding to linear dimensions of $\sim 8 \cdot 10^3 \text{ cm}$ (at a distance of 3.5 kpc) and implying lower limits to the brightness temperatures of $\sim 5 \cdot 10^5 \text{ K}$. The small source sizes and the extremely high brightness temperatures strongly suggest a maser interpretation for the 6 cm H_2CO emission near NGC 7538–IRS 1.

In the direction of many dark clouds the 6 cm line of H_2CO is observed in absorption against the 2.7 K cosmic background radiation (Minn and Greenberg, 1973; Dieter, 1973). This anomalous absorption is best explained by a collisional pump model originally suggested by Townes and Cheung (1969) and later confirmed by accurate numerical calculations of $\text{H}_2-\text{H}_2\text{CO}$ cross sections by Garrison et al. (1975). Since the circumstances in the molecular cloud in which NGC 7538–IRS 1 is embedded are presumably not very different from those in dark clouds the anomalous emission in the 6 cm H_2CO line will not be easily explained by collisional pumping.

Led by the fact that the level population of the H_2CO molecule is extremely sensitive to deviations from blackbody radiation at millimeter wavelengths (Thaddeus, 1972) and in view of the proximity of the ultra-compact H II region we propose in this paper that the H_2CO maser near NGC 7538–IRS 1 is pumped by the free-free continuum radiation of the H II region.

2. The Radiative Transfer Problem

We consider Formaldehyde molecules in a molecular Hydrogen cloud in the presence of a strong radio continuum radiation field. To find the H_2CO level populations we solve the equations of radiative transfer assuming that the thermal velocities of the molecules are much smaller than the large scale velocities in the cloud (cf. de Jong et al., 1975 and references therein).

Let $n_i(r)$ be the density of molecules in level i at distance r from the centre of a spherical cloud. In statistical equilibrium we have

$$\sum_j P_{jk} n_j = n_k \sum_j P_{ji} \quad (1)$$

where

$$\begin{aligned} P_{jk} &= A_{jk}(\beta_{jk}(1 + H_{jk}) + D_{jk}) + C_{jk} \quad (j > k) \\ &= A_{jk}(\beta_{jk} H_{jk} + D_{jk}) \frac{g_k}{g_j} + C_{jk} \quad (j < k). \end{aligned} \quad (2)$$

The transition probabilities P_{jk} are expressed in terms of the spontaneous radiative transition probability A_{jk} , the collision induced transition probabilities C_{jk} and the net radiative transition rates $\beta_{jk} A_{jk}$ where β_{jk} is the so called escape probability (cf. de

Send offprint requests to: W. Boland

Table 1. Frequencies and spontaneous emission rates of rotational transitions in ortho-H₂CO

| Transition | ν_{ij} (GHz) | A_{ij} (s ⁻¹) |
|----------------------------------|------------------|-----------------------------|
| 1 ₁₀ -1 ₁₁ | 4.83 | 3.59 10 ⁻⁹ |
| 2 ₁₂ -1 ₁₁ | 140.84 | 5.34 10 ⁻⁵ |
| 2 ₁₁ -1 ₁₀ | 150.50 | 6.52 10 ⁻⁵ |
| 2 ₁₁ -2 ₁₂ | 14.49 | 3.23 10 ⁻⁸ |
| 3 ₁₃ -2 ₁₂ | 211.23 | 2.29 10 ⁻⁴ |
| 3 ₁₂ -2 ₁₁ | 225.73 | 2.79 10 ⁻⁴ |
| 3 ₁₂ -3 ₁₃ | 28.98 | 1.29 10 ⁻⁷ |
| 4 ₁₄ -3 ₁₃ | 281.53 | 5.93 10 ⁻⁴ |
| 4 ₁₃ -3 ₁₂ | 300.84 | 7.24 10 ⁻⁴ |
| 4 ₁₃ -4 ₁₄ | 48.28 | 3.59 10 ⁻⁷ |
| 5 ₁₅ -4 ₁₄ | 355.71 | 1.25 10 ⁻³ |
| 5 ₁₄ -4 ₁₃ | 379.83 | 1.52 10 ⁻³ |
| 5 ₁₄ -5 ₁₅ | 72.41 | 8.09 10 ⁻⁷ |
| 6 ₁₆ -5 ₁₅ | 423.5 | 2.17 10 ⁻³ |
| 6 ₁₅ -5 ₁₄ | 452.5 | 2.65 10 ⁻³ |
| 6 ₁₅ -6 ₁₆ | 101.4 | 1.59 10 ⁻⁶ |
| 7 ₁₇ -6 ₁₆ | 494.0 | 3.51 10 ⁻³ |
| 7 ₁₆ -6 ₁₅ | 527.6 | 4.28 10 ⁻³ |
| 7 ₁₆ -7 ₁₇ | 135.0 | 2.82 10 ⁻⁶ |

Jong et al., 1975)

$$\beta_{jk}(r) = \frac{1}{2} \int_{-1}^{+1} \beta_{jk}(r, \mu) d\mu = \frac{1}{2} \int_{-1}^{+1} \frac{1 - \exp\{-\tau_{jk}(r, \mu)\}}{\tau_{jk}(r, \mu)} d\mu. \quad (3)$$

The optical depth

$$\tau_{jk}(r, \mu) = \frac{\lambda_{jk}^3 A_{jk}}{8\pi} \frac{r}{|1 - \alpha\mu^2|v(r)} \left(n_k \frac{g_l}{g_k} - n_j \right) \quad (4)$$

is defined along a direction which makes an angle $\arccos(\mu)$ with the radius vector. The parameter α is related to the gradient of the large scale velocity field in the gas

$$\alpha = 1 - \frac{r}{v(r)} \frac{dv(r)}{dr}. \quad (5)$$

The quantities $\beta_{jk} H_{jk}$ and $D_{jk}(r)$ are the so-called photon occupation numbers of the jk line (Ehitzur et al., 1976) of the blackbody background radiation field ($T_{bb} = 2.7$ K)

$$H_{jk} = \left(\exp\left(\frac{hc}{\lambda_{jk} k T_{bb}}\right) - 1 \right)^{-1} \quad (6)$$

and of the free-free continuum radiation field

$$D_{jk}(r) = \frac{1 - \exp\{-\tau_c(\lambda_{jk})\}}{\exp\left(\frac{hc}{\lambda_{jk} k T_e}\right) - 1} W(r) \quad (7)$$

where

$$W(r) = \frac{1}{2} \int_{|1 - \alpha\mu^2|^{1/2}}^1 \beta(r, \mu) d\mu \quad (8)$$

is a factor with which the continuum radiation field at distance r from the centre of the H II region ($r > R_{II}$) is diluted. If the continuum radiation is not absorbed in the surrounding neutral gas ($\beta = 1$) $W(r)$ reduces to the usual geometrical dilution factor.

The free-free continuum optical depth of an H II region with electron temperature $T_e = 10^4$ K may be written (cf. Spitzer, 1978)

$$\tau_c(\lambda) = (\lambda/\lambda_0)^2 \{1 + 0.311 \log(\lambda)\} \quad (9)$$

where λ and λ_0 are expressed in cm. The wavelength λ_0 where the H II region becomes optically thick equals

$$\lambda_0 = 6.35 \cdot 10^4 E_m^{-0.5} \text{ cm} \quad (10)$$

where

$$E_m = 4/3 n_e^2 R_{II} \quad (11)$$

is the emission measure of a spherical H II region.

The spontaneous radiative transition probabilities between the lowest 14 rotational levels of ortho-Formaldehyde, listed in Table 1, are calculated from the relation

$$A_{jk} = \frac{64\pi^3}{3h} E_{jk}^3 \mu_{jk}^2 \quad (12)$$

where the wavenumbers E_{jk} of the lower transitions are taken from measurements of Nerf (1972) and Winniewisser et al. (1974) and of the higher transitions from Townes and Schawlow (1955). The matrix elements $(\mu_{jk}/\mu)^2$ are interpolated from tables of Schwendemann and Laurie (1958) for an asymmetry factor $\kappa = -0.961$ using a dipole moment of $\mu = 2.34$ Debye (Townes and Schawlow, 1955). Hyperfine splitting of the rotational levels of H₂CO is neglected throughout our calculations.

The H₂-H₂CO collision rates are taken from Green et al. (1978) but multiplied by a factor 1.57 to correct for the difference in reduced mass of H₂ and He using a 80% H₂-20% He gas mixture.

3. Pump Mechanism

To illustrate the pump mechanism we approximate ortho-H₂CO by its four lowest rotational levels. Since we shall argue that the H₂CO maser is radiatively pumped by the radio continuum radiation of the H II region we consider only radiative transitions. Solving the equations of statistical equilibrium for this system we find

$$\frac{n_2}{n_1} = \frac{p P_{12} + P_{33} P_{34} F_{32}}{p P_{21} + P_{24} P_{33} P_{31}} \quad (13)$$

where $p = P_{31}(P_{32} + P_{33}) + P_{32} P_{34}$ and the transition rates P_{ij} are defined by Eq. (2) (with $C_{ij} = 0$). From Eq. (13) we derive the following condition for population inversion in the groundstate doublet

$$\frac{P_{34} P_{32}}{P_{33} P_{31}} \frac{P_{13} - q(1 + P_{31}/P_{34})}{P_{24} + q} > 1 \quad (14)$$

where $q = P_{21} - P_{12} = \beta_{21} A_{21}$. For illustrative purposes we further assume that all lines are optically thin ($\beta_{ij} = 1$) and that the 2.7 K background radiation field can be neglected ($H_{ij} = 0$) so that Eq. (2) reduce to

$$P_{jk} = A_{jk}(1 + D_{jk}) \quad j > k \quad (15)$$

$$= A_{jk} D_{jk} \frac{g_l}{g_j} \quad j < k$$

The only free parameters in Eqs. (14) and (15) are the wavelength λ_0 where the H II region becomes optically thick and the geometrical dilution factor W ($\beta_{ij} = 1$) which depends on the distance of the masing gas to the H II region. Inserting molecular parameters (see Table 2) one easily verifies that $P_{24} \gg q$ and $P_{13} \gg q(1 + P_{31}/P_{34})$ for $\lambda_0 \ll 10^3$ cm and $W \geq 0.1$ so that the population inversion condition may be written

$$\frac{F_{32}}{F_{31} F_{33}} > 1 \quad (16)$$




Fig. 1. Population ratios in the ground state doublet levels of 4-level ortho-Formaldehyde molecules near an H II region as a function of the radio continuum spectrum (λ_0 is the wavelength where the H II region becomes optically thick) and the distance to the H II region ($W=0.5$ corresponds to $r=R_{II}$ and $W=0.1$ to $r=5/3 R_{II}$). Collisions and radiation trapping are neglected




Fig. 2. Population ratios in the $J=1$ and $J=2$ doublet levels of 14-level ortho-Formaldehyde molecules near an H II region as a function of the radio continuum spectrum. Curves (1)–(3) are for the $1_{10} \rightarrow 1_{11}$ transition without collisions and radiation trapping (curve 1), with collisions but no radiation trapping (curve 2), with collisions and radiation trapping included (curve 3). Curve 4 gives the population ratio in the $2_{11} \rightarrow 2_{12}$ transition for the case that collisions and radiation trapping are included. The parameters adopted for this calculation are given in the text

Table 2. Parameters for NGC 7538 – IRS 1

| | | References |
|-----------------------|------------------------------------|-------------------------|
| Electron temperature | $T_e = 10^4$ K | |
| Total angular area | $\Delta\Omega = 0.67$ arc s^2 | Harris and Scott (1976) |
| Emission measure | $E_m = 6 \cdot 10^9$ cm $^{-6}$ pc | Harris and Scott (1976) |
| Distance | $d = 3.5$ kpc | Israel et al. (1973) |
| Radius | $R_{II} = 2.4 \cdot 10^{16}$ cm | |
| Mean electron density | $n_e = 2 \cdot 10^5$ cm $^{-3}$ | |
| Turn-over wavelength | $\lambda_0 = 2.7$ cm | |

where

$$F_{ij} = \frac{1 + D_{ij}}{D_{ij}} \quad (17)$$

is the ratio of the downward to the upward radiative transition rates between levels i and j . If spontaneous radiative transitions dominate $F_{ij} \gg 1$ and if induced radiative transitions dominate $F_{ij} \rightarrow 1$. Condition (16) has a simple physical explanation. Population inversion of the ground state $J=1$ doublet of H_2CO occurs if the free-free continuum radiation field induces relatively more downward transitions from level 4 to level 2 than from level 3 to

level 1 ($F_{42} > F_{31}$) while the populations of levels 4 and 3 are kept approximately equal by rapid radiative exchange ($F_{43} \approx 1$). This is numerically illustrated in Fig. 1 where n_2/n_1 is plotted against λ_0 for two values of W . For $W=0.5$ the masing gas is situated at the boundary R_{II} of the H II region, for $W=0.1$ the inversion layer is located at a distance $5/3 R_{II}$ from the centre of the H II region. If $W=0.5$ inversion occurs for $5 \text{ mm} \lesssim \lambda_0 \lesssim 50 \text{ cm}$ and if $W=0.1$ for $2 \text{ mm} \lesssim \lambda_0 \lesssim 20 \text{ cm}$. From this illustrative example we conclude that the $J=1$ doublet of H_2CO can be inverted by the free-free radio continuum radiation of an H II region if it is very compact with emission measures in the range $10^9 - 10^{10}$ cm $^{-6}$ pc.




Fig. 3. Schematic picture of our model of the compact H II region NGC 7538-IRS 1 and its surrounding molecular cloud




Fig. 4. The relative population inversion of the ground state doublet levels of ortho-Formaldehyde as a function of distance from the centre of NGC 7538-IRS 1 for different Formaldehyde abundances. The molecular gas has a density $n(\text{H}_2) = 10^4 \text{ cm}^{-3}$ and a temperature $T_k = 20 \text{ K}$

In Fig. 2 we show the results of a more realistic calculation where we have included 14 levels of ortho- H_2CO and where the effects of collisions with H_2 molecules and of radiation trapping are also taken into account. The parameters characterizing this calculation are chosen as follows: a dilution factor $W = 0.5$, a H_2 density $n(\text{H}_2) = 3 \cdot 10^4 \text{ cm}^{-3}$, a gas kinetic temperature $T_k = 20 \text{ K}$,

a Formaldehyde abundance $x(\text{H}_2\text{CO}) = n(\text{H}_2\text{CO})/n(\text{H}_2) = 4 \cdot 10^{-7}$ and a velocity gradient $dv/dr = 1280 \text{ km s}^{-1} \text{ pc}^{-1}$ ($\alpha = 0$, see Eq. (5)) corresponding to the observed line separation over the distance between the two maser spots.

From a comparison of Figs. 1 and 2 (curve 1) it is clear that including more Formaldehyde levels increases the population inversion of the ground state doublet and shifts the maximum inversion towards smaller values of λ_0 (more compact H II regions). Both effects are caused by the contributions to the pumping of the higher mm transitions in the H_2CO ladder. Including the effect of H_2 collisions narrows the range of λ_0 where population inversion occurs (curve 2). Taking in addition radiation trapping into account somewhat reduces the population inversion (curve 3). It also follows from Fig. 2 that for $\lambda_0 \leq 3 \text{ cm}$ both the $1_{10}-1_{11}$ transition (curve 3) and the $2_{11}-2_{12}$ transition (curve 4) are inverted while for $3 \text{ cm} \leq \lambda_0 \leq 10 \text{ cm}$ only the $1_{10}-1_{11}$ transition is inverted.

4. A Model for the Formaldehyde Maser near NGC 7538

We suppose that the H_2CO maser near NGC 7538 is located in dense cold molecular gas and that it is pumped by the free-free continuum radiation field of the compact H II region NGC 7538-IRS 1. Assuming spherical geometry one finds from its observed radio spectrum the physical parameters listed in Table 2. The free-free continuum optical depth becomes unity at $\lambda_0 = 2.7 \text{ cm}$ ($E_m = 6 \cdot 10^6 \text{ cm}^{-6} \text{ pc}$) so that we expect our radio continuum pump mechanism to operate (see Fig. 2).

The observed similarity of the 1720 MHz OH emission spectrum and the H_2CO emission spectrum suggests that both masers originate in the same gas cloud (cf. Forster et al., 1980). According to Elitzur (1976) the 1720 MHz maser is collision-dominated and it is most efficiently pumped for molecular Hydrogen densities in the range $10^3-10^5 \text{ cm}^{-3}$ and gas temperatures in the range 15-200 K. This and the fact that the lowest rotational levels of H_2CO thermalize at densities $n(\text{H}_2) \geq 3 \cdot 10^2 \text{ cm}^{-3}$ suggest that both masers are located in a molecular cloud outside the thin dense shell of compressed gas at the interface of the H II region and the molecular cloud (Mathews and O'Dell, 1969). Because this shell is very thin ($\sim 0.01 R_{II}$) and transparent to radio continuum radiation its presence is neglected in our model. The far-infrared radiation field generated in the shell and in the molecular cloud surrounding NGC 7538-IRS 1 (Willner, 1976; Werner et al., 1979) is also neglected because the free-free radio continuum radiation field dominates at all wavelengths of interest.

The H II region is optically thick at both 6 cm and at 18 cm so that the H_2CO and the OH lines must originate in molecular gas in front of the H II region and the observed high brightness temperatures are due to amplification of the radio continuum radiation of the H II region.

A comparison of the radial velocities of the H 110 α recombination line ($-66 \pm 5 \text{ km s}^{-1}$, Forster et al., 1980), of the H_2CO maser lines (-57.6 and -59.8 km s^{-1} , Forster et al., 1980) and of several other molecular lines ($\sim -57 \text{ km s}^{-1}$, CO observed by Wilson et al., 1974 and HCN, CN, HCO^+ , and N_2H^+ by Turner and Thaddeus, 1977) indicates that NGC 7538-IRS 1 is moving through the molecular cloud with a velocity of about 10 km s^{-1} towards us.

Led by the observations discussed above we assume that the compact H II region NGC 7538-IRS 1 moves through or expands into a molecular cloud with a velocity $V \approx 10 \text{ km s}^{-1}$. We further assume that it moves towards us and that the molecular gas in front of the H II region is pushed sideways. The velocity field in the




Fig. 5. Same as Fig. 4 for $n(\text{H}_2) = 3 \cdot 10^4 \text{ cm}^{-3}$

gas may then be approximated by

$$\frac{d\eta}{ds} = \frac{V}{R_H} (1 - \mu^2) \quad (18)$$

where μ is the cosine of the angle between line element ds and the line of sight. An attempt to give a pictorial representation of our model is shown in Fig. 3. The velocity gradient is zero (coherence) along the line of sight ($\mu=1$) and is maximal (V/R_H) perpendicular to the line of sight ($\mu=0$) in the direction in which the molecular gas is streaming away. The fact that the local approximation that we have used to solve the radiative transfer (cf. Sect. 2) breaks down in the radial direction does not invalidate our results because the solid angle over which $V(1-\mu^2) \leq v_{th}$ is small (~ 0.1 steradian), much smaller than the solid angle subtended by the H II region (cf. de Jong et al., 1975).

We have solved the line transfer problem for 14-level ortho- H_2CO molecules as outlined in Sect. 2 by putting $r=R_H$, $v(r)=V$ and $\alpha=1$ in Eq. (4). In Fig. 4 we present the variation of the relative population inversion

$$\eta = (n_3 g_2 / g_1 - n_2) / n(\text{H}_2\text{CO}) \quad (19)$$

with distance r from the centre of the H II region for several values of the H_2CO abundance $x(\text{H}_2\text{CO}) = n(\text{H}_2\text{CO}) / n(\text{H}_2)$. The results are obtained for $n(\text{H}_2) = 10^4 \text{ cm}^{-3}$ and $T_k = 20 \text{ K}$ using the parameters of NGC 7538-IRS 1 listed in Table 2 to characterize the continuum radiation field. The relative population inversion decreases with increasing distance because the radio continuum pumping becomes less efficient. It decreases with increasing H_2CO abundance because optical depth effects in the millimeter lines reduce the pump rate. Figure 5 illustrates the effects of thermalization by collisions. It shows the results for the same parameters but for a molecular Hydrogen density $n(\text{H}_2) = 3 \cdot 10^4 \text{ cm}^{-3}$. Compared with Fig. 4 the population inversion and the thickness of the inversion layer are both smaller by about a factor 2.

The brightness temperature of the H_2CO line can be calculated from

$$T_b = T_c \{ \exp(-\tau) - 1 \} \quad (20)$$

where $T_c = 10^4 \text{ K}$ is the brightness temperature of the radio continuum radiation of the H II region and

$$\tau = 3.4 \cdot 10^{-8} \bar{\eta} n(\text{H}_2\text{CO}) \frac{dl}{v_{th}} \quad (21)$$

is the 6 cm H_2CO line optical depth [see Eq. (4)]. In Eq. (21) $\bar{\eta}$ is the relative population difference [Eq. (19)] averaged over the thickness of the inversion layer dl and $v_{th} = 0.1 \text{ km s}^{-1}$ is the thermal velocity of H_2CO molecules. To reproduce the observed brightness temperatures in the H_2CO maser lines of $\sim 10^4 \text{ K}$ a line optical depth $\tau \approx -5$ is required. According to Fig. 4 this can be achieved for $n(\text{H}_2) = 10^4 \text{ cm}^{-3}$ where we find $\bar{\eta} \approx 0.006$ and $dl \approx 1.2 R_H$ if $x(\text{H}_2\text{CO}) = 8 \cdot 10^{-7}$.

To show that this Formaldehyde abundance can be obtained in dense dark clouds we now derive an analytic expression for the H_2CO abundance from a discussion of its chemical formation and destruction processes. For more details of dark cloud chemistry the reader is referred to the paper by de Jong et al. (1980).


The formation of H_2CO is initiated by the radiative association reaction



leading to CH_2^+ after collisions with H_2 . At large densities $n(\text{H}_2) \geq 10^4 \text{ cm}^{-3}$ CH_2^+ is removed from the gas by



rather than by recombination. CH_3^+ is predominantly destroyed by dissociative recombination leading to CH_3 and CH_4 upon recombination. CH_3 is destroyed by collisions with C^+ and H_2^+ and by the reaction



We assume that Formaldehyde is predominantly removed from the gas by collisions with HCO^+ , H_3^+ , and C^+ . At large densities and for $A_{ul} \geq 2$ the formation of C^+ is initiated by cosmic ray ionization of He atoms followed by the reaction



and the C^+ ions are mainly destroyed by reaction (a) as long as $k_2 > 1 \cdot 10^{-9} n(\text{O}_2) / n(\text{H}_2)$.

From our models of molecular clouds (de Jong et al., 1980) we found that in dense dark clouds all Hydrogen is in H_2 , that all Carbon is in the form of CO and that approximately 1/3 of the remaining Oxygen atoms are in the form of atomic Oxygen and 2/3 in the form of O_2 . Assuming that electrons are predominantly produced by cosmic ray ionization of molecular Hydrogen and removed from the gas by dissociative recombination with molecular ions like HCO^+ and H_3^+ one finds

$$n(e) = 1.6 \cdot 10^{-4} \left(\frac{\zeta}{5 \cdot 10^{-18} \text{ s}^{-1}} \right)^{1/2} \left(\frac{n(\text{H}_2)}{10^4 \text{ cm}^{-3}} \right)^{1/2} \text{ cm}^{-3} \quad (22)$$

for a gas temperature of $T=20 \text{ K}$ where ζ is the cosmic ray ionization rate of H_2 . Using a H_2CO destruction rate due to collisions with molecular ions of $2 \cdot 10^{-9} n(e) \text{ s}^{-1}$ we finally obtain the following expression for the Formaldehyde abundance

$$\frac{n(\text{H}_2\text{CO})}{n(\text{H}_2)} = 8 \cdot 10^{-7} \left(\frac{\zeta}{5 \cdot 10^{-18} \text{ s}^{-1}} \right)^{1/2} \left(\frac{10^4 \text{ cm}^{-3}}{n(\text{H}_2)} \right)^{1/2} \quad (23)$$

where we have assumed that CH_3 is predominantly destroyed by reaction (c). This is the case as long as Carbon and Oxygen are not more depleted than about a factor 100 compared to their

cosmical abundances. For our chemical scheme to work we need a reaction rate coefficient of reaction (a): $k_a \geq 5 \cdot 10^{-15} \text{ cm}^3 \text{ s}^{-1}$ in good agreement with a theoretical calculation of Herbst et al. (1977).

Sofar we have neglected photodestruction of Formaldehyde. Because the extinction in the compressed shell between the shock front and the ionization front around the compact H II region is quite large the UV radiation of the O6 star exciting NGC 7538-IRS 1 is completely absorbed in the shell (Willner, 1976). The O7 star exciting the optical nebula NGC 7538 (cf. Israel et al., 1973) at a projected distance of about 4 pc from NGC 7538-IRS 1 is another possible source of UV radiation. From Kurucz (1979) we estimate that the radiation field of an O7 star with a radius of $10 R_\odot$ at 4 pc distance is about 300 times as intense as the average interstellar radiation field at 1300 \AA (Habing, 1968). For photodestruction to be negligible (less than 10^{-13} s^{-1}) we require that the maser is formed at depth $A_v \geq 10$ into the molecular cloud (cf. de Jong et al., 1980).

We conclude that our model can explain the maser amplification required to reproduce the observed brightness temperatures. Our model predicts a maser spot size with typical dimensions $\sim 2 R_{II} \text{ arc cos} \{(1 - v_{rel}/V)^{1/2}\} \approx 5 \cdot 10^{15} \text{ cm}$ consistent with the observed upper limit. From our calculations we further find that the induced emission rate in the maser transition is approximately equal to the rate of population exchange in the millimeter lines so that the maser is close to saturation. This explains why both maser features have about the same intensity. Our model does not account for the fact that there are two maser spots separated by about $0.1 (5 \cdot 10^{15} \text{ cm})$ in position and by about 2 km s^{-1} in velocity. Special circumstances presumably due to gas dynamical effects (fluctuations in the gas density and/or the velocity field) are required to explain these facts.

Our model can also be used to predict a brightness temperature of the $2_{11}-2_{12}$ line of formaldehyde. From our numerical results we obtain an optical depth in that line of ~ -1 so that we predict a brightness temperature of $\sim 2 \cdot 10^6 \text{ K}$ consistent with the upper limit of $5 \cdot 10^6 \text{ K}$ that can be derived from the $\text{H}_2\text{CO } 2 \text{ cm}$ line observations of NGC 7538-IRS 1 by Evans et al. (1975) for a source size of 0.1 .

Acknowledgements. We would like to thank Dr. J. R. Foster and Dr. W. M. Goss for communicating their results prior to publication and for several enlightening discussions. The research of W. B. is supported by the Netherlands Foundation for Astronomical Research (ASTRON) with financial aid from the Netherlands Organization for the Advancement of Pure Research.

References

- de Jong, T., Chu, S.I., Dalgarno, A.: 1975, *Astrophys. J.* **199**, 69
 de Jong, T., Dalgarno, A., Boland, W.: 1980, *Astron. Astrophys.* **91**, 68
 Dieter, N.H.: 1973, *Astrophys. J.* **183**, 449
 Elitzur, M.: 1976, *Astrophys. J.* **203**, 124
 Elitzur, M., Goldreich, P., Scoville, N.: 1976, *Astrophys. J.* **205**, 384
 Evans, N.J., Zuckerman, B., Morris, G., Sato, T.: 1975, *Astrophys. J.* **196**, 433
 Forster, J.R., Goss, W.M., Wilson, T.L., Downes, D., Dickel, H.R.: 1980, *Astron. Astrophys.* **84**, L1
 Garrison, B.J., Lester, W.A., Miller, W.H., Green, S.: 1975, *Astrophys. J. Letters* **200**, L175
 Green, S., Garrison, B.J., Lester, W.A., Miller, W.H.: 1978, *Astrophys. J. Suppl. Ser.* **37**, 321
 Habing, H.J.: 1968, *Astron. Inst. Neth.* **19**, 421
 Harris, S., Scott, P.F.: 1976, *Monthly Notices Roy. Astron. Soc.* **175**, 371
 Herbst, E., Schubert, J.G., Certain, P.R.: 1977, *Astrophys. J.* **213**, 696
 Israel, F.P., Habing, H.J., de Jong, T.: 1973, *Astron. Astrophys.* **27**, 143
 Kurucz, R.L.: 1979, *Astrophys. J. Suppl. Ser.* **40**, 1
 Mathews, W.G., O'Dell, C.R.: 1969, *Ann. Rev. Astron. Astrophys.* **7**, 67
 Minn, Y.K., Greenberg, J.M.: 1973, *Astron. Astrophys.* **22**, 13
 Nerf, R.B.: 1972, *Astrophys. J.* **174**, 467
 Rots, A.H., Dickel, H.R., Forster, J.R., Goss, W.M.: 1980 (in preparation)
 Spitzer, L.: 1978, *Physical processes in the interstellar medium*, Wiley, New York
 Schwendemann, R.H., Laurie, V.W.: 1958, *Tables of line strength for rotational transitions of asymmetric rotor molecules*, Pergamon Press, New York
 Thaddeus, P.: 1972, *Astrophys. J.* **173**, 317
 Townes, C.H., Cheung, A.C.: 1969, *Astrophys. J. Letters* **157**, L103
 Townes, C.H., Schawlow, A.L.: 1955, *Microwave Spectroscopy*, McGraw-Hill, New York
 Turner, B.E., Thaddeus, P.: 1977, *Astrophys. J.* **211**, 755
 Werner, M.W., Becklin, E.E., Gatley, I., Mathews, K., Neugebauer, G., Wynn-Williams, C.G.: 1979, *Monthly Notices Roy. Astron. Soc.* **188**, 463
 Wilson, W.J., Schwartz, P.R., Epstein, E.E., Johnson, W.A., Etcheverry, R.D., Mori, T.T., Berry, G.G., Dyson, H.B.: 1974, *Astrophys. J.* **191**, 357
 Winnewisser, G., Mezger, P.G., Breuer, H.D.: 1974, *Topics in current chemistry*, Vol. 44, Springer, Berlin, Heidelberg, New York