

## Supporting Information

# Oxidative modifications of proteins and lipids of dried semen, urine and saliva stains as a function of age in forensic context

Nihad Achetib <sup>1,2†</sup>, Rosa E. Otto <sup>1†</sup>, Maurice C.G. Aalders <sup>1,2,3</sup> and Annemieke van Dam<sup>1,2,4,\*</sup>

<sup>1</sup> Amsterdam University Medical Centers (UMC), location AMC, University of Amsterdam, Biomedical Engineering and Physics, Meibergdreef 9, Amsterdam, The Netherlands.

<sup>2</sup> Amsterdam Public Health, Methodology, Amsterdam, The Netherlands.

<sup>3</sup> Co van Ledden Hulsebosch Center (CLHC), University of Amsterdam, Amsterdam, The Netherlands.

<sup>4</sup> Amsterdam University of Applied Science, Forensic Science, Tafelbergweg 51, Amsterdam, The Netherlands.

<sup>†</sup> Both authors contributed equally

\* **Correspondence:** [Annemiekevandam@amsterdamumc.nl](mailto:Annemiekevandam@amsterdamumc.nl), Tel.: +31 20 566 4390.




Figure S1: Distribution of the fluorescent ratios (.1) and the protein carbonyls concentrations (.2) of each donor for semen, urine and saliva. The whiskers indicate the range from the smallest to the largest value.