



UvA-DARE (Digital Academic Repository)

Draft genome of the marine bacterium *Alteromonas gracilis* strain J4 isolated from the green coenocytic alga *Caulerpa prolifera*

van Duijnhoven, H.J.; Dombrowski, N.; Kuperus, P.; Aires, T.; Coelho, M.A.G.; Silva, J.; Muyzer, G.; Engelen, A.H.

DOI

[10.1128/mra.00184-24](https://doi.org/10.1128/mra.00184-24)

Publication date

2024

Document Version

Final published version

Published in

Microbiology Resource Announcements

License

CC BY

[Link to publication](#)

Citation for published version (APA):

van Duijnhoven, H. J., Dombrowski, N., Kuperus, P., Aires, T., Coelho, M. A. G., Silva, J., Muyzer, G., & Engelen, A. H. (2024). Draft genome of the marine bacterium *Alteromonas gracilis* strain J4 isolated from the green coenocytic alga *Caulerpa prolifera*. *Microbiology Resource Announcements*, 13(7), Article e0018424. <https://doi.org/10.1128/mra.00184-24>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.
UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Draft genome of the marine bacterium *Alteromonas gracilis* strain J4 isolated from the green coenocytic alga *Caulerpa prolifera*

Hannah J. van Duijnhoven,¹ Nina Dombrowski,² Peter Kuperus,² Tânia Aires,¹ Márcio A.G. Coelho,¹ João Silva,¹ Gerard Muyzer,² Aschwin H. Engelen¹

AUTHOR AFFILIATIONS See affiliation list on p. 3.

ABSTRACT Here, we present the draft genome sequence of *Alteromonas gracilis* strain J4, isolated from the green macroalga *Caulerpa prolifera*. The draft genome is 4,492,914 bp in size and contains 4,719 coding DNA sequences, 67 tRNAs, and 16 rRNA-coding genes. Strain J4 may exhibit host growth-promoting properties.

KEYWORDS plant-microbe interactions, whole-genome sequencing, marine microbiology, growth-promoting bacteria, interspecific competition

To understand *Caulerpa* host-microbe interaction(1,2), we isolated bacteria from *C. prolifera* rhizoid tissue and conducted whole-genome sequencing. Here, we present the draft genome sequence of *Alteromonas gracilis* strain J4.

Strain J4 was isolated from rhizoids of hand-collected *C. prolifera* in the Ria Formosa lagoon (37°00'22.7"N 7°58'00.3"W, Faro, Portugal) and stored in a cooling box. Rhizoids were ground with mortar and pestle, and the lysate was plated on Difco Marine Agar 2216 and incubated in the dark at room temperature (20–25°C). After 3 days, individual colonies were replated and re-incubated. Isolate J4 was identified as an *Alteromonas* sp. by comparative full-length 16S rRNA gene Sanger sequencing (Applied BioSystems 3130xl Genetic Analyzer) analysis using primers 27F/1492R against the NCBI database (3). The 16S rRNA gene showed 96.80% sequence identity to *A. gracilis* strain 9a2 341. Genomic DNA was extracted using the peqGOLD Bacterial DNA Mini Kit (VWR). Genome sequencing was conducted on a MinION Mk1C device (Oxford Nanopore Technologies), using the Ligation Sequencing Kit (SQK-LSK110) and a Flongle Flow Cell (R9.4.1). Base-calling was performed using Guppy v6.2.7 (<https://community.nanopore-tech.com/downloads>). 225,773 reads passed quality control with a mean Q-score of 12 and an N50 of 3.71 kb.

Trimming residual sequencing adaptors and splitting chimeric reads [Porechop v0.2.4 (4)] resulted in 211,623 reads. Filtlong v0.2.1 was used to remove small (<1,000 bp) and poor quality (<5%) reads (<https://github.com/rrwick/Filtlong>). For the remaining 172,018 reads, 12 subsamples were generated at 55× depth using Tricycler v0.5.5 (5). Three subsamples were each assembled with (i) Flye v2.9.3, (ii) Miniasm v0.3 & Minipolish v0.1.3, (iii) Raven v1.8.3, and (iv) Unicycler v0.5.0 (6–10). The consensus assembly was further generated using Tricycler v0.5.5 and polished with Homopolish v0.4.1 (11), resulting in one circular contig with a total size of 4,492,914 bp (131× coverage) and a GC content of 44.0%. The genome was reoriented using Dnaaplcr chromosome v0.7.0 (12) by identifying *dnaA* as the replication initiator gene. The genome had a completeness of 98% and contamination of 5.1% (CheckM2 v1.0.2) (13) and contained 4,719 protein-coding genes, 67 tRNA, and 16 rRNA coding genes [Prokka v1.14.6 (14); Fig. 1]. 903 genes were identified as potential pseudogenes (<https://github.com/>

Editor Frank J. Stewart, Montana State University, Bozeman, Montana, USA

Address correspondence to Hannah J. van Duijnhoven, juliavanduijnhoven@msn.com.

The authors declare no conflict of interest.

Received 24 February 2024

Accepted 24 May 2024

Published 11 June 2024

Copyright © 2024 van Duijnhoven et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).




FIG 1 The genome map of *Alteromonas gracilis* strain J4. Each circle from inner to outer indicates potential plant growth-promoting (PGP) genes, coding sequences (CDS) in the leading strand, CDS in the lagging strand, tRNA and rRNA, GC content, GC skew+, and GC skew-. Potential PGP genes are indicated in red and labeled with EC numbers. If not labeled, tryptophan halogenases are represented.

[ndombrowski/j4_assembly](#)) using Pseudofinder v1.1.0 with the UniProtKB/Swiss-Prot database as reference (15, 16). J4 likely belongs to an uncharacterized species within the genus *Alteromonas*. The genome exhibits 85.3% average nucleotide identity (ANI) with *Alteromonas* sp009811495 ([GCF_016756315.1](#)) based on a comparison with the GTDB r214 database using GTDB-Tk v2.3.2 (17, 18).

The annotation analysis identified two genes encoding 4-hydroxyphenylpyruvate dioxygenase (EC 1.13.11.27), pivotal in melanin catalysis (19). Melanin plays a role in ensuring survival during symbiotic interactions (20). Protein genes *cspD*, *dnaJ*, *dnaK*, and

grpE, described to protect against cold/heat and oxidative stress, were detected (21), as well as 12 genes related to sulfur metabolism (22). Four indole-3-glycerol phosphate synthases (EC 4.1.1.48) and two tryptophan 2,3-dioxygenases (EC 1.13.11.11), both key precursors in indole-3-acetic acid biosynthesis were found. In all, 24 putative genes encoding tryptophan halogenases and six tryptophan synthases were found, suggesting potential growth-promoting properties in strain J4 with biotechnological applications (23, 24).

ACKNOWLEDGMENTS

The research leading to the results presented in this publication was financially supported by CCMAR under reference number CCMAR/BD/07/2022 for H.D., CEE-CINST/00114/2018 for AE, and funded by The BlueForests project of EEA Grants (PT-INNOVATION-0081). T.A. was supported by the fellowship reference—SFRH/BPD/116774/2016 from FCT and M.C. was supported by FCT (DivRestore/0013/2020). G.M. and P.K. were supported by the 2020–2021 Biodiversa+ and water JPI joint call for research projects, under the BiodivRestore ERA-NET Cofund (GA N°101003777) with the EU and the Dutch Ministry of Agriculture, Nature and Food Quality. This study received Portuguese national funds from FCT—Foundation for Science and Technology through projects UIDB/04326/2020 (DOI:10.54499/UIDB/04326/2020), UIDP/04326/2020 (DOI:10.54499/UIDP/04326/2020), and LA/P/0101/2020 (DOI:10.54499/LA/P/0101/2020).

AUTHOR AFFILIATIONS

¹Centre of Marine Sciences (CCMAR/CIMAR LA), Universidade do Algarve, Campus de Gambelas, Faro, Portugal

²Microbial Systems Ecology, Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, the Netherlands

AUTHOR ORCIDs

Hannah J. van Duijnhoven  <http://orcid.org/0000-0003-1519-6889>

Nina Dombrowski  <http://orcid.org/0000-0003-1917-2577>

Tânia Aires  <http://orcid.org/0000-0002-1964-6819>

Márcio A.G. Coelho  <http://orcid.org/0000-0001-7832-5241>

João Silva  <http://orcid.org/0000-0002-7211-1661>

Gerard Muyzer  <http://orcid.org/0000-0002-2422-0732>

Aschwin H. Engelen  <http://orcid.org/0000-0002-9579-9606>

DATA AVAILABILITY

The 16S rRNA sequence data, raw Nanopore sequence reads, and the assembled genome sequence have been deposited in GenBank under BioProject number [PRJNA1077798](https://ncbi.nlm.nih.gov/bioproject/PRJNA1077798), with BioSample accession numbers [PP541516](https://ncbi.nlm.nih.gov/biosample/PP541516), [SAMN40604929](https://ncbi.nlm.nih.gov/biosample/SAMN40604929), and [SAMN39982826](https://ncbi.nlm.nih.gov/biosample/SAMN39982826), respectively, and the reported genome is the second version, [CP145482.2](https://ncbi.nlm.nih.gov/biosample/CP145482.2).

REFERENCES

- Alexandre A, Santos R. 2020. High nitrogen and phosphorous acquisition by belowground parts of *Caulerpa prolifera* (Chlorophyta) contribute to the species' rapid spread in Ria Formosa lagoon, Southern Portugal. *J Phycol* 56:608–617. <https://doi.org/10.1111/jpy.12988>
- Parreira F, Martinez-Crego B, Lourenço Afonso CM, Machado M, Oliveira F, Manuel dos Santos Gonçalves J, Santos R. 2021. Biodiversity consequences of *Caulerpa prolifera* takeover of a coastal lagoon. *Estuar Coast Shelf Sci* 255:107344. <https://doi.org/10.1016/j.ecss.2021.107344>
- Johnson M, Zaretskaya I, Raytselis Y, Merezuk Y, McGinnis S, Madden TL. 2008. NCBI BLAST: a better web interface. *Nucleic Acids Res* 36:W5–9. <https://doi.org/10.1093/nar/gkn201>
- Wick R. 2017. Porechop: adapter trimmer for Oxford Nanopore reads. Available from: <https://github.com/rrwick/Porechop/>
- Wick RR, Judd LM, Cerdeira LT, Hawkey J, Méric G, Vezina B, Wyres KL, Holt KE. 2021. Tricycler: consensus long-read assemblies for bacterial genomes. *Genome Biol* 22:266. <https://doi.org/10.1186/s13059-021-02483-z>
- Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol* 37:540–546. <https://doi.org/10.1038/s41587-019-0072-8>

7. Li H. 2016. Minimap and miniasm: fast mapping and *de novo* assembly for noisy long sequences. *Bioinformatics* 32:2103–2110. <https://doi.org/10.1093/bioinformatics/btw152>
8. Wick RR, Holt KE. 2019. Benchmarking of long-read assemblers for prokaryote whole genome sequencing. *F1000Res* 8:2138. <https://doi.org/10.12688/f1000research.21782.4>
9. Vaser R, Šikić M. 2021. Time-and memory-efficient genome assembly with Raven. *Nat Comput Sci* 1:332–336. <https://doi.org/10.1038/s43588-021-00073-4>
10. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>
11. Huang YT, Liu PY, Shih PW. 2021. Homopolish: a method for the removal of systematic errors in nanopore sequencing by homologous polishing. *Genome Biol* 22:95. <https://doi.org/10.1186/s13059-021-02282-6>
12. Bouras G, Grigson SR, Papudeshi B, Mallawaarachchi V, Roach MJ. 2024. Dnaapler: a tool to reorient circular microbial genomes. *J Open Source Softw* 9:5968. <https://doi.org/10.21105/joss.05968>
13. Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2023. CheckM2: a rapid, scalable and accurate tool for assessing microbial genome quality using machine learning. *Nat Methods* 20:1203–1212. <https://doi.org/10.1038/s41592-023-01940-w>
14. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>
15. Syberg-Olsen MJ, Garber AI, Keeling PJ, McCutcheon JP, Husnik F. 2022. Pseudofinder: detection of pseudogenes in prokaryotic genomes. *Mol Biol Evol* 39:msac153. <https://doi.org/10.1093/molbev/msac153>
16. UniProt Consortium. 2023. Uniprot: the universal protein knowledge-base in 2023. *Nucleic Acids Res* 51:D523–D531. <https://doi.org/10.1093/nar/gkac1052>
17. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify genomes with the genome taxonomy database. *Bioinformatics* 36:1925–1927. <https://doi.org/10.1093/bioinformatics/btz848>
18. Parks DH, Chuvochina M, Chaumeil PA, Rinke C, Mussig AJ, Hugenholtz P. 2020. A complete domain-to-species taxonomy for bacteria and archaea. *Nat Biotechnol* 38:1079–1086. <https://doi.org/10.1038/s41587-020-0501-8>
19. Chen J, Wang X, Zhu S, Chen Y, Yang J. 2016. Complete genome sequence of *Alteromonas stellipolaris* LMG 21856, a budding brown pigment-producing oligotrophic bacterium isolated from the Southern Ocean. *Genome Announc* 4:10–1128. <https://doi.org/10.1128/genomeA.00137-16>
20. Pavan ME, López NI, Pettinari MJ. 2020. Melanin biosynthesis in bacteria, regulation and production perspectives. *Appl Microbiol Biotechnol* 104:1357–1370. <https://doi.org/10.1007/s00253-019-10245-y>
21. Torres M, Hong KW, Chong TM, Reina JC, Chan KG, Dessaux Y, Llamas I. 2019. Genomic analyses of two *Alteromonas stellipolaris* strains reveal traits with potential biotechnological applications. *Sci Rep* 9:1215. <https://doi.org/10.1038/s41598-018-37720-2>
22. Guédon E, Martin-Verstraete I. 2006. Cysteine metabolism and its regulation in bacteria, p 195–218. In *Amino acid biosynthesis~pathways, regulation and metabolic engineering*. Springer, Berlin, Heidelberg.
23. Gago JF, Viver T, Urdiain M, Pastor S, Kämpfer P, Ferreira E, Rossello-Mora R. 2021. Description of three new *Alteromonas* species *Alteromonas antoniana* sp. nov., *Alteromonas lipotruaeae* sp. nov. and *Alteromonas lipotruaeiana* sp. nov. isolated from marine environments, and proposal for reclassification of the genus *Salinimonas* as *Alteromonas*. *Syst Appl Microbiol* 44:126226. <https://doi.org/10.1016/j.syapm.2021.126226>
24. Navarro-Torre S, Carro L, Rodríguez-Llorente ID, Pajuelo E, Caviedes MÁ, Igual JM, Klenk H-P, Montero-Calasanz MDC. 2020. *Pseudoalteromonas rhizosphaerae* sp. nov., a novel plant growth-promoting bacterium with potential use in phytoremediation. *Int J Syst Evol Microbiol* 70:3287–3294. <https://doi.org/10.1099/ijsem.0.004167>