



UvA-DARE (Digital Academic Repository)

Harnessing the potential of the microbial sulfur cycle for environmental biotechnology

Gupta, S.; Plugge, C.M.; Muyzer, G.; Sánchez-Andrea, I.

DOI

[10.1016/j.copbio.2024.103164](https://doi.org/10.1016/j.copbio.2024.103164)

Publication date

2024

Document Version

Final published version

Published in

Current Opinion in Biotechnology

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Gupta, S., Plugge, C. M., Muyzer, G., & Sánchez-Andrea, I. (2024). Harnessing the potential of the microbial sulfur cycle for environmental biotechnology. *Current Opinion in Biotechnology*, 88, Article 103164. <https://doi.org/10.1016/j.copbio.2024.103164>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Review

Harnessing the potential of the microbial sulfur cycle for environmental biotechnology

Suyash Gupta^{1,2,*}, Caroline M Plugge^{1,3,#}, Gerard Muyzer^{2,\$} and Irene Sánchez-Andrea^{4,†}



The sulfur cycle is a complex biogeochemical cycle characterized by the high variability in the oxidation states of sulfur. While sulfur is essential for life processes, certain sulfur compounds, such as hydrogen sulfide, are toxic to all life forms. Micro-organisms facilitate the sulfur cycle, playing a prominent role even in extreme environments, such as soda lakes, acid mine drainage sites, hot springs, and other harsh habitats. The activity of these micro-organisms presents unique opportunities for mitigating sulfur-based pollution and enhancing the recovery of sulfur and metals. This review highlights the application of sulfur-oxidizing and -reducing micro-organisms in environmental biotechnology through three illustrative examples. Additionally, it discusses the challenges, recent trends, and prospects associated with these applications.

Addresses

¹ Wetsus, European Centre of Excellence for Sustainable Water Technology, Leeuwarden, the Netherlands

² Microbial Systems Ecology, Department of Freshwater and Marine Ecology, Institute of Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, the Netherlands

³ Laboratory of Microbiology, Wageningen University & Research, Wageningen, the Netherlands

⁴ Environmental Science for Sustainability Department, IE Universidad, Segovia, Spain

Corresponding author: Muyzer, Gerard (g.muijzer@uva.nl)

* <https://orcid.org/0000-0002-7774-4636>

<https://orcid.org/0000-0002-3391-7742>

\$ <https://orcid.org/0000-0002-2422-0732>

† <https://orcid.org/0000-0001-6977-3026>

Current Opinion in Biotechnology 2024, 88:103164

This review comes from a themed issue on **Environmental Biotechnology**

Edited by **Mari Winkler** and **George Wells**

For complete overview of the section, please refer to the article collection, "[Environmental Biotechnology \(2024\)](#)"

Available online 2 July 2024

<https://doi.org/10.1016/j.copbio.2024.103164>

0958-1669/© 2024 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

The sulfur cycle involves a series of oxidized and reduced sulfur states that can occur through chemical and biological processes. The sulfur atom cycles between the oxidation states +6 in sulfate and -2 in sulfide (Box 1). Sulfur molecules act as electron donors for aerobic respiration, electron acceptors for anaerobic respiration and anaerobic light-dependent CO₂ reduction, and fermentation substrates [1]. In natural environments, sulfate-reducing micro-organisms (SRM) contribute 29% of organic matter sequestration and play an important role in the biogeochemical cycling of sulfur [2]. SRM use sulfate as an electron acceptor in utilizing both organic and inorganic compounds [3].

Approximately 80–90% of sulfide formed can be re-oxidized to various intermediates or completely oxidized to sulfate by sulfide- and sulfur-oxidizing micro-organisms (SOM). The remaining sulfide is deposited as iron complexes (e.g. FeS, FeS₂) [2]. The intermediates, such as elemental sulfur, sulfite, and thiosulfate, can undergo further oxidation to sulfate by SOM, reduction to sulfide by thiosulfate reducing, sulfur-reducing and sulfite-reducing micro-organisms or disproportionation to sulfide and sulfate by sulfur-disproportionating micro-organisms (SPM). Although organo-sulfur compounds are prevalent in natural environments, this review focuses on inorganic sulfur compounds.

In man-made systems, such as wastewater treatment plants processing industrial waters, the mineralization or anaerobic digestion of organic compounds by micro-organisms is an important process. For industries that utilize sulfate or sulfuric acid, the water treatment leads to sulfide formation via the activity of SRM. Sulfide is toxic and highly corrosive because it and other reduced sulfur compounds, such as bisulfide and persulfide, react with metals, forming metal sulfides, consequently leading to corrosion [4•].

Sulfur-related micro-organisms have been successfully employed to mitigate industrial and environmental issues [5•]. Specific micro-organisms have been isolated from diverse environments, physiologically characterized, and further studied to understand the molecular mechanisms involved

in sulfur conversions [6,7]. Additionally, applying state-of-the-art omics techniques has identified novel taxa of sulfate reducers and sulfide oxidizers from different environments and revealed insights into their genetic diversity, novel functional genes, and metabolic pathways [8–12•].

In this review, we present three examples of sulfur bacteria's application in environmental biotechnology (Box 1) and discuss the challenges and prospects associated with these applications.

Example 1: biodesulfurization of sulfide-containing gas streams

Haloalkaline biodesulfurization (BD) is an eco-friendly, cost-effective process that converts toxic sulfide gas to elemental sulfur using haloalkaliphilic SOM under microoxic conditions [8,13]. The microoxic conditions inhibit the complete oxidation of sulfide to sulfate (Box 1), which is energetically preferred by chemolithoautotrophic SOMs under ambient oxygen supply and control the formation of sulfur (Box 1). Some notable SOMs include obligated chemolithoautotrophs such as *Thiomicrospira*, *Thioalkalivibrio*, *Thioalkalibacter*, *Thiohalophilus*, and facultative autotrophs such as *Alkalilimnicola*, *Alkalispirillum* [14,15].

The generated sulfur is a nontoxic compound with several downstream applications [16]. Commercially known as 'Thiopaq', the process removes >99% of sulfide from sour gas streams [17]. The process involves (1) an absorber column for absorbing sulfide gas in a haloalkaline solution, (2) a bioreactor for oxidizing sulfide to elemental sulfur, and (3) a sulfur recovery section for retrieving the elemental sulfur particles from the process solution [17]. The process converts ~90% of sulfide to sulfur and the rest to sulfate and thiosulfate. Unlike sulfur, these latter compounds are undesirable as they decrease the alkalinity, increasing caustic consumption and bleed water formation.

A recent innovation, 'Thiopaq Ultra', is being studied at the pilot scale ([18]), and it incorporates an additional anoxic bioreactor between the absorber and oxic bioreactor (Figure 1, [18]). The new configuration has a higher selectivity for sulfur formation (~97%) and depicts a ~90% reduction in caustic, make-up water consumption and bleed formation [13,18]. This leads to a significant decrease in the operational costs of the BD process. In addition, the operational stability is enhanced due to a better ability for sulfur to settle, resulting in easier separation and, consequently, a better sulfur recovery [18,19••].

Molecular biological analyses have elucidated the presence and activity of a shared microbial community in both full- and pilot-scale Thiopaq Ultra BD systems [8,20••]. *Thioalkalivibrio sulfidophilus* was the most

dominant sulfide-oxidizing bacterium (SOB) in the full-scale reactors [8]. The pilot-scale Thiopaq Ultra, however, was dominated by *Tv. sulfidophilus* (5–54%) and *Alkalilimnicola ehrlichii* (8–63%) [20••]. While *Thioalkalivibrio* is a well-known SOB, the prevalence of *Alkalilimnicola* is noteworthy as it has not been previously associated with sulfide oxidation under (micro)oxic conditions.

Moreover, SOMs in haloalkaline BD systems were electroactive in a bioelectrochemical (BES)-based BD system [21,22]. As such, they generated current while converting sulfide to sulfur (44%) and sulfate (56%). A combined BES-haloalkaline BD system could significantly enhance energy efficiency by eliminating the need for aeration of the oxic part while recovering energy from sulfide oxidation. Furthermore, the sulfur uptake by bacteria can also decrease the risk of electrode passivation due to sulfur deposition [22]. To date, BES systems for resource recovery from wastewater are under investigation to be used on a larger scale [23].


Significant progress has been made in the field of anoxic BD, which employs nitrate or nitrite as electron acceptors (as oxygen substitutes) to oxidize sulfide to sulfur or sulfate. One such advancement involves the replacement of traditional biotrickling filters (BTF) with continuous stirred tank bioreactors (CST-BRs) [24••]. In BTFs, the sulfur accumulates on the filters, thereby reducing their effectiveness. On the other hand, CST-BRs have shown promise, achieving a removal efficiency of 98.6% and recovering 88% of elemental sulfur [25]. A drawback is the increased operational costs due to the use of commercial chemicals as electron acceptors. However, this issue can be mitigated by using effluents from nitrification processes. For instance, in one anoxic BD system, ammonium-rich water is first nitrified to nitrate/nitrite, which is then used as an electron acceptor to oxidize sulfide into N₂ and elemental sulfur [26].

The composition of microbial communities in anoxic BD systems is largely unexplored. Current research primarily focuses on process development and stability, with only a handful of studies delving into the biodiversity of these systems. Dominant SOMs are affiliated to the genera *Thiobacillus*, *Sedimenticola*, *Sulfurimonas*, and *Rhodanobacter* [24••]. A more comprehensive understanding of the presence and activity of the microbial communities in the anoxic BD systems is needed, warranting further research in this area.

Example 2: biomining – mining of metals with microbes

Traditional mining, which generates a significant amount of waste in tailings (fine ground rock particles formed after ore processing), poses a severe challenge to land use and is energy intensive [27]. Biomining, on the

Figure 1



Schematic representation of the improved haloalkaline BD process (Thiopaq Ultra). Dashed lines indicate the flow of gas streams; solid lines indicate liquid circulation.

other hand, offers a less harmful alternative. This method involves the direct use of microbes or microbially produced lixivants (liquid media) to leach or extract metals, such as iron, copper, zinc, gold, and uranium, reducing waste generation [28••].


Bioleaching (BL) is primarily employed to extract metals from metal sulfides, utilizing chemolithotrophic iron/sulfur (Fe/S)-oxidizing microbes such as *Acidithiobacillus ferrooxidans*, *Acidithiobacillus caldus*, *Leptospirillum*, *Sulfolobus*, and *Ferroplasma* [29•–31]. These acidophilic micro-organisms thrive in highly acidic (pH < 3) and metal-rich environments, and by oxidizing ferrous iron (Fe^{2+}), they produce a BL lixiviant containing high concentrations of ferric iron (Fe^{3+}). The Fe^{3+} is involved in the conversion of sulfide to sulfate, not merely mediating but actively participating in the reaction via the formation of either thiosulfate (for acid-insoluble minerals) or polysulfides (for acid-soluble minerals) [29•,30,32]. The key reactions involved are shown in Box 1.

Despite its energy efficiency and a 70% lower carbon footprint than traditional methods, biomining may not be the most cost-effective option for high-quality ores, given the rate of BL is very slow. However, it presents an appealing approach for extracting metals from lower-grade ores, residual ores postmining (tailings), historical tailings, and electronic waste [27,28••].

Recently, *in situ* BL has been tested at a pilot scale for copper extraction from chalcopyrite (CuFeS_2) ores located more than 1 km deep via a process known as Deep *In Situ* Bioleaching (DISB; Figure 2). In this concept, an Fe^{3+} -rich bio-lixiviant is generated above ground in ferric iron-generating bioreactors (FIGB) and pumped into the underground chalcopyrite deposits via drilled channels. Acting as an electron acceptor, Fe^{3+} mediates the biotic/abiotic dissolution of Cu^{2+} in the acidic lixiviant, which is then pumped up from the other side of the ore and collected above ground in a recovery unit. Pure copper is extracted using solvent extraction and/or electrowinning, while the metal-depleted stream is recirculated back to the FIGB [28,33•]. In one DISB study, the FIGB was dominated by bacteria such as *Thiobacillus*, *Thiovirga*, *Alishewanella*, and a native fungal species [33•]. Even though these bacteria are not typically described as acidophiles, the overall consortium could maintain the desirable acidic conditions for leaching. Furthermore, iron-oxidizing bacteria might have been protected within biofilms and, therefore, not present in the sampled stream [33•].

BL has also been applied and studied in CST-BRs and heap-leaching reactors, extracting metals from tailings [34,35]. However, the extraction of chalcopyrite faces challenges, such as slow kinetics, passivation of ore due to the formation of a layer of sulfur compounds or decreased dissolution because of its inherent n-type electronic structure [36–39]. Ríos et al. [10•] proposed using moderate thermophilic acidophiles and controlling the

Figure 2



Schematic representation of DISB of metal sulfides using ferric iron (Fe^{3+})-generating bioreactors (FIGB) [28,72]. This is an example of indirect BL, which occurs differently for acid-soluble and nonacid-soluble metal sulfides. The sulfur conversions can occur both chemically and mediated by bacteria.

oxidation–reduction potential to improve BL. Their study revealed that consortia comprising *Sulfobacillus thermotolerans*, *Sulfobacillus acidophilus*, and *Ferroplasma acidiphilum* could maintain a desired redox potential below 680 mV (SHE), yielding maximum copper recovery. The study also highlighted synergistic interactions in consortia with weak iron oxidizers and *At. caldus* yielding more Fe^{3+} , thereby increasing the redox potential above 680 mV (SHE) and forming an undesired sulfur layer. Therefore, careful design of an effective synthetic community is necessary [10•]. Another study explored enhancing chalcopyrite dissolution by decoupling the growth and BL activity of the acidophiles by increasing the temperature of the consortium at the end of the growth phase with only 4°C [40].

Example 3: biosulfidogenesis for metal removal from metalliferous waters


The aforementioned metalliferous waters require treatment, which has traditionally involved indiscriminate neutralization with gypsum, followed by storage in tailings ponds. Unfortunately, this approach squanders valuable metals, relegating them to a toxic waste status. A more sustainable strategy, however, emerges in removing metals as metal sulfides via hydrogen sulfide [41,42]. While off-site hydrogen sulfide production and transport pose safety and economic challenges, onsite

sulfide generation offers a preferable alternative. Biosulfidogenesis (BS), the dissimilatory reduction of oxidized sulfur molecules to sulfide as an end product, has garnered significant attention [43–45]. It enables onsite sulfide production, facilitating the selective recovery of metals (e.g. Cu, Zn, Ni, Co) as high-purity metal sulfides suitable for recycling. Achieving this selectivity involves adjusting sulfide concentrations, metal content, and pH based on the corresponding pK values of desired metal sulfides [46]. For instance, copper sulfides will precipitate at lower pH values, while other metals, such as zinc or iron, will precipitate at increasing values [47]. Figure 3 shows different configurations of BS.

Sulfide production through biological sulfate reduction has been so far the focus of most investigations for several reasons: (1) sulfate is a readily available electron acceptor in acid mine drainage (AMD) being at concentrations up to molar amounts in some cases and, therefore, a cost-effective electron acceptor; (2) sulfate stability and solubility at low pH is high; (3) sulfate reduction at low pH consumes acidity and increases pH [48].

However, BS can be achieved from several sulfur-based electron acceptors such as sulfite, tetrathionate or elemental sulfur. Johnson and Sánchez-Andrea [34] high-

Figure 3



BS reactor and different configurations for metal precipitation from AMD.

light elemental sulfur reduction as the most promising technology, combining criteria such as economic efficiency per sulfide produced, solubility and stability. Elemental sulfur's efficiency requiring four times less electron donor for equivalent sulfide production significantly impacts process economics because adding electron donors is a large portion of the overall operating costs. Moreover, elemental sulfur, often a waste product, proves more cost-effective than traditional electron donors [49••,50]. Finally, a yet technologically underexplored metabolism for metal precipitation is elemental sulfur disproportionation, eliminating the need for external electron donors [51].

Biosulfidogenic technology, while promising, faces limitations due to the toxicity of metal-containing waters and their low pH under actual AMD conditions. Existing systems necessitate neutral conditions for sulfide production, followed by stripping in separate reactors. Recent investigations, however, have unveiled acidophilic sulfidogenic micro-organisms capable of resiliently precipitating metals. Notable species of SRMs include *Archaeoglobus profundus* and *Vulcanisaeta moutnovskia* (Archaea), alongside members of the bacterial phylum *Bacillota* (formerly *Firmicutes*) such as *Thermodesulfobium*, *Desulfosporosinus*, and *Acididesulfobacillus* [52–60••]. Additionally, members of the *Desulfobacterota* (formerly known as the *Deltaproteobacteria*), such as *Desulfothermobacter acidiphilus*, contribute to BS. Beyond sulfate reducers, sulfur reducers, such as *Desulfurella amikii* or sulfur disproportionators such as *Acidianus* strain AS80 offer alternative pathways [61,62] for sulfide generation at low pH. The application of micro-organisms naturally adapted to these extremely acidic conditions will broaden the application range and robustness of biosulfidogenic processes for metal precipitation and recovery.

Challenges and prospects

The microbial sulfur cycle plays a crucial role in many biotechnological applications. However, its complexity, given by the redox activity, presents significant engineering challenges and opportunities. Understanding the microbial communities and underlying mechanisms is essential for improving bioprocesses.

While most studies rely on 16S rRNA gene amplicon sequencing to assess microbial community composition, a deeper understanding of the genes, transcript, proteins, and metabolic levels is vital. This knowledge unravels functionality and mechanisms, enabling targeted bioprocess enhancements.

Direct application of pure and mixed cultures with genetically engineered strains offers promising avenues. For instance, CRISPR-Cas modifications have boosted sulfur formation by *Thioalkalivibrio versutus* by 24.7% while reducing sulfate formation by 15.2% compared to the wild type. Genetic modifications also shed light on sulfur metabolism [63]. However, using mixed or pure cultures requires careful consideration: (i) containment of genetically modified organisms, (ii) maintaining process robustness despite contaminants in the feed, and (iii) ensuring reactor sterility.

Physiology-based modeling has proven valuable in predicting bioprocess performance [64,65]. However, in recent times, data-driven technologies, such as artificial intelligence (AI), machine learning, and 'digital twins' have been widely adopted across industries, including biomanufacturing and mining. These technologies hold significant promise for bioprocesses, particularly in design, control, and scalability [66,67].

Scaling up remains a significant hurdle in biotechnology. Maintaining consistent process conditions across diverse environmental contexts is far from trivial. Therefore, achieving a robust, functionally redundant, and reproducible bioprocess becomes paramount.

In situ processes, such as DISB, present their own challenges. One critical issue is how to effectively halt the process once the desired ore has been sufficiently leached [33•]. If left unchecked, the ongoing process may lead to AMD, contaminating the surrounding area and groundwater. Consequently, it is essential to incorporate a BL cessation strategy into the design and operation of such *in situ* processes from the outset.

Biotechnological processes have evolved beyond product recovery, aiming for multifaceted goals and value-added outcomes. Energy efficiency, cost-effectiveness, and self-sufficiency are now the driving forces. One example is a microbial electrosynthesis system in which methane production from CO₂ is directly coupled with sulfide oxidation to sulfate, creating an eco-friendly system [68]. Another example is microbial protein formation from sulfide-rich biogas by methane- and sulfide-oxidizing bacteria [69]. A growing trend involves leveraging sulfur oxidation and reduction for extracting rare Earth metals from e-waste and lithium batteries [27,70]. In addition to new products, there is immense value in exploring creative ways to utilize abundant end products such as sulfur [16]. Likewise, BS can yield sulfide nanoparticles and contribute to a 'metal sulfides-microbe' biohybrid system. These systems find applications in environmental contexts, bridging the gap between biology and materials science [71].

Conclusion

Micro-organisms within the sulfur cycle present remarkable opportunities for tackling sulfur pollution while recovering valuable metals and elemental sulfur. However, while harnessing these bacteria is very attractive, managing toxic sulfur compounds under extreme conditions poses safety and operational efficiency challenges. Researchers are exploring to obtain a deeper understanding of the crucial micro-organisms involved in this process and seeking better consortia, substrates, and operational conditions to enhance process performance. Looking ahead, AI-based technologies could simulate process/application conditions for predicting and improving bioprocess efficiency. Overall, where environmental challenges meet biotechnological innovation, sulfur cycle-based micro-organisms continue to shape biotechnological applications for a more resilient future.

Funding

This work was performed in the co-operation framework of Wetsus, the European Centre of Excellence for Sustainable Water Technology (www.wetsus.nl),

University of Amsterdam, and Wageningen University and Research. Wetsus is co-funded by the Dutch Ministry of Economic Affairs and Ministry of Infrastructure and Environment, the European Union Regional Development Fund, the Province of Fryslan, and the Northern Netherlands Provinces. This work is part of a project that has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement no. 665874. The study was co-financed by Paqell B.V.

CRedit authorship contribution statement

Suyash Gupta wrote the draft version of the manuscript. All authors contributed to the final version of the manuscript.

Data Availability

No data were used for the research described in the article.

Declaration of Competing Interest

There is no conflict of interest.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Canfield DE, Raiswell R: **The evolution of the sulfur cycle.** *Am J Sci* 1999, **299**:697-723.
 2. Wasmund K, Mußmann M, Loy A: **The life sulfuric: microbial ecology of sulfur cycling in marine sediments.** *Environ Microbiol Rep* 2017, **9**:323-344.
 3. Muyzer G, Stams AJM: **The ecology and biotechnology of sulphate-reducing bacteria.** *Nat Rev Microbiol* 2008, **45**:187-192.
 4. Fan K, Wang W, Xu X, Yuan Y, Ren N, Lee DJ, Chen C: **Recent advances in biotechnologies for the treatment of environmental pollutants based on reactive sulfur species.** *Antioxidants* 2023, **12**:767.
- Notable review on prevalence and importance of reactive sulfur species in environmental technology combined with metabolic mechanisms.
5. Nguyen PM, Do PT, Pham YB, Doan TO, Nguyen XC, Lee WK, Nguyen DD, Vadiveloo A, Um MJ, Ngo HH: **Roles, mechanism of action, and potential applications of sulfur-oxidizing bacteria for environmental bioremediation.** *Sci Tot Environ* 2022, **852**:158203.
- Notable review on the roles and mechanisms of sulfur-oxidizing micro-organisms with critical discussion.
6. Ibáñez A, Garrido-Chamorro S, Coque JJR, Barreiro C: **From genes to bioleaching: unraveling sulfur metabolism in *Acidithiobacillus* genus.** *Genes* 2023, **14**:1772.
 7. Sorokin DY, Lysenko AM, Mityushina LL, Tourova TP, Jones BE, Rainey FA, Robertson LA, Kuenen GJ: ***Thioalkalimicrobium aerophilum* gen. nov., sp. nov. and *Thioalkalimicrobium sibericum* sp. nov., and *Thioalkalivibrio versutus* gen. nov., sp. nov., *Thioalkalivibrio nitratis* sp. nov. and *Thioalkalivibrio denitrificans* sp. nov., novel obligately alkaliphilic.** *Int J Syst Evol Microbiol* 2001, **51**:565-580.

8. Gupta S, Plugge CM, Klok JBM, Muyzer G: **Comparative analysis of microbial communities from different full-scale haloalkaline biodesulfurization systems.** *Appl Microbiol Biotechnol* 2022, **106**:1759-1776.
9. Buetti-Dinh A, Herold M, Christel S, Hajjami MEI, Bellenberg S, Ilie O, Wilmes P, Poetsch A, Sand W, Vera M, et al.: **Systems biology of acidophile biofilms for efficient metal extraction.** *Sci Data* 2020, **7**:215.
10. Ríos D, Bellenberg S, Christel S, Lindblom P, Giroux T, Dopson M: **Potential of single and designed mixed cultures to enhance the bioleaching of chalcopyrite by oxidation-reduction potential control.** *Hydrometallurgy* 2024, **224**:106245.
- A remarkable study investigating pure and mixed cultures and associated operational conditions to enhance their BL potential.
11. Vavourakis CD, Mehrshad M, Balkema C, Van Hall R, Andrei AŞ, Ghai R, Sorokin DY, Muyzer G: **Metagenomes and metatranscriptomes shed new light on the microbial-mediated sulfur cycle in a Siberian soda lake.** *BMC Biol* 2019, **17**:69.
12. Li M, Wen J: **Recent progress in the application of omics technologies in the study of bio-mining microorganisms from extreme environments.** *Micro Cell Fact* 2021, **20**:178.
13. Klok J, van Heeringen G, de Rink R, Wijnbelt J: **Introducing the next generation of the Thiopaq O&G process for biotechnological gas desulfurization: Thiopaq-SQ.** *Sulphur* 2017, **18260**:1-2.
14. Muyzer G, Kuenen JG, Robertson LA: **Colorless sulfur Bacteria.** In *The Prokaryotes: Prokaryotic Physiology and biochemistry*. Edited by Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F. Springer Berlin Heidelberg; 2013:555-588.
15. Sorokin DY, Zhilina TN, Lysenko AM, Tourova TP, Spiridonova EM: **Metabolic versatility of haloalkaliphilic bacteria from soda lakes belonging to the *Alkalispirillum-Alkalilimnicola* group.** *Extremophiles* 2006, **10**:213-220.
16. Pudi A, Rezaei M, Signorini V, Andersson MP, Baschetti MG, Mansouri SS: **Hydrogen sulfide capture and removal technologies: a comprehensive review of recent developments and emerging trends.** *Sep Purif Technol* 2022, **298**:121448.
17. Cline C, Hoksberg A, Abry R, Janssen A: Biological process for H₂S removal from gas streams: the Shell-Paques / Thiopaq™ gas desulfurization process. In *53rd Laurance Reid Gas Conditioning Conference*, 2003, 1-18. Norman (Oklahoma), USA.
18. de Rink R, Klok JBM, van Heeringen GJ, Sorokin DY, ter Heijne A, Zeijlmaker R, Mos YM, de Wilde V, Keesman KJ, Buisman CJN: **Increasing the selectivity for sulfur formation in biological gas desulfurization.** *Environ Sci Technol* 2019, **53**:4519-4527.
19. de Rink R, Gupta S, Piccioli de Carolis F, Liu D, ter Heijne A, Klok • JBM, Buisman CJN: **Effect of process conditions on the performance of a dual-reactor biodesulfurization process.** *J Environ Chem Eng* 2021, **9**:106450.
- Effects of process factors, hydraulic retention time, sulfide concentration, and pH were studied in detail to optimize further the sulfur-selectivity in a pilot dual line BD installation.
20. Gupta S, de Rink R, Klok JBM, Muyzer G, Plugge CM: **Process conditions affect microbial diversity and activity in a haloalkaline biodesulfurization system.** *Appl Environ Microbiol* 2024, **90**:e01864-23.
- First time that the active microbial community in haloalkaline BD installations has been identified. It reveals *Tv. sulfidophilus* and *A. ehrlichii* as the 'work horses' in these systems.
21. Ter Heijne A, De Rink R, Liu D, Klok JBM, Buisman CJN: **Bacteria as an electron shuttle for sulfide oxidation.** *Environ Sci Technol Lett* 2018, **5**:495-499.
22. de Rink R, B. Lavender M, Liu D, Klok JBM, Sorokin DY, ter Heijne A, Buisman CJN: **Continuous electron shuttling by sulfide oxidizing bacteria as a novel strategy to produce electric current.** *J Hazard Mater* 2022, **424**:127358.
23. Zhang Z, Sarkar D, Li L, Datta R: **Contaminant removal and resource recovery in bioelectrochemical wastewater treatment.** *Curr Pollut Rep* 2022, **8**:159-176.
24. Almenglo F, González-Cortés JJ, Ramírez M, Cantero D: **Recent advances in biological technologies for anoxic biogas desulfurization.** *Chemosphere* 2023, **321**:138084.
- State-of-the-art review on advances made in anoxic biogas BD using fixed biomass and suspended growth bioreactors since the first studies in this field in 2008.
25. González-Cortés JJ, Torres-Herrera S, Almenglo F, Ramírez M, Cantero D: **Anoxic biogas biodesulfurization promoting elemental sulfur production in a continuous stirred tank bioreactor.** *J Hazard Mater* 2021, **401**:123785.
26. González-Cortés JJ, Almenglo F, Ramírez M, Cantero D: **Simultaneous removal of ammonium from landfill leachate and hydrogen sulfide from biogas using a novel two-stage oxic-anoxic system.** *Sci Tot Environ* 2021, **750**:141664.
27. Dunbar WS: **Biotechnology and the mine of tomorrow.** *Trends Biotechnol* 2017, **35**:79-89.
28. Roberto FF, Schippers A: **Progress in bioleaching: part B, applications of microbial processes by the minerals industries.** *Appl Microbiol Biotechnol* 2022, **106**:5913-5928.
- Excellent review updating on the applications of microbial processes in BL and bio-oxidation.
29. Jones S, Santini JM: **Mechanisms of bioleaching: iron and sulfur oxidation by acidophilic microorganisms.** *Essays Biochem* 2023, **67**:685-699.
- Notable review exploring the mechanisms underlying BL and summarizes current knowledge on the enzymes driving microbial sulfur and iron oxidation in acidophiles.
30. Vera M, Schippers A, Hedrich S, Sand W: **Progress in bioleaching: fundamentals and mechanisms of microbial metal sulfide oxidation – part A.** *Appl Microbiol Biotechnol* 2022, **106**:6933-6952.
31. Moya-Beltrán A, Beard S, Rojas-Villalobos C, Issotta F, Gallardo Y, Ulloa R, Giaveno A, Degli Esposti M, Johnson DB, Quatrini R: **Genomic evolution of the class *Acidithiobacillia*: deep-branching Proteobacteria living in extreme acidic conditions.** *ISME J* 2021, **15**:3221-3238.
32. Schippers A, Sand W: **Bacterial leaching of metal sulfides proceeds by two indirect mechanisms via thiosulfate or via polysulfides and sulfur.** *Appl Environ Microbiol* 1999, **65**:319-321.
33. Bomberg M, Miettinen H, Hajdu-Rahkama R, Lakaniemi AM, Anacki W, Witecki K, Puhakka JA, Ineich T, Slabbert W, Kinnunen P: **Indirect in situ bioleaching is an emerging tool for accessing deeply buried metal reserves, but can the process be managed? – A case study of copper leaching at 1 km depth.** *Environ Technol Innov* 2023, **32**:103375.
- Notable study to upscale laboratory studies to field study monitoring development of iron oxidizing potential in *situ* enrichment of intrinsic micro-organisms and decommission procedures to inactivate the microbial population at the end of the operation.
34. Kremser K, Thalner S, Schoen H, Weiss S, Hemmelmaier C, Schnitzhofer W, Aldrian A, Guebitz GM: **Stirred-tank and heap-bioleaching of shredder-light-fractions (SLF) by acidophilic bacteria.** *Hydrometallurgy* 2020, **193**:105315.
35. Zhang R, Schippers A: **Stirred-tank bioleaching of copper and cobalt from mine tailings in Chile.** *Min Eng* 2022, **180**:107514.
36. O'Connor GM, Eksteen JJ: **A critical review of the passivation and semiconductor mechanisms of chalcopyrite leaching.** *Min Eng* 2020, **154**:106401.
37. Gericke M, Muller HH, van Staden PJ, Pinches A: **Development of a tank bioleaching process for the treatment of complex Cu-polymetallic concentrates.** *Hydrometallurgy* 2008, **94**:23-28.
38. Stott MB, Watling HR, Franzmann PD, Sutton D: **The role of iron-hydroxy precipitates in the passivation of chalcopyrite during bioleaching.** *Min Eng* 2000, **13**:1117-1127.
39. Ren Z, Chao C, Krishnamoorthy P, Asselin E, Dixon DG, Mora N: **The overlooked mechanism of chalcopyrite passivation.** *Acta Mater* 2022, **236**:118111.
40. Hedrich S, Joulain C, Graupner T, Schippers A, Guézennec AG: **Enhanced chalcopyrite dissolution in stirred tank reactors by**

- temperature increase during bioleaching. *Hydrometallurgy* 2018, **179**:125-131.
41. Santos AL, Johnson DB: **Design and application of a low pH upflow biofilm sulfidogenic bioreactor for recovering transition metals from synthetic wastewater at a Brazilian copper mine.** *Front Microbiol* 2018, **9**:2051.
 42. Li Y, Zhao Q, Liu M, Guo J, Xia J, Wang J, Qiu Y, Zou J, He W, Jiang F: **Treatment and remediation of metal-contaminated water and groundwater in mining areas by biological sulfidogenic processes: a review.** *J Hazard Mater* 2023, **443**:130377.
 43. Florentino AP, Sánchez-España J, Sánchez-Andrea I: **Editorial: advances in biosulfidogenic processes.** *Front Bioeng Biotechnol* 2022, **10**:1025093.
 44. Estay H, Barros L, Troncoso E: **Metal sulfide precipitation: recent breakthroughs and future outlooks.** *Minerals* 2021, **11**:1385.
 45. Johnson DB, Sánchez-Andrea I: **Dissimilatory reduction of sulfate and zero-valent sulfur at low pH and its significance for bioremediation and metal recovery.** In *Advances in Microbial Physiology*. Edited by Poole RK. Academic Press; 2019:205-231.
 46. Hidalgo-Ulloa A, Buisman C, Weijma J: **Metal sulfide precipitation mediated by an elemental sulfur-reducing thermoacidophilic microbial culture from a full-scale anaerobic reactor.** *Hydrometallurgy* 2022, **213**:105950.
 47. Weijma J, Klok JBM, Dijkman H, Jansen G, Sánchez-Andrea I, Buisman CJN, van Hullebusch ED, Hennebel T, du Laing G, Cruz H, et al.: **Established technologies for metal recovery from industrial wastewater streams.** In *Resource Recovery from Water*. Edited by Pikaar I, Guest J, Ganigué R, Jensen P, Rabaey K, Seviour T, Trimmer J, van der Kolk O, Vaneekhaute C, Verstraete W. IWA Publishing; 2022:295-317.
 48. Ayangbenro AS, Olanrewaju OS, Babalola OO: **Sulfate-reducing bacteria as an effective tool for sustainable acid mine bioremediation.** *Front Microbiol* 2018, **9**:1986.
 49. Hidalgo-Ulloa A, van der Graaf CM, Sanchez-Andrea I, Buisman CJN: **Continuous single-stage elemental sulfur reduction and copper sulfide precipitation under thermoacidophilic conditions.** *Water Res* 2023, **236**.
- Study investigating and optimizing single-stage metal sulfide precipitation process with elemental sulfur under thermoacidophilic conditions.
50. Sun R, Zhang L, Wang X, Ou C, Lin N, Xu S, Qiu YY, Jiang F: **Elemental sulfur-driven sulfidogenic process under highly acidic conditions for sulfate-rich acid mine drainage treatment: performance and microbial community analysis.** *Water Res* 2020, **185**:116230.
 51. Zou J, Qiu YY, Li H, Jiang F: **Sulfur disproportionation realizes an organic-free sulfidogenic process for sustainable treatment of acid mine drainage.** *Water Res* 2023, **232**:119647.
 52. Burggraf S, Jannasch HW, Nicolaus B, Stetter KO: ***Archaeoglobus profundus* sp. nov., represents a new species within the sulfate-reducing *Archaeobacteria*.** *Syst Appl Microbiol* 1990, **13**:24-28.
 53. Chernyh NA, Neukirchen S, Frolov EN, Sousa FL, Miroshnichenko ML, Merkel AY, Pimenov NV, Sorokin DY, Ciordia S, Mena MC, et al.: **Dissimilatory sulfate reduction in the archaeon '*Candidatus Vulcanisaeta moutnovskia*' sheds light on the evolution of sulfur metabolism.** *Nat Microbiol* 2020, **5**:1428-1438.
 54. Mori K, Kim H, Kakegawa T, Hanada S: **A novel lineage of sulfate-reducing microorganisms: *Thermodesulfobiaceae* fam. nov., *Thermodesulfobium narugense*, gen. nov., sp. nov., a new thermophilic isolate from a hot spring.** *Extremophiles* 2003, **7**:283-290.
 55. Frolov EN, Kublanov IV, Toshchakov SV, Samarov NI, Novikov AA, Lebedinsky AV, Bonch-Osmolovskaya EA, Chernyh NA: ***Thermodesulfobium acidiphilum* sp. nov., a thermoacidophilic, sulfate-reducing, chemoautotrophic bacterium from a thermal site.** *Int J Syst Evol Microbiol* 2017, **67**:1482-1485.
 56. Alazard D, Joseph M, Battaglia-Brunet F, Cayol JL, Ollivier B: ***Desulfosporosinus acidiphilus* sp. nov.: A moderately acidophilic sulfate-reducing bacterium isolated from acid mining drainage sediments.** *Extremophiles* 2010, **14**:305-312.
 57. Sánchez-Andrea I, Stams AJM, Hedrich S, Nancucheo I, Johnson DB: ***Desulfosporosinus acididurans* sp. nov.: an acidophilic sulfate-reducing bacterium isolated from acidic sediments.** *Extremophiles* 2015, **19**:39-47.
 58. Panova IA, Ikkert O, Avakyan MR, Kopitsyn DS, Mardanov AV, Pimenov NV, Shcherbakova VA, Ravin NV, Karnachuk OV: ***Desulfosporosinus metallidurans* sp. nov., an acidophilic, metal-resistant sulfate-reducing bacterium from acid mine drainage.** *Int J Syst Evol Microbiol* 2021, **71**:004876.
 59. Sánchez-Andrea I, van der Graaf CM, Hornung B, Bale NJ, Jarzembowska M, Sousa DZ, Rijpstra WIC, Sinnighe Damsté JS, Stams AJM: **Acetate degradation at low pH by the moderately acidophilic sulfate reducer *Acididesulfobacillus acetoxydans* gen. nov. sp. nov.** *Front Microbiol* 2022, **13**:816605.
- Notable study showing novel moderately acidophilic sulfate reducing bacteria which can oxidize organic acids to CO₂.
60. Egas RA, Sahonero-Canavesi DX, Bale NJ, Koenen M, Yildiz Ç, Villanueva L, Sousa DZ, Sánchez-Andrea I: **Acetic acid stress response of the acidophilic sulfate reducer *Acididesulfobacillus acetoxydans*.** *Environ Microbiol* 2024, **26**:e16565.
- First paper on transcriptomic study of acidophilic sulfate reducing bacteria showing genes involved in the response to acetic acid.
61. Florentino AP, Brienza C, Stams AJM, Sánchez-Andrea I: ***Desulfurella amilsii* sp. nov., a novel acidotolerant sulfur-respiring bacterium isolated from acidic river sediments.** *Int J Syst Evol Microbiol* 2016, **66**:1249-1253.
 62. Amenabar MJ, Urschel MR, Boyd ES: **Metabolic and taxonomic diversification in continental magmatic hydrothermal systems.** In *Microbial Evolution under Extreme Conditions*. Edited by Bakermans C. De Gruyter; 2015:57-96.
 63. Hao X, Mu T, Sharshar MM, Jia Y, Zhong W, Chen Z, Wen Q, Yang M, Wang C, Xing J: **CRISPR/Cas12a-mediated genome editing in *Thioalkalivibrio versutus*.** *ACS Synth Biol* 2023, **12**:1204-1215.
 64. Kiragosyan K, Klok JBM, Keesman KJ, Roman P, Janssen AJH: **Development and validation of a physiologically based kinetic model for starting up and operation of the biological gas desulfurization process under haloalkaline conditions.** *Water Res X* 2019, **4**:100035.
 65. Klok JBM, de Graaff M, van den Bosch PLF, Boelee NC, Keesman KJ, Janssen AJH: **A physiologically based kinetic model for bacterial sulfide oxidation.** *Water Res* 2013, **47**:483-492.
 66. Herwig C, Pörtner R, Möller J: **Digital Twins.** Springer International Publishing; 2021.
 67. Singh M, Srivastava R, Fuenmayor E, Kuts V, Qiao Y, Murray N, Devine D: **Applications of digital twin across industries: a review.** *Appl Sci* 2022, **12**:5727.
 68. Kambara H, Dinh HTT, Matsushita S, Aoi Y, Kandaichi T, Ozaki N, Ohashi A: **New microbial electrosynthesis system for methane production from carbon dioxide coupled with oxidation of sulfide to sulfate.** *J Environ Sci* 2023, **125**:786-797.
 69. Areniello M, Matassa S, Esposito G, Lens PNL: **Microbial protein production from sulfide-rich biogas through an enrichment of methane- and sulfur-oxidizing bacteria.** *Bioresour Technol* 2023, **383**:129237.
 70. Martínez-Bellange P, von Bernath D, Navarro CA, Jerez CA: **Biomining of metals: new challenges for the next 15 years.** *Micro Biotechnol* 2022, **15**:186-188.
 71. Su Z, Li X, Xi Y, Xie T, Liu Y, Liu B, Liu H, Xu W, Zhang C: **Microbe-mediated transformation of metal sulfides: mechanisms and environmental significance.** *Sci Tot Environ* 2022, **825**:153767.
 72. Johnson DB: **Biomining-biotechnologies for extracting and recovering metals from ores and waste materials.** *Curr Opin Biotechnol* 2014, **30**:24-31.