



UvA-DARE (Digital Academic Repository)

Biopsychosocial aspects of sleep bruxism in children

Restrepo Serna, C.C.

[Link to publication](#)

Citation for published version (APA):

Restrepo Serna, C. C. (2018). Biopsychosocial aspects of sleep bruxism in children

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



1

Chapter

I n t r o d u c t i o n

Bruxism was defined by a panel of experts, as a repetitive jaw-muscle activity characterized by clenching or grinding of the teeth and/or by bracing or thrusting of the mandible¹. Although bruxism is not necessarily a treatment-demanding condition^{2,3}, it remains a common concern for parents, who are usually worried when their children gnash and grind their teeth during sleep⁴.

Over the years, several studies have been performed in adults to try explaining the pathophysiology of sleep bruxism (SB) and its clinical correlates. As a result, researchers working on adults agree on some basic concepts. Among those, there is consensus that SB is of central origin^{5,6,7} and that it should be assessed within a multiple-variable framework^{1,8}, due to the multiplicity of biological, psychological, and social factors that interact and lead to its occurrence⁹. In addition, the common belief that tooth wear has a linear relationship with SB, has been discarded in adults⁸.

On the other hand, investigations on bruxism in children are scarce, in part because of potential poor compliance with research protocols, as well as ethical and legal issues that make it difficult to perform researches in children. In addition, validated criteria for the assessment of SB in children are lacking.

The literature suggests that polysomnography (PSG) should be considered the reference standard for a quantitative assessment of SB¹. Such technique monitors multiple physiologic parameters during sleep and it is useful in the evaluation of sleep disorders (i.e., treatment-demanding conditions), as well as their discrimination from sleep behaviors (i.e., physiological phenomena)¹⁰. The electroencephalogram, eye movements, chin and masticatory muscles electromyogram, nasal airflow, thoracic and abdominal respiratory effort, electrocardiogram, and oxygen saturation; are among the parameters that are recorded continuously during sleep in a laboratory setting and are displayed on a computerized system or on paper^{10,11}.

The widespread use of PSG is limited by the high costs, limited availability, and comprehensive technical requirements^{12,13}. Unfortunately, those issues are even more critical in children than adults, thus further limiting the diffusion of PSG for SB assessment in children. Therefore, available investigations have forcedly adopted parental (proxy) report as the main strategy to gather data on SB in children.

Whilst such an approach seems reasonable from a cost-effectiveness ratio, the literature suggests that any self- or parental-reported approach to the assessment of SB may have its limits, if used alone^{12,13}. In particular, there are ongoing debates on the validity of self-report in adults^{2,3}. Thus, researches are currently performed to identify alternative options to PSG and to determine

the correlation between the various approaches to SB assessment (viz., self-report, clinical assessment, and PSG measurement)^{8,14}.

In children, as stated above, the most common method is the use of questionnaires answered by the parents, which is particularly useful when evaluating large numbers of participants^{15,16}. Nevertheless, the main concern regarding the use questionnaires for SB data collection in children, is that their agreement with PSG findings has not been assessed.

Due to the potential shortcomings of self-reported approaches and the technical limitations that prevent the large-scale usage of PSG, alternative methods are being explored. Among those, the use of portable electromyography (EMG) devices, which are available to measure the masticatory muscle activity (MMA) during sleep time at home in adults^{17,18}, could be useful for children as well. Portable EMG recorders would allow keeping a child in his/her habitual sleep environment, but the correlation between portable EMG findings and PSG measurements has never been evaluated in children.

The diagnostic difficulties to approach SB and to provide a definite assessment in children, reflect in the inconclusive knowledge that is available on some etiological and clinical aspects. For decades, the search for an association of a tense personality and/or a stressful life with SB in children has been the main target of etiological studies^{19,20,21,22,23}, while tooth wear was the general focus for the evaluation of SB consequences^{24,25}. Notwithstanding, findings on those two topics deserve further investigation, to provide a comprehensive assessment of SB in children.

From the above, it emerges that evidence on SB in children is still scarce and fragmental, and that getting deeper into the study of several biopsychological issues may provide further insights into the pathophysiology of SB in children.

Aims and outline of the thesis

The general aim of this thesis is to obtain a deeper insight into the mutual interactions of the biopsychological aspects of SB in children. Below, the specific aims are formulated for each chapter.

The starting point of the project is a systematic literature review on the epidemiology of SB in children. The review, described in **Chapter 2** of this thesis, aims to summarize all the available information concerning SB prevalence in children.

In the study described in **Chapter 3**, the correlation between proxy-reported SB and a “definite” SB diagnosis via PSG, using PSG/SB adult criteria

as the gold standard¹, will be explored. In addition, due to the difficulties of using PSG in children, a study on the correlation between the measurements of MMA, using a portable single-channel EMG device and PSG, will be presented in **Chapter 4**.

The next studies aim to investigate the psychosocial issues that may be linked to the central origin of SB in children. Due to the psychological and behavioral nature of the variables, as well as the need to recruit large samples for epidemiological purposes, the studies described in **Chapter 5 and 6** are questionnaire-based. The investigations assess the associations of SB with quality of life (QoL) variables and sleep behaviors, in children belonging to different social layers. Validated psychological instruments will be used, and both studies will be performed on representative samples, to increase the internal and external validity of the results and to explore possible associations to be refined in future studies.

Finally, the investigation presented in **Chapter 7** evaluates one of the clinical signs that is commonly considered a consequence of SB, viz., tooth wear. Specifically, SB and dietary-habits will be evaluated as comorbidities and associated factors to the presence and severity of tooth wear in the mixed dentition.

References

1. Lobbezoo F, Ahlberg J, Glaros AG, Kato T, Koyano K, Lavigne GJ et al. Bruxism defined and graded: an international consensus. *J Oral Rehabil.* 2013; 402-404.
2. Raphael KG, Santiago V, Lobbezoo F. Is bruxism a disorder or a behaviour? Rethinking the international consensus on defining and grading of bruxism. *J Oral Rehabil.* 2016;43:791-798.
3. Manfredini D, De Laat A, Winocur E, Ahlberg J. Why not stop looking at bruxism as a black/white condition? Aetiology could be unrelated to clinical consequences. *J Oral Rehabil.* 2016;43:799-801.
4. Meltzer LJ, Johnson C, Crosette J, Ramos M, Mindell JA. Prevalence of Diagnosed Sleep Disorders in Pediatric Primary Care Practices. *Pediatrics.* 2010;125:1410-1418.
5. Lavigne GJ, Kato T, Kolta A, Sessle BJ. Neurobiological mechanisms involved in sleep bruxism. *Crit Rev Oral Biol Med.* 2003;14:30-46.

6. Lobbezoo F, Naeije M. Bruxism is mainly regulated centrally, not peripherally. *J Oral Rehabil.* 2001;28:1085-1091.
7. Seraidarian P, Seraidarian PI, das Neves Cavalcanti B, Marchini L, Claro Neves AC. Urinary levels of catecholamines among individuals with and without sleep bruxism. *Sleep Breath.* 2009;13:85-88.
8. Casett E, Réus JC, Stuginski-Barbosa J, Porporatti AL, Carra MC, Peres MA, de Luca Canto G, Manfredini D. Validity of different tools to assess sleep bruxism: a meta-analysis. *J Oral Rehabil.* 2017;44:722-734.
9. Carra MC, Huynh N, Fleury B, Lavigne G. Overview on Sleep Bruxism for Sleep Medicine Clinicians. *Sleep Med Clin.* 2015;10:375-384.
10. Morgenthaler TI, Owens J, Alessi C, Boehlecke B, Brown TM, Coleman J Jr, Friedman L, Kapur VK, Lee-Chiong T, Pancer J, Swick TJ; American Academy of Sleep Medicine. Practice parameters for behavioral treatment of bedtime problems and night wakings in infants and young children. *Sleep.* 2006;29:1277-1281.
11. Khoury S, Carra MC, Huynh N, Montplaisir J, Lavigne GJ. Sleep Bruxism-Tooth Grinding Prevalence, Characteristics and Familial Aggregation: A Large Cross-Sectional Survey and Polysomnographic Validation. *Sleep.* 2016; 39:2049-2056
12. Huynh NT, Desplats E, Bellerive A. Sleep bruxism in children: sleep studies correlate poorly with parental reports. *Sleep Med.* 2016;19:63-68.
13. Aurora RN, Lamm CI, Zak RS, Kristo DA, Bista SR, Rowley JA, Casey KR. Practice parameters for the non-respiratory indications for polysomnography and multiple sleep latency testing for children. *Sleep.* 2012;35:1467-1473.
14. Manfredini D, Ahlberg J, Castroflorio T, Poggio CE, Guarda-Nardini L, Lobbezoo F. Diagnostic accuracy of portable instrumental devices to measure sleep bruxism: a systematic literature review of polysomnographic studies. *J Oral Rehabil.* 2014;41:836-842.
15. Drumond CL, Souza DS, Serra-Negra JM, Marques LS, Ramos-Jorge ML, Ramos-Jorge J. Respiratory disorders and the prevalence of sleep bruxism among schoolchildren aged 8 to 11 years. *Sleep Breath.* 2017;21:203-208.

16. Emodi Perlman A, Lobbezoo F, Zar A, Friedman Rubin P, van Selms MK, Winocur E. Self-Reported bruxism and associated factors in Israeli adolescents. *J Oral Rehabil.* 2016;43:443-450.
17. Castroflorio T, Bargellini A, Rossini G, Cugliari G, Deregibus A, Manfredini D. Agreement between clinical and portable EMG/ECG diagnosis of sleep bruxism. *J Oral Rehabil.* 2015;42:759-764.
18. Stuginski-Barbosa J, Porporatti AL, Costa YM, Svensson P, Conti PC. Diagnostic validity of the use of a portable single-channel electromyography device for sleep bruxism. *Sleep Breath.* 2016;20:695-702.
19. Restrepo CC, Vásquez LM, Alvarez M, Valencia I. Personality traits and temporomandibular disorders in a group of children with bruxing behaviour. *J Oral Rehabil.* 2008; 35:585-593.
20. Oliveira MT, Bittencourt ST, Marcon K, Destro S, Pereira JR. Sleep bruxism and anxiety level in children. *Braz Oral Res.* 2015; 29: 11-15.
21. Ferreira-Bacci Ado V, Cardoso CL, Díaz-Serrano KV. Behavioral problems and emotional stress in children with bruxism. *Braz Dent J.* 2012;23:246-251.
22. Serra-Negra JM, Paiva SM, Flores-Mendoza CE, Ramos-Jorge ML, Pordeus IA. Association among stress, personality traits, and sleep bruxism in children. *Pediatr Dent.* 2012;34:30-34.
23. Serra-Negra JM, Ramos-Jorge ML, Flores-Mendoza CE, Paiva SM, Pordeus IA. Influence of psychosocial factors on the development of sleep bruxism among children. *Int J Paediatr Dent.* 2009;19:309-317.
24. Restrepo C, Peláez A, Alvarez E, Paucar C, Abad P. Digital imaging of patterns of dental wear to diagnose bruxism in children. *Int J Paediatr Dent.* 2006;16:278- 285.
25. Vélez AL, Restrepo CC, Peláez-Vargas A, Gallego GJ, Alvarez E, Tamayo V, Tamayo M. Head posture and dental wear evaluation of bruxist children with primary teeth. *J Oral Rehabil.* 2007;34:663-670.