



UvA-DARE (Digital Academic Repository)

Soft, absorbed X-ray spectra of the new transient IGR J17451-3022

Heinke, C.O.; Bahramian, A.; Altamirano, D.; Wijnands, R.

Publication date

2014

Document Version

Final published version

Published in

The astronomer's telegram

[Link to publication](#)

Citation for published version (APA):

Heinke, C. O., Bahramian, A., Altamirano, D., & Wijnands, R. (2014). Soft, absorbed X-ray spectra of the new transient IGR J17451-3022. *The astronomer's telegram*, 6459. <http://www.astronomerstelegam.org/?read=6459>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Outside
 GCN
 IAUCs

Other
 ATel on [Twitter](#) and [Facebook](#)
 ATELstream
 ATel Community Site
 MacOS: [Dashboard Widget](#)

This space for free for your conference.



[[Previous](#) | [Next](#) | [ADS](#)]

Soft, absorbed X-ray spectra of the new transient IGR J17451-3022

ATel #6459; *C. O. Heinke, A. Bahramian (Alberta), D. Altamirano (Southampton), R. Wijnands (Amsterdam)*
 on 10 Sep 2014; 21:47 UT
 Credential Certification: *Craig Heinke (cheinke@virginia.edu)*

Subjects: X-ray, Neutron Star, Transient

Referred to by ATel #: [6469](#), [6486](#), [6501](#), [6533](#), [7028](#), [7361](#)



We report Swift/XRT followup observations of the new transient IGR J17451-3022 discovered in JEM-X observations performed Aug. 22-24 (Atel #[6451](#)). Swift observed with XRT in PC mode on Sept. 5 (1648 seconds; position reported in Atel #[6451](#)), Sept. 9 (1950 s, starting at UT 10:00), and on Sept. 10 (856 s, starting at UT 2:12). IGR J17451-3022 is clearly detected in each observation, with moderate pileup.

We extracted source spectra (excluding the central 8-10" to avoid the piled-up region) and background spectra from the 3 XRT PC mode observations, constructed effective area files, and used the appropriate response matrices. We grouped the data to 50 counts/bin in the first two observations, and 25 counts in the third observation, and fit the spectra simultaneously in XSPEC with absorbed (TBABS, using Wilms et al. (2000, ApJ, 542, 914) abundances) power-law, blackbody (bbodyrad), or bremsstrahlung models. An absorbed power-law model gave a poor fit (reduced chi-squared of 1.34 for 59 degrees of freedom), with photon indices greater than 3.3 (best-fit around 3.7). Bremsstrahlung models gave a better fit (reduced chi-squared of 1.16 for 59 dof), but with low temperatures around 2 keV, which seem physically implausible. Fits with a blackbody allowing the absorption, temperature, and radius to vary provided a good fit (reduced chi-squared of 1.16 for 59 dof). In this case, the temperature remains essentially constant at 0.82 (± 0.06 , at 90% confidence) keV, while the absorption increased ($N_H = 5.2 \pm 0.8 \times 10^{22} \text{ cm}^{-2}$ on Sept. 5; $7.2 \pm 1.0 \times 10^{22}$ on Sept. 9; $6.4 \pm 1.5 \times 10^{22}$ on Sept. 10). The bbodyrad normalization may increase (from 33 to 50, then down to 44), but the variations are not significant. The 2-10 keV unabsorbed flux increases from $1.2(\pm 0.1) \times 10^{-10} \text{ ergs cm}^{-2} \text{ s}^{-1}$ to $1.9(\pm 0.2) \times 10^{-10} \text{ ergs cm}^{-2} \text{ s}^{-1}$, then to $1.6(\pm 0.2) \times 10^{-10} \text{ ergs cm}^{-2} \text{ s}^{-1}$. We attempted blackbody fitting with the absorption tied (best-fit at 6×10^{22}). In this case, the temperature varies (from 0.77 ± 0.04 , to 0.88 ± 0.05 , then to 0.83 ± 0.06), but the fit is markedly poorer (reduced chi-squared of 1.25 for 61 dof).

The nature of IGR J17451-3022 remains unclear. The high absorption is consistent with a location near (or beyond) the Galactic Center. The rather soft spectrum seems to rule out cataclysmic variables and high-mass X-ray binaries, and appears consistent with either a low-mass X-ray binary in the soft state, or perhaps a magnetar. If it is an LMXB located in the Galactic Center, the luminosity (2-10 keV $L_x = 1 \times 10^{36} \text{ erg/s}$) would be unusually low for a soft state. This might be explained if the system is nearly edge-on, reducing our visibility of the inner

Related

- 7361** Discovery of eclipses in the X-ray transient IGR J17451-3022
- 7096** INTEGRAL detection of the on-going outbursts from 1RXS J180408.9-342058 and GRO J1750-27
- 7039** Swift observations of 1RXS J180408.9-342058
- 7028** Continuing outburst of Galactic transient IGR J17451-3022
- 7008** MAXI/GSC observation of 1RXS J180408.9-342058 in outburst
- 6997** Swift/BAT detects an outburst from the neutron star binary 1RXS J180408.9-342058
- 6839** New Outburst of the Be/X-ray Transient GRO J1750-27 Detected with Fermi/GBM
- 6602** INTEGRAL/JEM-X sees enhanced activity in the Galactic center region: SAX J1747.0-2853 and IGR J17454-2919
- 6574** Hard X-ray spectral and timing properties of IGR J17454-2919 consistent with a black hole in the hard state
- 6533** Chandra Localization of IGR J17451-3022
- 6530** IGR J17454-2919: a new X-ray transient found by INTEGRAL/JEM-X close to the Galactic Center
- 6501** New Galactic transient IGR J17451-3022 still soft
- 6486** Detection of spectral hardening in IGR J17451-3022; evidence for a LMXB
- 6469** Swift observations of the ongoing outburst of IGR J17451-3022
- 6459** Soft, absorbed X-ray spectra of the new transient IGR J17451-3022
- 6451** A new X-ray transient, IGR J17451-3022, discovered by INTEGRAL/JEM-X near the Galactic Centre
- 1400** Further observations of GRO J1750-27 (AX J1749.1-2639) with INTEGRAL
- 1385** INTEGRAL Galactic bulge monitoring observations of GRO J1750-27 (AX J1749.1-2639), H1743-322 and SLX 1746-331

disk, which could also explain the varying N_H . Alternatively, the system may be located well beyond the Galactic Center; a distance of 20 kpc would bring the L_x up to 10^{37} ergs/s, reasonable for the soft state. The apparently varying N_H could in this case be due to variations in a more complex spectrum (e.g. a disk blackbody, plus comptonization, and perhaps another blackbody from a neutron star surface) that our limited spectra cannot reasonably constrain. Alternatively, IGR J17451-3022 may be a transient magnetar, in which case we should detect pulsations (in the 2-12 second range) in data of sufficient time resolution.

We thank the Swift team for their rapid scheduling of these observations.

[**Telegram Index**]

R. E. Rutledge, Editor-in-Chief

`rrutledge@astronomerstelegam.org`

Derek Fox, Editor

`dfox@astronomerstelegam.org`

Mansi M. Kasliwal, Co-Editor

`mansi@astronomerstelegam.org`