UvA-DARE (Digital Academic Repository)

Using a digital game for training desirable behavior in cognitive-behavioral therapy for burnout syndrome: a controlled study

Zielhorst, T.; van den Brule, D.; Visch, V.; Melles, M.; van Tienhoven, S.; Sinkbaek, H.; Schrieken, B.; Tan, E.S.H.; Lange, A.

DOI

[10.1089/cyber.2013.0690](https://doi.org/10.1089/cyber.2013.0690)

Publication date

2015

Document Version

Final published version

Published in

Cyberpsychology, Behavior, and Social Networking

[Link to publication](#)

Citation for published version (APA):

Zielhorst, T., van den Brule, D., Visch, V., Melles, M., van Tienhoven, S., Sinkbaek, H., Schrieken, B., Tan, E. S. H., & Lange, A. (2015). Using a digital game for training desirable behavior in cognitive-behavioral therapy for burnout syndrome: a controlled study. *Cyberpsychology, Behavior, and Social Networking*, 18(2), 101-111.
<https://doi.org/10.1089/cyber.2013.0690>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Using a Digital Game for Training Desirable Behavior in Cognitive–Behavioral Therapy of Burnout Syndrome: A Controlled Study

Thomas Zielhorst, MSc,^{1,*} Daphne van den Brule, MSc,^{2,†} Valentijn Visch, PhD,²
Marijke Melles, PhD,² Sam van Tienhoven, MSc,³ Helle Sinkbaek, MSc,³ Bart Schrieken, MSc,³
Eduard S.-H. Tan, Prof.,¹ and Alfred Lange, Prof.⁴

Abstract

Burnout is a globally increasing illness, and as a result, many forms of burnout therapy have arisen. The use of digital games can be psychotherapeutically effective because they can transform exercises that are by themselves unattractive into intrinsically motivated action. This pilot study aims to test whether a specially designed game contributes to patients learning desired behavior and achieving other specific therapeutic goals in an online cognitive–behavioral therapy (CBT)-based burnout treatment context. In total, 101 participants took part in the experiment, under four conditions: (a) Game + Therapy, (b) Therapy Only, (c) Game Only, and (d) No Game + No Therapy. Pre- and postmeasures were taken online. Results showed that the two therapy conditions (Game + Therapy and Therapy Only) showed a greater decrease in complaints and disengagement, and a stronger increase in coping skills than the nontherapy conditions (Game Only and No Game + No Therapy). As expected, the Game + Therapy condition outperformed the Therapy Only condition on combined improvement measures of burnout symptoms. However, analyses of individual measures showed no effects. It can be cautiously concluded that the therapeutic digital game may be a useful tool when embedded in a therapeutic burnout treatment program and is probably more efficient than CBT, as it is used in current practice.

Introduction

BURNOUT, ALSO REFERRED to as work-related stress, is a rapidly growing labor-related illness.¹ In the Netherlands, 12% of employees were reported to suffer from burnout in 2011, 2% more than in 2004.^{2,3} Burnout has also increased prevalence in other countries.^{4,5}

Burnout has been defined as “the occurrence of emotional exhaustion, unusual fatigue, a cynical or detached working attitude and a feeling of decreasing competence.”^{6(p14)} The Maslach Burnout Inventory (MBI) is the most commonly used self-report index, measuring three factors: emotional exhaustion, depersonalization, and decreasing personal accomplishment.^{7,8} In another recently proposed index, the latter factor was omitted.⁹ Burnout has a three-stage etiology: (a) imbalance of job demands and energy resources, (b) precursive symptoms, and (c) burn-

out syndrome.¹⁰ Precursive symptoms include stress,^{11–15} depression,^{11,12,15–17} and anxiety^{12,15,16,18–20} They tend to persist after the second stage.

Among the treatments for burnout syndrome,²¹ relaxation, stress management, and cognitive–behavioral therapy (CBT) have proven effective and are broadly used.¹¹ CBT was found to be effective for the specific burnout-related problems of stress,²² unexplained fatigue,²³ depression,²⁴ and anxiety disorder.²⁵ CBT hinges on the insight that psychological complaints are based on automatized behavior and cognition that can be altered through learning.²⁶ CBT aims to cure patients by training them to cope with stress. Favorable cognitive and behavioral habits are learnt, such as recognizing negative thoughts and setting personal boundaries.^{11,22}

Burnout therapy is increasingly administered online. Online platforms include e-mail, e-chat, Web sites, video conferencing, and mobile applications, either as (guided) self-help

¹Department of Communication Science, University of Amsterdam, Amsterdam, The Netherlands.

²Department of Industrial Design, Delft University of Technology, Delft, The Netherlands.

³Interapy, Amsterdam, The Netherlands.

⁴Department of Clinical Psychology, University of Amsterdam, Amsterdam, The Netherlands.

*Current affiliation: Global Electronic Operations, Elsevier B.V., Amsterdam, The Netherlands.

†Current affiliation: Push the Button, Venlo, The Netherlands.

therapy or as actual psychotherapy.²⁷ The Amsterdam-based firm Interyp was the first certified agency worldwide to offer a complete online burnout CBT program. Use is made of e-mail and a Web site for correspondence between the patient and the therapist. Figure 1 describes how the program functions. The Interyp program is an effective and easily accessible program.¹² This study investigates whether the program could be further improved through the use of serious games.

Some positive outcomes of serious gaming have been found in a number of fields.^{28–31} Serious games are not primarily designed for entertainment, but for purposes such as education, training, therapy, or behavioral change.³² Research on the effects of psychotherapeutic games is quite recent. Yet, support has been reported of beneficial effects in the treatment of a variety of complaints, including depression, divorce trauma, autism, and phobias.^{33–45} This article reports a pilot study testing the incremental value in terms of user motivation and treatment effectiveness of “gamifying” an existing online CBT-based burnout module.

The psychotherapeutic effectiveness of games originates in their capability of transforming exercises that are in themselves unattractive into intrinsically motivated action. Playing games is consciously experienced as enjoyable, while, at the same time, players are also honing their skills and capacities, even though they are unaware of doing so.^{46,47} The game experience elements responsible for users’ engagement in what is an exercise include *challenge*^{48,49} and *flow*.⁵⁰ Appropriate game challenges are neither too difficult nor too easy to meet. Optimal challenges make for an exceptionally pleasurable state of immersion in activity re-

ferred to as flow.^{50–53} This experiential state requires a perfect balance between game challenges and user skills.^{53,54} Appropriate games link flow to two ends. First, optimal challenge and flow boost players’ self-esteem because not-too-easy successes are experienced and ascribed to the self. Second, their (cognitive) behavioral repertoire is expanded with new responses, which in turn boosts self-esteem,⁵² as demonstrated in players of therapeutic games.^{54–56}

Gamification and the Persuasive Game Design model

This study aims to test whether a game specially designed to present users with carefully planned challenges and flow experiences contributes to patients learning desired behavior in an online CBT-based burnout treatment context. The design philosophy is gamification based on existing psychotherapy, enriching the proven practice of burnout CBT with games as an added tool for the therapist. The Persuasive Game Design model (Fig. 2) provides the framework for the game’s design. Motivating game elements, such as challenges, first transport the player into a game world where equivalents of real world tasks are carried out. Persuasive games can be so structured as to achieve transfer of effects from the experienced game world to the real world.⁴⁸ In the present case, transfer effects would result in real world skills reducing burnout symptoms.

Research expectations and hypotheses

The reported experiment had a two factor design: therapy versus no therapy and gaming versus nongaming. As the

FIG. 1. The Interyp burnout program,¹² including protocol for participants in the Game + Therapy condition.

FIG. 2. Application of the Persuasive Game Design model on burnout therapy.⁵⁹

game was designed to enrich current therapy, it was anticipated that therapy plus game would outperform therapy without it. However, as a successful therapy was gamified, it was expected that the game without therapy would be less effective than therapy without the game. In contrast, using the game without therapy was expected to have significant but less than optimal effects. Finally, there was a control condition where patients received no treatment at all.

The tested hypothesis, then, is that the treatment's effectiveness would increase over the experimental conditions in the following order: (a) no therapy or game (control), (b) game without therapy, (c) therapy without game, and (d) therapy including game.

Method

Participants

The study was embedded in the online Interapy burnout treatment program (Fig. 1). The experiment enrolled 101 people, 85 of whom completed the entire procedure. All patients had symptoms of burnout, and 76% of the patients were medically diagnosed as being burnout patients. The ages of participants ranged between 24 and 63 years old, with an average of 40 years ($SD=9.09$); 78% had achieved higher education, and 51% were male.

Design and procedures

The study is a quasi-experiment, with a 2×2 pretest-posttest mixed design and conditions Game Only (GO, $n=27$), Game+Therapy (GT, $n=17$), Therapy Only (TO, $n=20$), and No Game+No Therapy (NGNT, $n=21$). Experimental manipulations came into effect right after the awareness phase (Fig. 1). In TO and GT conditions, premeasures were taken before the awareness phase; in the remaining conditions, this was done upon participants' enrolment in the study. Postmeasures were taken 10–15 days after the premeasure when TO and GT participants were at the end of the behavioral change and cognitive restructuring phase. These participants received enjoyment questionnaires upon completing the game.

Participants in the GT condition were recruited from another cohort than those in the TO condition. Due to a lack of patients, it was not possible to run four groups at once. For this reason, data were initially collected for GT, TO, and NGNT conditions. Due to changes in the program manage-

ment, the TO condition could only be run 18 months later, effectively rendering the study into a quasi-experiment. Participants to GT and TO conditions were Interapy burnout patients, who were invited by their therapist. Participants to GO and NGNT conditions were recruited from two online burnout forums and randomly assigned. Ethical procedures of the certified therapy institute were strictly followed. All participants signed a consent form before taking part. Participants could win a book in return for their participation. The researcher was aware of the assigned condition of each patient, whereas patients and therapists were not.

Experimental material: the game

Assignments that were part of the original therapy module formed the basis of the levels of the game. In the gamification process, six general game elements were applied to increase the players' engagement and enjoyment of the game world (see Table 1).

The module *Dealing with Others* aims to train patients in dealing assertively with job colleagues and entertaining positive thoughts about themselves or their social environment. Abstract stylization of the game world reduced the challenge set by the first level. Later levels increased the challenge by requiring new behaviors in more settings, consolidation of positive thoughts, and a positive mind-set. Positive feedback from the game reinforced positive thinking, also lowering anxiety and depression and contributing to self-esteem, ultimately improving the player's capability to deal with others in the real world. Figure 3 shows a description of the therapeutic game. Patients played the game online, and there was no supervision from the therapists.

Dependent variables

Pre- and postmeasures included the following variables: (a) (precursive) burnout symptoms, (b) self-esteem, (c) coping skills, and (in the GO and GT conditions) (d) game enjoyment. Burnout symptoms are defined as symptoms affecting body, mood, and thoughts and consist of depression, anxiety, stress, and disengagement. The first three complaints were measured by items from the Depression Anxiety Stress Scale that uses 21 variables with a 7-point rating scale.⁵⁷ It should be noted that depression, anxiety, and stress are considered by therapists as *precursive symptoms*—early symptoms exhibited by burnout patients in the process of developing the syndrome. Disengagement is defined as

TABLE 1. DESCRIPTION OF THE GAME ELEMENTS THAT SERVED AS THE BASIS FOR THE DIGITAL GAME

<i>Game element</i>	<i>Element description</i>	<i>Therapeutic goal</i>
Narrative	The game encloses the storyline of a fictional character, who encounters problems deeming to be similar to those that the player has	Supplies a safe virtual environment for self-monitoring and learning behavior
Metaphor	The game uses metaphors, such as the filing cabinet as the human brain, to provide a surrounding for the mini games	Aids in the comprehension of the player's illness
Fun	The balance between game challenge and player skill will determine the amount of fun that the player will have. The narrative and the game quality can also enhance this experience	Ensures enjoyment and motivation to continue
Control	Whether the player needs to link up positive and negative thoughts or shoot negative thoughts out of the sky, the player's actions determine whether the game progresses or not	Gives the player the opportunity to meet challenges
Simulation	From the beginning to the end the game turns from an abstract simulated experience (clouds with text) to a more concrete and easily recognizable situation (recorded boss interview)	Serves as the foundation for practicing skills to be used beyond the gaming situation
Feedback	Feedback could either be textual ("You've done a great job!") or via immediate score information ("10 points")	Helps the player to understand if challenges have been met and underlines this progress with background information

distancing oneself from one's work and experiencing negative attitudes toward it.¹⁰ It was measured by eight items on a 7-point Likert scale from the Oldenburg Burnout Inventory.^{10,58}

Self-esteem is defined as a feeling of confidence in one's competence. It was measured by a 7-point Likert item from the Single Item Self Esteem (SISE) scale.⁵⁹ Coping skills involve setting of personal boundaries, positive self-image thinking, and restraining perfectionism. These three variables are standard measures of progress in Interapy's burnout program and are derived from validated instruments such as the Depression Anxiety Stress Scale and the Maslach Burnout Inventory.¹² Each were measured by five option multiple choice questions that were used as a test variable to see if the patient learned the desired behavior.

The experience of enjoyment while playing the game was measured only in the TG and GO conditions, using 7-point rating scales: likeable, funny, interesting, exciting, challenging, and useful. Flow was measured by three 7-point scales for challenge, immersion, and autonomy from the EGameFlow scale.⁶⁰

Control variables were gender, age, education, Internet usage, game usage, game attitude, and game completion.

Results

Treatment effects

Table 2 lists the results of pre- and posttest of precursive symptoms and of disengagement.

Precursive symptoms of burnout—depression, anxiety, and stress—were combined into one measure ($\alpha=0.76$). A multivariate analysis of covariance (ANCOVA) of combined complaints and disengagement, with treatment conditions as the between-subjects factor and pre-posttest as the within-subjects factor, was performed with gender, age, and education as covariates. Main effects of pre-posttest, $F(1, 78)=5.17$,

$p<0.05$, and treatment conditions, $F(3, 82)=6.15$, $p=0.001$, were found to indicate that levels of precursive symptoms and disengagement across pre- and posttest differed from one condition to another. An interaction effect between the pretest and the posttest and treatment conditions was detected, $F(3, 82)=7.29$, $p<0.001$, $\eta_p^2=0.210$, observed power=0.980.

Precursive symptoms

A main effect of pre-posttest was seen, $F(1, 82)=9.97$, $p<0.005$,^a as well as a main effect in the treatment condition, $F(3, 82)=2.86$, $p<0.05$, indicating that the levels of precursive symptoms across pre- and posttest differed from one condition to another. Post hoc tests showed a significant difference between therapy and nontherapy conditions. However, it was found that the difference between playing the game and not playing the game was not significant. The effect for pre-posttest disappeared when controlling for age, gender, and education, $F_{\text{pre-posttest}}(3, 78)=2.55$, $p=\text{n.s.}$

An interaction effect between pre-posttest and treatment conditions was found, $F(3, 78)=8.59$, $p<0.001$, $\eta_p^2=0.248$. Precursive symptoms exhibited a decreasing trend for the GT, TO, and GO conditions, while the NGNT condition exhibited an increasing trend (Fig. 4). A repeated measures analysis of variance with only GT and TO conditions showed a significant interaction between pre-posttest and condition factors, $F(1, 43)=7.15$, $p<0.05$, $\eta_p^2=0.152$; the steeper decrease of complaints in the GT compared to the TO condition was reliable.

Disengagement

Disengagement scores differed between the pretest and the posttest, and point out a main effect for pre-posttest, $F_{\text{pre-posttest}}(3, 78)=7.17$, $p<0.001$. However, no main effect was found for treatment condition, meaning that treatment

Level	Description and illustrations of the level
Intro	<p>The introduction presents in four screens the story of “Cloudy”, acting as the player’s avatar. Cloudy was once happy in his job but over time got overloaded with work. He feels highly responsible for his work, but due to exhaustion and feelings of helplessness he ends up staying at home and develops a huge feeling of guilt towards his colleagues.</p>
1. Match	<p>In this level the patient has to match positive and negative statements by dragging the right combinations to the boxes and press the ‘check’ button. In case of a correct match the player receives two points. In case of an incorrect match the player’s score is reduced by 1 point. The level is finished when all 8 correct combinations are made.</p>
2. Battle	<p>This level has the player confronted with a negative statement by a devil-like creature in the air, the so-called “Negativo”. For example, the Negativo states: “My boss does not like me” and the player can choose between the arguments A) “I do not like him either, so we are even”, B) “I cannot read his mind” or C) “It is as it is. I do not mind”. The player has to select the matching positive statement out of three options, aim, and shoot the negative statement. In this case the correct answer is “B: I cannot read his mind”, as it is a common pitfall for burnout patients to fill in the thoughts of others. For every chosen argument, the player receives feedback on why this argument is right or wrong and 2 points for the right argument and minus 1 point for a wrong argument.</p>

FIG. 3. Description of the game.

conditions did not differ as to disengagement level, $F_{\text{condition}}(1, 78)=0.314$, $p=\text{n.s.}$ Post hoc tests again showed significant differences between the therapy and nontherapy conditions, but not between game and no game ones. No interaction was obtained between pre–posttest and treatment condition ($p=0.18$).

Separate multivariate analyses of variance for depression, anxiety, stress, and disengagement showed differences between treatment conditions and between pre- and posttest, but no interaction effect of the two factors.

<p>3. Recognition</p>	<p>In this level the player is confronted with an animated typical work-related scenario. During the conversation negative and positive thoughts pop up at one of the animated figures, with whom the player identifies himself with. The thoughts appear for 5 seconds. If the player sees a negative thought appear, he should shoot this thought down by clicking on it. Five positive alternative thoughts pass by and the player should drag and drop the best one into the empty thinking cloud. For every choice, he will receive feedback. One point is lost when the answer is false and the player gets the chance to see the scenario again and choose the right answer.</p>
<p>4. Facing the Boss</p>	<p>During this level the player has to react to a request from a person that talks to him on the screen. This person is a real person acting out a work related situation that is recorded on video in advance. The player's response will be recorded via his webcam. The exercise can be compared with a role playing game, but then in a virtual world. After recording, he is able to review his recording and reflect on the exercise by answering questions like 'how did it go?' and 'did you recognize stress symptoms?'. At the end the player can choose to save the reflection text file and recording on his computer and decide whether he wants to share this recording with his therapist in order to receive feedback.</p>

Visual feedback was shaped using the metaphor of a filing cabinet. Burnout patients recognize themselves as struggling against a cabinet overloading with unfinished files. In the game successful performance filled the cabinet with compliments. Players received 2 credit points for a right and minus 1 point for a wrong answer. Resits were given for every failure. Entrance to the next level was granted upon reaching the designated score.

FIG. 3. (Continued).

Coping skill improvement and self-esteem

A main effect of pre–posttest for coping skill improvement, $F(1, 81)=31.60$, $p<0.001$, was found. Here, the treatment condition also had a main effect, $F(3, 81)=3.96$, $p<0.05$, while an interaction effect between pre–posttest and

treatment conditions was present as well, $F(3, 81)=4.96$, $p<0.005$, $\eta^2_p=0.155$. Post hoc tests for complaints and disengagement pointed out similar results. However, this time there was no difference between the TO condition and the GO condition. Finally, self-esteem was found to be stable from pre- to posttest and invariant across treatment conditions.

TABLE 2. MEANS AND STANDARD DEVIATIONS FOR PRECURSIVE SYMPTOMS AND DISENGAGEMENT PER TREATMENT CONDITION

	N	Pretest precursive symptoms		Posttest precursive symptoms		Pretest disengagement		Posttest disengagement	
		M	SD	M	SD	M	SD	M	SD
GT condition	17	4.21	0.94	3.44	0.88	4.30	0.86	3.78	1.20
TO condition	20	3.94	1.24	3.50	0.79	4.24	1.40	4.13	1.38
GO condition	27	4.52	1.58	4.43	1.49	5.16	1.80	5.16	1.63
NGNT condition	21	4.38	1.30	4.69	1.06	5.37	1.29	5.79	1.15

GT, Game + Therapy; TO, Therapy Only; GO, Game Only; NGNT, No Game + No Therapy.

FIG. 4. Plots for interaction between pre-posttest and treatment condition. Estimated marginal means for precursive symptoms.

The role of enjoyment and flow

Descriptive statistics for the experience of the game as a whole and for separate levels can be found in Table 3. A between subjects ANCOVA showed no differences in enjoyment of the game between GO and GT conditions, $F(1,43) = 1.99$, $p = n.s.$

In order to check the assumption that flow causes enjoyment, a regression analysis was performed on joint data of GT and GO conditions with flow, game behavior, game attitude, gender, age, and education as predictors.^b Flow predicted enjoyment ($\beta = 0.868$, $p < 0.001$), as did age ($\beta = -0.188$, $p < 0.01$). The effects of game enjoyment on improvement were explored (Table 4). A combined measure for game enjoyment did not predict any of the three reductions in complaints. However, enjoyment did predict improvement of coping skills, while skill improvement in turn predicted decrease of stress and depression symptoms.

Discussion

Findings

This pilot study aimed to test whether a therapeutic digital game is an efficient and suitable tool to aid patients in achieving therapeutic goals in an online CBT-based burnout treatment environment. As predicted, the two therapy conditions (GT and TO) showed a higher decrease of precursive complaints and disengagement as well as a stronger increase of coping skills than the nontherapy conditions (GO and

NGNT). As also expected, the GT condition outperformed the TO condition on improvement measures of precursive symptoms (taken as a combination of disengagement, depression, anxiety, and stress). However, analyses of separate measures did not reveal differential decreases across treatment conditions. Finally, skill improvement was more pronounced in the two therapy conditions, but no more so in the GT than the TO condition.

Differences in progress between therapeutic and non-therapeutic conditions can be interpreted to mean that participating in a CBT-based burnout program results in a reduction of the (precursive) burnout symptoms—depression, anxiety, stress, and disengagement—superior to not participating in such a program. In addition, it can be concluded that the therapeutic digital game may be an effective tool when embedded in a therapeutic burnout treatment program and is more effective than CBT alone as it is used in the current practice of Interapy. Caution is needed, however, because significantly better progress was found on the combined measures only. More importantly, the game as a standalone burnout treatment is not as successful as participating in a burnout treatment program. This lends support to a gamification approach that takes existing therapeutic practice as its starting point. That approach also contributed to the ecological validity of the results because measures could be used that have served as treatment outcomes for more than a decade.^c Major insights acquired in the therapeutic contribution of game play include (a) that experiencing the flow of playing the game boosted enjoyment and

TABLE 3. MEANS AND STANDARD DEVIATIONS FOR ENJOYMENT PER CONDITION ON A 9-POINT LIKERT SCALE

	Enjoyment		Enjoyment Level 1		Enjoyment Level 2		Enjoyment Level 3		Enjoyment Level 4		Flow	
	M	SD	M	SD	M	SD	M	SD	M	SD	M	SD
GT group	4.94	0.78	4.76	0.95	4.56	1.24	5.13	0.96	4.30	0.88	4.75	1.16
GO group	4.48	1.60	4.52	1.45	3.45	1.57	4.93	1.30	4.57	1.69	4.33	0.88

TABLE 4. OVERVIEW OF THE PERFORMED REGRESSION ANALYSES WITH MULTIPLE DEPENDENT VARIABLES

<i>Dependent variable</i>	<i>Predictor</i>	β	t	p	R^2
Disengagement (diff.)	Enjoyment	-0.137	-0.872	0.388 (n.s.)	0.289 (adj. R^2 =0.147)
	Relevant skill improvement (diff.)	-0.058	-0.343	0.733 (n.s.)	
	Self-esteem (diff.)	-0.112	-0.802	0.427 (n.s.)	
	Depression (diff.)	-0.328	-1.717	0.093 (n.s.)	
	Anxiety (diff.)	0.055	0.321	0.749 (n.s.)	
	Stress (diff.)	0.659	3.687	0.001	
	Gender	-0.067	-0.491	0.626 (n.s.)	
	Age	-0.014	-0.014	0.919 (n.s.)	
	Education	-0.078	-0.557	0.581 (n.s.)	
Stress (diff.)	Enjoyment	0.100	0.780	0.439 (n.s.)	0.506 (adj. R^2 =0.420)
	Coping skill improvement (diff.)	0.294	2.234	0.030	
	Self-esteem (diff.)	0.216	1.949	0.057 (n.s.)	
	Depression (diff.)	0.481	3.419	0.001	
	Anxiety (diff.)	0.311	2.323	0.025	
	Gender	0.162	1.463	0.150 (n.s.)	
	Age	0.071	0.657	0.514 (n.s.)	
	Education	-0.321	-1.156	0.254 (n.s.)	
Anxiety (diff.)	Enjoyment	-0.156	-1.176	0.246 (n.s.)	0.465 (adj. R^2 =0.372)
	Coping Skill Improvement (diff.)	-0.200	-1.417	0.163 (n.s.)	
	Self-esteem (diff.)	0.036	0.300	0.766 (n.s.)	
	Depression (diff.)	0.350	2.255	0.029	
	Stress (diff.)	0.337	2.323	0.025	
	Gender	0.018	0.154	0.879 (n.s.)	
	Age	-0.031	-0.276	0.784 (n.s.)	
	Education	0.122	1.017	0.315 (n.s.)	
Depression (diff.)	Enjoyment	0.137	1.147	0.257 (n.s.)	0.566 (adj. R^2 =0.491)
	Coping Skill Improvement (diff.)	-0.272	-2.205	0.033	
	Self-esteem (diff.)	-0.251	-2.469	0.017	
	Anxiety (diff.)	0.284	2.255	0.029	
	Stress (diff.)	0.422	3.419	0.001	
	Gender	-0.191	-1.864	0.069 (n.s.)	
	Age	0.055	0.540	0.592 (n.s.)	
	Education	0.192	1.823	0.075 (n.s.)	
Self-esteem (diff.)	Enjoyment	-0.071	-0.428	0.671 (n.s.)	0.064 (adj. R^2 =-0.031)
	Coping Skill Improvement (diff.)	0.216	1.329	0.190 (n.s.)	
	Gender	0.071	0.488	0.627 (n.s.)	
	Age	0.108	0.759	0.452 (n.s.)	
	Education	0.088	0.616	0.541 (n.s.)	
Coping skill improvement (diff.)	Enjoyment	0.494	3.959	0.000	0.301 (adj. R^2 =0.230)
	Self-esteem (diff.)	0.161	1.329	0.190 (n.s.)	
	Gender	-0.029	-0.234	0.816 (n.s.)	
	Age	-0.024	-0.196	0.845 (n.s.)	
	Education	0.119	0.961	0.341 (n.s.)	

Bold indicates significant results.

hence learning motivation, and (b) that enjoying the game indirectly targets burnout by facilitating acquisition of skills for coping with its symptoms.

Related work and future research

In the last decade, research on computerized CBT (cCBT) has been plentiful. Most studies imply that digitizing CBT in one way or another has great potential.^{61–64} Also, as mentioned earlier, many games and “playful interventions” have been developed for therapeutic purposes.^{33–45} However, the use of games as a tool for cCBT is quite uncommon. In fact, only one study of a game fully integrated into a CBT surrounding was found.⁶⁵ This study found high usability and acceptance among therapists and patients of a game under development, while the present study

tested treatment outcomes of a novel game integrated into actual therapy. More research should go toward the exploration of “gamified CBT” (gCBT), a term introduced now on the fly.

In further research, additional therapy modules may be gamified, possibly enhancing the present game’s potential. The design is on the right track; witness the finding that enjoyment is positively associated with practicing skills. Another improvement might be to grant the therapist a role in the currently unsupervised game, in order to reinforce desirable behavior in combination with the game’s feedback.

Limitations

A major setback to this study as a one module treatment manipulation is that it can only capture changes in a limited

number of outcome variables, leaving out, for example, emotional exhaustion symptoms and growth of self-esteem, which could not be established in any of the therapy conditions. This requires a longitudinal design across subsequent CBT modules.

Other shortcomings include:

- As is common in clinical treatment effect studies, assignment to conditions could not be completely random. Differences between conditions prior to treatment prevented direct comparison of postmeasures. The number of patients willing to participate was low. The number of patients in the GT condition ($n=17$) in particular may have reduced power to detect differences in comparison to the TO condition.
- The TO condition was carried out almost 18 months after the other conditions. This may have resulted in historical bias.
- Of all the patients, 76% were medically diagnosed as burnout patients. This statistic could only be established at the point of pretest, after the conditions were already assigned. In spite of burnout patients being randomly assigned, they were not equally divided across the conditions.
- The two therapy conditions may have developed treatment sensitization^{66,67} with unknown consequences.

The results of this study cannot be generalized to all burnout types and severities, and current game levels and situations may not fit with different treatment modules. Tailoring the game toward the specifics of other stressors than used here is a direction for future research to explore.

Conclusion

This study showed that people who experience the burnout symptoms of depression, anxiety, stress, and disengagement can be treated at least as effectively—and probably more effectively—when the therapist deploys a specially designed digital game compared with conventional therapy in a CBT-based burnout treatment program. It was found that experiencing flow and enjoyment, due to playing a properly designed therapeutic game, may ultimately benefit recovery from burnout symptoms and complaints. Thus, the research underlines the promise of gamifying mental health practices for improved patient experience.

Notes

- a. For all repeated measures analyses of variance, a GLM procedure (MANOVA) was used in SPSS.
- b. The adjusted R^2 for this model was 0.789.
- c. In addition to this, solid Cotan (Dutch commission of test affairs) scores were administered to the test scales that form the basis of the treatment.

Author Disclosure Statement

Sam van Tienhoven, Helle Sinkbaek, and Bart A.L. Schrieken are employees of Interapy and receive a regular salary from Interapy according to their labor contract. Alfred Lange is the founder of Interapy. When Interapy was sold to five mental health organizations in 2011, the financial rela-

tionship between Alfred Lange and Interapy ended. No other competing financial interests exist.

References

1. Maslach C, Leiter MP. (1998) *Burnout: oorzaken, gevolgen, remedies*. Amsterdam: Contact.
2. Centraal Bureau voor de Statistiek. (2005) Burn-out en psychische belasting. www.cbs.nl/nl-NL/menu/themas/gezondheid-welzijn/publicaties/artikelen/archief/2005/2005-1738-wm.htm (accessed Mar. 14, 2011).
3. Centraal Bureau voor de Statistiek. (2012) Burn-outklachten bij hoge werkdruk en bij weinig sociale steun. www.cbs.nl/nl-NL/menu/themas/arbeid-sociale-zekeerheid/publicaties/artikelen/archief/2012/2012-3712-wm.htm (accessed Dec. 9, 2012).
4. Korunka C, Tement S, Zdrehus C, et al. (2010) *Burnout: definition, recognition and prevention approaches*. Vienna: BOIT.
5. Goutas LJ. Burnout. *The Write Stuff* 2008; 17:135–136.
6. Karsten C. (2000) *Omgaan met burnout: preventie, hulp en reïntegratie*. Rijswijk: Elmar B.V.
7. Halbesleben JRB, Demerouti E. The construct validity of an alternative measure of burnout investigating the English translation of the Oldenburg Burnout Inventory. *Work & Stress* 2005; 19:208–220.
8. Kalliath TJ, O'Driscoll MP, Gillespie DF, et al. A test of the Maslach Burnout Inventory in three samples of health-care professionals. *Work & Stress* 2000; 14:35–50.
9. Lee RT, Ashforth BE. A meta-analytic examination of the correlates of the three dimensions of job burnout. *Journal of Applied Psychology* 1996; 81:123–133.
10. Demerouti E, Bakker AB, Vardakou I, et al. The convergent validity of two burnout instruments: a multitrait-multimethod analysis. *European Journal of Psychological Assessment* 2002; 19:12–23.
11. Hoogduin CAL, Schaufeli WB, Schaap CPDR, et al. (2001) *Behandelingsstrategieën bij burnout*. Houten: Bohn Stafleu Van Loghum.
12. Lange A, Van de Ven J-P, Schrieken B, et al. Interapy-burnout: preventie en behandeling van burnout via het internet. *Directieve Therapie* 2003; 23:121–145.
13. Deary IJ, Watson R, Hogston R. A longitudinal cohort study of burnout and attrition in nursing students. *Journal of Advanced Nursing* 2003; 43:71–81.
14. Etzion D. Moderating effect of social support on the stress–burnout relationship. *Journal of Applied Psychology* 1984; 69:615–622.
15. Rada RE, Johnson-Leong C. Stress, burnout, anxiety and depression among dentists. *Journal of the American Dental Association* 2004; 135:788–794.
16. Jeanneau M, Armelius K. Self-image and burnout in psychiatric staff. *Journal of Psychiatric & Mental Health Nursing* 2000; 7:399–406.
17. McKnight JD, Glass DC. Perceptions of control, burnout, and depressive symptomatology: a replication and extension. *Journal of Consulting & Clinical Psychology* 1995; 63:490–494.
18. Clarke LA, Watson D. Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications. *Journal of Abnormal Psychology* 1991; 100:316–336.
19. Endler NS. Stress, anxiety and coping: the multidimensional interaction model. *Canadian Psychology* 1997; 38: 136–153.

20. Slivar B. The syndrome of burnout, self-image, and anxiety with grammar school students. *Psihološka Obzorja* 2001; 10:21–32.
21. Farber BA. Treatment strategies for different types of teacher burnout. *Psychotherapy in Practice* 2000; 56:675–689.
22. Granath J, Ingvarsson S, Von Thiele U, et al. Stress management: a randomized study of cognitive behavioural therapy and yoga. *Cognitive Behaviour Therapy* 2006; 35:3–10.
23. Huibers MJH, Beurskens AJHM, Van Schayck CP, et al. Efficacy of cognitive-behavioural therapy by general practitioners for unexplained fatigue among employees. *The British Journal of Psychiatry* 2004; 184:240–246.
24. Fava GA, Rafanelli C, Grandi S, et al. Prevention of recurrent depression with cognitive behavioral therapy. *Archives of General Psychiatry* 1998; 55:816–820.
25. Borkovec TD, Costello E. Efficacy of applied relaxation and cognitive-behavioral therapy in the treatment of generalized anxiety disorder. *Journal of Consulting & Clinical Psychology* 1993; 61:611–619.
26. Korrelboom K, Ten Broeke E. (2004) *Geïntegreerde cognitieve gedragstherapie. Handboek voor theorie en praktijk*. Bussum: Uitgeverij Coutinho.
27. Ruwaard J. (2013) *The efficacy and effectiveness of online CBT*. Enschede: IpskampDrukkers BV.
28. Durkin K. Videogames and young people with developmental disorders. *Review of General Psychology* 2010; 14:122–140.
29. Gee JP. (2003) *What video games have to teach us about learning and literacy*. New York: Palgrave Macmillan.
30. Kato PM. Video games in health care: closing the gap. *Review of General Psychology* 2010; 14:113–121.
31. Squire K. Video games and education: designing learning systems for an interactive age. *Educational Technology Magazine* 2008; 48:17–26.
32. Abt CC. (1975) *Serious games*. New York: Viking Press.
33. Robillard G, Bouchard S, Fournier T, et al. Anxiety and presence during VR immersion: a comparative study of the reactions of phobic and non-phobic participants in therapeutic virtual environments derived from computer games. *CyberPsychology & Behavior* 2003; 6:467–476.
34. Bouchard S, Côte S, St-Jacques J, et al. Effectiveness of virtual reality exposure in the treatment of arachnophobia using 3D games. *Technology & Healthcare* 2006; 14: 19–27.
35. Walshe DG, Lewis EJ, Kim SI, et al. Exploring the use of computer games and virtual reality in exposure therapy for fear of driving following a motor vehicle accident. *CyberPsychology & Behavior* 2003; 6:329–334.
36. Rizzo A, Kim GJ. A SWOT analysis of the field of virtual reality rehabilitation and therapy. *Presence* 2005; 14:119–146.
37. Gardner JE. Can the Mario Bros. help? Nintendo games as an adjunct in psychotherapy with children. *Psychotherapy* 1991; 28:667–670.
38. Ceranoglu TA. Video games in psychotherapy. *Review of General Psychology* 2010; 14:141–146.
39. Ceranoglu TA. Star Wars in psychotherapy: video games in the office. *Academic Psychiatry* 2010; 34:233–236.
40. Barker J, Brinkman L, Deardorff M. Computer intervention for adolescent children of divorce. *Journal of Divorce & Remarriage* 1995; 23:197–213.
41. Strickland D, Marcus LM, Mesibov GB, et al. Brief report: two case studies using virtual reality as a learning tool for autistic children. *Journal of Autism & Developmental Disorders* 1996; 26:651–659.
42. Parsons S, Mitchell P, Leonard A. The use and understanding of virtual environments by adolescents with autistic spectrum disorders. *Journal of Autism & Developmental Disorders* 2004; 34:449–466.
43. Coyle D, Matthews M, Sharpy J, et al. Personal Investigator: a therapeutic 3D game for adolescent psychotherapy. *Interactive Technology & Smart Education* 2005; 2:73–88.
44. Coyle D, Doherty G, Sharpy J. An evaluation of a solution focused computer game in adolescent interventions. *Clinical Child Psychology & Psychiatry* 2009; 14:345–360.
45. Brezinka V, Hovestadt L. Serious games can support psychotherapy of children and adolescents. *Lecture Notes in Computer Science* 2007; 4799:357–364.
46. Tan ES-H. Entertainment is emotion: the functional architecture of the entertainment experience. *Media Psychology* 2008; 11:28–51.
47. Steen FF, Owens SA. Evolution's pedagogy: an adaptationist model of pretense and entertainment. *Journal of Cognition & Culture* 2001; 1:289–321.
48. Visch V, Vegt N, Anderiesen H, et al. Persuasive game design: a model and its definitions. *CHI Conference 2013*, Paris, France.
49. Tan E, Jansz J. (2008) The game experience. In Schifferstein R, Hekkert P, eds. *Product experience*. Amsterdam: Elsevier, pp. 531–556.
50. Csikszentmihalyi M. (2008) *Flow: the psychology of optimal experience*. New York: Harper Perennial.
51. Csikszentmihalyi M, Csikszentmihalyi IS, eds. (1988) *Optimal experience: psychological studies of flow in consciousness*. New York: Cambridge University Press.
52. Csikszentmihalyi M, LeFevre J. Optimal experience in work and leisure. *Journal of Personality & Social Psychology* 1989; 56:815–822.
53. Sherry JL. Flow and media enjoyment. *Communication Theory* 2004; 14:328–347.
54. Vorderer P, Hartmann T, Klimmt C. (2003) Explaining the enjoyment of playing video games: the role of competition. *ICEC '03 Proceedings of the Second Conference on Entertainment Computing*. Pittsburgh, PA: Carnegie Mellon University, pp. 1–9.
55. Deci EL, Koestner R, Ryan RM. A meta-analytic review of experiments examining the effects of extrinsic rewards on intrinsic motivation. *Psychological Bulletin* 1999; 125: 627–668.
56. Ryan RM, Rigby CS, Przybylski A. The motivational pull of video games: a self-determination theory approach. *Motivation & Emotion* 2006; 30:347–363.
57. De Beurs E, Van Dyck R, Marquenie LA, et al. De DASS: een vragenlijst voor het meten van depressie, angst en stress. *Gedragstherapie* 2001; 34:35–53.
58. Bakker AB, Heuven E. Emotional dissonance, burnout, and in-role performance among nurses and police officers. *International Journal of Stress Management* 2006; 13: 423–440.
59. Robins RW, Hendin HM, Trzesniewski KH. Measuring global self-esteem; construct validation of a single-item measure and the Rosenberg Self-Esteem Scale. *Personality & Social Psychology Bulletin* 2001; 27:151–161.
60. Fu F-L, Su R-C, Yu S-C. EGameFlow: a scale to measure learners' enjoyment of e-learning games. *Computers & Education* 2009; 52:101–112.

61. Baer L, Greist J, Marks IM. Computer-aided cognitive behavior therapy. *Psychotherapy & Psychosomatics* 2007; 76:193–195.
62. Vincent N, Walsh K, Lewycky S. Sleep locus of control and computerized cognitive-behavioral therapy (cCBT). *Behaviour Research & Therapy* 2010; 48:779–783.
63. Elsegood K, Powell D. Computerized cognitive-behaviour therapy (cCBT) and older people: a pilot study to determine factors that influence willingness to engage with cCBT. *Counseling & Psychotherapy Research: Linking Research with Practice* 2008; 8:189–192.
64. Titov N, Andrews G, Choi I, et al. Randomized controlled trial of web-based treatment of social phobia without clinician guidance. *Australian & New Zealand Journal of Psychiatry* 2009; 43:913–919.
65. Coyle D, McGlade N, Doherty G, et al. (2011) Exploratory evaluations of a computer game supporting cognitive behavioural therapy for adolescents. CHI '11 Proceedings of the SIGCHI Conference on Human Factors in Computing Systems, Vancouver, BC, Canada.
66. Bergin AE, Garfield SL. (1994) *Handbook of psychotherapy and behavior change*. Singapore: John Wiley.
67. Lana RE. (1969) Pretest sensitization. In Rosenthal R, Rosnow RL, eds. *Artifact in behavioral research*. New York: Academic Press, pp. 119–141.

Address correspondence to:
 Thomas Zielhorst
 Elsevier B.V.
 Global Electronic Operations
 Radarweg 29
 1043 NX Amsterdam
 The Netherlands

E-mail: thomas_zielhorst@hotmail.com