



UvA-DARE (Digital Academic Repository)

Hipparcos data on Herbig Ae/Be stars: an evolutionary scenario

van den Ancker, M.E.; The, P.S.; Tjin a Djie, H.R.E.; Catala, C.; de Winter, D.; Blondel, P.F.C.; Waters, L.B.F.M.

Published in:
Astronomy & Astrophysics

[Link to publication](#)

Citation for published version (APA):

van den Ancker, M. E., The, P. S., Tjin a Djie, H. R. E., Catala, C., de Winter, D., Blondel, P. F. C., & Waters, L. B. F. M. (1998). Hipparcos data on Herbig Ae/Be stars: an evolutionary scenario. *Astronomy & Astrophysics*, 324, L33-L36.

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*Letter to the Editor***Hipparcos data on Herbig Ae/Be stars: an evolutionary scenario** ^{*}M.E. van den Ancker¹, P.S. Thé¹, H.R.E. Tjin A Djie¹, C. Catala², D. de Winter³, P.F.C. Blondel¹, and L.B.F.M. Waters^{1,4}¹ Astronomical Institute “Anton Pannekoek”, University of Amsterdam, Kruislaan 403, NL–1098 SJ Amsterdam, The Netherlands² Laboratoire d’Astrophysique et Unité de Recherche associée au CNRS 285, Observatoire Midi-Pyrénées, France³ Dpto. Física Teórica C–XI, Facultad de Ciencias, Universidad Autónoma de Madrid, Cantoblanco, E–28049 Madrid, Spain⁴ SRON Space Research Laboratory, P.O. Box 800, 9700 AV Groningen, The Netherlands

Received 24 April 1997 / Accepted 30 May 1997

Abstract. Fundamental astrophysical parameters (distance, temperature, luminosity, mass, age) of a sample of 10 Herbig Ae/Be candidates and 3 non-emission line A and B stars in star forming regions were computed combining Hipparcos parallaxes with data from literature. All genuine Herbig stars in our sample are located between the birthline and the zero-age main sequence (ZAMS) in the Hertzsprung-Russell diagram (HRD), in accordance with what is expected for pre-main sequence stars. The region in the HRD close to the birthline is relatively devoid of stars when compared to the region closer to the ZAMS, in agreement with the expected evolutionary time scales. The Herbig Ae/Be stars not associated with star forming regions were found to be located close to the ZAMS. Additionally we discuss a possible evolutionary scenario for the circumstellar environment of Herbig stars.

Key words: Circumstellar matter – Stars: Distances – HR diagram – Stars: Pre-main sequence – Stars: Variables

1. Introduction

In a historical paper Herbig (1960) was the first to realize that the “Be and Ae stars associated with nebulosity” are in fact stars of intermediate mass still in their pre-main sequence (PMS) phase of evolution: stars which have lost most of their envelope of infalling gas and dust, but are not yet fusing hydrogen into helium and whose energy is mainly supplied by gravitational contraction. Criteria for membership of this class of stars, now more commonly known as Herbig Ae/Be (HAeBe) stars, are (Thé et al. 1994): (1) spectral type earlier than F8; (2) presence of emission lines; (3) presence of IR-excess in the spectral energy

distribution; (4) location in or near a probable star formation region.

Later, stars were found which are not near any plausible star formation region, but show all other characteristics of HAeBe stars. Whether these so-called isolated Herbig Ae/Be stars are young stellar objects remains controversial. Moreover, the membership criteria outlined above do not uniquely select intermediate-mass pre-main sequence stars, but will also select evolved massive stars (Davies et al. 1990) which are still located in the region where they formed. Therefore one has to be careful in identifying Herbig stars with pre-main sequence objects (Herbig 1992).

In this *letter* we will derive astrophysical parameters of a sample of 10 Herbig Ae/Be candidates as well as 3 non emission-line A and B-type stars in star formation regions for which the Hipparcos mission resulted in 3σ detections of the parallax. An attempt will be made to construct an evolutionary scenario for the circumstellar environment of HAeBe stars. In a forthcoming paper (van den Ancker et al. 1997b; Paper II) we will study the photometric behaviour of the Herbig Ae/Be stellar group using the photometric data on all Herbig stars provided by Hipparcos.

2. Data Analysis

Accurate absolute astrometry as well as photometry was obtained by Hipparcos on a sample of 38 A- and B-type stars in nearby star forming regions. This sample is a subset of a list of objects (Thé et al. 1982) which was accepted in 1982 by the Hipparcos consortium and which was released to the P.I. prior to the publication of the Hipparcos Catalogue (ESA 1997), in which a full description of the data products will be given. Table 1 lists the stars for which a 3σ detection of the parallax was achieved. A full analysis of the complete sample will be given in Paper II.

The first two columns in Table 1 give the Hipparcos number and the name of each star. The third column lists the trigono-

Send offprint requests to: M.E. van den Ancker (mario@astro.uva.nl)

* Based on data from the Hipparcos astrometry satellite.

Table 1. Astrophysical parameters of programme stars.

Name		π	f_{rej}	d	Association			d_{lit}	Ref.	Sp. Type	Ref.	A_V	$\log T_{\text{eff}}$	$\log L_*/L_\odot$	M	Age
HIP	Other	[mas]	[%]	[pc]	D.C.	R.N.	S.A.	[pc]			[m]			[M_\odot]	[Myr]	
22910	AB Aur	6.9 ± 1.0	0	144^{+23}_{-17}	L1519,(L1517)		Tau R2	140	(1)	A0Ve	(11)	0.50	4.00	$1.72^{+0.13}_{-0.11}$	2.4 ± 0.2	2.5 ± 1.0
34116	HD 53367	4.1 ± 1.4	8	250^{+120}_{-60}	L1657	S292	CMa R1	1150	(2)	B0IVe	(12)	2.23	4.50	$4.04^{+0.34}_{-0.26}$	13 ± 3	i 0.05
54413	HD 97048	5.7 ± 0.8	0	180^{+30}_{-20}	DC297.2-15.6	S135	CED 111	160	(3)	B9-A0ep+sh	(3)	1.24	4.00	$1.61^{+0.13}_{-0.10}$	2.5 ± 0.2	i 2
54557	HD 97300	5.3 ± 1.0	1	190^{+40}_{-30}			CED 112	160	(3)	B9V	(13)	1.33	4.02	$1.54^{+0.17}_{-0.15}$	2.5 ± 0.3	i 3
56379	HD 100546	9.7 ± 0.6	0	103^{+7}_{-6}	(DC296.2-7.9)			170	(4)	B9Vne	(14)	0.28	4.02	$1.51^{+0.06}_{-0.05}$	2.4 ± 0.1	i 10
58520	HD 104237	8.6 ± 0.5	0	116^{+8}_{-7}			Cha III	84	(4)	A0Vpe	(4)	0.71	3.98	$1.77^{+0.06}_{-0.06}$	2.5 ± 0.1	2.0 ± 0.5
79080	HR 5999	4.8 ± 0.9	0	210^{+50}_{-30}	DC339.7+9.2	S14	Lupus 3	140	(5)	A5-7III/IVe+sh	(12)	0.37	3.88	$1.89^{+0.17}_{-0.15}$	3.1 ± 0.5	0.6 ± 0.4
80462	HD 147889	7.4 ± 1.2	0	140^{+30}_{-20}	L1687	IC 4603	Sco R1	160	(6)	B2V	(15)	3.32	4.34	$3.32^{+0.16}_{-0.13}$	7.5 ± 1.0	i 0.1
81624	HD 150193	6.7 ± 1.7	3	150^{+50}_{-30}	L1729		Sco R1	160	(6)	A1Ve	(16)	1.61	3.97	$1.47^{+0.23}_{-0.19}$	2.3 ± 0.2	i 2
87819	HD 163296	8.2 ± 1.0	0	122^{+17}_{-13}				150	(7)	A1Ve	(16)	0.25	3.97	$1.48^{+0.12}_{-0.10}$	2.3 ± 0.1	i 2.5
93425	HD 176386	7.4 ± 1.2	0	140^{+30}_{-20}	DC359.8-17.9	NGC 6727		130	(8)	B9IV	(17)	0.62	4.03	$1.69^{+0.14}_{-0.13}$	2.7 ± 0.2	i 2
100289	BD+40 4124	9.3 ± 2.2	18	110^{+30}_{-20}	L888/L895	NGC 6910	Cyg R1	980	(9)	B2Ve	(18)	3.16	4.34	$2.04^{+0.23}_{-0.19}$		
103763	HD 200775	2.3 ± 0.6	0	430^{+160}_{-90}	L1174	NGC 7023	Cep R2	440	(10)	B2.5IVe	(19)	1.92	4.31	$3.89^{+0.26}_{-0.21}$	10 ± 2	0.02 ± 0.01

References to Table 1: (1) Elias (1978); (2) Herbst et al. (1982); (3) Whittet et al. (1997); (4) Hu et al. (1989); (5) Hughes et al. (1993); (6) Whittet (1974); (7) Thé et al. (1985); (8) Marraco & Rydgren (1981); (9) Shevchenko et al. (1991); (10) Whitcomb et al. (1981); (11) Böhm & Catala (1993); (12) Finkenzeller (1985); (13) Rydgren (1980); (14) Houk et al. (1975); (15) Cohen (1973); (16) Houk et al. (1988); (17) Houk et al. (1982); (18) Hillenbrand et al. (1995); (19) Rogers et al. (1995).

metric parallax measured by Hipparcos and the 1σ uncertainty in this (both in milliarcseconds). The fourth column lists the percentage of measurements that had to be rejected to arrive at the solution for the parallax given in the third column. The next column in Table 1 gives the distance, with its 1σ uncertainty, computed from the parallax. Columns 6–8 contain names of the star forming region in which the star is located. In column 6 an association with a dark cloud from the catalogue of Lynds (1962), or its extension to the southern hemisphere (Hartley et al. 1986) is given. When the name of the dark cloud is in parentheses, this indicates that the star is located near the edge, so the association of the star with the cloud might be doubtful. The next column in Table 1 lists the reflection nebula with which the star is associated, whereas the 8th column in Table 1 gives the name of a stellar aggregate with which the star is associated. A literature distance and a reference to the paper in which this was derived is provided in columns 9 and 10.

3. Astrophysical Parameters

In order to compute the total luminosity of the sample stars, photometric data, from the ultraviolet (ANS, TD1, IUE), through the optical (Walraven *WULBV*, Johnson/Cousins *UBVRI*) to the infrared (*JHKLMNQ*, IRAS), were collected from literature. Since many of the stars show strong variations in brightness, only optical and UV data obtained near maximum brightness were used. The stellar luminosity, extinction and effective temperature were computed following the method outlined in van den Ancker et al. (1997a). This method includes a correction for a possible anomalous extinction law towards the object. Since in H Ae Bes one often overestimates stellar temperatures from UV spectra due to the presence of heated layers in the immediate surroundings of the stars' photospheres (e.g. Blondel & Tjin A Djie 1994), and photometric methods to estimate stellar temperatures often yield erroneous results when the extinction law

for the circumstellar material is anomalous, as is often the case for such stars (e.g. Thé et al. 1996), we only use spectral types from literature based on optical spectra (Table 1). The computed visual extinction at maximum brightness A_V , effective temperature T_{eff} and stellar luminosity L_* are listed in columns 13–15 of Table 1.

In Fig. 1, we plot the programme stars in the HR diagram. The error in $\log T_{\text{eff}}$ is about 0.05 (or one subclass in spectral type), but individual data points may have larger errors. The error in luminosity is dominated by the error in the distance. Also shown in Fig. 1 are the pre-main sequence evolutionary tracks and the birthline (i.e. the line where a star first becomes optically visible on its evolution to the zero-age main sequence) computed by Palla & Stahler (1993). Using these evolutionary tracks and the isochrones given by the same authors, we can make an estimate of the masses and ages of our programme stars. These are listed in the last two columns of Table 1. Note that for HD 200775 the given age, based on it being a pre-main sequence star, is rather dubious. Given the short main sequence life time of a $10 M_\odot$ star and the fact that the post-main sequence lifetime of such a star is much longer than its pre-main sequence contraction time, it seems more likely that HD 200775 is already evolving away from the ZAMS, in which case its age would be around 20 Myr.

4. Discussion and Conclusions

The observed distribution of stars fits well with what is expected for pre-main sequence stars: they are almost all located between (or at) the zero-age main sequence (ZAMS) and the birthline. There is a good correlation between the luminosity class (Table 1) and the position in the HRD: All giants are located to the right of the ZAMS. The clustering of stars around the evolutionary track of mass $2.5 M_\odot$ in Fig. 1 is probably due to a selection effect: There are no Herbig Be stars in the nearest star forming




Fig. 1. Hertzsprung-Russell diagram of Herbig Ae/Be stars with parallaxes measured by Hipparcos. Open plot symbols indicate stars with $f_{\text{rej}} > 3\%$. Also shown are the theoretical pre-main sequence evolutionary tracks (solid lines and dashed lines) and the birthlines for 10^{-4} (upper dotted line) and $10^{-5} M_{\odot} \text{ yr}^{-1}$ (lower dotted line) by Palla & Stahler (1993).

regions. Since the pre-main sequence evolution of B-type stars proceeds much faster than those of their A-type counterparts this could reflect a difference in age between nearby and more distant star forming regions.

Peculiar is the position of BD+40°4124 in Fig. 1, far below the ZAMS. Such a position seems impossible for a young stellar object. BD+40°4124 illuminates a reflection nebula, in which many other young stellar objects can be found (Hillenbrand et al. 1995), so the possibility of this being an extremely rare evolved object seen projected towards a star forming region can be excluded. Therefore one or more of our assumptions used for deriving this position in the HRD must be wrong. Different authors in literature agree well on the spectral type of BD+40°4124, B2–3e. Also, a spectral type which would place it near the zero-age main sequence is not compatible with the observed spectral energy distribution. However, the distance as measured by Hipparcos (110 ± 30 pc) differs greatly from the literature value of the distance to the star forming region in which BD+40°4124 is located (1000 pc). Therefore we conclude that, although formally the Hipparcos data products give a 4σ detection for the parallax of BD+40°4124, this must be wrong. We note that for BD+40°4124 and HD 53367 the percentage of measurements that was rejected to arrive at the solution for the trigonometric parallax given in the Hipparcos Catalogue is very high, $f_{\text{rej}} > 3\%$ (indicated by open circles in Fig. 1). Furthermore, BD+40°4124 was flagged as a variability induced mover




Fig. 2. Observed (squares) and extinction-corrected (circles) SEDs for (a) HD 104237, (b) AB Aur, (c) HD 100546, (d) β Pic. Also shown are Kurucz (1991) models fitted to the extinction-corrected SEDs.

in the Hipparcos Catalogue, most probably indicating problems with the astrometric solution. This case may reflect the difficulty of parallax determinations by Hipparcos in nebulous regions. For HD 53367 the discrepancy between the literature and Hipparcos distance that can be seen in Table 1 is statistically hardly significant.

A fair number of stars in our sample cluster around the $2.5 M_{\odot}$ track (Fig. 1). This allows us to construct an evolutionary sequence for their circumstellar environment. We select the objects HD 104237, AB Aur and HD 100546 for which reliable ages were estimated (Table 1), and plot their spectral energy distributions (SEDs) in Fig. 2. Included in Fig. 2 is the SED of β Pic which is even older than HD 100546 (Crifo et al. 1997), but has a slightly lower mass. The younger stars in our sample have SEDs that tend to be more or less flat in the infrared, whereas for the more evolved ones the dust excess tends to have a double-peaked structure, with a broad dip in the energy distribution around $10 \mu\text{m}$. Extrapolating such a behaviour, we would expect to end up with a higher-mass equivalent of a star like β Pic. Interestingly, Waelkens et al. (1992) showed that the SEDs of Herbig stars can be classified as “flat” or “double-peaked” in the infrared and proposed an evolutionary scenario in which a broad dip around $10 \mu\text{m}$ develops with time. The ages derived from the Hipparcos data are consistent with such a scenario.

Spectroscopic evidence for evolution of the circumstellar dust component comes from observations with the Infrared Space Observatory ISO (Waelkens et al. 1996): HD 100546 shows a strong crystalline silicate component, similar to those seen in the ground-based $10 \mu\text{m}$ spectra of β Pic (Knacke et al. 1993) and of comets (Hanner et al. 1994). A recent ISO spectrum of the comet Hale-Bopp also shows strong crystalline silicates (Crovisier et al. 1997). In contrast, the object HD 104237 shows little evidence for such a crystalline dust component (Malfait, private communication). The occurrence of crystalline dust is an

indication for the degree of “processing” which the circumstellar material underwent after accretion from the molecular cloud, and is thus a measure of age. We find that for the small sample discussed here, the age derived from Hipparcos is consistent with the degree of processing seen in the ISO spectra.

We stress that the timescale for the evolution of the central star and the circumstellar material are probably not well coupled: studies of very young open clusters show that only a small percentage of the cluster stars, many of which are undoubtedly pre-main sequence in nature, show circumstellar gas and dust (e.g. van den Ancker et al. 1997a). Furthermore, it is known that some giants, with presumably lower stellar ages, do show a double-peaked infrared excess, making the construction of evolutionary sequences difficult. Provided that the initial configuration and spatial distribution of the circumstellar dust envelopes of the Herbig Ae/Be stars are similar, some unknown physical mechanism regulates the timescale at which the circumstellar matter is dispersed. So, while the *timescales* may be very different for individual stars, it is quite possible that the *sequence* is as proposed in Fig. 2.

It is tempting to speculate about the origins of the formation of a 10 μm dip. From the presence of 10 μm amorphous silicate emission in most Herbig Ae/Be stars, we know that the dust envelopes surrounding these stars must be optically thin, and that the presence of a broad dip around 10 μm must correspond to a physical gap in the radial distribution of dust around H AeBes. Such a gap can only be caused by perturbations by another body surrounding the central star. This could either be a low-mass star outside the dust shell, clearing the region between the inner and outer Lindblad resonances through long-range resonance effects (Lin & Papaloizou 1979), or a Jupiter-sized planet forming inside the dust envelope, clearing $2\sqrt{3}$ Roche lobe radii on either side of the planet (Artymowicz 1987). Since many of the H AeBes with faint companions detected in recent years by speckle interferometry as well as by Hipparcos (paper II) do not show a dip in their energy distributions, we consider the latter explanation more likely.

Acknowledgements. The authors would like to thank K. Malfait, C. Waelkens and P.R. Wesselius for many useful discussions on the ISO spectra of Herbig Ae/Be stars. DDW is supported in part by Spanish grant DGICYT PB94-0165. LBFMW acknowledges financial support from the Royal Netherlands Academy of Arts and Sciences, and through an NWO *Pionier* grant. This research has made use of the Simbad data base, operated at CDS, Strasbourg, France.

References

- van den Ancker, M.E., Thé, P.S., Feinstein, A., Vázquez, R.A., de Winter, D., Pérez, M.R. 1997a, A&AS, in press
- van den Ancker, M.E., Thé, P.S., Tjin A Djie, H.R.E., Catala, C., de Winter, D. 1997b, A&A, to be submitted (Paper II)
- Artymowicz, P. 1987, *Icarus* 70, 303
- Blondel, P.F.C., Tjin A Djie, H.R.E. 1994, ASP Conf. Ser. 62, 211
- Böhm, T., Catala, C. 1993, A&AS 101, 629
- Cohen, J.G. 1973, ApJ 186, 149
- Crifo, F., Vidal-Madjar, A., Lallement, R., Ferlet, R., Gerbaldi, M. 1997, A&A 320, L29
- Crovisier, J., Leech, K., Bockelée-Morvan, D., Brooke, T.Y., Hanner, M.S., Altieri, B., Keller, H.U., Lellouch, E. 1997, *Science* 275, 1904
- Davies, J.K., Evans, A., Bode, M.F., Whittet, D.C.B. 1990, MNRAS 247, 517
- Elias, J.H. 1978, ApJ 224, 857
- ESA, 1997, The Hipparcos Catalogue, ESA SP-1200
- Finkenzeller, U. 1985, A&A 151 340
- Hanner, M.S., Lynch, D.K., Russell, R.W. 1994, ApJ 425, 274
- Hartley, M., Manchester, R.N., Smith, R.M., Tritton, S.B., Goss, W.M. 1986, A&AS 63, 27
- Herbig, G.H. 1960, ApJS 4, 337
- Herbig, G.H. 1992, ASP Conf. Ser. 62, 3
- Herbst, W., Miller, D.P., Warner, J.W., Herzog, A. 1982, AJ 87, 98
- Hillenbrand, L.A., Meyer, M.R., Strom, S.E., Skrutskie, M.F. 1995, AJ 109, 280
- Houk, N., Cowley, A.P. 1975, Michigan Spectral Survey, Vol. 1, Univ. Michigan Press
- Houk, N. 1982, Michigan Spectral Survey, Vol. 3, Univ. Michigan Press
- Houk, N., Smith-Moore, A. 1988, Michigan Spectral Survey, Vol. 4, Univ. Michigan Press
- Hu, J.Y., Thé, P.S., de Winter, D. 1989, A&A 208, 213
- Hughes, J., Hartigan, P., Clampitt, L. 1993, AJ 105, 571
- Knacke, R.F., Fajardo-Acosta, S.B., Telesco, C.M., Hackwell, J.A., Lynch, D.K., Russell, R.W. 1993, ApJ 418, 440
- Kurucz, R.L. 1991, Precision photometry: Astrophysics of the galaxy (Eds. A.G. Davis Philip, A.R. Upgren, K.A. Janes), L. Davis press, Schenectady, New York
- Lin, D.N.C., Papaloizou, J. 1979, MNRAS 188, 191
- Lynds, B.T. 1962, ApJS 7, 1
- Marraco, H.G., Rydgren, A.E. 1981, AJ 86, 62
- Palla, F., Stahler, S.W. 1993, ApJ 418, 414
- Rogers, C., Heyer, M.H., Dewdney, P.E. 1995, ApJ 442, 694
- Rydgren, A.E. 1980, AJ 85, 444
- Shevchenko, V.S., Ibragimov, M.A., Chernysheva, T.L. 1991, SvA 35, 229
- Thé, P.S., Felenbok, P., Cuypers, H., Tjin A Djie, H.R.E. 1985, A&A 149, 429
- Thé, P.S., Tjin A Djie, H.R.E., Praderie, F., Catala, C. 1982, ESA SP-177, 105
- Thé, P.S., de Winter, D., Pérez, M. 1994, A&AS 104, 315
- Thé, P.S., Pérez, M.R., Voshchinnikov, N.V., van den Ancker, M.E. 1996, A&A 314, 233
- Waelkens, C., Bogaert, E., Waters, L.B.F.M. 1992, ASP Conf. Ser. 62, 405
- Waelkens, C., Waters, L.B.F.M., de Graauw, M.S., et al. 1996, A&A 315, L245
- Whitcomb, S.E., Gatley, I., Hildebrand, R.H., Keene, J., Sellgren, K., Werner, M.W. 1981, ApJ 246, 416
- Whittet, D.C.B. 1974, MNRAS 168, 371
- Whittet, D.C.B., Prusti, T., Franco, G.A.P., Gerakines, P.A., Kilkenny, D., Larson, K.A., Wesselius, P.R. 1997, A&A, in press