UvA-DARE (Digital Academic Repository)

Working memory and cognitive flexibility-training for children with an autism spectrum disorder: a randomized controlled trial

de Vries, M.; Prins, P.J.M.; Schmand, B.A.; Geurts, H.M.

DOI

[10.1111/jcpp.12324](https://doi.org/10.1111/jcpp.12324)

Publication date

2015

Document Version

Final published version

Published in

Journal of Child Psychology and Psychiatry

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

de Vries, M., Prins, P. J. M., Schmand, B. A., & Geurts, H. M. (2015). Working memory and cognitive flexibility-training for children with an autism spectrum disorder: a randomized controlled trial. *Journal of Child Psychology and Psychiatry*, 56(5), 566-576.
<https://doi.org/10.1111/jcpp.12324>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Working memory and cognitive flexibility-training for children with an autism spectrum disorder: a randomized controlled trial

Marieke de Vries,^{1,2} Pier J.M. Prins,¹ Ben A. Schmand,^{1,3} and Hilde M. Geurts^{1,2,4}

¹Univeristy of Amsterdam, Amsterdam; ²Dutch Autism and ADHD Research Center, Amsterdam; ³Academic Medical Center Amsterdam, Amsterdam; ⁴Dr. Leo Kannerhuis, Amsterdam, the Netherlands

Background: People with autism spectrum disorders (ASDs) experience executive function (EF) deficits. There is an urgent need for effective interventions, but in spite of the increasing research focus on computerized cognitive training, this has not been studied in ASD. Hence, we investigated two EF training conditions in children with ASD.

Methods: In a randomized controlled trial, children with ASD ($n = 121$, 8–12 years, $IQ > 80$) were randomly assigned to an adaptive working memory (WM) training, an adaptive cognitive flexibility-training, or a non-adaptive control training (mock-training). Braingame Brian, a computerized EF-training with game-elements, was used. Outcome measures (pretraining, post-training, and 6-week-follow-up) were near-transfer to trained EFs, far-transfer to other EFs (sustained attention and inhibition), and parent's ratings of daily life EFs, social behavior, attention deficit hyperactivity disorder (ADHD)-behavior, and quality of life. **Results:** Attrition-rate was 26%. Children in all conditions who completed the training improved in WM, cognitive flexibility, attention, and on parent's ratings, but not in inhibition. There were no significant differential intervention effects, although children in the WM condition showed a trend toward improvement on near-transfer WM and ADHD-behavior, and children in the cognitive flexibility condition showed a trend toward improvement on near-transfer flexibility. **Conclusion:** Although children in the WM condition tended to improve more in WM and ADHD-behavior, the lack of differential improvement on most outcome measures, the absence of a clear effect of the adaptive training compared to the mock-training, and the high attrition rate suggest that the training in its present form is probably not suitable for children with ASD.

Keywords: Executive functioning, working memory, flexibility, cognitive training, autism.

Introduction

Executive functions (EFs) are cognitive functions needed for goal directed behavior that is adaptive to the environment. EFs are important in many aspects of life, such as school success, and well-developed EFs can predict lifelong achievement (Diamond, 2013). A growing body of evidence suggests EFs can be trained (e.g., Melby-Lervåg & Hulme, 2013; Zinke, Einert, Pfennig, & Kliegel, 2012), although generalization to other functions and to daily life (Holmes, Gathercole, & Dunning, 2009), and long term effects (Roughan & Hadwin, 2011) are not well established. EF training has been studied in typically developing (TD) people (e.g., Alloway, Bibile, & Lau, 2013) and clinical groups, such as children with Attention Deficit Hyperactivity Disorder (ADHD: e.g., Klingberg et al., 2005). One of the mainstream cognitive theories on autism spectrum disorders (ASD) postulates that EF deficits are the underlying problem (Russell, 1997). Moreover, EF deficits in ASD are related to adaptive behavior (Gilotty, Kenworthy, Sirian, Black, & Wagner, 2002), which in turn is associated with quality of life (Chiang & Wineman, 2014). However, only a few studies focused on EF interventions in ASD (Baltruschat et al., 2011; Fisher & Happé, 2005; Kenworthy

et al., 2014). ASDs are characterized by difficulties in social interaction and communication, and restricted and repetitive behavior and interests (American Psychiatric Association, 2000). These difficulties interfere with daily life, and may partly be caused by EF deficits (Damasio & Maurer, 1978; Geurts, Vries, & van den Bergh, 2014; Russell, 1997). Hence, an EF intervention may be fruitful for ASD.

Some intervention studies in ASD included EF as an outcome measure. Although a Theory of Mind (ToM)-training did not influence EF (Fisher & Happé, 2005), positive reinforcement (Baltruschat et al., 2011) and a social competence intervention (Stichter et al., 2010) did show positive effects on EF, although studied without a control group. A Social Skills training also improved EF, but to a lesser extent than a direct EF intervention (Kenworthy et al., 2014). The effects of direct EF interventions in ASD have hardly been studied, and findings are mixed; both positive (Kenworthy et al., 2014) and null effects (Fisher & Happé, 2005) have been reported. Children with ASD do show development of EFs (Pellicano, 2010a), which implies some plasticity. In sum, there are tentative indications that targeting EFs in interventions for ASD is promising, and that the best approach is to target EFs directly.

Working memory (WM)-training has been studied quite extensively in both clinical and TD groups (e.g.,

Conflict of interest statement: No conflicts declared.

Melby-Lervåg & Hulme, 2013), in particular, the computerized WM-training Cogmed (Chacko et al., 2013; Shipstead, Hicks, & Engle, 2012). Findings are inconsistent. Both near-transfer (improvement of the trained function) and far-transfer (improvement of other functions) have been reported, even when compared to an active control group, who performed a nonadaptive version of the training task (e.g., Holmes et al., 2009; Klingberg et al., 2005; Thorell, Lindqvist, Bergman Nutley, Bohlin, & Klingberg, 2009). However, recent reviews and meta-analyses are critical. Short term near-transfer to verbal and visual-spatial WM is reported, but long term and far-transfer (to attention, ADHD-behavior, and reasoning) is small and non-significant (Melby-Lervåg & Hulme, 2013). Hence, the effects are not (yet) evident, also due to lack of replication and caveats in the studies (Shipstead, Redick, & Engle, 2012). Clearly, more methodologically sound WM-training research is needed. Moreover, there is a striking absence of studies on WM-training in ASD.

Although flexibility deficits in ASD are not consistently reported in the research setting, inflexible behavior in daily life is evident (Geurts, Corbett, & Solomon, 2009; Kenworthy, Yerys, Anthony, & Wallace, 2008; Leung & Zakzanis, 2014). Therefore, a flexibility-training might be particularly fruitful. The few studies that focused on flexibility-training reported improved flexibility, interference control, and verbal WM in children with ADHD (Kray, Karbach, Haenig, & Freitag, 2011) and TD people (Karbach & Kray, 2009). Hence, flexibility seems trainable.

In spite of evident EF deficits in ASD and positive results of WM and flexibility-training in related disorders, studies on EF training in ASD are lacking. Therefore, we studied the effect of a WM and a

flexibility-training compared to an active control condition (mock-training) in a large randomized controlled trial of children (8–12 years) with ASD. Repetition of monotonous tasks in EF training reduces motivation and increases attrition-rates, and adding game-elements enhances motivation, and may improve the training effect (Prins, Dovis, Ponsioen, Ten Brink, & Van der Oord, 2011). We, therefore, used an EF training with game-elements: Braingame Brian (Prins et al., 2013), that showed promising pilot-results in ADHD (Oord, Ponsioen, Geurts, Brink, & Prins, 2014). We expected near-transfer to WM and flexibility respectively, and far-transfer to other EF tasks (inhibition, sustained attention), and to parent's ratings of daily life EFs and behavior (Figure S1).

Methods

Participants

Families were approached through mental health care institutions and internet advertisements. Of the initially 166 signed up families, 34 did not participate because of the required time investment (see Figure 1). The remaining 132 children were screened for (a) a prior independent clinical ASD diagnosis according to the Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV: American Psychiatric Association, 2000), diagnosed by a multidisciplinary team specialized in ASD, (b) age 8–12 years, (c) IQ above 80, and (d) absence of a seizure disorder. First, the diagnosis was verified if the score on the Social Responsiveness Scale parent report (SRS: Constantino et al., 2003; Roeyers, Thys, Druart, De Schryver, & Schittekatte, 2011) was above the specified Dutch clinical ASD cut-off of 57 (raw score), if Autism Diagnostic Interview Schedule-Revised (ADI-R: De Jonge & de Bildt, 2007; Lord, Rutter, & Couteur, 1994) cut-offs were reached on two out of the three domains, and if ASD was recognized before the age of 36 months (Gray, Tonge, & Sweeney, 2008). Second, IQ was

Figure 1 Flow chart of inclusion for RCT (conforming to CONSORT)

estimated with two subtests of the Dutch version of the Wechsler Intelligence Scale for Children (WISC-III: Kort et al., 2002; Sattler, 2001). Eleven children were excluded after screening (2 ADI-R, 6 IQ, 3 personal circumstances). One hundred and twenty-one children were randomly (double blind) assigned to the three intervention-conditions. Six children dropped out after pretraining. Finally, 115 children performed the training (40 WM, 37 flexibility, 38 mock-training). Twenty-five children did not complete 25 sessions (9 WM, 10 flexibility, and 6 mock-training; similar in the three conditions [$\chi^2(2) 1.3, p = .52$]). Forty children used psychotropic medication (17 abstained during test-appointments, 23 continued). Of the 90 children who completed the training, 24 children used medication (10 abstained, 14 continued). Years since diagnosis was 2.1 ($SD = 1.7, N = 84$, as six parents did not indicate the date of diagnosis).

Tasks and questionnaires

Measures include screening and four steps of outcome measures: (a) near-transfer to WM and flexibility tasks resembling the training tasks, (b) near-transfer to WM and flexibility tasks differing from the training tasks, (c) far-transfer to inhibition and sustained attention tasks, and (d) far-transfer to daily life (see Figure 2).

Inclusion/screening. *IQ* (Kort et al., 2002). The WISC-III subtests Vocabulary and Block Design, used to estimate IQ, correlate highly with Full Scale IQ, and have good reliability (Legerstee, van der Reijden-Lakeman, Lechner-van der Noort, & Ferdinand, 2004; Sattler, 2001).

Diagnosis: *The SRS* (Constantino et al., 2003; Dutch version: Roeyers et al., 2011) is a reliable and valid questionnaire measuring autism characteristics (Bolte, Poustka, & Constantino, 2008). The ADI-R (Lord et al., 1994; Rutter, Lord, & LeCouteur, 2003) is an extensive semi-structured interview including three domains of autism symptoms (social, communication, and repetitive behavior). Reaching specified cut-offs on these domains, and clear indications of abnormalities in development before 36 months, suggests a DSM-IV (American Psychiatric Association, 2000) or ICD-10 (World Health Organisation, 1992) diagnosis of autism (Gray et al., 2008; Rutter et al., 2003).

WM and flexibility tasks resembling the training tasks. *WM:* The Corsi block tapping task (Corsi-BTT: Corsi, 1972) measures visual-spatial WM (WM-training near-transfer, flexibility-training far-transfer). In the currently used

computerized version, nine blocks on a computer screen light up (500 ms each) in a certain sequence, which has to be repeated with mouse-clicks. The initial sequence length is three blocks, and after every four trials, the sequence length increases with one, up to a maximum of eight blocks. The task ends after four incorrectly repeated sequences of the same length. The outcome measure is the longest correctly repeated sequence length.

Flexibility: The Gender-emotion switch-task (de Vries & Geurts, 2012) is an adaptation of the classical switch-task (Rogers & Monsell, 1995; flexibility-training near-transfer, WM-training far-transfer). Pictures of easy to recognize happy or angry male and female faces (Lundqvist, Flykt, & Öhman, 1998) are displayed on the computer screen, and have to be sorted, alternatingly, on gender or emotion by pressing designated keys on the keyboard (de Vries & Geurts, 2012). The sorting rule changes randomly after two, three, or four trials (30% switch-trials). Children with ASD are quite capable to recognize these basic emotions (Geurts, Luman, & van Meel, 2008), hence emotion recognition most likely does not influence performance. Outcome measures are error rates (ER) and reaction time (RT) switch costs (switch-trial minus repeat-trial scores).

WM and flexibility tasks different from the training tasks. *WM:* In the N-back task (Casey et al., 1995; Smith & Jonides, 1999) participants have to remember pictures that are presented on a computer screen and indicate if the current picture matches the picture N trials before. Three levels with increasing WM-load were included: 0-back (baseline, indicate if the picture is a dog or not), 1-back, and 2-back (de Vries & Geurts, 2014). The outcome measures were the ER and RT on the 2-back, and 1-back task (corrected by subtraction for ER and RT on the 0-back task).

Flexibility: In the Number-gnome switch-task, an adaptation of the number-switch-task (Cepeda, Cepeda, & Kramer, 2000) pictures of one, two or three gnomes (Geirnaert, 2010) with a speech bubble with the number '1', '2', or '3' is displayed. Participants have to report the number of gnomes, or the number in the speech bubble, by pressing the 1, 2, or 3 on the keyboard. The task design is identical to the gender-emotion switch-task. Outcome measures were ER, and RT switch costs.

Far-transfer to inhibition and sustained attention tasks. *Inhibition:* In the current adaptation of the classical Stop-task (Logan, 1994) a picture of a yellow dog facing

Figure 2 Inclusion and outcome measures. ADI-R, Autism Diagnostic Interview revised; BRIEF, Behavioral Rating Inventory of Executive Functioning; Corsi-BTT, Corsi block tapping task; CSBQ Children's Social Behavior Questionnaire; DBDRS, Disruptive Behavior Disorders Rating Scale; PedsQL, Pediatric Quality of Life questionnaire; SART, Sustained Attention Response Task; SRS, Social Responsiveness Scale, WISC-III, Wechsler Intelligence Scale for Children, 3rd version

left or right appears on the computer screen. Participants have to press a corresponding left or right key on the keyboard as fast as possible. On 30% of the trials, the dog turns red (stop-signal), and the response has to be inhibited (de Vries & Geurts, 2014). The outcome measure was the stop-signal reaction time (SSRT: Logan & Cowan, 1984).

Sustained attention: In the Sustained attention response task (SART: Robertson, Manly, Andrade, Baddeley, & Yiend, 1997) numbers 1–9 appear on the computer screen (250 ms) followed by a cross (1000 ms or until response). After each number, except '3', when the cross appears, the spacebar has to be pressed. The outcome measures were ER and RT.

Far-transfer to daily life. EF: The Behavior Rating Inventory of Executive Function (BRIEF: Gioia, Isquith, Guy, & Kenworthy, 2000; Dutch Version: Smidts & Huizinga, 2009; 75 items, 3-point Likert-scale) has high to very high internal consistency and test-retest stability (Huizinga & Smidts, 2010). The subdomains Shift and WM, and the total scale were the outcome measures.

Social behavior: The Children's Social Behavior Questionnaire (CSBQ, Dutch version: Hartman, Luteijn, Moorlag, De Bildt, & Minderaa, 2007; 49 items, 3-point Likert-scale) has good internal consistency and test-retest stability (Hartman, Luteijn, Serra, & Minderaa, 2006). The total scale was the outcome measure.

ADHD-behavior: The Dutch parent version of the Disruptive Behavior Disorders Rating Scale (Oosterlaan, Scheres, Antrop, Roeyers, & Sergeant, 2000; DBDRS: Pelham, Gnagy, Greenslade, & Milich, 1992; 42 items, 4-point Likert-scale) has good psychometric properties (Oosterlaan et al., 2000). The subscales inattention, and hyperactivity/impulsivity combined (DBDRS-ADHD) were used as the outcome measure.

Quality of life: The Pediatric Quality of Life Inventory (PedsQL: Bastiaansen, Koot, Bongers, Varni, & Verhulst, 2004; Varni, Seid, & Kurtin, 2001; 23 items, 5-point Likert-scale) has satisfactory reliability and validity (Varni et al., 2001). The total scale was used as the outcome measure.

Intervention

Each child performed one of three intervention-conditions of 'Braingame Brian' (Prins et al., 2013): a WM-training, a flexibility-training, or a mock-training (Appendix S1 and Figure S3). In each intervention-condition, all EF tasks were performed but whether the level was adaptive differed per intervention-condition. The WM-training included five adaptive WM-training tasks with increasing difficulty, and the other tasks remained at a low level. The flexibility-training included an adaptive flexibility task, and the other tasks remained at a low level. In the mock-training all tasks remained at a low, nonadaptive level. In the flexibility-training and mock-training only the first most basic task of the five WM-training tasks was performed.

Procedure

First, written informed consent, SRS, and screening questionnaires were obtained. Next, training and test-appointments [screening, pretraining, post-training (after 6 weeks), and follow-up (FU, after 6 more weeks)] were scheduled. During screening, the ADI-R was administered to the parents, and the WISC-III subtests to the children. Parents, children, and assessors were kept blind to intervention-condition until after

follow-up. At pretraining, post-training, and FU (at the University of Amsterdam) all outcome tasks were administered to the children, and parents filled out the questionnaires. Pretraining a laptop with Braingame Brian installed was handed over and children and parents were instructed. Children and parents were told that there were three intervention-conditions, but not how these conditions were designed. All received similar basic training instructions and were told that tasks could change over time, but that this was not always noticeable. During the six training-weeks, parents were contacted weekly about the child's progress. In these weekly telephone calls, parents were asked about the current progress in the training, motivation of the child, and possible reward systems used. After the training, parents were asked about the amount of parental support their child received during the training. The number of completed sessions was retrieved from the automatically saved log-files of the training. The test-appointments lasted 50–150 min (as additional, for the current study irrelevant, tasks and questionnaires were administered). Children received a small gift for participating. Parents received a report with the ADI-R results. Travel expenses were largely covered.

Data analysis

The main (per-protocol) analyses included children who finished 25 sessions. If applicable, inverse efficiency scores (IES = mean RT/(1-ER; Bruyer & Brysbaert, 2011) were used as an outcome measure, to take possible speed-accuracy trade-off into account. IES are reliable when RT and ER are correlated, and ER <10% (Bruyer & Brysbaert, 2011), which was satisfied for the SART.

First, *t*-tests and chi-square tests were conducted on screening and pretraining measures to test if children who finished the training differed from the drop-outs, and a MANOVA was conducted to test if children in the three intervention-conditions differed. Second, two repeated measures ANOVAs were conducted to test whether children in the WM, and flexibility intervention-condition improved on the within-training tasks (WM: mean final reached sequence length on the five training tasks, flexibility: final level reached). Third, training effects were examined with four (to increase power, missing data-points 1.47%) repeated measures MANOVA's with intervention (WM, flexibility, and mock-training) as between subject variable, and time (pretraining, post-training, and FU) as within subject variable. Outcome measures were (a) Corsi-BTT and n-back task (near-transfer WM), (b) gender-emotion switch-task and number-gnome switch-task (near-transfer flexibility), (c) SART, and stop task, and (d) questionnaires (far-transfer). Additional intention-to-treat analyses (last observation carried forward) were conducted: the four MANOVA's on intervention effects were repeated including all participants, with number of completed sessions as covariate. Finally, we explored in children who performed all sessions if within-training-improvement (mean of five WM-training tasks, and reached level of flexibility-training task) was correlated with near-transfer training-outcome (Corsi-BTT and gender-emotion task improvement between pretraining and post-training).

Bonferroni corrections for multiple testing were applied (screening and pretraining: $0.05/25 = 0.002$, training effects: $0.05/4 = 0.0125$). Analyses were conducted both with and without children who used medication during test-appointments. Due to missing data, numbers differed slightly for different measures (see Table 1).

Results

Screening and pretraining

The drop-outs of the intervention used medication slightly more often [$\chi^2(1) = 6.5, p = .04$], and showed

Table 1 Mean (standard deviations) and MANOVAs for near-transfer and far-transfer measures

Measure	Group							Comparison ^a					
	Pre-training		Post-training		FU		Time	Time*Intervention					
	Pre-training	Post-training	Pre-training	Post-training	Pre-training	Post-training			Time				
	WM-training <i>n</i> = 31		Flexibility-training <i>n</i> = 26		Mock- training <i>n</i> = 29								
Corsti-BTT	5.2 (0.7)	6.3 (1.0)	6.2 (0.8)	5.5 (0.9)	6.0 (1.0)	5.8 (0.8)	5.1 (1.2)	5.7 (1.2)	5.8 (1.1)	26.6***	.24	2.6*	.06
N-back ^{bc}													
ER1-back	9.0 (12.1)	1.8 (5.3)	0.8 (6.2)	4.3 (7.9)	1.9 (6.8)	1.0 (4.8)	8.7 (11.8)	3.7 (6.9)	3.0 (7.5)	17.1***	.17	1.2	.03
ER2-back	18.1 (12.6)	12.2 (8.5)	11.5 (8.1)	16.3 (7.7)	12.0 (8.8)	7.8 (6.0)	17.5 (11.9)	12.5 (8.4)	10.1 (7.7)	22.6***	.21	0.4	.01
RT1-back	67.6 (41.4)	43.8 (54.6)	47.5 (44.4)	65.2 (39.5)	51.1 (41.9)	38.0 (32.9)	91.5 (46.2)	62.7 (55.2)	52.3 (46.2)	11.8***	.12	0.7	.02
RT2-back	77.3 (60.5)	71.3 (64.9)	72.0 (39.8)	79.0 (61.6)	89.1 (51.4)	59.4 (62.2)	113.0 (74.7)	103.2 (65.5)	85.4 (59.0)	2.9	.03	0.8	.02
		<i>n</i> = 30			<i>n</i> = 26			<i>n</i> = 29		<i>F</i> _(2, 164)	η^2_p	<i>F</i> _(2, 236)	η^2_p
Switch costs on the gender-emotion switch- task													
RT	109.3 (73.0)	96.7 (58.2)	91.6 (57.1)	107.1 (81.3)	88.5 (68.7)	91.1 (77.9)	108.0 (63.9)	62.7 (42.5)	79.6 (67.4)	5.0**	.06	0.7	.02
ER	5.2 (8.0)	8.8 (6.9)	6.3 (5.8)	8.2 (6.6)	6.9 (7.5)	6.2 (5.1)	6.1 (7.2)	7.6 (6.4)	5.9 (5.5)	1.6	.02	1.1	.03
Switch costs on the number-gnome switch-task													
RT	64.0 (79.3)	56.7 (58.0)	74.0 (67.0)	67.0 (79.7)	52.3 (51.8)	70.7 (64.7)	71.9 (63.2)	46.9 (37.6)	54.7 (59.0)	2.5	.03	0.6	.01
ER	6.8 (6.4)	7.2 (7.1)	7.7 (7.3)	9.1 (5.6)	8.4 (6.2)	6.3 (5.4)	6.2 (7.4)	7.8 (5.7)	7.6 (6.5)	0.2	.00	1.1	.03
		<i>n</i> = 30			<i>n</i> = 25			<i>n</i> = 25		<i>F</i> _(2, 154)	η^2_p	<i>F</i> _(2, 154)	η^2_p
SSART-IES	233.3 (61.3)	216.8 (61.6)	226.6 (75.8)	233.0 (52.1)	197.1 (53.2)	203.9 (55.3)	251.9 (91.2)	203.3 (61.6)	230.1 (80.5)	10.4***	.12	1.1	.03
SSSRT ^c	269.5 (79.6)	242.7 (66.9)	254.2 (48.5)	240.1 (50.9)	249.0 (74.9)	252.9 (64.5)	239.9 (63.8)	248.5 (81.1)	248.2 (78.6)	0.2	.00	1.0	.03
		<i>n</i> = 30			<i>n</i> = 25			<i>n</i> = 31		<i>F</i> _(2, 168)	η^2_p	<i>F</i> _(4, 168)	η^2_p
BRIEF-shift	18.4 (3.0)	17.3 (3.1)	16.9 (2.9)	18.3 (3.4)	17.4 (3.3)	16.5 (4.0)	17.2 (2.5)	16.5 (2.8)	16.5 (2.7)	16.5***	.17	1.2	.03
BBRIEF-WM	24.3 (5.3)	23.3 (5.0)	22.7 (5.4)	23.2 (4.7)	23.1 (4.4)	22.4 (5.4)	23.0 (4.1)	22.3 (4.7)	22.4 (4.3)	5.1**	.06	0.7	.02
BRIEF-total	164.6 (22.2)	157.6 (22.1)	154.0 (27.3)	164.7 (17.8)	160.9 (19.5)	155.2 (27.5)	163.0 (17.3)	156.7 (18.6)	157.2 (17.9)	17.9***	.18	1.0	.02
CSBQ	47.8 (14.1)	44.2 (15.3)	42.8 (16.2)	46.0 (14.1)	42.2 (14.9)	41.9 (17.0)	45.9 (14.3)	41.6 (12.5)	42.5 (13.6)	9.1***	.10	0.2	.01
DBDRS-ADHD	28.6 (10.8)	27.0 (10.8)	24.7 (9.9)	25.6 (9.6)	25.5 (10.3)	24.6 (10.9)	27.0 (10.3)	24.2 (8.6)	25.6 (10.0)	6.4***	.07	2.5*	.06
PedsQI	34.2 (12.0)	29.8 (12.8)	29.0 (11.4)	31.5 (10.4)	28.5 (11.2)	28.0 (10.6)	30.7 (12.1)	27.4 (11.1)	28.7 (12.6)	8.6***	.09	0.5	.01

BRIEF, Behavior Rating Inventory of Executive Function; Corsi-BTT, Corsi block tapping task; CSBQ, Children Social Behavior Questionnaire; DBDRS-ADHD, inattention and hyperactivity/impulsivity subscales Disruptive Behavior Disorders Rating Scale; ER, % error; FU, 6-week-follow-up; PedsQL, Pediatric Quality of Life Inventory; RT, reaction time ms; SART-IES, sustained attention response task, inverse efficiency score; WM, working memory.

^a*df* were corrected (Greenhouse-Geisser) if assumption of sphericity was violated (Mauchly's test).

^bCorrected for 0-back.

^cAnalyses without outliers did not change the main findings.

p* < .05; *p* < .01; ****p* < .001.

slightly less flexible behavior according to parents ($T_{(1, 118)} = 2.3$, $p = .02$, $r = .21$, NS after Bonferroni correction, Table S1). The RTs on the 1-back task were slightly higher in the mock-training condition ($F_{(2, 87)} = 3.4$, $p = .04$, NS after Bonferroni correction, see Table S2). There were no further differences between the three intervention-conditions on screening or pretraining variables.

Within training

Children in the WM and flexibility intervention-conditions improved significantly in sequence lengths in the WM-training tasks, and level in the flexibility-training task (p 's $< .001$, see Figure 3, and Table S3). Hence, the adaptive function in the training was effective.

Training effects

Near-transfer. WM: In all three intervention-conditions there was a significant effect of time on the Corsi-BTT ($p < .001$, see Table 1). Post hoc analysis revealed a significant increase between pretraining and post-training scores ($p < .001$), but not between post-training and FU scores ($p > .05$). There was a time $\times$ intervention interaction trend ($p = .04$, NS after Bonferroni correction). Children in the WM-training condition tended to improve more than children in the flexibility and mock-training conditions.

In all intervention-conditions there was a significant effect of time on ER and RT on the 1-back task, and ER on the 2-back task (see Table 1). Post hoc analysis revealed significant differences between pretraining and both post-training and FU (p 's $< .01$), but not between post-training and FU. There were no

significant time $\times$ intervention interactions. Children in the adaptive-intervention-conditions did not improve more in performance than children in the mock-training condition (see Table 1, and Figure S2).

Flexibility: There was a significant effect for time on the RT switch costs on the gender-emotion switch-task ($p < .01$). Post hoc analysis revealed a significant decrease in RT switch costs between pretraining and post-training ($p < .01$), but not between post-training and FU ($p > .01$). There was no effect of time on ER switch costs on both switch-tasks. Children in the adaptive intervention-conditions did not increase more in performance than children in the mock-training-condition (see Table 1).

Far-transfer. Sustained attention and inhibition: There was a significant effect for time on the SART-IES. Post hoc analysis revealed only a significant decrease in IES between pretraining and post-training ($p < .001$). There was no significant time $\times$ intervention interaction. Hence, sustained attention improved in all intervention-conditions, but did not improve more in the adaptive intervention-conditions than in the mock-training condition.

There was no significant time effect, nor time $\times$ intervention interaction in SSRT on the stop task (see Table 1). Inhibition did not improve.

Far-transfer to daily life. There was a significant effect for time on all questionnaires (see Table 1).¹ Post hoc analysis revealed a significant difference between pretraining and post-training (p 's $< .01$, except for BRIEF-WM: $p = .03$), but not between post-training and FU (p 's $> .01$). There was a time $\times$ intervention interaction trend on DBDRS-ADHD ($p = .05$, NS after Bonferroni correction).

Figure 3 Mean and standard errors of (A) the sequence length on the five training tasks in the WM-training and (B) level of cognitive flexibility-training

Hence, EFs, social behavior, and quality of life improved equally in all three intervention-conditions, but ADHD-behavior improved slightly more in the WM intervention-condition.

Rerunning the MANOVAs on training effect without children who used medication during testing did not change the main findings, except that the time $\times$ intervention interaction trend on DBDRS-ADHD was not significant anymore, probably due to a lack of power.

Intention-to-treat analysis

Intention-to-treat analyses [WM $N = 41$, flexibility $N = 40$, mock-training $N = 40$, $M(SD)$ performed sessions: WM: 21 (7), flexibility: 20 (9), mock: 22 (7)] revealed that overall children did not improve in Corsi-BTT performance, but there was a marginal time $\times$ number of sessions ($F_{(2, 228)} = 4.3$, $p = .02$, $\eta_p^2 = .04$, NS after Bonferroni correction), and a significant time $\times$ intervention interaction ($F_{(4, 228)} = 4.4$, $p = .002$, $\eta_p^2 = .07$). Corsi-BTT performance was positively related to the number of finished sessions, and improved the most in the WM intervention-condition between pre- and post-training ($F_{(2, 119)} = 5.7$, $p = .004$, $\eta_p^2 = .09$), but not between post-training and FU ($p > .01$). Children in the WM intervention-condition improved more in Corsi-BTT performance than children in the flexibility and mock-training, and remained at a higher level at FU.

There was a time $\times$ intervention interaction trend on the switch costs in ER in the gender-emotion switch-task ($F_{(4, 234)} = 2.9$, $p = .02$, $\eta_p^2 = .05$, NS after Bonferroni correction). Switch costs in ER improved more in the flexibility intervention-condition than in the WM and mock-training condition between pre- and post-training ($F_{(2, 117)} = 4.6$, $p = .01$, $\eta_p^2 = .07$), but not between post-training and FU ($p > .05$).

On the BRIEF Shift, there was a marginal time $\times$ number of sessions interaction ($F_{(2, 230)} = 3.2$, $p = .04$, $\eta_p^2 = .03$, NS after Bonferroni correction). Children who performed more sessions improved slightly more on the BRIEF-shift scale.

On the DBDRS-ADHD, the time effect was not significant ($p > .05$), but there was a time $\times$ intervention interaction trend ($F_{(3.7, 212.2)} = 2.6$, $p = .04$, $\eta_p^2 = .04$, NS after Bonferroni correction). Post hoc analyses revealed that between pretraining and post-training, there was neither time nor a time $\times$ intervention interaction ($p > .05$), and between post-training and FU there was no significant time effect ($F_{(1, 116)} = 0.1$, $p = .80$, $\eta_p^2 = .00$), but a significant time $\times$ intervention interaction ($F_{(2, 117)} = 5.3$, $p = .006$, $\eta_p^2 = .08$). Overall, children in the WM intervention-condition decreased more in their parent rated ADHD-behavior than children in the other intervention-conditions, and after the post-training this improvement continued only in the WM intervention-condition.

In line with per-protocol analysis, there were no time, nor interaction effects on the number-gnome

switch-task and the stop task. In contrast to per-protocol analyses, there were no significant time nor interaction effects on the n-back task, SART, BRIEF-WM, BRIEF-total, CSBQ, and PedsQL.

Relation between within training improvement and training effect

The improvement in the training tasks was not significantly correlated to the primary outcome measures (WM $r = -.09$ to $.16$, flexibility $r = -.33$ to $.06$). Although children did improve within the training, this improvement was not significantly related with performance on similar tasks (Corsi-BTT, and gender-emotion switch-task; see Table S4).

Discussion

Compared to a mock-training, a WM-training and flexibility-training did not induce near-transfer nor far-transfer in children with ASD in this randomized controlled trial. All children improved in WM, flexibility, attention, and parent rated EF, social behavior, ADHD-behavior, and quality of life, but the adaptive-intervention-conditions did not result in a larger improvement than the mock-training. Overall, differently measured flexibility and inhibition did not improve. The WM-training did tend to induce more improvement in WM (near-transfer) and ADHD-behavior (far-transfer) than the other intervention-conditions. The medium effect sizes and the significant findings in intention-to-treat analyses suggest a trend, whereas the non-significant treatment $\times$ time interaction in the per-protocol analyses might have resulted from a lack of power. Moreover, intention-to-treat analyses suggested a trend toward near-transfer of the flexibility-training, and a trend that children who performed more sessions, improved more in daily life flexibility.

The trend of near-transfer of the WM-training is consistent with studies in children with ADHD (e.g., Chacko et al., 2013). However, the lack of generalization to the broader WM construct and other EFs makes the trend in parent reported improvement in ADHD-behavior difficult to interpret. Although WM is related to ADHD symptoms (Oord et al., 2012; Rapport et al., 2009), training induced improvements in WM have not unequivocally shown improvement in ADHD symptoms (Melby-Lervåg & Hulme, 2013; Rapport, Orban, Kofler, & Friedman, 2013). Moreover, WM improvement was found in a recent study on Braingame Brian in children with ADHD (S. DAVIS, S. Oord, R.W. Wiers & P.J.M. Prins, unpublished data), but far-transfer to ADHD-behavior was not found. Hence, the currently found trend of improved WM and ADHD-behavior in children with ASD should be further examined in future research.

The trend of near-transfer of the flexibility-training in the intention-to-treat analysis is consistent with previous studies reporting positive effects of this type

of training (Karchach & Kray, 2009; Kray et al., 2011). However, this small effect was not revealed in the main analyses. As children with poor flexibility seem to benefit more from flexibility-training (Karchach & Kray, 2009), and as not all children with ASD show flexibility deficits (Pellicano, 2010b; de Vries & Geurts, 2012), there might have been too little room for improvement. Moreover, improvement may relapse when discontinuing a task-switching training (Karchach & Kray, 2009). Hence, including FU might reveal a less strong effect as compared to a pretraining/post-training design. In sum, although the current findings suggest that our computerized flexibility-training is ineffective for ASD, there is limited evidence of a near-transfer effect.

In line with previous research (Thorell et al., 2009), we found no improvement in inhibition. Recent research suggests that a combined EF intervention might be more successful to improve inhibition (S. Dövis, S. Oord, R.W. Wiers & P.J.M. Prins, unpublished data). The equal improvement in all intervention-conditions indicated that both the WM-training and the flexibility-training did not induce far-transfer to attention, daily EFs, social behavior, and quality of life. The overall improvement might be due to test-retest-, practice-, and expectancy effects, or regression to the mean (Melby-Lervåg & Hulme, 2013). However, nonspecific training aspects, such as the needed sustained attention to perform low level EF tasks over 25 sessions (Rueda, Checa, & Cómbita, 2012); and interacting with characters in the virtual world (Best, 2014) might have induced improvement in all intervention-conditions. Moreover, parents likely give their child extra attention during the training by jointly visiting the test-sessions, and motivating and supporting their child to complete the training. This treatment related parental involvement may have ameliorated behavioral problems (Forehand, Jones, & Parent, 2013).

The finding that the WM-training showed more effect than the flexibility and mock-training may be due to the fact that this training included five different WM-training tasks, while the other conditions only included one WM-training task, and the flexibility-training only included one adaptive flexibility-training task. Performing various tasks may be more demanding, but also more motivating, and importantly, more effective (Karchach & Kray, 2009). The WM-training includes remembering, manipulating, and updating of information. Targeting a broader range of functions, even though all part of WM, may contribute to generalization of the training effect (Buitenvweg, Murre, & Ridderinkhof, 2012; Karchach & Kray, 2009). In contrast, the solely increasing speed in the flexibility-training may be more difficult to generalize to daily life. Moreover, the gap between cognitive flexibility in the research setting and in daily life may be too large. A cognitive

behavioral approach may be more appropriate to improve cognitive flexibility (Kenworthy et al., 2014), while a computerized WM-training may be promising to improve WM. Moreover, it may be fruitful to train more EFs simultaneously, while training a single EF is perhaps too limited to result in far-transfer.

Attrition was relatively high (26%), compared to previous studies (5%–10%; Kenworthy et al., 2014; Melby-Lervåg & Hulme, 2013; Stichter et al., 2010). Children who dropped out used medication more often, indicating that these children possibly experienced more severe problems. There were no differences on other measures, but uncontrolled factors might be relevant to attrition. We unfortunately did not quantify parental support, or children's attitude toward the training-game, such as motivation to succeed on the training tasks, or to complete the training. These factors may have contributed to success or failure to complete the training, or benefit from the training. The weekly phone calls did indicate more enthusiasm about the game world of the training than about the training tasks, and integrating the training tasks more thoroughly in the game world might make the training more appealing and rewarding to the children.

One might argue that the currently used mock-training also trained EFs to a certain extent. However, the EF demand in this mock-training was very low, and undoubtedly much smaller than in the other training conditions. Completely ruling out EFs may be difficult, if not impossible, in virtually any task, and including a completely different task is hard to compare to a computer training. Thus, we consider our mock-training a proper control condition, but a challenge for future research may be to design an active, though non-EF applying control condition.

The current study has some limitations. Firstly, some children used psychotropic medication. However, medication-use was similar in all conditions, and analyses without these children gave similar results. Secondly, we did not use teacher ratings. However, teacher ratings would probably not have shown larger effects (e.g., Klingberg et al., 2005). Thirdly, although on a group level EF deficits are present in children with ASD, not every individual within this group experiences EF deficits (de Vries & Geurts, 2014). The training may be effective only for children with ASD who show EF deficits. It will be an important next step to identify specific subgroups that may benefit from this training.

In sum, the current study indicates that WM-training shows only marginal effects on WM and ADHD-behavior, and effects of the flexibility-training are even less clear, though we found limited evidence for near-transfer. The training in its present form is not a feasible treatment for all children with ASD. However, given the major individual differences in ASD (Pellicano, 2010b; de Vries & Geurts, 2012,

2014), pursuing whether the training is useful for specific subgroups is a worthwhile enterprise.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Description of Braingame Brian.

Table S1. Mean (*M*) and SD of screening and pretraining measures for children who did, and did not finish the training.

Table S2. Mean (*M*) and SD of screening and pretraining measures for children in the three training groups.

Table S3. Within training: Mean (*M*), and SD of the number of sequences at the end of each training task in the working memory training, and reached level in the cognitive flexibility-training.

Table S4. Correlation between within training improvement, and training effect.

Appendix S2. Expected and actual pattern of findings and visual representation of WM tasks.

Figure S1. Expected pattern of findings.

Figure S2. Pattern of findings.

Figure S3. Visual representation of the WM-training and cognitive flexibility tasks.

Acknowledgements

The study was approved by the ethics committee of the Academic Medical Center Amsterdam (METC 2010-185), and registered in the Dutch trial register (NL32255.018.10). Study funding was provided by the University of Amsterdam. P.J.M.P. is member of the foundation Gaming and Training, a nonprofit organization that develops and evaluates online interventions, such as EF training, for youth mental health care. The other authors have declared that they have no competing or potential conflicts of interest. The authors thank all the children and their parents for participating in the study; the students for testing the participants; and the mental health care institutions for helping with patient recruitment.

Correspondence

Marieke de Vries, Brain & Cognition, Psychology Department, University of Amsterdam, Weesperplein 4, 1018 XA Amsterdam, the Netherlands; Email: m.devries@uva.nl

Key points

- Children with autism spectrum disorders (ASD) show executive functioning (EF) deficits.
- EF training seems promising in other populations.
- A working memory (WM) and cognitive flexibility-training may be promising for children with ASD.
- WM-training shows marginal effects on WM and ADHD-behavior in children with ASD, and marginal effects of flexibility-training on flexibility are indicated.
- The training in its current form is no feasible treatment for children with ASD.

Note

1. RCI's (Jacobson & Truax, 1991) based on test-retest reliability (BRIEF: Smidts & Huizinga, 2009; CSBQ: Hartman et al., 2007; DBD-ADHD: Oosterlaan et al., 2008), and Crohnbachs α (PedsQL: Varni, Burwinkle, Seid, & Skarr, 2003) indicated that between pretraining and post-training relatively few children improved significantly (one sided 95% confidence interval; BRIEF 12.4%, CSBQ 14.6%, DBD-ADHD 7.9%, PedsQL 28.1%).

References

- Alloway, T.P., Bibile, V., & Lau, G. (2013). Computerized working memory training: Can it lead to gains in cognitive skills in students? *Computers in Human Behavior*, 29, 632–638.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders*. Washington, DC: American Psychiatric Association.
- Baltruschat, L., Hasselhorn, M., Tarbox, J., Dixon, D.R., Najdowski, A.C., Mullins, R.D., & Gould, E.R. (2011). Addressing working memory in children with autism through behavioral intervention. *Research in Autism Spectrum Disorders*, 5, 267–276.
- Bastiaansen, D., Koot, H.M., Bongers, I.L., Varni, J.W., & Verhulst, F.C. (2004). Measuring quality of life in children referred for psychiatric problems: Psychometric properties of the PedsQLTM 4.0 generic core scales. *Quality of Life Research*, 13, 489–495.
- Best, J.R. (2014). Relations between video gaming and Children's executive functions. In F.C. Blumberg (Ed.), *Learning by playing video gaming in education* (p. 42). Oxford: Oxford University Press.
- Bolte, S., Poustka, F., & Constantino, J.N. (2008). Assessing autistic traits: Cross-cultural validation of the social responsiveness scale (SRS). *Autism Research*, 1, 354–363.
- Bruyer, R., & Brysbaert, M. (2011). Combining speed and accuracy in cognitive psychology: Is the inverse efficiency score (IES) a better dependent variable than the mean reaction time (RT) and the percentage of errors (PE)? *Psychologica Belgica*, 51, 5–13.
- Buitenweg, J.I., Murre, J.M., & Ridderinkhof, K.R. (2012). Brain training in progress: A review of trainability in healthy seniors. *Frontiers in Human Neuroscience*, 6, 183.
- Casey, B.J., Cohen, J.D., Jezzard, P., Turner, R., Noll, D.C., Trainor, R.J., ... & Rapoport, J.L. (1995). Activation of prefrontal cortex in children during a nonspatial working memory task with functional MRI. *Neuroimage*, 2, 221–229.

- Cepeda, N.J., Cepeda, M.L., & Kramer, A.F. (2000). Task switching and attention deficit hyperactivity disorder. *Journal of Abnormal Child Psychology*, 28, 213–226.
- Chacko, A., Feisen, N., Bedard, A., Bedard, A.C., Uderman, J.Z., & Chimiklis, A. (2013). Cogmed working memory training for youth with ADHD: A closer examination of efficacy utilizing evidence-based criteria. *Journal of Clinical Child and Adolescent Psychology*, 42, 769–783.
- Chiang, H.M., & Wineman, I. (2014). Factors associated with quality of life in individuals with autism spectrum disorders: A review of literature. *Research in Autism Spectrum Disorders*, 8, 974–986.
- Constantino, J.N., Davis, S.A., Todd, R.D., Schindler, M.K., Gross, M.M., Brophy, S.L., ... & Reich, W. (2003). Validation of a brief quantitative measure of autistic traits: Comparison of the social responsiveness scale with the autism diagnostic interview-revised. *Journal of Autism and Developmental Disorders*, 33, 427–433.
- Corsi, P.M. (1972). Human memory and the medial temporal region of the brain. *Dissertation Abstracts International*, 34, 819B.
- Damasio, A.R., & Maurer, R.G. (1978). A neurological model for childhood autism. *Archives of Neurology*, 35, 777–786.
- De Jonge, M., & de Bildt, A. (2007). *Nederlandse bewerking van de ADI-R*. Amsterdam: Hogrefe Uitgevers B.V.
- Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64, 135–168.
- Fisher, N., & Happé, F. (2005). A training study of theory of mind and executive function in children with autistic spectrum disorders. *Journal of Autism and Developmental Disorders*, 35, 757–771.
- Forehand, R., Jones, D.J., & Parent, J. (2013). Behavioral parenting interventions for child disruptive behaviors and anxiety: What's different and what's the same. *Clinical Psychology Review*, 33, 133–145.
- Geirnaert, J. (2010). *Het grote kabouter wesley boek*. Amsterdam: Uitgeverij de Harmonie.
- Geurts, H.M., Corbett, B., & Solomon, M. (2009). The paradox of cognitive flexibility in autism. *Trends in Cognitive Sciences*, 13, 74–82.
- Geurts, H.M., Luman, M., & van Meel, C.S. (2008). What's in a game: The effect of social motivation on interference control in boys with ADHD and autism spectrum disorders. *Journal of Child Psychology and Psychiatry*, 49, 848–857.
- Geurts, H.M., deVries, M., & van den Bergh, S.F.W.M. (2014). Executive functioning theory and autism. *Handbook of executive functioning* (pp. 121–141) New York, NY: Springer.
- Gilotty, L., Kenworthy, L., Sirian, L., Black, D.O., & Wagner, A.E. (2002). Adaptive skills and executive function in autism spectrum disorders. *Child Neuropsychology*, 8, 241–248.
- Gioia, G.A., Isquith, P.K., Guy, S.C., & Kenworthy, L. (2000). Test review behavior rating inventory of executive function. *Child Neuropsychology*, 6, 235–238.
- Gray, K.M., Tonge, B.J., & Sweeney, D.J. (2008). Using the autism diagnostic interview-revised and the autism diagnostic observation schedule with young children with developmental delay: Evaluating diagnostic validity. *Journal of Autism and Developmental Disorders*, 38, 657–667.
- Hartman, C.A., Luteijn, E., Moorlag, A., De Bildt, A., & Minderaa, R.B. (Eds.). (2007). *Manual for the CSBQ [handleiding voor de VISK]*. Amsterdam: Harcourt.
- Hartman, C.A., Luteijn, E., Serra, M., & Minderaa, R. (2006). Refinement of the children's social behavior questionnaire (CSBQ): An instrument that describes the diverse problems seen in milder forms of PDD. *Journal of Autism and Developmental Disorders*, 36, 325–342.
- Holmes, J., Gathercole, S.E., & Dunning, D.L. (2009). Adaptive training leads to sustained enhancement of poor working memory in children. *Developmental Science*, 12, F9–F15.
- Huizinga, M., & Smidts, D.P. (2010). Age-related changes in executive function: A normative study with the dutch version of the behavior rating inventory of executive function (BRIEF). *Child Neuropsychology*, 17, 51–66.
- Jacobson, N.S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12.
- Karbach, J., & Kray, J. (2009). How useful is executive control training? age differences in near and far transfer of task-switching training. *Developmental Science*, 12, 978–990.
- Kenworthy, L., Anthony, L.G., Naiman, D.Q., Cannon, L., Wills, M.C., Luong-Tran, C., ... & Bal, E. (2014). Randomized controlled effectiveness trial of executive function intervention for children on the autism spectrum. *Journal of Child Psychology and Psychiatry*, 55, 274–383.
- Kenworthy, L., Yerys, B.E., Anthony, L.G., & Wallace, G.L. (2008). Understanding executive control in autism spectrum disorders in the lab and in the real world. *Neuropsychology Review*, 18, 320–338.
- Klingberg, T., Fernell, E., Olesen, P.J., Johnson, M., Gustafsson, P., Dahlstrom, K., ... & Westerberg, H. (2005). Computerized training of working memory in children with ADHD – A randomized, controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44, 177–186.
- Kort, W., Schittekatte, M., Compaa, E.L., Bosmans, M., Bleichrodt, N., & Vermeir, G., ... & Verhaeghe, P. (2002). *WISC-III NL. handleiding. nederlandse bewerking*. London: The Psychological Corporation.
- Kray, J., Karbach, J., Haenig, S., & Freitag, C. (2011). Can task-switching training enhance executive control functioning in children with attention deficit/-hyperactivity disorder? *Frontiers in Human Neuroscience*, 5, 1–9.
- Legerstee, J.S., van der Reijden-Lakeman, I.A., Lechner-van der Noort, M.G., & Ferdinand, R.F. (2004). Bruikbaarheid verkorte versie WISC-RN in de kinderpsychiatrie. *Kind En Adolescent*, 25, 178–182.
- Leung, R.C., & Zakzanis, K.K. (2014). Brief report: Cognitive flexibility in autism spectrum disorders: A quantitative review. *Journal of Autism and Developmental Disorders*, 1–18.
- Logan, G.D. (1994). On the ability to inhibit thought and action: A users' guide to the stop signal paradigm. In D. Dagenbach & T.H. Carr (Eds.), *Inhibitory processes in attention, memory, and language*. (pp. 189–239). San Diego, CA: Academic Press.
- Logan, G.D., & Cowan, W.B. (1984). On the ability to inhibit thought and action: A theory of an act of control. *Psychological Review*, 91, 295–327.
- Lord, C., Rutter, M., & Couteur, A. (1994). Autism diagnostic interview-revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 24, 659–685.
- Lundqvist, D., Flykt, A., & Öhman, A. (1998). *The karolinska directed emotional faces (KDEF)*. (91–630 ed.). Stockholm: Department of Clinical Neuroscience, Psychology Section, Karolinska Institutet.
- Melby-Lervåg, M., & Hulme, C. (2013). Is working memory training effective? A meta-analytic review *Developmental Psychology*, 49, 270.
- Oord, S.V., Ponsioen, A.J., Geurts, H.M., Brink, E.L., & Prins, P.J.M. (2014). A pilot study of the efficacy of a computerized executive functioning remediation training with game elements for children with ADHD in an outpatient setting: Outcome on parent- and teacher-rated executive functioning and ADHD behavior. *Journal of Attention Disorders*, 18, 699–712.
- Oosterlaan, J., Bayens, D., Scheres, A., Antrop, J., Roeyers, H., & Sergeant, J.A. (2008). *Handleiding van de VvGK6-16 vragenlijst voor gedragsproblemen bij kinderen van 6 tot en met 16 jaar [manual for the DBDRS disruptive behavior*

- disorder rating scale for children from 6 to 16 years old]. Amsterdam: Harcourt Assessment BV.
- Oosterlaan, J., Scheres, A., Antrop, I., Roeyers, H., & Sergeant, J.A. (2000). *Handleiding bij de vragenlijst voor gedragsproblemen bij kinderen [manual of the rating scale for disruptive behavior disorders in Children—DBDRS]*. Lisse: Swets & Zeitlinger.
- Pelham, W.E., Gnagy, E.M., Greenslade, K.E., & Milich, R. (1992). Teacher ratings of dsm-iii-R symptoms for the disruptive behavior disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31, 210–218.
- Pellicano, E. (2010a). The development of core cognitive skills in autism: A 3-Year prospective study. *Child Development*, 81, 1400–1416.
- Pellicano, E. (2010b). Individual differences in executive function and central coherence predict developmental changes in theory of mind in autism. *Developmental Psychology*, 46, 530–544.
- Prins, P.J.M., Brink, E.T., DAVIS, S., Ponsioen, A., Geurts, H.M., de Vries, M., & van der Oord, S. (2013). “Braingame brian”: Toward an executive function training program with game elements for children with ADHD and cognitive control problems. *Games for Health Journal*, 2, 44–49.
- Prins, P.J.M., DAVIS, S., Ponsioen, A., Ten Brink, E., & Van der Oord, S. (2011). Does computerized working memory training with game elements enhance motivation and training efficacy in children with ADHD? *Cyberpsychology, Behavior, and Social Networking*, 14, 115–122.
- Rapport, M.D., Bolden, J., Kofler, M.J., Sarver, D.E., Raiker, J.S., & Alderson, R.M. (2009). Hyperactivity in boys with attention-deficit/hyperactivity disorder (ADHD): A ubiquitous core symptom or manifestation of working memory deficits? *Journal of Abnormal Child Psychology*, 37, 521–534.
- Rapport, M.D., Orban, S.A., Kofler, M.J., & Friedman, L.M. (2013). Do programs designed to train working memory, other executive functions, and attention benefit children with ADHD? A meta-analytic review of cognitive, academic, and behavioral outcomes. *Clinical Psychology Review*, 33, 1237–1252.
- Robertson, I.H., Manly, T., Andrade, J., Baddeley, B.T., & Yiend, J. (1997). Oops!': Performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*, 35 (6), 747–758.
- Roeyers, H., Thys, M., Druart, C., De Schryver, M., & Schittekatte, M. (2011). *Screeninglijst voor autismespectrumstoornissen (SRS) handleiding*. Amsterdam: Hogrefe uitgevers bv.
- Rogers, R.D., & Monsell, S. (1995). Costs of a predictable switch between simple cognitive tasks. *Journal of Experimental Psychology: General*, 124, 207.
- Roughan, L., & Hadwin, J.A. (2011). The impact of working memory training in young people with social, emotional and behavioural difficulties. *Learning and Individual Differences*, 21, 759–764.
- Rueda, M.R., Checa, P., & Cómbita, L.M. (2012). Enhanced efficiency of the executive attention network after training in preschool children: Immediate changes and effects after two months. *Developmental Cognitive Neuroscience*, 2, S192–S204.
- Russell, J. (1997). *Autism as an executive disorder*. New York: Oxford University Press.
- Rutter, M., Lord, C., & LeCouteur, A. (2003). *ADI-R autism diagnostic interview—revised. manual*. Los Angeles: Western Psychological Services.
- Sattler, J. (2001). *Assessment of children: Cognitive applications*. San Diego, CA: Jerome Sattler Publisher Inc.
- Shipstead, Z., Hicks, K.L., & Engle, R.W. (2012). Cogmed working memory training: Does the evidence support the claims? *Journal of Applied Research in Memory and Cognition*, 1, 185–193.
- Shipstead, Z., Redick, T.S., & Engle, R.W. (2012). Is working memory training effective? *Psychological Bulletin*, 138, 628.
- Smids, D., & Huizinga, M. (Eds.) (2009). *BRIEF executieve functies gedragsvragenlijst: Handleiding*. Amsterdam: Hogrefe Uitgevers.
- Smith, E.E., & Jonides, J. (1999). Storage and executive processes in the frontal lobes. *Science*, 283, 1657–1661.
- Stichter, J.P., Herzog, M.J., Visovsky, K., Schmidt, C., Randolph, J., Schultz, T., & Gage, N. (2010). Social competence intervention for youth with asperger syndrome and high-functioning autism: An initial investigation. *Journal of Autism and Developmental Disorders*, 40, 1067–1079.
- Thorell, L.B., Lindqvist, S., Bergman Nutley, S., Bohlin, G., & Klingberg, T. (2009). Training and transfer effects of executive functions in preschool children. *Developmental Science*, 12, 106–113.
- Varni, J.W., Burwinkle, T.M., Seid, M., & Skarr, D. (2003). The PedsQL™ 4.0 as a pediatric population health measure: Feasibility, reliability, and validity. *Ambulatory Pediatrics*, 3, 329–341.
- Varni, J.W., Seid, M., & Kurtin, P.S. (2001). PedsQL (TM) 4.0: Reliability and validity of the pediatric quality of life inventory (TM) version 4.0 generic core scales in healthy and patient populations. *Medical Care*, 39, 800–812.
- de Vries, M., & Geurts, H.M. (2012). Cognitive flexibility in ASD; task switching with emotional faces. *Journal of Autism and Developmental Disorders*, 42, 2558–2568.
- de Vries, M., & Geurts, H.M. (2014). Beyond individual differences: Are working memory and inhibition informative specifiers within ASD?. *Journal of Neural Transmission*, 121, 1183–1198.
- World Health Organisation. (1992). *The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and diagnostic guidelines*. Geneva: World Health Organization.
- Zinke, K., Einert, M., Pfennig, L., & Kliegel, M. (2012). Plasticity of executive control through task switching training in adolescents. *Frontiers in Human Neuroscience*, 6, 41.

Accepted for publication: 11 August 2014

Published online: 26 September 2014