



UvA-DARE (Digital Academic Repository)

Sulfate-dependent acetate oxidation under extremely natron-alkaline conditions by syntrophic associations from hypersaline soda lakes

Sorokin, D.Y.; Abbas, B.; Tourova, T.P.; Bumazhkin, B.K.; Kolganova, T.V.; Muyzer, G.

DOI

[10.1099/mic.0.075093-0](https://doi.org/10.1099/mic.0.075093-0)

Publication date

2014

Document Version

Final published version

Published in

Microbiology (United Kingdom)

[Link to publication](#)

Citation for published version (APA):

Sorokin, D. Y., Abbas, B., Tourova, T. P., Bumazhkin, B. K., Kolganova, T. V., & Muyzer, G. (2014). Sulfate-dependent acetate oxidation under extremely natron-alkaline conditions by syntrophic associations from hypersaline soda lakes. *Microbiology (United Kingdom)*, *160*(4), 723-732. <https://doi.org/10.1099/mic.0.075093-0>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, P.O. Box 19185, 1000 GD Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Sulfate-dependent acetate oxidation under extremely natron-alkaline conditions by syntrophic associations from hypersaline soda lakes

Dimitry Y. Sorokin,^{1,2} Ben Abbas,² Tatjana P. Tourova,¹
Boris K. Bumazhkin,³ Tatjana V. Kolganova³ and Gerard Muyzer⁴

Correspondence

Dimitry Y. Sorokin
soroc@inmi.ru or
d.sorokin@tudelft.nl

¹Winogradsky Institute of Microbiology, Russian Academy of Sciences,
Prospect 60-let Otyabrya 7/2, 117811 Moscow, Russia

²Department of Biotechnology, Delft University of Technology, Delft, The Netherlands

³Bioengineering Centre, Russian Academy of Sciences, Moscow, Russia

⁴Department of Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics,
University of Amsterdam, Amsterdam, The Netherlands

So far, anaerobic sulfate-dependent acetate oxidation at high pH has only been demonstrated for a low-salt-tolerant syntrophic association of a clostridium '*Candidatus Contubernalis alkalaceticum*' and its hydrogenotrophic sulfate-reducing partner *Desulfonatronum cooperativum*. Anaerobic enrichments at pH 10 inoculated with sediments from hypersaline soda lakes of the Kulunda Steppe (Altai, Russia) demonstrated the possibility of sulfate-dependent acetate oxidation at much higher salt concentrations (up to 3.5 M total Na⁺). The most salt-tolerant purified cultures contained two major components apparently working in syntrophy. The primary acetate-fermenting component was identified as a member of the order *Clostridiales* forming, together with '*Ca. Contubernalis alkalaceticum*', an independent branch within the family *Syntrophomonadaceae*. A provisional name, '*Ca. Syntrophonatronum acetioxidans*', is suggested for the novel haloalkaliphilic clostridium. Two phylotypes of extremely haloalkaliphilic sulfate-reducing bacteria of the genus *Desulfonatronospira* were identified as sulfate-reducing partners in the acetate-oxidizing cultures under extreme salinity. The dominant phylotype differed from the two species of *Desulfonatronospira* described so far, whilst a minor component belonged to *Desulfonatronum thiodismutans*. The results proved that, contrary to previous beliefs, sulfate-dependent acetate oxidation is possible, albeit very slowly, in nearly saturated soda brines.

Received 14 November 2013

Accepted 24 January 2014

INTRODUCTION

Soda lakes are naturally occurring highly alkaline and saline habitats containing sodium carbonate at high concentrations, maintaining a stable high pH of the brines. Despite these extreme conditions, soda lakes usually have high primary production. This, together with high sulfate concentrations, is the reason why the microbiological sulfur cycle is highly active in soda lakes. Soda lake sediments usually contain millimolar concentrations of free sulfide and FeS in the top 20 cm (Sorokin *et al.*, 2011a). Detailed studies of the reductive sulfur cycle in soda lakes, including

Abbreviations: DGGE, denaturing gradient gel electrophoresis; SRB, sulfate-reducing bacteria.

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA and *dsrAB* gene sequences determined in this study are KF588515–KF588528 and KF835251–KF835261, respectively.

One supplementary figure is available with the online version of this paper.

measurements of sulfate reduction rates, revealed active sulfate, thiosulfate and sulfur/polysulfide reduction mostly stimulated by formate/H₂ (Gorlenko *et al.*, 1999; Kulp *et al.*, 2006, 2007; Sorokin *et al.*, 2004, 2010a). However, only sulfur reduction was apparently active with acetate as electron donor in a relatively short-term incubation experiment. This was corroborated by the results of a cultivation approach, which demonstrated a domination of lithotrophic sulfate-reducing bacteria (SRB) using H₂/formate, lactate and ethanol as electron donors and disproportionating thiosulfate/sulfite, such as in the genera *Desulfonatronum*, *Desulfonatronovibrio* and *Desulfonatronospira* (Pikuta *et al.*, 1998, 2003; Zhilina *et al.*, 1997, 2005a; Sorokin *et al.*, 2008, 2011b). Heterotrophic SRB isolated from soda lakes are either 'incomplete oxidizers', producing acetate as a final product during oxidation of simple organic compounds (*Desulfobotulus alkaliphilus* and *Desulfobulbus alkaliphilus*; Sorokin *et al.*, 2010b, 2012) or 'complete oxidizers', which cannot use external acetate (*Desulfonatronobacter*

acidivorans; Sorokin *et al.*, 2012). However, a specific group of acetate- and propionate-oxidizing sulfur/polysulfide respiring natronophiles able to grow in concentrated soda brines can be isolated easily from hypersaline soda lakes (*Desulfurispira natronophila*; Sorokin & Muyzer, 2010).

So far, there has been only a single report on the possibility of indirect sulfate-dependent acetate oxidation at high pH by a syntrophic association of a novel obligate syntrophic endospore-forming clostridium '*Candidatus Contubernalis alkalaceticum*' and a hydrogenotrophic SRB, *Desulfonatronum cooperativum* (Zhilina *et al.*, 2005b). The association was enriched from the low-salt alkaline lake Hadyn in Tuva Republic (Russia) as a part of a low-salt anaerobic cellulose-degrading community (total Na⁺ ~0.25 M, pH 10) selected by using acetate as electron donor and sulfate as electron acceptor (Kevbrin *et al.*, 1999). The association oxidized ~25 mM acetate in 30 days, producing up to 20 mM sulfide. However, neither the pH nor salt ranges were reported for the acetate-oxidizing association. Meanwhile, the question of the salt limit for the low-energy-yielding anaerobic acetate oxidation with sulfate as final electron acceptor is fundamental for understanding thermodynamic limits of microbial life. Growth at high salt demands high extra energy input for the synthesis of osmolytes and, therefore, low-energy-generating metabolism is compromised under these conditions (Oren, 2011). So far, the salt limit for direct oxidation of acetate by halophilic SRB is 2 M NaCl, which is not surprising taking into account the very low energy yield of the conversion: CH₃COO⁻ + SO₄²⁻ → HS⁻ + 2 HCO₃⁻ [$\Delta G^{\circ} = -56$ kJ (pH 7; Schink & Stams, 2006); $\Delta G^{\circ} = -47$ kJ (pH 10; Oren, 2011)].

The results described in this paper present, to the best of our knowledge, the first proof of anaerobic acetate oxidation with sulfate as electron acceptor by natronophilic syntrophic associations at much higher salt concentrations than believed previously. The acetate-oxidizing association consisted of a member of a novel branch within the class *Clostridia* and its extremely natronophilic lithoautotrophic SRB partner from the genus *Desulfonatronospira*, which, working together, oxidized acetate in nearly saturated soda brines.

METHODS

Samples. Anaerobic sediment cores (5–15 cm depth) were obtained from the following soda lakes of the Kulunda Steppe (south-eastern Siberia, Altai, Russia): (i) the moderately saline soda lake Cock Soda Lake (salinity 70 g l⁻¹, pH 10.1, carbonate alkalinity 0.7 M), and (ii) the hypersaline soda lakes Tanatar-5 (salinity 170 g l⁻¹, pH 9.9, carbonate alkalinity 1.9 M) and Bitter-1 (salinity 400 g l⁻¹, pH 10.65, carbonate alkalinity 4.4 M) in July 2010, and Bitter-1 in July 2011 (salinity 400 g l⁻¹, pH 10.1, carbonate alkalinity 4.0 M).

Enrichment and cultivation conditions. Anaerobic acetate-dependent sulfate-reducing cultures were enriched from soda lake sediments at 30 °C in a mineral medium based on sodium carbonate/bicarbonate buffer with stable pH 10 containing in total 0.6–3 M Na⁺, 0.1–0.3 M NaCl and 1 g K₂HPO₄ l⁻¹. After sterilization,

the medium was supplemented with 4 mM NH₄Cl, 1 mM MgSO₄, 20 mg yeast extract l⁻¹ and 1 ml l⁻¹ of each solution of acidic trace metals and vitamins (Pfennig & Lippert, 1966), and 1 ml basic filter-sterilized Se/W solution (Plugge, 2005). Acetate and sulfate at 20 mM were used as electron donor and electron acceptor, respectively; 1 mM HS⁻ was added as a reductant. The SRB partners were sub-cultured from the acetate-utilizing association in a medium specific for *Desulfonatronospira*, containing 3 M total Na⁺ with formate and sulfite (Sorokin *et al.*, 2008). Routine cultivation was performed in 18 ml Hungate tubes with 10 ml medium made anoxic by several cycles of flushing with argon and evacuation. Growth was monitored by sulfide production and measurements of OD₆₀₀. When the sulfide concentration in the enrichments exceeded 5 mM, the cultures were transferred into new medium at 1:100 dilution. After two or three successful transfers, the enrichments were serially diluted up to 10⁻¹⁰. Growth on solid medium was not observed. One of the probable reasons was a problem with clarity and solidification of agar at high sodium carbonate concentrations.

The pH dependence was examined at 2 M Na⁺ using the following filter-sterilized buffers: for pH 7–8, 0.1 M HEPES and NaCl; for pH 8.5–11, a mixture of sodium bicarbonate/sodium carbonate. Final pH values were taken to indicate a suitable range for growth, because of the pH shift at pH extremes. To study the influence of salt concentration on growth, mineral sodium carbonate bases at pH 10 containing 0.6 and 4.0 M total Na⁺ were mixed in different proportions.

Analyses. Sulfide was precipitated in 10% (w/v) zinc acetate and analysed by the methylene blue method after separation from the supernatant as ZnS (Trüper & Schlegel, 1964). The amount of cell protein was measured by the Lowry method (Lowry *et al.*, 1951) after removal of interfering FeS from the cell pellet by a double wash with 1 M NaCl, pH 4. Acetate in the supernatant was analysed by GC after removal of sulfide and acidification to pH 5 (Chromoteq-Crystall 5000.2; column Sovpol-5, 1 m, 180 °C; detector PID, 30 °C). Phase-contrast photomicrographs were obtained with an Axioplan Imaging 2 microscope (Zeiss).

Genetic and phylogenetic analysis. DNA was extracted from the cells using the UltraClean Microbial DNA Isolation kit (MoBio Laboratories) following the manufacturer's instructions. The nearly complete 16S rRNA gene was obtained from finally diluted associations by standard molecular cloning procedures using general bacterial primers 11f/1492r (Lane, 1991). The PCR products were purified using the Wizard SV-gel and PCR Clean-Up System (Promega). The purified fragments were ligated into plasmids using the pGEM-T Easy Vector System (Promega) and the plasmids were then electroporated into the competent cells of *Escherichia coli* strain DH10B. DNA from positive clones ($n=25$) was extracted with Wizard MiniPreps (Promega). Community analysis of syntrophic associations was performed by using 16S-rRNA-gene-based and *dsrB*-based denaturing gradient gel electrophoresis (DGGE) according to Schäfer & Muyzer (2001). For the 16S rRNA analysis, the primer pair was bacterial 341f + GC clamp/907r and the gel gradient was from 20 to 70%. For the *dsrB* analysis, the primer pair was DSRp2060f + GC clamp/DSR4r (Geets *et al.*, 2006; Wagner *et al.*, 1998) with the gel gradient from 30 to 65%.

For the phylogenetic analysis of the 16S rRNA gene, the obtained sequences were first compared to all sequences stored in GenBank using the BLAST algorithm and were consequently aligned using CLUSTALW. The phylogenetic trees were reconstructed using TREECON W and the neighbour-joining algorithm. Phylogeny of the *dsrB* fragments was reconstructed using ARB software. The sequences were aligned using Codoncode Aligner. Sequences were aligned with complete-length sequences of closest relatives from the order *Desulfovibrionales* and the family *Desulfobacteraceae* obtained from the updated *dsrAB* database (Loy *et al.*, 2009) using the ARB 'fast

aligner' utility. The maximum likelihood method, RAxML (implemented in ARB), was used to calculate the resulting phylogenetic tree.

RESULTS AND DISCUSSION

Enrichment and isolation of syntrophic acetate-oxidizing sulfidogenic associations from soda lake sediments

In contrast to our previous negative short-term experiments (Sorokin *et al.*, 2010a), prolonged incubations of sediment slurries amended with acetate and sulfate showed sulfide formation after 3–6 months in samples taken in 2010 at 0.6–2.0 M total Na^+ , pH 10. After several subcultures, a maximum of 10 mM sulfide was produced in 3 months. In serial dilutions, cultures with reduced morphological diversity were obtained from three lakes. The final diluted culture at the lowest salinity was dominated by two clearly different phenotypes (Fig. S1a, available in the online Supplementary Material), whilst the other two cultures only included short rods (Fig. S1b, c). In 2011, the experiment with the most saline soda lake, Bitter-1 (sample 3KL-011), was repeated to determine the salt limit of the process by slurry incubations at 2–4 M Na^+ , pH 10. Sulfide formation was observed at sodium carbonate concentrations up to 3 M total Na^+ after 6 months of incubation, with the purified culture dominated by short rods (Fig. S1d). None of the acetate-oxidizing cultures obtained contained endospore-forming clostridia observed previously in the low-salt alkaliphilic acetate-oxidizing association (Zhilina *et al.*, 2005a, b). These results demonstrated the extraordinary possibility of a very slow, but reproducible microbial sulfate-dependent oxidation of acetate in soda lake sediments at moderate to extremely high salt concentrations.

Growth dynamics and influence of sodium on acetate-dependent sulfidogenesis in cultures from the hypersaline soda lake Bitter-1

In maximally purified culture from 3KL-011 grown at 2 M total Na^+ , pH 10, full oxidation of 18 mM acetate was

observed in 3 months (Fig. 1a). The oxidation was accompanied by biomass growth and formation of 16 mM sulfide, which corresponded well to the eight-electron stoichiometry 1:1 of the donor (acetate) and acceptor (sulfide) assuming that part of the acetate was assimilated. The experimental maximum specific growth yield was 1.2 mg cell protein (mmol acetate)⁻¹. Cultures from both 2010 and 2011 were obligately alkaliphilic, growing within a range from pH 8.7 to 10.5 with an optimum around pH 10 (Fig. 1b). Both cultures grew in sodium carbonate brines containing at least 1 M total Na^+ with an optimum at 1.5–2 M. There was a slight difference between the upper salt limit, which was higher for the culture from 2011 which grew in nearly saturated sodium carbonate brines containing 3.5 M Na^+ (Fig. 1c). However, full oxidation of 18 mM acetate under such extreme conditions was at least three times slower than under the optimal salt concentration.

Analysis of the composition of acetate-oxidizing syntrophic cultures

DGGE analysis of the 16S rRNA genes of three acetate-oxidizing cultures from the 2010 samples showed a domination of two closely related clostridial phylotypes (Fig. 2a); one presented at a low-to-moderate salinity (3KL-010 and 7KL-010) and another presented at high salinity (5KL-010). Together with the two clones detected directly in anaerobic sediments of one of the Bitter lake systems (T. P. Tourova, unpublished), they formed a novel lineage within the family *Syntrophomonadaceae* (class *Clostridia*) distantly related to '*Ca. Contubernalis alkalaceticum*' (Fig. 3). The SRB partner at the lowest salinity was identified as a member of the genus *Desulfonatronum* and the SRB partner at the highest salinity was identified as a novel representative of the extremely natronophilic genus *Desulfonatronospira* (Sorokin *et al.*, 2008). A more specific analysis of the functional gene *dsrB* showed the presence of a novel *Desulfonatronospira* in moderate- and high-salt cultures, and confirmed the presence of *Desulfonatronum* in the low-salt culture (Fig. 2b, Table 1). Several *Bacteroidetes* phylotypes present in cultures




Fig. 1. (a) Growth dynamics at 2 M Na^+ and pH 10, (b) pH profile at 2 M Na^+ , and (c) salt profile at pH 10 in natronophilic syntrophic acetate-oxidizing culture 3KL-011. (a) ●, Biomass; ▲, sulfide; ◆, acetate; (b, c) ▲, culture AAS1; ●, culture 3KL-011. Means from two independent experiments.




Fig. 2. (a) 16S rRNA gene and (b) *dsrB* DGGE analysis of syntrophic acetate-oxidizing sulfate-reducing enrichment cultures from Kulunda soda lakes, 2010, grown at pH 10 and variable Na^+ concentrations. 3KL-010: Cock Soda Lake, dilution -4; 7KL-010: Tanatar-5, dilutions -4 and -5; 5KL-010: Bitter-1, dilution -4; AAS1: Bitter-1, final dilution culture -8. Band identification in (a): 1, 2, 7-9, 13-15, 20, uncultured bacteroidetes from soda lakes; 3, 4, 11, 16-19, 21-25, *Clostridiales*, novel branch; 5, 6, *Desulfonatronum thioautotrophicum* (99%); 12, *Desulfonatronospira* spp. (97-98%). Band identification in (b): 1, 18, *Desulfobacterium anilini* (93%); 2-4, 35, *Desulfonatronum lacustre* (96%); 5, 6, 10, 13, 20, 22, 23, 24, 26, *Desulfonatronospira thiodismutans* (89-92%). Bands 11 and 12 gave bad quality sequences and are not included.

at low dilutions (Fig. 2a) disappeared at high dilutions, indicating their irrelevance for the main process.

The comparison of the enrichment culture from Bitter-1 lake (2011) at 3 M Na^+ with the 'AAS1' culture from the same lake (2010) showed the presence of the same key players, i.e. a novel member of the class *Clostridia* and a novel *Desulfonatronospira* (Fig. 4). Subculturing on a medium specific for *Desulfonatronospira* with formate and sulfite resulted in the domination of a rod-shaped organism and, in addition, a spiral-shaped *Desulfonatronospira* proliferated in the offshoots of the 3KL-011 culture. Both types were highly enriched and identified by cloning. The rod-shaped phylotype was distant from the described *Desulfonatronospira* by its full-length 16S rRNA gene and probably represents a novel species within this genus, whilst

the spiral morphotype belonged to the type species *Desulfonatronospira thiodismutans* (Fig. 5a). It seems likely that the novel rod-shaped *Desulfonatronospira* may have been selected in the high-salt acetate-oxidizing syntrophic cultures on the basis of its better adaptation to grow with sulfate as electron acceptor than the species described previously (Sorokin *et al.*, 2008).

Interestingly, in the 3KL-011 culture, another SRB was detected that belonged to the order *Desulfobacterales* with *Desulfonatronobacter acidivorans* as the closest relative (Figs 4 and 5b, Table 1). The latter is the only known 'complete oxidizing' natronophilic SRB found so far in soda lakes, but it cannot utilize external acetate and cannot grow lithotrophically with formate. As this organism was absent in the 2010 culture, we can only conclude that it was not




Fig. 3. Phylogenetic position based on 16S rRNA gene sequence analysis of the acetate-oxidizing clostridial members in syntrophic associations from soda lakes. The clones directly detected in lake Bitter-3 are in bold type. Bar, 5 nt changes per 100 nt. The percentage of bootstraps was derived from 1000 resampling iterations using the neighbour-joining algorithm; only values >70% are indicated. GenBank accession numbers are given in parentheses.

essential for the syntrophic acetate oxidation. The same applies to the presence of '*Halanaerobium hydrogeniformans*' in the formate + sulfite offshoots from the 3KL-011 culture. This haloalkaliphilic clostridium is a fermentative saccharolytic (Brown *et al.*, 2011). However, it must be pointed out that some members of the order *Halanaerobiales* have the potential to use sulfur-dependent respiration as an additional metabolism (Sorokin *et al.*, 2011a).

Attempts to find other electron donors for the highly purified syntrophic cultures from Bitter-1 lake were complicated by the fact that *Desulfonatrosipira* is more diverse physiologically than the SRB partner (genus *Desulfonatrosipira*) in the low-salt association described previously. The latter was reported to be able to oxidize 1- and 2-propanol, *iso*-butyrate, serine and fructose, apart from acetate (Zhilina *et al.*, 2005a). The high salt cultures actively grew with ethanol, propanol and butanol, the substrates also utilized by *Desulfonatrosipira* alone. Other donors, positive for the '*Ca. Contubernalis*'/*Desulfonatrosipira* association, did not support growth of the high-salt cultures, as well as the range of substrates tested (negative for *Desulfonatrosipira* alone), including propionate, malate, fumarate, succinate, valerate, caproate and peptone. Also, the acetate-oxidizing clostridium alone

did not grow on substrates employed commonly for the cultivation of syntrophs (McInerney *et al.*, 2008; Stams & Plugge, 2009), such as fumarate, pyruvate, crotonate and ethylene glycol.

In summary, the results presented here demonstrate anaerobic acetate oxidation with sulfate as electron acceptor at extremely natron-alkaline conditions of hypersaline soda lakes. This process is driven by syntrophic interaction between a novel and apparently obligately syntrophic clostridial lineage in the family *Syntrophomonadaceae* and a novel member of the extremely natronophilic lithotrophic SRB of the genus *Desulfonatrosipira*. The clostridial member is suggested to be accommodated into a novel candidate taxon '*Candidatus Syntrophonatronum acetioxidans*'. Despite a very slow growth, the mere fact that such a low-energy-yielding catabolic reaction can support growth at extreme salt and pH is an important finding for understanding the limits of microbial life and its biogeochemical role. That such a process has been detected so far only in hypersaline soda lakes, and not in hypersaline habitats with neutral pH (Oren, 2011), indicates that sodium carbonates may create an environment fundamentally different from the conditions present in a high-NaCl world.

Table 1. Dsr(AB) sequences identified in syntrophic acetate-oxidizing cultures from hypersaline soda lakes (DGGE band sequences)

The reference *dsrAB* gene sequences of *Desulfonatrobacter acidivorans* APT2^T have been deposited in GenBank under accession numbers KF835252 and KF835254.

Culture				Sequence source*	Closest culturable relative in GenBank	Domination	Similarity of translated amino acids (%)	GenBank accession no. of the representative sequence
Lake	Salinity (M Na ⁺)	Dilution	Substrate					
3KL-010	0.6	-4	Acetate/SO ₄ ²⁻	DGGE1_bands 2-4, 35 DGGE1_band 1	<i>Desulfonatronum lacustre</i> <i>Desulfobacterium anilini</i>	Dominant Minor	96 92	Band (1)-4: KF835261 No
7KL-010	1.0	-4/-5	Acetate/SO ₄ ²⁻	DGGE1_bands 20, 22-24, 26	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	89-92	Band (1)-20: KF835263
5KL-010 (AAS1)	2.0	-4	Acetate/SO ₄ ²⁻	DGGE1_bands 5, 6, 10-13	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	89-92	Band (1)-13: KF835262
	2.0	-8	Acetate/SO ₄ ²⁻	DGGE2_band 3	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	91	Band (2)-3: KF835255
	3.0	-8	Formate/SO ₃ ²⁻	DGGE2_bands 14, 15	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	91-92	Band (2)-15: KF835257
	3.0	-8	Formate/SO ₃ ²⁻	Direct sequencing	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	90	Full sequence: KF835251; KF835253
3KL-011	3.0-3.5	-8	Acetate/SO ₄ ²⁻	DGGE2_band 8	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	91-92	Band (2)-8: KF835256
	3.0	-8	Formate/SO ₃ ²⁻	DGGE2_bands 22, 23	<i>Desulfonatronospira thiodismutans</i>	Dominant (phylotype 1)	92	Band (2)-22: KF835259
				DGGE2_bands 26, 27	<i>Desulfonatronospira thiodismutans</i>	Minor (phylotype 2)	100	Band (2)-26: KF835260
				DGGE2_bands 20, 21, 24, 25	<i>Desulfonatrobacter acidivorans</i>	Minor	94	Band (2)-20: KF835258

*DGGE1 corresponds to Fig. 2(b); DGGE2 corresponds to Fig. 4(b).




Fig. 4. Comparative (a) 16S RNA gene and (b) *dsrB* DGGE analysis of two extremely natronophilic anaerobic syntrophic cultures oxidizing acetate with sulfate as *electron acceptor* enriched from hypersaline soda lake Bitter-1. AAS1, final dilution culture from 2010 sample; 3KL-011, final dilution culture from 2011 sample. Left part of the gels shows cultures grown with acetate + sulfate; right part shows the transfer to formate + sulfite. Band identification in (a): 0, *Bacteroidetes* (87%); 1, *Clostridiales*, novel branch; 3, 5, *Halanaerobium hydrogenotiformans* (99%); 2, 4, *Desulfonatronospira* spp. (98%); 6, *Desulfonatronospira thiodismutans* (99%); 7, *Desulfonatronobacter acidivorans* (95%). Band identification in (b): 3–6, 8, 11, 14, 15, 17–19, 22, 23, 32, *Desulfonatronospira thiodismutans* (89–92%); 26, 27, *Desulfonatronospira thiodismutans* (100%); 20, 21, 24, 25, 31, *Desulfonatronobacter acidivorans* (94%). Band 9 gave a bad quality sequence and is not included.

Description of ‘*Candidatus Syntrophonatronum*’

‘*Candidatus Syntrophonatronum*’ (Syn.tro.pho.natro’n.um. Gr. prep. *syn* in company with, together with; Gr. n. *trophos* one who feeds; L. n. *natron* soda; M.L. neutr. n. *Syntrophonatronum* syntrophic natronophile).

Gram-positive rods. Obligately anaerobic, oxidizing fatty acids in syntrophy with the H₂-consuming partner. Obligate natronophiles. Habitat: anaerobic sediments of hypersaline soda lakes. Members of the family *Syntrophomonadaceae* (order *Clostridiales*, class *Clostridia*). Type species: ‘*Candidatus Syntrophonatronum acetioxidans*’.

Description of ‘*Candidatus Syntrophonatronum acetioxidans*’

‘*Candidatus Syntrophonatronum acetioxidans*’ [acet.i.’oxi.dan.s. L. n. *acetum* vinegar; N.L. n. *acidum aceticum* acetic acid; N.L. v. *oxido* (from Gr. adj. *oxus* acid or sour

and in combined words indicating oxygen) to make acid, oxidize; N.L. part. adj. *acetioxidans* oxidizing acetate].

Cells are non-spore-forming and non-motile Gram-positive rods, 0.5 × 2–3 µm. Obligately anaerobic, oxidizing acetate in the presence of a lithoautotrophic sulfate-reducing partner. Other possible substrates are not known. Does not grow alone on pyruvate, crotonate, fumarate or ethylene glycol. Obligately natronophilic with the ability to grow in syntrophic culture within the range pH 8.5–10.5 (optimum at pH 10) and sodium carbonate concentrations from 1.0 to 3.5 M total Na⁺ (optimum 1.5–2.0 M). Growth is optimal at 35 °C and possible up to 42 °C.

The type strain ASS1^T exists in a syntrophic co-culture with the extremely natronophilic SRB partner from the genus *Desulfonatronospira* (the GenBank accession number of the 16S rRNA gene sequence is KF588524). It is deposited in the UNIQEM culture collection (Institute of Microbiology, Russian Academy of Sciences) under the



Fig. 5. Phylogenetic position based on 16S rRNA gene sequence analysis of the SRB present in acetate-oxidizing syntrophic associations from soda lakes. (a) Sequences in the order *Desulfovibrionales*. (b) Sequences in the family *Desulfobacteraceae*, order *Desulfobacterales*. Bars, 5 nt changes per 100 nt. The percentage of bootstraps was derived from 1000 resampling iterations using the neighbour-joining algorithm; only values >70% are indicated. GenBank accession numbers are given in parentheses.

number U977. The culture was obtained from anaerobic sediments of the hypersaline soda lake Bitter-1 in Kulunda Steppe (Altai, Russia). The GenBank accession number of the 16S rRNA gene sequence of the type strain AAS1^T is KF588515.

ACKNOWLEDGEMENTS

This work was supported by the Russian Foundation for Basic Research (grants 13-04-00049 and 13-04-40405 comfi) to D. S. and an ERC Advanced Grant to G. M.

REFERENCES

- Brown, S. D., Begemann, M. B., Mormile, M. R., Wall, J. D., Han, C. S., Goodwin, L. A., Pitluck, S., Land, M. L., Hauser, L. J. & Elias, D. A. (2011). Complete genome sequence of the haloalkaliphilic, hydrogen-producing bacterium *Halanaerobium hydrogenoformans*. *J Bacteriol* **193**, 3682–3683.
- Geets, J., Borremans, B., Diels, L., Springael, D., Vangronsveld, J., van der Lelie, D. & Vanbroekhoven, K. (2006). *DsrB* gene-based DGGE for community and diversity surveys of sulfate-reducing bacteria. *J Microbiol Methods* **66**, 194–205.
- Gorlenko, V. M., Namsaraev, B. B., Kulyrova, A. V., Zavarzina, D. G. & Zhilina, T. N. (1999). Activity of sulfate-reducing bacteria in the sediments of the soda lakes in south-east Transbaikal area. *Microbiology (English translation of Mikrobiologiya)* **68**, 580–586.
- Kezbrin, V. V., Zhilina, T. N. & Zavarzin, G. A. (1999). Decomposition of cellulose by the anaerobic alkaliphilic microbial community. *Microbiology (English translation of Mikrobiologiya)* **68**, 601–609.
- Kulp, T. R., Hoefft, S. E., Miller, L. G., Saltikov, C., Murphy, J. N., Han, S., Lanoil, B. & Oremland, R. S. (2006). Dissimilatory arsenate and sulfate reduction in sediments of two hypersaline, arsenic-rich soda lakes: Mono and Searles Lakes, California. *Appl Environ Microbiol* **72**, 6514–6526.
- Kulp, T. R., Han, S., Saltikov, C. W., Lanoil, B. D., Zargar, K. & Oremland, R. S. (2007). Effects of imposed salinity gradients on dissimilatory arsenate reduction, sulfate reduction, and other microbial processes in sediments from two California soda lakes. *Appl Environ Microbiol* **73**, 5130–5137.
- Lane, D. J. (1991). 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial Systematics*, pp. 115–177. Edited by E. Stackebrandt & M. Goodfellow. Chichester: Wiley.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *J Biol Chem* **193**, 265–275.
- Loy, A., Duller, S., Baranyi, C., Musmann, M., Ott, J., Sharon, I., Béjà, O., Le Paslier, D., Dahl, C. & Wagner, M. (2009). Reverse dissimilatory sulfite reductase as phylogenetic marker for a subgroup of sulfur-oxidizing prokaryotes. *Environ Microbiol* **11**, 289–299.
- McInerney, M. J., Struchtemeyer, C. G., Sieber, J., Mouttaki, H., Stams, A. J. M., Schink, B., Rohlin, L. & Gunsalus, R. P. (2008). Physiology, ecology, phylogeny, and genomics of microorganisms capable of syntrophic metabolism. *Ann N Y Acad Sci* **1125**, 58–72.
- Oren, A. (2011). Thermodynamic limits to microbial life at high salt concentrations. *Environ Microbiol* **13**, 1908–1923.
- Pfennig, N. & Lippert, K. D. (1966). Über das Vitamin B₁₂ – bedürfnis phototropher Schwefel bacterien. *Arch Mikrobiol* **55**, 245–256.
- Pikuta, E. V., Zhilina, T. N., Zavarzin, G. A., Kostrikina, N. A., Osipov, G. A. & Rainey, F. A. (1998). *Desulfonatronum lacustre* gen. nov., sp. nov.: a new alkaliphilic sulfate-reducing bacterium utilizing ethanol. *Microbiology (English translation of Mikrobiologiya)* **67**, 105–113.
- Pikuta, E. V., Hoover, R. B., Bej, A. K., Marsic, D., Whitman, W. B., Cleland, D. & Krader, P. (2003). *Desulfonatronum thiodis-mutans* sp. nov., a novel alkaliphilic, sulfate-reducing bacterium capable of lithoautotrophic growth. *Int J Syst Evol Microbiol* **53**, 1327–1332.
- Plugge, C. M. (2005). Anoxic media design, preparation, and considerations. *Methods Enzymol* **397**, 3–16.
- Schäfer, H. & Muyzer, G. (2001). Denaturing gradient gel electrophoresis in marine microbial ecology. *Methods Microbiol* **30**, 425–468.
- Schink, B. & Stams, A. J. M. (2006). Syntrophism among Prokaryotes. In *The Prokaryotes*, 3rd edn, vol. 2, pp. 309–335. Edited by M. Dworkin, S. Falkow, E. Rosenberg, K. H. Schleifer & E. Stackebrandt. New York: Springer.
- Sorokin, D. Y. & Muyzer, G. (2010). *Desulfurispira natronophila* gen. nov. sp. nov.: an obligately anaerobic dissimilatory sulfur-reducing bacterium from soda lakes. *Extremophiles* **14**, 349–355.
- Sorokin, D. Y., Gorlenko, V. M., Namsaraev, B. B., Namsaraev, Z. B., Lysenko, A. M., Eshinimaev, B. T., Khmelenina, V. N., Trotsenko, Y. A. & Kuenen, J. G. (2004). Prokaryotic communities of the north-eastern Mongolian soda lakes. *Hydrobiologia* **522**, 235–248.
- Sorokin, D. Y., Tourova, T. P., Henstra, A. M., Stams, A. J. M., Galinski, E. A. & Muyzer, G. (2008). Sulfidogenesis under extremely haloalkaline conditions by *Desulfonatronospira thiodis-mutans* gen. nov., sp. nov., and *Desulfonatronospira delicata* sp. nov. – a novel lineage of *Deltaproteobacteria* from hypersaline soda lakes. *Microbiology* **154**, 1444–1453.
- Sorokin, D. Y., Rusanov, I. I., Pimenov, N. V., Tourova, T. P., Abbas, B. & Muyzer, G. (2010a). Sulfidogenesis under extremely haloalkaline conditions in soda lakes of Kulunda Steppe (Altai, Russia). *FEMS Microbiol Ecol* **73**, 278–290.
- Sorokin, D. Y., Detkova, E. N. & Muyzer, G. (2010b). Propionate and butyrate dependent bacterial sulfate reduction at extremely haloalkaline conditions and description of *Desulfobotulus alkaliphilus* sp. nov. *Extremophiles* **14**, 71–77.
- Sorokin, D. Y., Kuenen, J. G. & Muyzer, G. (2011a). The microbial sulfur cycle at extremely haloalkaline conditions of soda lakes. *Front Microbiol* **2**, 44.
- Sorokin, D. Y., Tourova, T. P., Kolganova, T. V., Detkova, E. N., Galinski, E. A. & Muyzer, G. (2011b). Culturable diversity of lithotrophic haloalkaliphilic sulfate-reducing bacteria in soda lakes and the description of *Desulfonatronum thioautotrophicum* sp. nov., *Desulfonatronum thiosulfatophilum* sp. nov., *Desulfonatronovibrio thiodis-mutans* sp. nov., and *Desulfonatronovibrio magnus* sp. nov. *Extremophiles* **15**, 391–401.
- Sorokin, D. Y., Panteleeva, A. N., Tourova, T. P. & Muyzer, G. (2012). *Desulfonatronobacter acidivorans* gen. nov., sp. nov. and *Desulfobulbus alkaliphilus* sp. nov., haloalkaliphilic heterotrophic sulfate-reducing bacteria from soda lake. *Int J Syst Evol Microbiol* **62**, 2107–2113.
- Stams, A. J. M. & Plugge, C. M. (2009). Electron transfer in syntrophic communities of anaerobic bacteria and archaea. *Nat Rev Microbiol* **7**, 568–577.
- Trüper, H. G. & Schlegel, H. G. (1964). Sulfur metabolism in *Thiorhodaceae*. 1. Quantitative measurements on growing cells of *Chromatium okenii*. *Antonie van Leeuwenhoek* **30**, 225–238.
- Wagner, M., Roger, A. J., Flax, J. L., Brusseau, G. A. & Stahl, D. A. (1998). Phylogeny of dissimilatory sulfite reductases supports an early origin of sulfate respiration. *J Bacteriol* **180**, 2975–2982.
- Zhilina, T. N., Zavarzin, G. A., Rainey, F. A., Pikuta, E. N., Osipov, G. A. & Kostrikina, N. A. (1997). *Desulfonatronovibrio hydrogenovorans* gen. nov., sp. nov., an alkaliphilic, sulfate-reducing bacterium. *Int J Syst Bacteriol* **47**, 144–149.
- Zhilina, T. N., Zavarzina, D. G., Kuever, J., Lysenko, A. M. & Zavarzin, G. A. (2005a). *Desulfonatronum cooperativum* sp. nov., a novel

hydrogenotrophic, alkaliphilic, sulfate-reducing bacterium, from a syntrophic culture growing on acetate. *Int J Syst Evol Microbiol* **55**, 1001–1006.

Zhilina, T. N., Zavarzina, D. G., Kolganova, T. V., Turova, T. P. & Zavarzin, G. A. (2005b). [*Candidatus* Contubernalis alkalaceticum],

an obligately syntrophic alkaliphilic bacterium capable of anaerobic acetate oxidation in a coculture with *Desulfonatronum cooperativum*.] *Mikrobiologiya* **74**, 800–809 (in Russian).

Edited by: C. Dahl