UvA-DARE (Digital Academic Repository)

Treatment effect on recidivism for juveniles who have sexually offended

A multilevel meta-analysis

ter Beek, E.; Spruit, A.; Kuiper, C.H.Z.; van der Rijken, R.E.A.; Hendriks, J.; Stams, G.J.J.M.

DOI

[10.1007/s10802-017-0308-3](https://doi.org/10.1007/s10802-017-0308-3)

Publication date

2018

Document Version

Final published version

Published in

Journal of Abnormal Child Psychology

License

Article 25fa Dutch Copyright Act

[Link to publication](#)

Citation for published version (APA):

ter Beek, E., Spruit, A., Kuiper, C. H. Z., van der Rijken, R. E. A., Hendriks, J., & Stams, G. J. J. M. (2018). Treatment effect on recidivism for juveniles who have sexually offended: A multilevel meta-analysis. *Journal of Abnormal Child Psychology*, 46(3), 543-556. <https://doi.org/10.1007/s10802-017-0308-3>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Treatment Effect on Recidivism for Juveniles Who Have Sexually Offended: a Multilevel Meta-Analysis

Ellis ter Beek¹ · Anouk Spruit² · Chris H. Z. Kuiper¹ · Rachel E. A. van der Rijken³ · Jan Hendriks^{2,4} · Geert Jan J. M. Stams²

Published online: 25 May 2017
© Springer Science+Business Media New York 2017

Abstract The current study investigated the effect on recidivism of treatment aimed at juveniles who have sexually offended. It also assessed the potential moderating effect of type of recidivism, and several treatment, participant and study characteristics. In total, 14 published and unpublished primary studies, making use of a comparison group and reporting on official recidivism rates, were included in a multilevel meta-analysis. This resulted in the use of 77 effect sizes, and 1726 participants. A three-level meta-analytic model was used to calculate the combined effect sizes (Cohens *d*) and to perform moderator analyses. Study quality was assessed with the EPHPP Quality Assessment Tool for

Quantitative Studies. A moderate effect size was found ($d = 0.37$), indicating that the treatment groups achieved an estimated relative reduction in recidivism of 20.5% as compared to comparison groups. However, after controlling for publication bias, a significant treatment effect was no longer found. Type of recidivism did not moderate the effect of treatment, indicating that treatment groups were equally effective for all types of recidivism. Also, no moderating effects of participant or treatment characteristics were found. Regarding study characteristics, a shorter follow up time showed a trend for larger effect sizes, and the effect size calculation based on proportions yielded larger effect sizes than calculation via mean frequency of offending. Implications for future research and clinical practice are discussed.

✉ Ellis ter Beek
Ellis.terBeek@horizon.eu

Anouk Spruit
A.Spruit@uva.nl

Chris H. Z. Kuiper
Chris.Kuiper@horizon.eu

Rachel E. A. van der Rijken
Rachel.van.der.Rijken@deviersprong.nl

Jan Hendriks
JHendriks@dewaagnederland.nl

Geert Jan J. M. Stams
G.J.J.M.Stams@uva.nl

Keywords Juveniles · Sex offending · Treatment effect · Recidivism · Meta-analysis

An interest in treatment for juveniles who have sexually offended was first sparked in the late 1970's. Research on adults who had sexually offended showed that they had often started their abusing careers in adolescence (Groth et al. 1982). In accordance with findings from research on adult sex offenders, it was assumed that dynamic risk factors explaining juvenile sex offending would differ from dynamic risk factors explaining non-sexual juvenile delinquency, and that specific treatment for juveniles who had sexually offended would be needed to target these dynamic risk factors (Barbaree and Marshall 2006). Recent research, however, shows that juveniles who have sexually offended differ psychologically from adult sex offenders (Adjorlolo and Egbenya 2016). And that notably, in general, dynamic risk factors for juvenile sex-offense recidivism do not always differ from risk factors

¹ Horizon Youth Care and Education, Mozartlaan 150, 3055 KM Rotterdam, The Netherlands

² University of Amsterdam, Amsterdam, The Netherlands

³ Viersprong Institute for Studies on Personality Disorders, Halsteren, The Netherlands

⁴ Forensic Psychotherapeutic Centre de Waag, Amsterdam, The Netherlands

for juvenile general offense recidivism (Carpentier and Proulx 2011; Christiansen and Vincent 2013; Worling and Långström 2003).

Also, sexual (re)offending by juveniles remains mainly adolescence limited (Beaudry-Cyr et al. 2015; Loeber et al. 2012; Lussier et al. 2012). In 2016, Caldwell conducted a meta-analysis, and found a weighted mean of 5% official recidivism by juveniles. This probably is an underestimation of the true recidivism rate due to the rarity of documented sexual offending (Wittebrood 2006). Interestingly, however, the reoffending base rate has significantly declined to a weighted mean of 3% in the last 15 years. Improved quality of treatment was hypothesized to be the most likely explanation (Caldwell 2016). Research on recidivism reduction through treatment of juveniles who have sexually offended, however, is highly diverse, differing in study design, outcome, treatment and participant characteristics, which is thought to affect study findings (Dopp et al. 2015; Hanson et al. 2009). The current study aims to synthesize the results of earlier studies by testing the hypothesis that specialized treatment for juveniles is effective.

Treatment Components

The Risk-Need-Responsivity (RNR) model (Bonta and Andrews 2007), prominently states that treatment allocation and delivery should be guided by specific criminogenic treatment needs and recidivism risk levels, and should be delivered responsively (i.e., attuned to personal life goals, learning and motivational style) in order to be effective. As a consequence, aiming treatment at established risk factors for (sexual) reoffending is considered to be imperative (Ter Beek et al. 2016).

Juveniles who generally display antisocial tendencies, who have also displayed other forms of offending behavior, seem most at risk for recidivism, including sexual reoffending (Drew 2013; Hendriks and Bijleveld 2005; Lawing et al. 2010; Van Wijk et al. 2007). Overall, juveniles who have sexually offended display a fair amount of non-sexual recidivism (Caldwell 2016; Lobanov-Rostovsky 2015). Specialized juvenile treatment may therefore need to address risk factors for general delinquency, such as immaturity, impulsiveness, and antisocial beliefs, to effectively reduce sexual as well as general recidivism risk. Seto and Lalumière (2010) and Fanniff and Kimonis (2014), comparing juveniles who have sexually offended with juveniles who have committed non-sexual offenses only, found both similarities (mainly their level of antisocial thoughts and behavior) and differences. Juveniles who had sexually offended showed more atypical sexual interests, more social and emotional problems, and more experiences of sexual victimization, abuse and neglect than did non-sex offenders. These findings suggest that general etiological theories of

antisocial behavior might only partially explain juvenile sexual misconduct, and that specific or additional explanations remain relevant.

Christiansen and Vincent (2013), in their study of 39,249 juveniles, of whom 695 juveniles had offended sexually, showed prior non-sexual offending, non-contact sexual offenses, offending against a child, early age of onset, poor school performance, and non-school attendance to be predictive of sexual recidivism in juveniles. In 2011, Carpentier and Proulx examined 351 male adolescents who had sexually offended, and found paternal abandonment, childhood sexual victimization, association with younger children, and having victimized a stranger to be associated with a higher risk for sexual recidivism. Prior to these studies, Worling and Långström (2003) conducted a literature review of juvenile sex offending, which provided empirical evidence for the following risk factors: sexual deviation (e.g., interest in prepubescent children or sexual violence), prior criminal sanctions for sexual assault(s), having two or more victims, victimization of strangers, lack of intimate peer relationships/social isolation, and incomplete offense-specific treatment. Notably, the dynamic (changeable) risk factors, such as sexual deviation, are of most interest for recidivism risk reduction through treatment (Beggs and Grace 2011; Hanson and Morton-Bourgon 2005; Pedersen et al. 2010).

Although support is found for dynamic risk factors that need to be addressed during treatment, most of the empirically established risk factors for juvenile sexual recidivism (e.g., number of previous victims, prior offending behavior, having offended against a child or a stranger, early age of onset) are considered static (unchangeable). Successful treatment however, may also focus on reduction of negative consequences of static risk factors (for example, by means of trauma treatment to alleviate the consequences of earlier victimization). Also, non-criminogenic (mental health) treatment needs should be addressed (Lord 2016).

Studies into treatment needs of juveniles who have sexually offended, have substantiated the existence of several psychological typologies. Studies into specific mental illnesses, executive functioning, and personality profiles have, for example, concluded that juveniles with child victims display different treatment needs than do juvenile peer abusers (Drew 2013; Glowacz and Born 2012; Hart-Kerkhoffs et al. 2009; Hendriks 2006; Kjellgren et al. 2006). Also, Butler and Seto (2002), and Hissel et al. (2006) have found sex-only offenders to differ from mixed offenders (juveniles having also committed non-sexual offenses). Hendriks (2006) also differentiated group offenders from solo offenders. The diversity in the population of juveniles who have sexually offended has become increasingly clear, especially because to date no psychological typology has been found to represent a subgroup at increased risk for sexual recidivism (Cale et al. 2016; Lussier et al. 2012).

Effectiveness of Treatment for Juveniles Who Have Sexually Offended

Treatments for juveniles who have sexually offended make use of different treatment strategies (i.e., wilderness therapy, cognitive and/or behavioral based group therapy, cognitive and/or behavioral based individual therapy, family therapy and psycho-education) and are delivered in both residential and community settings (Ryan et al. 2010; Veneziano and Veneziano 2002). Meta-analyses, predominantly including studies that combine adult and juvenile samples, have shown cognitive behavioral based treatments and multisystemic therapy to be most effective in reducing recidivism (Dopp et al. 2015; Hanson et al. 2009; Lösel and Schmucker 2005; Schmucker and Lösel 2015; Walker et al. 2004). Treatment types that incorporate these standards are, therefore, generally considered established treatment.

To date, only two meta-analyses have been performed that focus specifically on treatment for juveniles. Walker et al. (2004) performed a meta-analysis of ten published and unpublished studies; two studies with and eight without a comparison group. Reitzel and Carbonell (2006) based their findings on nine studies, all using a comparison or control group, and including five unpublished studies. These meta-analyses demonstrated a moderate effect of treatment on sexual recidivism by juveniles. Moderator analyses showed that outcomes measured via official recidivism data (rearrests or convictions) resulted in smaller effect sizes compared to measurement via self-report or penile plethysmography (i.e., the measurement of physical arousal to deviant stimuli). Furthermore, random assignment to treatment and comparison groups yielded larger effect sizes. The hypotheses that cognitive behaviorally based treatments are more effective than other treatment types, was not substantiated in this meta-analysis (Reitzel and Carbonell 2006).

Reitzel and Carbonell (2006), as well as Walker et al. (2004), assessed the effects of treatment on sexual recidivism only, and based their findings on a restricted number of studies. In the last decade more and more robust studies (making use of a comparison or control group and examining the effectiveness of treatment for multiple types of recidivism) have been conducted. In addition, multilevel statistical techniques have become more refined, making it possible to analyze moderators with a restricted number of studies, accounting for both within and between study variability in effect sizes. A multilevel approach enables comprehensive moderator analyses to examine the influence of study, sample, and intervention characteristics (Van den Noortgate et al. 2013; Van Den Noortgate and Onghena 2003). This may be useful in the complex field of juvenile sex offending, where multiple causes may contribute to recidivism. Furthermore, multilevel meta-analytical techniques enable the use of all available effect sizes in the primary studies, thus preserving all

information and generating maximum statistical power (Assink et al. 2015). This is especially important when examining the results of studies with small samples, as is the case with many of the primary studies in this field.

Study aim

The aim of this study is to review contemporary research on the effect on officially measured recidivism (re-arrest or conviction) of treatment for juveniles who have sexually offended. In addition, the potential moderating effects of participant, treatment, and study characteristics were investigated. This provided an opportunity to detect factors that may influence the effect of treatment on the reduction of recidivism.

Method

To assess the effect of treatment on recidivism, and the factors moderating this effect, a multilevel meta-analysis was carried out. The term meta-analysis refers to a stepwise procedure and a set of statistical techniques, combining results of independent primary studies into effect sizes, so that overall conclusions can be drawn. An important requirement for traditional univariate meta-analytic approaches is that no dependency between effect sizes is allowed, so that only one effect size per primary study can be included. By stepping away from the traditional univariate approach, it becomes possible to deal with dependency of effect sizes, so all information can be preserved and a maximum of statistical power is achieved. In this study, applying a three-level structure statistically considers the three different variance components distributed over three levels: differences among all effect sizes or random sampling error (level 1), differences in effect sizes within studies (level 2), and differences in effect sizes between studies (level 3). If there is evidence for heterogeneity in effect sizes, moderator analyses can be conducted to test variables that may explain within-study or between-study heterogeneity. For these analyses, the three-level random effects model can be extended with study and effect size characteristics, making the model a three-level mixed effects model (Assink and Wibbelink 2016).

Inclusion Criteria

Multiple inclusion criteria were formulated to select the studies. First, the treatment condition had to be aimed at treating sexually offending behavior. Second, the study sample had to exclusively contain juveniles. Therefore, the mean age of the researched group had to lie between 12 and 18 years of age, and/or the study had to specifically report on juveniles or

adolescents. Third, the studies had to report treatment and comparison group recidivism rates via officially obtained data for sexual and/or other types of recidivism. And finally, the comparison group had to comprise juveniles who had sexually offended as to ensure basic comparability between groups.

Selection of Studies

All studies published before February 2016 that met the inclusion criteria were to be included in the current meta-analysis. Firstly, several electronic databases were searched, including Campbell library, PubMed, OVID (Medline, PsycINFO, ERIC), and Proquest (Sociological Abstracts, Social Services Abstracts, Proquest Dissertations). Secondly, Google Scholar was searched. The following English search string was used: (sex* offender) AND (juvenile OR adolescent) AND (treatment OR therapy) AND (effect* OR efficacy) AND (recidivism). Finally, the references of other meta-analyses and reviews were checked for eligible studies, and authors of non-published work were contacted. Regrettably the contacted authors did not respond, so a few non-published studies could not be included. A flow chart is presented as Fig. 1 in the results section.

The initial search and screening resulted in 23 studies that met the basic criterion of examining the effectiveness of an intervention for juveniles who have sexually offended with the use of a comparison group. After exclusion, mainly because of the use of other outcome variables (see also Table 2), 14 studies remained, with 77 effect sizes, 1726 participants, and 13 independent samples. Table 1 presents the study characteristics of the included studies. Table 2 specifies the excluded studies, and our reasons for excluding them in italic.

Coding the Studies

The dependent variable in this meta-analysis was recidivism. Three types of recidivism were distinguished: any recidivism, sexual recidivism, and non-sexual recidivism. In all studies, recidivism was measured by means of official data as to ensure comparability among the effect sizes of the primary studies, because the type of measurement used may influence the effect measured (Reitzel and Carbonell 2006).

The independent variable was the treatment offered. Type of recidivism, participant, treatment, and study characteristics were coded to assess whether treatment effects varied among the possible moderator variables. In order to reduce the problem of multiple testing (Tabachnik and Fidell 2013), only moderators of possible theoretical importance were used. Studies with overlapping samples (Worling and Curwen 2000; Worling et al. 2010) were coded with corresponding study identification numbers.

Firstly, **outcome characteristics** were coded as the type of recidivism reported; recidivism in general (including all types of recidivism), sexual recidivism, or non-sexual recidivism.

The following **participant characteristics** were coded. The criminal status of the clients was coded as all adjudicated or as also suspected of a sex crime, as not all studies only included formally adjudicated offenders. The cultural background of the juveniles was coded as the percentage of Caucasians in the researched group. Furthermore, the percentage of juveniles with child victims (< 12 years of age and ≥5 years younger than their assailant) was coded, as was the percentage of contact offenders (physical sexual offending, such as rape or sexual assault; not including, for example, exhibitionism). In most reports information on the estimated recidivism risk level of the respondents was available (groups were designated as relatively high at risk or low at risk), so also the mean estimated recidivism risk level of clients was coded as high or low. Finally, the living condition of participants was coded as living at home versus living in a residential facility or mixed (both types of living situation present in the researched group).

Several **treatment characteristics** were coded: firstly, whether the type of treatment was cognitive behavioral, systemic, or another form of (non -established) treatment as usual. Mean treatment duration was coded in months. Finally, the exclusion of clients with a lower IQ was coded as yes or no, for this may influence learnability of the researched group.

As for **study characteristics**, we coded whether the authors were independent researchers or whether they were involved in the development or implementation of the intervention. It was also coded whether the study was peer reviewed. Further, the design of the study (randomized controlled trial versus quasi-experimental) was coded. The type of comparison treatment used was coded as treatment as usual (TAU) or established treatment (incorporating cognitive behavioral treatment and/or systemic therapy). The comparability of the treatment and comparison group(s) was coded as high (including statistically controlling for any potential influential differences found, for example, differences in group recidivism risk levels) or low (when significant differences were reported between groups and this was not controlled for in results). The mean length of follow up was coded in months. The type of effect size was coded as proportion based (the difference in proportion of reoffenders in the treatment and comparison group) or based on mean number of re-offenses in each group. Type of recidivism measured was coded as re-arrest or reconviction. Lastly, study quality was coded by use of the EPHPP Quality Assessment Tool for Quantitative Studies (<http://www.ehphp.ca/tools.html>). This tool assesses the quality of a study as weak, moderate or strong, providing a comprehensive and structured assessment of the concept of study quality (Armijo-Olivo et al. 2012). The EPHPP was developed by the Effective Public Health Practice Project in Canada. It is a generic tool used to evaluate a variety of intervention study designs. It has been judged suitable to be used in systematic reviews of effectiveness (Deeks et al. 2003) and has been reported to have content

and construct validity (Jackson and Waters 2005; Thomas et al. 2004). The tool assesses six domains: selection bias, study design (including appropriateness of the design), confounders, blinding, data collection methods, and withdrawals and drop-outs, see Table 1 for the results of the assessment.

The first author of this article coded the included studies according to the suggestions of Lipsey and Wilson (2001). All studies were double coded by the second author. Following the guidelines by Shrout (1989), for the continuous variables single ICCs were calculated for the two-way random effects model, with absolute agreement as a criterion. On all 18 variables single measures showed an ICC of > 0.99 (i.e., substantial reliability). For 23 categorical variables kappa was calculated (Landis and Koch 1977). Two variables reached substantial interrater reliability (kappa = 0.61–0.80), all others reached almost perfect interrater reliability (kappa > 0.81).

Calculations and Analyses

Effect sizes were transformed into Cohen's d by using the calculator of Wilson (2013) and formulas of Lipsey and Wilson (2001). Most d -values were calculated based on reported means and percentages of recidivism. A positive effect size indicated that the treatment group benefited more than the comparison group, whereas a negative effect size indicated that the comparison group benefitted more than the treatment group. If a study only mentioned that an effect was not significant (as was the case in 9% of all effect sizes), the effect size was coded as zero (Lipsey and Wilson 2001). The continuous variables (percentage immigrants, percentage with child victims, percentage contact, percentage mixed offenders, and mean treatment length) were centered around their mean, and all other (categorical) variables were recoded into dummy variables. We checked for the presence of extreme outliers using box plots (Tabachnik and Fidell 2013); no extreme outliers were identified. Standard errors were estimated using formulas of Lipsey and Wilson (2001).

In all studies, we were able to calculate more than one effect size. Most studies reported on multiple outcome variables (i.e., all types of recidivism, sex only recidivism, non-sex recidivism, and sometimes violent non-sexual recidivism and non-violent non-sexual recidivism). Effect sizes from the same study may prove more alike than effect sizes from different studies. Therefore, the assumption of statistical independency, which underlies classical meta-analytic strategies, was violated (Hox 2002; Lipsey and Wilson 2001). In line with recently conducted meta-analyses, we applied a multilevel approach in order to deal with the interdependency of effect sizes (Assink et al. 2015; Houben et al. 2015; Spruit et al. 2016; Weisz et al. 2013). The multilevel approach accounts for the hierarchical structure of the data in which effect sizes are nested within the studies (Van Den Noortgate and Onghena 2003).

We used a three-level meta-analytic model to calculate the combined effect sizes and to perform moderator analyses. Three sources of variance were modelled, including the sampling variance for the observed effect sizes (level 1), the variance between effect sizes from the same study (level 2), and the variance between the studies (level 3) (Cheung 2014; Van den Noortgate et al. 2013; Wibbelink and Assink 2015). The sampling variance of observed effect sizes (level 1) was estimated by using the formula of Cheung (2014). Log-likelihood-ratio-tests were performed to compare the deviance of the full model to the deviance of the models excluding one of the variance parameters, making it possible to determine whether significant variance is present at the second and third level (Wibbelink and Assink 2015). Significant variance at level 2 or 3 indicates a heterogeneous effect size distribution, meaning that the effect sizes cannot be treated as estimates of a common effect size. In that case, we proceed to moderator analyses, because the differences between the effect sizes may be explained by outcome, study, sample, and/or intervention characteristics. Moderator analyses were only performed when each category of the potential moderator was filled with at least three studies. As a result, Study Quality was collapsed into a dichotomous variable (weak versus medium to strong) because only two studies were considered to be of strong quality. Exclusion of IQ could not be tested because only two studies reported on the exclusion of lower IQ respondents.

The multilevel meta-analysis was conducted in R (version 3.2.0) with the metafor-package, using a multilevel random effects model (Viechtbauer 2010; Wibbelink and Assink 2015). The restricted maximum likelihood estimate was used to estimate all model parameters, and the Knapp and Hartung (2003) was used for testing individual regression coefficients of the meta-analytic models and for calculating the corresponding confidence intervals (see also Assink et al. 2015; Houben et al. 2015; Spruit et al. 2016; Wibbelink and Assink 2015).

Publication Bias

In systematic reviews, the aim is to include all studies previously conducted that meet the inclusion criteria (Lipsey and Wilson 2001). However, a common problem is that studies may not have been published due to non-significant or unfavorable findings, and, therefore, are difficult to locate. Not including these studies may lead to an overestimation of the true effect size, the so called “publication bias” (Rosenthal 1979). In order to check the presence of publication bias in our meta-analysis, we performed a trim and fill procedure (Duval and Tweedie 2000) by drawing a trim-and-fill plot in R (Version 3.2.0) using the function “trimfill” of the metafor package (Viechtbauer 2010). We tested whether effect sizes were missing on the left side of the distribution, since

publication bias would only be likely to occur in case of non-significant or unfavorable (i.e., negative) results.

Results

The results of the literature search are outlined in Fig. 1. For details on the excluded studies, see Table 1. The most important characteristics of the included studies are presented in Table 2.

Overall, a significant, moderate effect, $d = 0.37$, $p < 0.001$, of treatment on recidivism was found, indicating that the treatment groups achieved an estimated reduction in recidivism of 20.5% as compared to comparison groups. The likelihood ratio test comparing models with and without between-study variance (level 3) showed that significant variance was present at the between-study level, $\sigma^2_{\text{level } 3} = 0.11$, $\chi^2(1) = 37.12$; $p < 0.000$. The variance between the effect sizes within studies (level 2) was not significant, $\sigma^2_{\text{level } 2} = 0.01$, $\chi^2(1) = 3.32$; $p = 0.068$. About 24% of the total effect size variance was accounted for by the sampling variance (level 1), 7% for the variance between effect sizes within studies (level 2), and 69% for the variance between studies (level 3). Because the variance on the third level was significant, we proceeded to moderator analyses to assess factors that could possibly explain variance in treatment effects (see Table 3).

As presented in the last column of Table 3, only the calculation type of effect size moderated the effect of treatment on recidivism in adolescents who have sexually offended. Stronger treatment effects were found for effect sizes that were proportion-based compared to effect sizes based on the mean number of re-offenses. The proportion of recidivism was significantly lower in the treatment groups compared to the comparison groups, but the mean number of offenses for every reoffender in the two groups did not significantly differ. Furthermore, a moderating trend was found for the follow up period, indicating that stronger treatment effects were

found in shorter follow up periods. None of the participant characteristics and none of the treatment characteristics moderated the effect of treatment on recidivism in adolescents who have sexually offended.

The trim-and-fill plot imputed estimations of effect sizes of missing studies, indicating the presence of publication bias (see Fig. 2). We included the imputed estimations (the open circles) and performed the meta-analysis again to compute an overall effect size that takes the influence of publication bias into account (Duval and Tweedie 2000). After controlling for publication bias, no significant treatment effect was found ($d = 0.15$, $p = 0.176$).

Discussion

A multilevel meta-analysis was performed to assess the strength of the effect on the reduction of recidivism of specific treatment for juveniles who have sexually offended, and to assess what variables have a moderating influence on the effect measured. An overall significant and moderate effect ($d = 0.37$) was found, indicating treatment to be more effective in reducing recidivism than comparison or control conditions, a finding similar to that of previous meta-analyses (Reitzel and Carbonell 2006; Walker et al. 2004). However, indications of publication bias were present, and after statistically controlling for this bias, the effect of treatment on recidivism was no longer significant. Studies with less favorable outcomes may indeed not have been published, and the moderate effect may therefore represent an overestimation of the true overall effect size. Thus, in the current meta-analysis, after correcting for publication bias, an effect of treatment on recidivism, as compared to comparison or control conditions, was not demonstrated. The treatment conditions in the primary studies, however, were mainly compared to other forms of treatment ($N = 10$), or to groups of assessment-only juveniles, who may have received treatment elsewhere ($N = 5$). Therefore,

Fig. 1 Flowchart of study selection

Table 1 Characteristics of Included Studies

Study	Study characteristics				Sample characteristics					Treatment characteristics			
	Treat. coh.	N	# r (M)	Pub	Study quality	Design	Indepauth.	Type of rec.	Mean age	% Male	% Caucasian	% Child victim	Contact
Barlow 1998	1995	44	2 (0.328)	No	Weak	QE	Yes	1/2	15.2	100	86	-	-
Borduin et al., 1990	1983–1985	16	6 (1.130)	Yes	Weak	RCT	No	1/2/3	14.0	100	63	50	94
Borduin et al., 2009	1990–2001	48	5 (0.915)	Yes	Moderate	RCT	No	1/2/3	14.0	96	71	-	-
Gillis and Gass 2010	1990–2005	190	6 (0.341)	Yes	Moderate	QE	Yes	1/2/3	-	100	65	-	-
Guarino-Ghezzi and Kimball 1998	1993–1994	75	5 (0.683)	Yes	Moderate	QE	Yes	1/2/3	-	100	53	46	83
Hendriks and Bijleveld 2005	1988–2003	325	3 (0.256)	Yes	Weak	QE	No	1/2/3	14.4	100	62	44	100
Holly 2000	1990–1998	64	14 (0.583)	No	Moderate	QE	Yes	1/2/3	14.5	98	-	-	100
Johnson Erickson 2008	2000–2007	78	4 (–0.367)	No	Strong	QE	Yes	1/2/3	15.3	99	78	87	92
Lab et al. 1993	1988–1991	155	2 (0.248)	Yes	Weak	QE	Yes	1/2	14.5	98	57	66	24
Letourneau et al. 2013	2004–2006	131	2 (0.051)	Yes	Strong	RCT	No	1	14.7	100	44	55	86
Schram et al. 1991	1984–1991	196	6 (0.055)	No	Weak	QE	Yes	1	14.5	100	89	76	97
Waite et al. 2005	1992–2001	256	4 (0.316)	Yes	Weak	QE	No	1/2/3	16.8	100	44	43	31
Worling and Curwen 2000	1987–1995	148 ^a	12 (0.575)	Yes	Moderate	QE	No	1/2/3	15.6	94	-	65	98
Worling et al. 2010	1987–1995	148 ^a	6 (0.352)	Yes	Moderate	QE	No	1/2/3	15.6	94	-	55	98

Treat. coh. = Years in which treatment was delivered, N = number of participants, # r (M) = number of effect sizes (mean d), Pub. = published in peer reviewed article yes/no, Study Quality = strong/moderate/weak according to EPHPP quality assessment tool, Design = RCT (randomized controlled trial) or QE (quasi experimental), Independent author = yes/no

Type of recidivism measured: 1 = all recidivism, 2 = sexual, 3 = non-sexual; Age = mean age of sample; % Male = percentage of males in sample; % Caucasian = percentage of Caucasian ethnicity; % Contact = percentage contact offense index crime; % Child molest = percentage child molestation index crime; Setting = outpatient, community based, residential or a mix; EXP type = Type of experimental intervention; Experimental Condition = name of experimental intervention; Comparison or Control condition = Type of control intervention(s)

*the allocation of control group status was based on available information, sometimes control group juveniles may have still have received alternate forms of treatment elsewhere

^a studies using the same participants,

^b treatment comparison group,

^c no treatment control group

Table 2 Excluded Studies and Reasons for Exclusion

Study	Authors & year of publication	Treat. cohort	N	Setting	Outcome measure	Experimental treatment	Comparison ^a or Control ^b condition
Brannon and Troyer 1991	1987–1988	110	Residential	Official Recidivism Rates	Residential State Treatment	Residential State Treatment	<i>JNSO</i> ^a
				Psychosocial Functioning & Official Recidivism Rates	Play therapy (5 to 12 year olds)*	CBT (5 to 12 year olds) ^a	
Kahn and Chambers 1991	1984	221	Mixed	Official Recidivism Rates (total group)*	8 Outpatient programs	2 Residential programs ^a	
Seabloom et al. 2003	1977–1986*	122	Outpatient	Official Recidivism Rates	Personal Social Awareness Program completers	Personal Social Awareness Program withdrawers ^b	
Letourneau and Armstrong 2008	1995–2005	222	-	Official Recidivism Rates	Registration*	Non-Registration ^b	
Letourneau et al. 2009	2004–2006	127	Community	Psychosocial Functioning*	JSO Multi Systemic Therapy	JSO Usual Community Services ^a	
Henggeler et al. 2009	2004–2006	127	Community	Psychosocial Functioning*	JSO Multi Systemic Therapy	JSO Usual Community Services ^a	
Van Outsem 2009	-	113	Outpatient	Psychological Functioning*	Treatment completers	Treatment non-completers and 'normal' controls ^b	
Weinrott et al. 1997	NR	69	Outpatient	Phallometric measurement and Selfreport*	Vicarious Sensitization	Cognitive Therapy ^a	

JSO Juvenile Sex offender, *JNSO* Juvenile Non Sex Offender, *CBT* Cognitive Behavioral Therapy

*The characteristic in italic font specifies reason(s) for exclusion. Mostly the outcome variable was other than a measure of recidivism (Letourneau et al., Henggeler et al., Van Outsem, and Weinrott et al.). In the Brannon and Troyer study a comparison group of non-sex offenders was used. In the Carpentier et al. study the respondents were too young to be included, in the Kahn and Chambers study the official recidivism rates were not available for the experimental and control group, and in the Letourneau and Armstrong study registration was researched which, to our judgement, is not a form of treatment. The Seabloom et al. study included transgressive acts (i.e., homosexuality, transvestitism) that were then considered to be a sexual transgression, but are now considered a variation in normal human sexuality, so their sample was deemed not comparable

Table 3 Overall Results and Moderator Effects of the Relation between Treatment and Recidivism

Moderator variables	# Studies	# ES	β_0 (mean d)	t_0	β_1	t_1	$F(df_1, df_2)$
Overall relation	13	77	0.367	3.713***			
After trim and fill	18	99	0.154	1.364	$(p = 0.176)$		
Outcome characteristics							
Type of Recidivism	13	77					$F(2,74) = 0.543$
Any offenses (RC)	13	29	0.401	3.764***			
Sex offenses	11	19	0.348	3.100**	−0.054	−0.735	
Non-sex offenses	9	28	0.332	3.012**	−0.069	−0.999	
Participant characteristics							
Criminal Status	13	77					$F(1,75) = 1.135$
All convicted (RC)	9	58	0.299	2.506*			
Also suspects	4	19	0.533	2.880**	0.235	1.065	
Percentage Caucasian	11	46	0.333	2.694*	0.406	0.473	$F(1,43) = 0.224$
Proportion Child Molestation	8	44	0.337	2.105*	0.395	0.451	$F(1,42) = 0.203$
Proportion Contact Offenses	10	64	0.332	2.554*	0.084	0.185	$F(1,62) = 0.034$
Recidivism Risk	12	72					$F(1,70) = 1.162$
High risk (RC)	8	49	0.423	3.284**			
Low risk	4	23	0.186	1.044	−0.237	−1.078	
Living Conditions	13	77					$F(2,74) = 0.085$
Residential (RC)	3	12	0.327	1.432			
Living at home	5	31	0.427	2.393*	0.100	0.345	
Mixed	5	34	0.339	1.971+	0.012	0.041	
Treatment characteristics							
Type of Treatment	13	77					$F(2,74) = 0.372$
TAU (RC)	3	14	0.213	0.993			
CBT	6	46	0.438	2.864**	0.226	0.857	
Systemic	4	17	0.384	1.961+	0.171	0.590	
Duration Treatment	9	49	0.400*	2.554	−0.002	−0.116	$F(1,47) = 0.014$
Study characteristics							
Authors	13	77					$F(1,75) = 1.262$
Dependent (RC)	6	38	0.488	3.342**			
Independent	7	39	0.265	1.969+	0.223	1.123	
Peer Reviewed	13	77					$F(1,75) = 2.456$
Yes (RC)	9	51	0.463	4.125***			
No	4	26	0.144	0.850	−0.319	−1.567	
Study Design	13	77					$F(1,75) = 2.379$
Randomized (RC)	3	13	0.653	3.131**			
Quasi-experimental	10	64	0.292	2.739**	−0.361	−1.542	
Type of Control Treatment	12	61					$F(1,55) = 0.057$
Treatment as usual (RC)	5	23	0.423	2.895**			
Established treatment	7	34	0.389	3.057**	−0.035	−0.233	
Comparability of Groups	10	56					$F(1,54) = 0.026$
High (RC)	5	35	0.374	2.867**			
Low	5	21	0.351	0.162	−0.023	−0.162	
Length Follow Up	10	68	0.336	2.460*	−0.002	−1.919+	$F(1,66) = 3.681+$
Type effect	13	77					$F(1,75) = 5.249*$
Proportion (RC)	13	68	0.396	3.728***			
Means	4	9	0.162	1.167	−0.234	−2.291*	
Recidivism Measure	13	77					$F(1,75) = 1.021$
Reconviction (RC)	3	20	0.178	0.841			
Rearrest	10	57	0.421	3.731***	0.243	1.011	
Study Quality	13	77					$F(1,75) = 0.018$
Strong/Moderate (RC)	7	54	0.380	2.739**			
Weak	6	23	0.353	2.270*	−0.028	−0.134	

studies = number of independent studies, #ES = number of effect sizes, t_0 = difference in mean r with zero, t_1 = difference in mean d with reference category, mean d = mean effect size (d), $F(df_1, df_2)$ = omnibus test, (RC) = reference category

+ = trend, significant at a 0.1 level, * = significant at a 0.05 level, ** = significant at a 0.01 level, *** = significant at a 0.001 level

the difference in effect on recidivism between the treatment conditions and the comparison conditions might have been reduced. Comparing treatment to a control condition (a group that receives no treatment at all), is

considered unethical in this population as it requires withholding treatment. Such research, therefore, will remain scarce. Few studies ($N = 3$) made use of drop out control groups. These studies, in contrast, may have

Note. The closed circles represent the primary studies included. The open circles represent forecasted 'missing' effect sizes, pointing out possible publication bias

Fig. 2 Results of trim-and-fill procedure testing for publication bias

resulted in an overestimation of the effect of treatment because drop outs are at higher risk for reoffending (Hendriks and Bijleveld 2005).

Significant variance was present at the between-study level. This indicates that the effect size distribution was heterogeneous, and that moderating variables might explain differences in strength of the effect sizes. Therefore, moderator analyses were performed.

The type of recidivism measured did not moderate the effects significantly; on all types of recidivism a moderate effect was achieved. This result could signify that, in juveniles, while treating sexual offending behavior, other offending behavior is also treated. This effect has also previously been established by Lösel and Schmucker (2005) and Schmucker and Lösel (2015). Treatment usually entails a confrontation with the sexual offending behavior, thoughts, and feelings prior to, during, and after the offense, the development of victim empathy, the development of anger management and stress coping skills, social skills training, treatment of the juveniles own victimization experiences, and relapse prevention strategies (Becker and Hunter 1997; Becker and Johnson 2001). Moreover, contemporary treatments have been developed to take into account broader issues. The Good Lives Model, for example, was presented as an addition to the relatively risk focused RNR model (Lord 2016). Contextual treatment forms, such as Multisystemic Therapy-Problem Sexual Behavior, are explicitly holistic in nature (Swenson et al. 2005). Thus, risk factors for juvenile delinquency in general also seem to be more and more addressed via treatment aimed at sex offending. This is

considered expedient, because while sexual recidivism amongst juveniles is rare, other types of recidivism are relatively frequent (Caldwell 2010; Caldwell 2016; Lobanov-Rostovsky 2015).

None of the tested participant characteristics moderated the effect of treatment on recidivism in juveniles. Thus, the currently coded subgroups within the sample did not account for differences in effects on reducing recidivism between the treatment and comparison or control groups. Notably, few studies defined their participants more specifically than juveniles who have sexually offended, and as a result, in the included studies the heterogeneous group of juveniles who commit sexual offenses was presented as relatively homogenous. This hampered research into moderating effects of participant characteristics. Many studies on specific mental illnesses, traits, executive functioning, and personality profiles of adolescents who have sexually offended have been conducted (Adjorlolo and Egbenya 2016; Glowacz and Born 2012; Hart-Kerkhoffs et al. 2009; Lawing et al. 2010; Margari et al. 2015; Purcel 2010; Seto and Lalumière 2010). None of the studies included in the current meta-analysis, however, reported on psychological profiles or the specific treatment needs juveniles presented at admission that deemed them eligible or in need of specific treatment (other than their type of offending behavior). Also, several typologies are commonly in use: non-contact versus contact offenders, sex-only offenders versus mixed or sex-plus offenders, peer abusers versus child abusers, and group offenders versus solo offenders (Butler and Seto 2002; Drew 2013; Hendriks 2006; Hissel et al. 2006; Kjellgren et al. 2006), and are described as in need of different treatment approaches. Of these typologies, only two (victim age and type of offending behavior) could be tested in this study. A more extensive moderator analysis could have been performed if studies had reported on treatment needs or typologies of their respondents.

Further, none of the coded treatment characteristics moderated the effect of treatment. The treatment conditions were grouped as predominantly cognitive behavioral based (e.g., SAFE-T, TBASOP), systemic (i.e., Multi Systemic Therapy and Functional Family Therapy), or as treatment as usual for juveniles who have sexually offended (TAU). The first two categories are established treatments, the third category contained a compilation of non-established treatment forms (i.e., adventure or wilderness therapy, individual treatment, or family treatment). The established treatments were expected to achieve better results in reducing recidivism in juveniles based on previous research (Hanson et al. 2009; Lösel and Schmucker 2005; Schmucker and Lösel 2015; Walker et al. 2004).

In study characteristics, one moderator significantly explained variance (calculation type of effect size), while for another (follow up time) a moderating trend was found, indicating stronger treatment effects in shorter follow up periods.

These results are in line with expectations. Calculation of Cohen's d via proportions (the relative number of recidivists per group) yielded a more pronounced effect (difference measured) than calculation via mean number of re-offenses (how many times recidivists reoffended per group). This indicates that specialist treatment showed a stronger reduction in the number of juveniles reoffending than comparison conditions did, but that reoffenders recidivated as many times in both groups. Regarding follow up time, most recidivism by juveniles who have sexually offended was reported in the first years after release (Hendriks and Bijleveld 2005), while the majority of adolescent offending proves to be adolescence limited (Lussier et al. 2012; Moffit 1993). Therefore, differences in percentages of recidivism were expected to be most pronounced in the first years after treatment. Lastly, comparing treatment with a non-established treatment comparison group did not generate larger effects than comparing treatment with an established treatment comparison group. The inclusion of drop-out respondents in an established treatment comparison group (by most of the studies), may have reduced their effect, since drop-outs were not able to profit fully from the established treatment administered. Drop-outs also may have been more distressed or less motivated than treatment completers; a reason for dropping out and achieving less positive results (Hendriks and Bijleveld 2005).

Notably, not many moderating influences were detected. However, non-significant or borderline significant results may still prove clinically relevant (Man-Son-Hing et al. 2002), especially when an analysis is conducted with a relatively small sample size. We refer to the limitations section for an overview of our borderline significant findings.

Limitations

This study was limited by only using studies that relied on official measures of recidivism. Thereby, undoubtedly, recidivistic behavior was missed that was not officially reported. Levels of unreported sexual problem behavior ('dark numbers') are known to be quite high (Wittebrood 2006). Also, official reports involve getting caught, which results in a specific subgroup of treated juveniles who were caught sexually offending, and again getting caught reoffending (Yun and Lee 2013). The use of a conservative measure, however, to our opinion generated greater comparability between studies and, therefore, more reliable results.

The restrictive inclusion criteria used, and, therefore, the relatively small number of included studies (mostly performed on small samples), presented us with underpowered analyses. However, to assess the effect of treatment the inclusion of studies with a comparison group is imperative (Weisburd et al. 2001). Some results previously detected in other reviews might have been replicated, if a larger sample size would have been available (i.e., the moderating effect of established

treatment, high at recidivism risk respondent groups, dependent authors, publication, study design, and re-arrest as outcome measure), because some of these variables did show rather pronounced, yet non-significant, differences in mean effect size (see Table 3).

Furthermore, there are methodological difficulties in the use of trim and fill procedures in multilevel meta-analytic data (Nakagawa and Santos 2012; Peters et al. 2007; Terrin et al. 2003). Therefore, the difference between the adjusted and observed mean effect size in this study should be interpreted as indicative of (the effects of) publication bias; the adjusted effect size is not to be considered the true effect size.

In the included studies, most likely due to ethical considerations, one type of treatment was usually compared to another type of treatment, that is, treatment as usual. Only one comparison of a treatment group to an explicit non-treatment (refuser) group was available (Worling and Curwen 2000). As a result, the moderating effect of no treatment could not be tested and no statement about the effects of therapy as compared to administering no therapy could be made. Also, we could not test whether treatment specifically aimed at juveniles who have sexually offended proved as effective or less effective as non-specific treatment (treatment aimed at juveniles with all types of behavioral problems) in reducing recidivism, for only one study addressed this issue (Guarino-Ghezzi and Kimball 1998).

Lastly, recidivism is just one aspect relevant to assess the quality of treatment. In addition to reduction of recidivism, treatment aims at improving the quality of life and psychosocial functioning of clients. This dual focus is important to bear in mind, since some treatments may excel in reducing recidivism (the treatment need of society), while others might more primarily serve treatment needs as formulated by their clients (quality of life). A meta-analysis on the effect of treatment on psychosocial improvement would, therefore, complement the findings of the current study.

Conclusion

Juveniles who have sexually offended constitute a notoriously heterogeneous group regarding treatment needs and offending behavior patterns. Various treatment forms are aimed at them, although sexual recidivism rates are generally low. Relatively low sexual recidivism rates, however, do not imply that juveniles who have sexually offended are not in need of therapy to reduce recidivism risk, if only because their general recidivism rates are relatively high. To date, recidivism risk factors that are specific to juveniles who have sexually offended mainly indicate at what specific criminogenic treatment needs treatment should be aimed (i.e., childhood sexual victimization, sexual deviation, association with younger children, poor school performance, not attending school and lack of intimate peer relationships/social isolation). Also, mixed offenders,

who display more antisocial tendencies, have been shown to be at higher risk for recidivism, and therefore, in greater need of treatment. The current study has shown sex offense specific treatment to have an equal effect on other types of recidivistic behavior. Thus, juveniles who have sexually offended and who display specific as well as general criminogenic needs may be best off in treatment aimed at reducing sexual recidivism. Not all juveniles who have sexually offended, however, display these features equally, so not all juveniles who have sexually offended may be in need of sex offense specific treatment as to reduce recidivism.

The authors would like to encourage more robust (independent, high quality) studies on the effect of treatment for juveniles who have sexually offended, possibly making use of a non-specific treatment comparison group. Reporting on specific treatment needs might also shed more light onto what treatment works, for what (type of) sexually offending juvenile. This could improve allocation to treatment and contribute to more positive treatment effects.

Compliance with Ethical Standards

Conflict of Interest The authors declare that they have no conflict of interest.

Ethical Approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent Informed consent was obtained from all individual participants included in the study.

Human Studies This article does not contain any studies with human participants performed by any of the authors.

References

The asterisks indicate original research included in the analysis.

- Adjorlolo, S., & Egbenya, D. L. (2016). Executive functioning profiles of adolescent and juvenile male sexual offenders: A systematic review. *The Journal of Forensic Psychiatry and Psychology*, 27, 349–375.
- Armijo-Olivo, S., Stiles, C. R., Hagen, N. A., Biondo, P. D., & Cummings, G. G. (2012). Assessment of study quality for systematic reviews: A comparison of the Cochrane collaboration risk of bias tool and the effective public health practice project quality assessment tool. *Journal of Evaluation in Clinical Practice*, 18, 12–18.
- Assink, M., & Wibbelink, C. M. (2016). Fitting three-level meta-analytic models in R: A step-by-step tutorial. *The Quantitative Methods for Psychology*, 12, 154–174.
- Assink, M., Van Der Put, C. E., Hoeve, M., De Vries, S. L., Stams, G. J. J., & Oort, F. J. (2015). Risk factors for persistent delinquent

- behavior among juveniles: A meta-analytic review. *Clinical Psychology Review*, 42, 47–61.
- Barbaree, H. E., & Marshall, W. L. (2006). *The juvenile sex offender*. New York: The Guilford Press.
- *Barlow, K. N. (1998). *Recidivism rates of level six residential programs for youthful male sexual offenders: 1995–1996*. Unpublished Master's thesis. Utah: Utah State University.
- Beaudry-Cyr, M., Jennings, W. G., Zgoba, K. M., & Tewksbury, R. (2015). Examining the continuity of juvenile sex offending into adulthood and subsequent patterns of sex and general recidivism. *International Journal of Offender Therapy and Comparative Criminology*, 9, 1–18.
- Becker, J. V., & Hunter, J. A. (1997). Understanding and treating child and adolescent sex offenders. In T. H. Ollendick & R. J. Prinz (Eds.), *Advances in clinical Psychology* (pp. 177–197). New York: Plenum.
- Becker, J. V., & Johnson, B. R. (2001). Treating juvenile sex offenders. In J. Ashford & B. Sales (Eds.), *Treating adult and juvenile sex offenders with special needs* (pp. 273–289). Washington DC: American Psychological Association.
- Beggs, S. M., & Grace, R. C. (2011). Treatment gain for sexual offenders against children predicts reduced recidivism: A comparative validity study. *Journal of Consulting and Clinical Psychology*, 79, 182–192.
- Bonta, J., & Andrews, D. A. (2007). *Risk-need-Responsivity model for offender assessment and rehabilitation*. Ottawa: Public Safety Canada.
- *Borduin, C. M., Henggeler, S. W., Blaske, D. M., & Stein, R. J. (1990). Multisystemic treatment of adolescent sexual offenders. *International Journal of Offender Therapy and Comparative Criminology*, 996, 105–113.
- *Borduin, C. M., Heiblum, N., & Schaeffer, C. M. (2009). A randomized clinical trial of Multisystemic therapy with juvenile sexual offenders: Effects on youth social ecology and criminal activity. *Journal of Consulting Psychology*, 77, 26–37.
- Brannon, J. M., & Troyer, R. (1991). Peer group counseling: A normalized residential alternative to the specialized treatment of adolescent sex offenders. *International Journal of Offender Therapy and Comparative Criminology*, 35, 225–234.
- Butler, S., & Seto, M. C. (2002). Distinguishing two types of adolescent sex offenders. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41, 83–90.
- Caldwell, M. F. (2010). Study characteristics and recidivism base rates in juvenile sex offender recidivism. *International Journal of Offender Therapy and Comparative Criminology*, 54, 197–211.
- Caldwell, M. F. (2016). Quantifying the decline in juvenile sexual recidivism rates. *Psychology, Public Policy and Law*, 22, 414–426.
- Cale, J., Smallbone, S., Rayment-McHugh, S., & Dowling, C. (2016). Offense trajectories, the unfolding of sexual and non-sexual criminal activity, and sex offense characteristics of adolescent sex offenders. *Sexual Abuse: A Journal of Research and Treatment*, 28, 791–812.
- Carpentier, J., & Proulx, J. (2011). Correlates of recidivism among adolescents who have sexually offended. *Sexual Abuse: A Journal of Research and Treatment*, 23, 434–455.
- Carpentier, M. Y., Silovsky, J. F., & Chaffin, M. (2006). Randomized trial of treatment for children with sexual behavior problems: Ten years later. *Journal of Consulting and Clinical Psychology*, 74, 482–488.
- Cheung, M. W. L. (2014). Modeling dependent effect sizes with three-level meta-analyses: A structural equation modeling approach. *Psychological Methods*, 19, 211–229.
- Christiansen, A. K., & Vincent, J. P. (2013). Characterization and prediction of sexual and nonsexual recidivism among adjudicated juvenile sex offenders. *Behavioural Sciences and the Law*, 31, 506–529.
- Deeks, J. J., Dinnes, J., D'Amico, R., Sowden, A. J., Sakarovich, C., Song, F., et al. (2003). Evaluating non-randomized intervention studies. *Health Technology Assessment*, 7, 3–173.

- Dopp, A. R., Borduin, C. M., & Brown, C. E. (2015). Evidence-based treatments for juvenile sexual offenders: Review and recommendations. *Journal of Aggression, Conflict and Peace Research*, 7, 223–236.
- Drew, C. H. (2013). Classification of juvenile sexual offenders by victim age based subgroups. *Electronic Theses, Treatises and Dissertations*. Paper 7359.
- Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56, 455–463.
- Fanniff, A. M., & Kimonis, E. R. (2014). Juveniles who have committed sexual offenses: A special group? *Behavioural Sciences and the Law*, 32, 240–257.
- *Gillis, H. L., & Gass, M. A. (2010). Treating juveniles in a sex offender program using adventure-based programming: A matched group design. *Journal of Child Sexual Abuse*, 19, 20–34.
- Glowacz, F., & Born, M. (2012). Do adolescent child abusers, peer abusers, and non-sex offenders have different personality profiles? *European Child and Adolescent Psychiatry*, 22, 117–225.
- Groth, N. A., Longo, R. E., & McFaddin, J. B. (1982). Undetected recidivism among rapists and child molesters. *Crime and Delinquency*, 128, 450–458.
- *Guarino-Ghezzi, S., & Kimball, L. M. (1998). Juvenile sex offenders in treatment. *Corrections Management Quarterly*, 2, 45–54.
- Hanson, R. K., & Morton-Bourgon, K. E. (2005). The characteristics of persistent sexual offenders: A meta-analysis of recidivism studies. *Journal of Consulting and Clinical Psychology*, 73, 1154–1163.
- Hanson, R. K., Bourgon, G., Helmus, L., & Hodgson, S. (2009). *A meta-analysis of the effectiveness of treatment for sexual offenders: Risk, need, and Responsivity*. Ottawa: Public Safety Canada.
- Hart-Kerkhoffs, L. A., Doreleijers, T. A. H., Jansen, L. M. C., Van Wijk, A. P. H., & Bullens, R. A. R. (2009). Offense related characteristics and psychosexual development of juvenile sex offenders. *Child and Adolescent Psychiatry and Mental Health*, 3, 1–10.
- Hendriks, J. (2006). *Jeugdige zedendelinquenten. Een studie naar subtypen en recidive. [Juvenile sex offenders. A study on subtypes and recidivism]*. Dissertation. Retrieved from <http://dare.uvu.vu.nl/handle/1871/15503>.
- *Hendriks, J., & Bijleveld, C. (2005). Recidive van jeugdige zedendelinquenten: poliklinisch behandelde versus niet-behandelde. [Recidivism by juvenile sexual offenders: Outpatient treatment versus non treatment]. *Tijdschrift voor Seksuologie*, 29, 150–160.
- Henggeler, S. W., Letourneau, E. J., Chapman, J. E., Borduin, C. M., Schewe, P. A., & McCart, M. R. (2009). Mediators of change for multisystemic therapy with juvenile sexual offenders. *Journal of Consulting and Clinical Psychology*, 77, 451–462.
- Hissel, S., Bijleveld, C., Hendriks, J., Jansen, B., & Collot d'Escury-Koenigs, A. (2006). Jeugdige zedendelinquenten: Specialisten, generalisten en first offenders [juvenile sex offenders: Specialists, generalists, and first offenders]. *Tijdschrift voor Seksuologie*, 30, 215–225.
- *Holly, M. C. (2000). *Long-term follow up of a community based treatment program for adolescent sex offenders*. Unpublished master's thesis. Thunder Bay: Lakehead University.
- Houben, M., Van Den Noortgate, W., & Kuppens, P. (2015). The relation between short-term emotion dynamics and psychological well-being: A meta-analysis. *Psychological Bulletin*, 141, 901–930.
- Hox, J. (2002). *Multilevel analysis: Techniques and applications*. Mahwah: Lawrence.
- Jackson, N., & Waters, E. (2005). Criteria for the systematic review of health promotion and public health interventions. *Health Promotion International*, 20, 367–374.
- *Johnson Erickson, C. (2008). *The effectiveness of functional family therapy in the treatment of juvenile sexual offenders*. Unpublished thesis. Bloomington: Indiana University.
- Kahn, T. J., & Chambers, H. J. (1991). Assessing re-offense risk with juvenile sexual offenders. *Child Welfare*, 70, 333–345.
- Kjellgren, C., Wassberg, A., Carlberg, M., Långström, N., & Svedin, C. G. (2006). Adolescent sexual offenders: A total survey of referrals to social services in Sweden and subgroup characteristics. *Sexual Abuse: A Journal of Research and Treatment*, 18, 357–372.
- Knapp, G., & Hartung, J. (2003). Improved tests for a random effects meta-regression with a single covariate. *Statistics in Medicine*, 22, 2693–2710.
- *Lab, S. P., Schields, G., & Schondel, C. (1993). Research note: An evaluation of juvenile sexual offender treatment. *Crime and Delinquency*, 39, 543–553.
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33, 159–174.
- Lawing, C., Frick, P. J., & Cruise, K. R. (2010). Differences in offending patterns between adolescent sex offenders high or low in callous-unemotional traits. *Psychological Assessment*, 22, 298–305.
- Letourneau, E. J., & Armstrong, K. S. (2008). Recidivism rates for registered and nonregistered juvenile sexual offenders. *Sexual Abuse*, 20, 393–408.
- Letourneau, E. J., Henggeler, S. W., Borduin, C. M., Schewe, P. A., McCart, M. R., Chapman, J. E., & Saldana, L. (2009). Multisystemic therapy for juvenile sexual offenders: 1 year results from a randomized effectiveness trial. *Journal of Family Psychology*, 23, 89–201.
- *Letourneau, E. J., Borduin, C. M., Henggeler, S. W., McCart, M. R., Schewe, P. A., & Armstrong, K. S. (2013). Two-year follow-up of a randomized effectiveness trial evaluating MST for juveniles who sexually offend. *Journal of Family Psychology*, 27, 978–985.
- Lipsey, M. W., & Wilson, D. B. (2001). *Practical meta-analysis*. Thousand Oaks: Sage publications.
- Lobanov-Rostovsky, C. (2015). *Recidivism of juveniles who commit sexual offenses*. SO+MAPI Research Brief. July 2015.
- Loeber, R., Hovee, M., Slot, N. W., & Van Der Laan, P. H. (2012). *Persisters and desisters in crime from adolescence into adulthood*. Surrey: Ashgate Publishing Limited.
- Lord, A. (2016). Integrating risk, the good lives model and recovery for mentally disordered sexual offenders. *Journal of Sexual Agression*, 22, 107–122.
- Lösel, F., & Schmucker, M. (2005). The effectiveness of treatment for sexual offenders: A comprehensive meta-analysis. *Journal of Experimental Criminology*, 1, 117–146.
- Lussier, P., Van Den Berg, C., Bijleveld, C., & Hendriks, J. (2012). A developmental taxonomy of juvenile sex offenders for theory, research and prevention. *Criminal Justice and Behaviour*, 39, 1559–1581.
- Man-Son-Hing, M., Laupacis, A., O'Rourke, K., Molnar, F. J., Mahon, J., Chan, K. B. Y., & Wells, G. (2002). Determination of the clinical importance of study results: A review. *Journal of General Internal Medicine*, 17, 469–476.
- Margari, F., Lecce, P. A., Craig, F., Laforzezza, E., Lisi, A., Pinto, F., Stallone, F., Pierri, G., Pisani, R., Zagaria, G., Margari, L., & Grattagliano, I. (2015). Juvenile sex offenders: Personality profile, coping styles and parental care. *Psychiatry Research*, 229, 82–88.
- Moffitt, T. E. (1993). Adolescence-limited and life-course persistent antisocial behaviour, a developmental taxonomy. *Psychological Review*, 100, 674–701.
- Nakagawa, S., & Santos, E. S. A. (2012). Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology*, 26, 1253–1274.
- Pedersen, L., Rasmussen, K., & Elsass, P. (2010). Risk assessment: The value of structured professional judgments. *International Journal of Forensic Mental Health*, 9, 74–81.
- Peters, J. L., Sutton, A. J., Jones, D. R., Abrams, K. R., & Rushton, L. (2007). Performance of the trim and fill method in the presence of

- publication bias and between-study heterogeneity. *Statistics in Medicine*, 26, 4544–4562.
- Purcel, M. (2010). A personality based classification of male adolescent sex offenders using the MACI (Thesis). Retrieved from <https://researchspace.auckland.ac.nz/bitstream/handle/2292/6538/01front.pdf?sequence=2>.
- Reitzel, L. R., & Carbonell, J. L. (2006). The effectiveness of sexual offender treatment for juveniles as measured by recidivism: A meta-analysis. *Sexual Abuse*, 18, 401–421.
- Rosenthal, R. (1979). The file drawer problem and tolerance for null results. *Psychological Bulletin*, 86, 638–641.
- Ryan, G., Laversee, T., & Lane, S. (2010). *Juvenile sexual offending, causes, consequences and correction*. Hoboken: John Wiley and Sons.
- Schmucker, M., & Lösel, F. (2015). The effects of sexual offender treatment on recidivism: An international meta-analysis of sound quality evaluations. *Journal of Experimental Criminology*, 11, 597–630.
- *Schram, D. D., Darling Milloy, C., & Rowe, W. E. (1991). *Juvenile sex offenders: a follow up study of reoffensive behavior*. Unpublished research report. Washington: Washington State institute for public policy.
- Seabloom, W., Seabloom, M. E., Seabloom, E., Barron, R., & Hendrickson, S. (2003). A 14 to 24 year longitudinal study of a comprehensive sexual health model treatment program for adolescent sex offenders. *International Journal of Offender Therapy and Comparative Criminology*, 47, 468–481.
- Seto, M. C., & Lalumière, M. L. (2010). What is so special about male adolescent sexual offending? A review and test of explanations through meta-analysis. *Psychological Bulletin*, 136, 526–575.
- Shrout, P. E. (1989). Measurement reliability and agreement in psychiatry. *Statistical Methods in Medical Research*, 7, 301–317.
- Spruit, A., Assink, M., Van Vugt, E. S., Van Der Put, C. E., & Stams, G. J. J. M. (2016). The effects of physical activity interventions on psychosocial outcomes in adolescents: A meta-analytic review. *Clinical Psychology Review*, 45, 56–71.
- Swenson, C. C., Henggeler, S. W., Taylor, I. S., & Addison, O. W. (2005). *Multisystemic Therapy and Neighborhood Partnerships*. New York: The Guildford Press.
- Tabachnik, B. G., & Fidell, L. S. (2013). *Using multivariate statistics* (6th ed.). Boston: Allynand Bacon.
- Ter Beek, E., Van Der Rijken, R. A. E., Kuiper, C. H. Z., Hendriks, J., & Stams, G. J. J. M. (2016). The allocation of sexually transgressive juveniles to intensive specialized treatment. *International Journal of Offender Therapy and Comparative Criminology*. Advance online publication.
- Terrin, N., Schmid, C. H., Lau, J., & Olkin, I. (2003). Adjusting for publication bias in the presence of heterogeneity. *Statistics in Medicine*, 22, 2113–2126.
- Thomas, B. H., Ciliska, D., Dobbins, M., & Micucci, S. (2004). A process for systematically reviewing the literature: Providing the research evidence for public health nursing interventions. *Worldviews on Evidence-Based Nursing*, 1, 176–184.
- Van Den Noortgate, W., & Onghena, P. (2003). Multilevel meta-analysis: A comparison with traditional meta-analytical procedures. *Educational and Psychological Measurement*, 6, 765–790.
- Van den Noortgate, W., López-López, J. A., Marín-Martínez, F., & Sánchez-Meca, J. (2013). Three-level meta-analysis of dependent effect sizes. *Behavior Research Methods*, 45, 576–594.
- Van Outsem, R. E. (2009). *Exploring psychological characteristics of sexually abusive juveniles*. Unpublished Thesis. Amsterdam: Vrije Universiteit Amsterdam.
- Van Wijk, A. P., Vreugdenhil, C., Van Horn, J., Vermeiren, R., & Doreleijers, T. A. H. (2007). Incarcerated Dutch juvenile sex offenders compared with non-sex offenders. *Journal of Child Sexual Abuse*, 16, 1–21.
- Veneziano, C., & Veneziano, L. (2002). Adolescent sex offenders: A review of the literature. *Trauma, Violence and Abuse*, 3, 247–260.
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36, 1–48.
- *Waite, D., Keller, A., McGarvey, E. L., Wieckowski, E., Pinkerton, R., & Brown, G. L. (2005). Juvenile sex offender re-arrest rates for sexual, violent nonsexual and property crimes: A 10-year follow-up study. *Sexual Abuse: A Journal of Research and Treatment*, 17, 313–331.
- Walker, D. F., McGovern, S. K., Poey, E. L., & Otis, K. E. (2004). Treatment effectiveness for male adolescent sexual offenders: A meta-analysis and review. *Journal of Child Sexual Abuse*, 13, 281–293.
- Weinrott, M. R., Riggan, M., & Frothingham, S. (1997). Reducing deviant arousal in juvenile sex offenders using vicarious sensitization. *Journal of Interpersonal Violence*, 12, 704–728.
- Weisburd, D., Lum, C., & Petrosino, A. (2001). Does research design affect study outcomes in criminal justice? *The Annals of the American Academy of Political and Social Science*, 578, 50–70.
- Weisz, J. R., Kuppens, S., Eckshtain, D., Ugueto, A. M., Hawley, K. M., & Jensen-Doss, A. (2013). Performance of evidence-based youth psychotherapies compared with usual clinical care. *JAMA Psychiatry*, 70, 750–761.
- Wibbelink, C. J. M., & Assink, M. (2015). *Manual for conducting a three-level meta-analysis in R*. Amsterdam: Universiteit van Amsterdam.
- Wilson, D. B. (2013). Practical meta-analysis effect size calculator. Retrieved from: <http://www.campbellcollaboration.org/escalc/html/EffectSizeCalculator-Home.php>.
- Wittebrood, K. (2006). *Slachtoffers van criminaliteit: Feiten en achtergronden. (Victims of crime: facts and backgrounds)*. Den Haag: Sociaal Cultureel Planbureau.
- *Worling, J. R., & Curwen, T. (2000). Adolescent sexual offender recidivism: Success of specialized treatment and implications of risk prediction. *Child Abuse and Neglect*, 24, 965–982.
- Worling, J. W., & Långström, N. (2003). Assessment of criminal recidivism risk with adolescents who have offended sexually: A review. *Trauma, Violence & Abuse*, 4, 341–362.
- *Worling, J. R., Littlejohn, M. A., & Bookalam, M. A. (2010). 20-year prospective follow-up study of specialized treatment for adolescents who offend sexually. *Behavioural Sciences and the Law*, 28, 46–57.
- Yun, I., & Lee, J. (2013). IQ and delinquency: The differential detection hypothesis revisited. *Youth Violence and Juvenile Justice*, 11, 196–211.