



## UvA-DARE (Digital Academic Repository)

### Chasing sympatric speciation: The relative importance and genetic basis of prezygotic isolation barriers in diverging populations of *Spodoptera frugiperda*

Hänniger, S.

**Publication date**

2015

**Document Version**

Final published version

[Link to publication](#)

**Citation for published version (APA):**

Hänniger, S. (2015). *Chasing sympatric speciation: The relative importance and genetic basis of prezygotic isolation barriers in diverging populations of Spodoptera frugiperda*. [Thesis, fully internal, Universiteit van Amsterdam].

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

**GEOGRAPHIC VARIATION IN SEXUAL  
ATTRACTION OF *SPODOPTERA FRUGIPERDA*  
CORN- AND RICE-STRAIN MALES TO  
PHEROMONE LURES**

**Melanie Unbehend<sup>1</sup>, Sabine Hänniger<sup>1</sup>, Gissella M. Vásquez<sup>2</sup>,  
María Laura Juárez<sup>3</sup>, Dominic Reisig<sup>4</sup>, Jeremy N. McNeil<sup>5</sup>,  
Robert L. Meagher<sup>6</sup>, David A. Jenkins<sup>7</sup>, David G. Heckel<sup>1</sup> &  
Astrid T. Groot<sup>1,8</sup>**

<sup>1</sup>Department of Entomology, Max Planck Institute for Chemical Ecology (MPICE), Hans-Knoell-Str. 8, 07745 Jena, Germany; <sup>2</sup>Department of Entomology, North Carolina State University, Raleigh, 27695 NC, USA; <sup>3</sup>Sección Zoología Agrícola, Estación Experimental Agroindustrial Obispo Colombres, Las Talitas (T4104AUD), Tucumán, Argentina; <sup>4</sup>North Carolina State University, Vernon James Research & Extension Center, 207 Research Station Road, Plymouth, 27962 NC, USA; <sup>5</sup>University of Western Ontario, Department of Biology, Biological & Geological Sciences Building, London, Ontario, N6A 5B7 Canada; <sup>6</sup>USDA-ARS, CMAVE, 1700 SW 23rd Dr., Gainesville, 32608 FL, USA; <sup>7</sup>Tropical Agriculture Research Station, 2200 Pedro A. Campos Ave., Suite 201, Mayaguez, 00680 Puerto Rico; <sup>8</sup>Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands

**Abstract**

The corn- and rice-strains of *Spodoptera frugiperda* exhibit several genetic and behavioral differences and appear to be undergoing ecological speciation in sympatry. Previous studies reported conflicting results when investigating male attraction to pheromone lures in different regions, but this could have been due to inter-strain and/or geographic differences. Therefore, we investigated whether corn- and rice-strain males differed in their response to different synthetic pheromone blends in different regions in North America, the Caribbean and South America. All trapped males were strain-typed by two strain-specific mitochondrial DNA markers. In the first experiment, we found a nearly similar response of corn- and rice-strain males to two different 4-component blends, resembling the corn- and rice-strain female blend we previously described from females in Florida. This response showed some geographic variation in fields in Canada, North Carolina, Florida, Puerto Rico, and South America (Peru, Argentina). In dose-response experiments with the critical secondary sex pheromone component (Z)-7-dodecenyl acetate (Z7-12:OAc), we found some strain-specific differences in male attraction. While the response to Z7-12:OAc varied geographically in the corn-strain, rice-strain males showed almost no variation. We also found that the minor compound (Z)-11-hexadecenyl acetate (Z11-16:OAc) did not increase attraction of both strains in Florida and of corn-strain males in Peru. In a fourth experiment, where we added the stereo-isomer of the critical sex pheromone component, (E)-7-dodecenyl acetate, to the major pheromone component (Z)-9-tetradecenyl acetate (Z9-14:OAc), we found that this compound was attractive to males in North Carolina, but not to males in Peru. Overall, our results suggest that both strains show rather geographic than strain-specific differences in their response to pheromone lures, and that regional sexual communication differences might cause geographic differentiation between populations.

**INTRODUCTION**

Geographic variation in the sexual communication signals of animals is a widespread phenomenon, being reported in frogs (Ryan et al. 1996; Bernal et al. 2005; Pröhl et al. 2006), birds (Slabbekoorn and Smith 2002; Tack et al. 2005; Podos and Warren 2007), fish (Gonzalez-Zuarth et al. 2011) and insects (Ackerman 1989; Miller et al. 1997; Zhu et al. 2009). This variation can be the result of isolation by distance, with a positive correlation between genetic dissimilarity and geographic distance (Balaban 1988; MacDougall-Shackleton and MacDougall-Shackleton 2001; Lampert et al. 2003), but this is not always the case (Seppä and Laurila 1999; Leblois et al. 2000; Kaefer et al. 2012). Furthermore, mating signals can be influenced by environmental factors such as temperature (Delisle and Royer 1994; Olvido et al. 2010; Roeser-Mueller et al. 2010; Green et al. 2012), humidity (Kumar and Saxena 1986; Royer and McNeil 1991, 1993), photoperiod length (Delisle and McNeil 1987; Gemeno and Haynes 2001), host plant volatiles (Landolt and Phillips 1997; Reddy and Guerrero 2004) or interspecific olfactory cues (Groot et al. 2010b) that vary geographically.

Geographic variation in sexual communication systems has been reported in several lepidopteran species (Toth et al. 1992; McElfresh and Millar 1999; Wu et al. 1999; Gemeno et al. 2000; Kawazu et al. 2005; Groot et al. 2009) and is of interest because changes in the sex pheromone signal and/or response to sex pheromones could result in reproductive isolation and subsequently may lead to speciation (Roelofs and Cardé 1974; Phelan 1992; Baker 2002; Smadja and Butlin 2009). Furthermore, geographically varying sexual communication is of interest for pest management, as many lepidopteran insects are pest species which are commonly monitored, disrupted or killed via pheromone-mediated methods (Cardé and Minks 1995; El-Sayed et al. 2009; Witzgall et al. 2010).

The fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae), consists of two genetically and behaviorally distinct strains, the corn- and rice-strain, occurring sympatrically throughout North- and South America (Pashley 1986). Both strains appear to be undergoing ecological speciation in sympatry and reveal several possible prezygotic isolation barriers (Groot et al. 2010c). These include differential host plant choice (Pashley 1986, 1989; Lu and Adang 1996; Nagoshi et al. 2006a; Machado et al. 2008), strain-specific mating times in the scotophase (Pashley et al. 1992; Schöfl et al. 2009), as well as differences in the female sex pheromone composition (Groot et al. 2008; Lima and McNeil 2009; Unbehend et al. 2013). Among these prezygotic mating barriers, the strain-specific timing of reproduction seems to be the most important one that differentiates both strains, as host plant preference is not as clear cut as previously thought (Nagoshi et al. 2006a, 2007; Groot et al. 2010c; Juárez et al. 2012), and no strain-specific mating based on sex pheromone differences could be shown so far (Pashley et al. 1992; Meagher and Nagoshi 2013; Unbehend et al. 2013). Additionally, a postzygotic mating barrier, i.e. reduced fertility of RC (rice-strain ♀ × corn-strain ♂) hybrid females, contributes to the divergence of the two strains and separates them in nature (Groot et al. 2010c). Besides being an excellent model to study the evolution of reproductive isolation (Groot et al. 2010c), *S. frugiperda* is a serious pest species that feeds on a large variety of agricultural crops (Pashley 1988), and can cause annual damages of up to ~300 million dollars in the United States (Sparks 1986).

The sex pheromone of *S. frugiperda* was identified by Tumlinson et al. 1986) to consists of (Z)-9-tetradecenyl acetate (Z9-14:OAc) as the major sex pheromone component, and (Z)-7-dodecenyl acetate (Z7-12:OAc) as critical secondary sex pheromone component. A number of other minor compounds like (Z)-11-hexadecenyl acetate (Z11-16:OAc) and (Z)-9-dodecenyl acetate (Z9-12:OAc) have also been identified from the gland (Tumlinson et al. 1986; Groot et al. 2008; Lima and McNeil 2009), but with unclear behavioral function so far (Unbehend et al. 2013). Analysis of sex pheromone gland extracts from females collected in Florida showed that corn-strain females contained significantly lower relative amounts of

Z7-12:OAc and Z9-12:OAc than rice-strain females (Groot et al. 2008; Unbehend et al. 2013). However, different male trapping experiments conducted in Louisiana and Florida showed no consistent attraction of males to females of their own strain (Pashley et al. 1992; Meagher and Nagoshi 2013; Unbehend et al. 2013), which suggests that differences in the female pheromone are not sufficient to cause assortative mating in the field.

There is evidence that there are geographic differences in this species in the female sex pheromone blend (Tumlinson et al. 1986; Batista-Pereira et al. 2006; Groot et al. 2008; Lima and McNeil 2009), as well as in the male response (Jones and Sparks 1979; Mitchell et al. 1985; Tumlinson et al. 1986; Andrade et al. 2000; Fleischer et al. 2005; Batista-Pereira et al. 2006; Unbehend et al. 2013). For example, while females from Brazil (Batista-Pereira et al. 2006) produce (E)-7-dodecenyl acetate (E7-12:OAc), those from Florida, Louisiana or French Guyana do not (Descoins et al. 1988; Groot et al. 2008; Lima and McNeil 2009). In addition, studies on females originating from Florida and Louisiana provide evidence of geographic variation in the production of sex pheromone by females of both strains (Groot et al. 2008; Lima and McNeil 2009; Unbehend et al. 2013). Numerous studies have shown that the number of males caught varies with the pheromone blend used. For example, while the minor compound Z11-16:OAc did not affect male attraction in Florida and Brazil (Tumlinson et al. 1986; Batista-Pereira et al. 2006; Unbehend et al. 2013), it did marginally increase capture rates in Costa Rica (Andrade et al. 2000), and addition of Z11-16:OAc and Z9-12:OAc to binary blends (of Z9-14:OAc and Z7-12:OAc) almost doubled the attraction of males in Pennsylvania (Fleischer et al. 2005). However, most of these studies did not determine the strain identity of the males captured. Consequently, the variation in male attraction observed in these studies could either be due to strain-specific and/or due to geographic differences.

To disentangle strain-specific variation from geographic variation in male response, we investigated the response of corn- and rice-strain males to different synthetic pheromone blends in six different countries in North America, the Caribbean and South America. We tested (A) two synthetic 4-component blends (Blend 1 and 2) in different fields in Canada, North Carolina, Florida, Puerto Rico, Peru and Argentina; (B) different doses of Z7-12:OAc in Florida, Puerto Rico and Peru; (C) different doses of Z11-16:OAc in Florida and Peru; and (D) different doses of E7-12:OAc and Z7-12:OAc in Peru and North Carolina. We found that corn- and rice-strain males showed some variation in their response to different synthetic pheromone blends in different geographic regions. Overall, our results suggest that there is less strain-specific than geographic variation in male response and that regional sexual communication differences might cause geographic differentiation between populations.

## MATERIALS AND METHODS

### *Ethics statement*

The examined species is neither endangered nor protected. Trapping experiments in Canada, North Carolina and Peru were conducted at experimental research stations of the University of Western Ontario, the North Carolina State University, and the Universidad Nacional Agraria La Molina, with permission of the local university members/experimenters (Table 1). Trapping experiments in Florida were conducted at a) the Everglades Research and Education Center of the University of Florida in Belle Glade with permission of Gregg S. Nuessly, b) a private field in Hague with permission obtained by RLM, and c) private fields at the Graham Farm in Moore Haven with permission of the farm manager Tommy Toms (Table 1). Field experiments in Puerto Rico were conducted at field sites of the seed companies Monsanto and 3<sup>rd</sup> Millenium Genetics with permission of the company employees Wilson Rivera González and Jose Santiago, respectively. In Argentina, experiments were conducted in a private field with permission obtained by MLJ (Table 1).

### *Male trapping experiments*

To test whether a certain synthetic pheromone blend is equally attractive for corn- and rice-strain males in different geographic regions, four different trapping experiments were conducted in six different regions in North America, the Caribbean and South America (Table 1). For experiment (A), we prepared two synthetic 4-component blends (Blend 1 and 2) based on strain-specific pheromone differences found in laboratory females from Florida by Groot et al. (2008). Both blends consisted of 100% Z9-14:OAc, but with different percentages of Z11-16:OAc, Z7-12:OAc and Z9-12:OAc (Table 2), as described by Unbehend et al. (Unbehend et al. 2013). Both blends were tested in Canada, the United States (North Carolina and Florida), Puerto Rico, Peru and Argentina (Table 1).

To evaluate the relative importance of Z7-12:OAc for male attraction in Florida, Puerto Rico and Peru (experiment B), different percentages of Z7-12:OAc (0, 2, 4, 10%) were added to the major sex pheromone component Z9-14:OAc alone (Table 2). The percentages used were chosen to examine whether Z7-12:OAc is necessary for male attraction in all regions and fields (0%, lures baited only with Z9-14:OAc), to test whether males can distinguish between 2 and 4% Z7-12:OAc, which is the difference that we found between corn- and rice-strain females (Groot et al. 2008), and to investigate a possible repellent effect of high dosages of Z7-12:OAc (10%).

To assess whether Z11-16:OAc would affect male attraction, we conducted experiment (C), in which different amounts of Z11-16:OAc (0, 8, 13, 18%) were added to a 'minimal blend', consisting of 100% Z9-14:OAc and 2% Z7-12:OAc (Table 2). The 'minimal blend' (0% Z11-16:OAc) was used as control, while 8 and 13% Z11-16:OAc reflect the percentages found in rice- and corn-strain females

from Florida, respectively (Groot et al. 2008). To test possible repellent effects, 18% was used as the highest concentration. Experiment C was conducted in Florida and Peru.

**TABLE 1.** *Spodoptera frugiperda* trapping experiments conducted in North America, the Caribbean and South America.

Country		Location & Coordinates	Field	Experiment <sup>1</sup>	Experimenter	Date
Canada	Ontario	+43°4'26.08", -81°20'21.81"	Corn	A	JMN	Sept. 2011
North Carolina	Plymouth	+35°50'46.19", -76°39'46.24"	Soybean	A, D	DR	Sept. 2011
		+35°51'01.93", -76°39'11.28"	Cotton; Grass			
		+35°51'48.80", -76°39'33.61"	Soybean; Corn			
Florida <sup>2</sup>	Belle Glade	+26°40'7.20", -80°37'57.63"	Corn A	A, B	MU, SH	April-May 2010
	Hague	+29°47'7.40", -82°25'3.66"	Corn B	C	RLM	Sept. 2011
	Moore Haven	+26°53'3.04", -81°7'21.17"	Grass	A, B, C	MU, SH	April-May 2010
Puerto Rico	Santa Isabel	+17°59'0.93", -66°23'29.88"	Corn A	A	MU,SH, ATG,DAJ	April 2010
		+17°57'30.65", -66°23'32.43"	Corn B	A, B		
Peru	Lima	-12°4'51.56", -76°57'9.14"	Corn	A, B, C, D	GV	May-July 2011
Argentina	El Molino	-27°20'11.1", -65°41'25.8"	Corn	A	MLJ	Dec. 2010-Jan. 2011

<sup>1</sup>Experiments: A) Test of strain-specific blends, B) Z7-12:OAc dose-response, C) Z11-16:OAc dose-response, D) Importance of E7-12:OAc.

<sup>2</sup>Data adapted from Unbehend et al. 2013.

To test the importance of the isomers Z7-12:OAc and E7-12:OAc in North and South America (experiment D), we added different doses of E7-12:OAc and Z7-12:OAc (0, 1, 2%) to 100% Z9-14:OAc (Table 2). The ‘minimal blend’ (2% Z7-12:OAc + 100% Z9-14:OAc) was used as control, and as an equivalent we prepared an E-blend with 2% E7-12:OAc and 100% Z9-14:OAc. To investigate a possible interaction effect of both isomers together, 1% as well as 2% of E- and Z7-12:OAc were added to 100% Z9-14:OAc. The fourth experiment (Exp. D) was carried out in North Carolina and Peru.

We were not able to conduct all four experiments in all countries, due to technical limitations (i.e. limited time availability of collaborators, limited access to infested field sites and variability of moth population densities). All data from trapping experiments in Florida were published previously (Unbehend et al. 2013) and were included in this study for comparison. In all experiments, the synthetic pheromone lures were placed in plastic green-yellow-white Unitraps (Pherobank, Wageningen, the Netherlands), which contained a Vaportape II insecticide strip (Hercon Environmental, Emigsville, PA, USA) to kill the males captured. At each site, traps were hung just above the crop canopy (1-2 m above the ground depending on crop phenology), spaced 15 m apart and at least 15 m from the edge of the field using a complete randomized block design. There were three replicates per treatment per field (n=3), except for experiments conducted in North Carolina, where each replicate was conducted in a different field (Table 1). Traps were rotated and emptied three or four times, depending on the number of treatments (Exp. A: n=3; Exp. B-D: n=4), and traps were rotated every 1-6 days, depending on the population density in the field. The males captured were stored at -20 °C until strain-identification in the laboratory (see below).

### ***Preparation of pheromone lures***

All pheromone compounds used to prepare lures were bought from Pherobank (Wageningen, the Netherlands), and had a purity of  $\geq 99\%$ . Red rubber septa (Thomas Scientific, Swedesboro, NJ, USA) were soaked in hexane for 24 hours and air dried before they were loaded with 100  $\mu$ l hexane containing 300  $\mu$ g of the major pheromone component, Z9-14:OAc, plus different amounts (relative to 300  $\mu$ g Z9-14:OAc) of the minor compounds Z11-16:OAc, Z7-12:OAc, Z9-12:OAc, and E7-12:OAc (Table 2). To avoid variable loading of our multi-component lures, all components of one specific lure were mixed together in hexane, formulated into one blend, and checked on the gas chromatograph before they were loaded onto a septum. All lures within one experiment were prepared using one master solution to avoid variation between replicates. The loaded relative amounts of each minor pheromone component reflected the relative amounts present in the female pheromone gland (Groot et al. 2008). To avoid variation in attraction due to differential amounts of the major component, the amount of Z9-14:OAc was set to

100% in all blends (Table 2), similar to studies conducted by Batista-Pereira et al. (2006), and Groot et al. (2007, 2010a). Therefore, the sum of all components in our multi-component blends was always larger than 100%, and the total amount of pheromone per lure ranged from 300 up to 360  $\mu\text{g}$  (Table 2). All prepared lures were stored in glass vials at  $-20\text{ }^{\circ}\text{C}$  until used 1-3 months later in the field. Each lure was only used once within one experiment (for  $\sim$ 1-3 weeks), and we did not observe a consistent decrease in lure-effectiveness at the end of an experiment (Figure S1). However, we cannot completely exclude a decrease in emission rates over time.

### Chemical analysis

The purity and composition of the prepared pheromone solutions were verified by gas chromatography (GC) analysis, using a HP7890 gas chromatograph with a 7683 automatic injector. A 2  $\mu\text{l}$  aliquot of each pheromone solution used for the preparation of the pheromone lures (Table 2) was injected into a splitless inlet attached to a polar capillary column (DB-WAXetr; 30 m  $\times$  0.25 mm  $\times$  0.5  $\mu\text{m}$ ) and a flame-ionization detector (FID). The GC program ran from 60  $^{\circ}\text{C}$ , with a 2 min hold, to 180  $^{\circ}\text{C}$  at 30  $^{\circ}\text{C}/\text{min}$ , 230  $^{\circ}\text{C}$  at 5  $^{\circ}\text{C}/\text{min}$  and to 245  $^{\circ}\text{C}$  at 20  $^{\circ}\text{C}/\text{min}$ , followed by a 15 min hold at 245  $^{\circ}\text{C}$  to clean the column for the next sample. The FID detector was held at 250  $^{\circ}\text{C}$ .

**TABLE 2.** Composition of pheromone lures to test the attraction of *Spodoptera frugiperda* males in the field.

Experiment and Lures <sup>1</sup>		Z9-14:OAc	Z11-16:OAc	Z7-12:OAc	E7-12:OAc	Z9-12:OAc
A	Blend 1	100%	13%	2%	-	1%
	Blend 2		8%	4%	-	2%
	Blank (Hexane)		-	-	-	-
B	0%	100%	-	-	-	-
	2%		-	2%	-	-
	4%		-	4%	-	-
	10%		-	10%	-	-
C	0%	100%	-	-	-	-
	8%		8%	-	-	-
	13%		13%	-	-	-
	18%		18%	-	-	-
D	2% Z7-12:OAc	100%	-	2%	-	-
	2% E7-12:OAc		-	-	2%	-
	1+1% Z/E7-12:OAc		-	1%	1%	-
	2+2% Z/E7-12:OAc		-	2%	2%	-

<sup>1</sup>Compound concentrations were as follows: 100% = 300  $\mu\text{g}$ , 18% = 54  $\mu\text{g}$ , 13% = 39  $\mu\text{g}$ , 10% = 30  $\mu\text{g}$ , 8% = 24  $\mu\text{g}$ , 4% = 12  $\mu\text{g}$ , 2% = 6  $\mu\text{g}$ , 1% = 3  $\mu\text{g}$ .

### ***Strain identification***


The strain identity of all trapped males was determined via two strain-specific markers, i.e. *MspI*- and *SacI*-digest of the mitochondrial COI gene, which are known to be diagnostic for strain-identification of the two fall armyworm strains in North and South America (Meagher and Gallo-Meagher 2003; Nagoshi et al. 2006a, 2007). DNA of all males captured was extracted as described by Unbehend et al. (Unbehend et al. 2013) using CTAB (Cetyltrimethylammonium bromide) and isopropanol for DNA precipitation. The extracted DNA was tested at MPICE for strain-specific polymorphisms at the mitochondrial COI gene by amplification and strain-specific digestion (Nagoshi et al. 2006a; Unbehend et al. 2013). The amplified part of the COI gene was digested with *MspI* as well as *SacI* and analyzed electrophoretically on a 1% agarose gel (Unbehend et al. 2013). *MspI* digestion detected corn-strain individuals, whereas *SacI* digestion proved rice-strain identity (Nagoshi et al. 2006a).

### ***Statistical analysis***

Data of each field site of one experiment (Exp. A-D) were analyzed separately, using a generalized linear model (GLM) with a Poisson distribution, or a quasi-Poisson distribution if the residual deviance of the data was larger than the residual degrees of freedom (over-dispersion), using the R software 2.11.1 (R Development Core Team 2007). To assess whether there was any effect of geographic location, field crop, and/or any strain-specific effect that influenced male attraction, data of experiment A and B were additionally analyzed with a multivariate analysis of variance (MANOVA) and a Wilks Lambda test. Untransformed data were used for all GLM analyses, while data were square root transformed for the MANOVA. Treatments that did not catch any moths within any rotation in any of the three biological replicates per field were excluded from the statistical analysis (e.g. blanks in Experiment A). Whenever a certain blend attracted one or more males in at least one trap within one biological replicate, possible zero catches of the other biological replicates were included in the analysis. In all graphs, we averaged the number of males of all rotations of one treatment, calculated one percentage value for each of the 3 biological replicates, and plotted the mean percentage of males per trap (i.e. the average of the 3 biological replicates). In the statistical analysis, only raw data (no means or percentages) were used.

## **RESULTS**

Overall, the field tests showed that *S. frugiperda* males of both strains exhibited some geographic variation in their attraction to two different synthetic 4-component-blends (Blends 1 and 2) in North America, the Caribbean and South America (Figure 1). We found a significant effect of geographic region, field crop, and an interaction effect between geographic region  $\times$  strain ( $P < 0.001$ , Table 3).



**FIGURE 1. Attraction of *Spodoptera frugiperda* corn- and rice-strain males to two 4-component blends in different regions.** Bars show the mean percentage of males caught per trap (Blend 1: 100% Z9-14:OAc + 13% Z11-16:OAc + 2% Z7-12:OAc + 1% Z9-12:OAc; Blend 2: 100% Z9-14:OAc + 8% Z11-16:OAc + 4% Z7-12:OAc + 2% Z9-12:OAc) and per biological replicate. There were three biological replicates per field (n=3), except for all fields in North Carolina (n=1) and for rice-strain males in Peru (n=1), where only one replicate caught males. The standard errors in all fields in North Carolina show the variation between rotations in one replicate (n=3), while all other error bars show the variation between biological replicates (n=3). Numbers in brackets/bars represent the total number of males caught, \* =  $P < 0.05$ , \*\* =  $P < 0.01$ , n.s. = not significant. Data from Florida are adapted from Unbehend et al. (2013).

**TABLE 3.** Test statistics on the *Spodoptera frugiperda* male trap catches of different experiments.


	EXPERIMENT A			EXPERIMENT B		
	Wilks	F	P	Wilks	F	P
Country	0.448	12.152	< <b>0.001</b>	0.196	26.711	< <b>0.001</b>
Strain	0.975	1.551	0.216	0.697	9.245	< <b>0.001</b>
Field	0.619	8.345	< <b>0.001</b>	0.907	2.181	0.078
Country:Strain	0.471	11.245	< <b>0.001</b>	0.425	11.333	< <b>0.001</b>
Strain:Field	0.899	1.680	0.104	0.787	5.743	< <b>0.001</b>

Experiments A (Test of two 4-component blends: Blend 1 and Blend 2) and B (Z7-12:OAc dose-response experiment) were analyzed individually using square root transformed data in a MANOVA and a Wilks' Lambda test. Bold P-values show a significant effect of geographic region, strain-identity of males, and/or the field crop, influencing the attraction of fall armyworm males to synthetic pheromone blends. Mean values and standard errors are shown in Figure 1 (Exp. A) and Figure 2 (Exp. B).

Corn-strain males showed a significantly higher attraction to Blend 1 than to Blend 2 in corn fields in Florida, Puerto Rico (field A), Peru and Argentina, but did not show a preference for any of the two blends in corn fields in Canada and Puerto Rico (field B), the mixed habitats in North Carolina or a grass field in Florida (Figure 1). Rice-strain males were equally attracted to Blend 1 and Blend 2 in all cases, with only one exception in a corn field in Florida, where Blend 1 was more attractive than Blend 2. Both strains only differentiated between the two blends when the experiment was conducted in a corn field, where males of both strains were more attracted to Blend 1 than to Blend 2 (Figure 1). Control traps baited with hexane were usually empty in all fields (data not shown), but did trap some males in Argentina (n=2) and in Puerto Rico (n=19 in field A, n=2 in field B). Interestingly, 18 out of the 19 males found in control traps in corn field A in Puerto Rico were caught during the first trap rotation at a time where male density was extremely high (over 50% of all males caught in this experiment were caught at the date of the first rotation, Figure S1).

In experiment B, the Z7-12:OAc dose-response experiment, we found a significant effect of geographic region and strain, as well as an interaction effect between geographic region  $\times$  strain, and field crop  $\times$  strain ( $P < 0.001$ ; Table 3). Interestingly, corn-strain males exhibited a greater differentiation in their response to Z7-12:OAc than rice-strain males (Figure 2), i.e. the highest number of corn-strain males was captured with lures containing 2% Z7-12:OAc, which was significantly different from all other ratios at three field sites in Florida and Puerto Rico. In Peru, lures with 2 and 4% Z7-12:OAc captured equal numbers of corn-strain males (Figure 2A). In Puerto Rico, lures containing no Z7-12:OAc, which is


the critical secondary sex pheromone component of this species, attracted 37 corn-strain males. In contrast to corn-strain males, rice-strain males showed a similar response to different concentrations of Z7-12:OAc in all regions tested, being equally attracted to blends containing 2 or 4% Z7-12:OAc, and showing some level of response to lures with 10% Z7-12:OAc (Figure 2B). No data of rice-strain males could be gathered in Peru, as only corn-strain males were found in this field (Figure 2A).




**FIGURE 2. Attraction of corn-strain (A) and rice-strain (B) males to different doses of Z7-12:OAc.** Bars represent the mean percentage of males caught per trap (0%, 2%, 4%, or 10% Z7-12:OAc + 100% Z9-14:OAc) and per biological replicate (n=3). Different letters above the bars indicate significant differences. Error bars show the variation between biological replicates (n=3). Numbers in brackets/bars represent the total number of males caught. Data from Florida are adapted from (Unbehend et al. 2013).

Testing different doses of Z11-16:OAc (experiment C) added to the minimal blend (i.e. 100% Z9-14:OAc and 2% Z7-12:OAc), revealed that corn-strain males were similarly attracted to binary blends as to three-component blends containing 8, 13 or 18% Z11-16:OAc (Figure 3A). This was true in Peru and Florida. Similarly, addition of Z11-16:OAc did not influence the attraction of rice-strain males in Florida (Figure 3B). In Peru, no rice-strain males were caught in the corn field.

In experiment D, when testing different doses (0, 1, 2%) of E7-12:OAc and Z7-12:OAc added to 100% Z9-14:OAc, we found that corn-strain males from Peru were not attracted to traps baited only with 2% E7-12:OAc added to Z9-14:OAc, but were equally attracted to all other blends tested (Figure 4). In North Carolina, *S. frugiperda* males were equally attracted to synthetic blends to which E7-12:OAc, Z7-12:OAc or E- and Z-7-12:OAc was added (Figure 4). Unfortunately, we were not able to identify the strain-type of any of the trapped males in North Carolina because of DNA degradation of the samples.



**FIGURE 3. Attraction of corn-strain (A) and rice-strain (B) males to different doses of Z11-16:OAc.** Bars represent the mean percentage of males caught per trap (0%, 8%, 13%, or 18% Z11-16:OAc + 100% Z9-14:OAc + 2% Z7-12:OAc) and per biological replicate (n=3). Error bars show the variation between biological replicates (n=3). Numbers in the bars represent the total number of males caught. n.s.=not significant. Data from Florida are adapted from (Unbehend et al. 2013).



**FIGURE 4. Attraction of *Spodoptera frugiperda* males to different doses of E7-12:OAc and Z7-12:OAc in different fields.** Bars show the mean percentage of males caught per trap (100% Z9-14:OAc + a: 2% E7-12:OAc, b: 2% Z7-12:OAc, c: 1% E- and Z-7-12:OAc, d: 2% E- and Z-7-12:OAc) and per biological replicate. The standard errors in all fields in North Carolina (NC) show the variation between rotations in one replicate (n=3), while error bars in Peru show the variation between biological replicates (n=3). Males from Peru are corn-strain males, while males from NC could not be strain-typed but probably belonged to both strains. Numbers in brackets/bars represent the total number of males caught. Different letters above the bars indicate significant differences. n.s.=not significant.

## DISCUSSION

We investigated the variation in attraction of *S. frugiperda* corn- and rice-strain males in North America, the Caribbean and South America, and found a) geographic variation in corn-strain male attraction to synthetic 4-component blends and to different doses of Z7-12:OAc, b) some geographic variation in rice-strain male attraction to different synthetic blends, c) no variation in male attraction to the minor compound Z11-16:OAc, and d) some evidence of geographic variation in response to E7-12:OAc, the compound identified in females from Brazil. Taken together our results indicate that both strains exhibit rather geographic than strain-specific variation in their response to synthetic pheromone lures.

We realize that our field data should be interpreted with some degree of caution for several reasons. Firstly, the composition of the lures of the first experiment (Blend 1 vs. Blend 2) was based on the relative amount of compounds observed in pheromone gland extracts (Groot et al. 2008), and not from airborne collections, which are somewhat different in composition (Tumlinson et al. 1986). We decided to load rubber septa with compound ratios and concentrations that occur in the female gland, assuming a similar emission of volatiles from the pheromone septum as from the female gland surface. Secondly, abiotic conditions may affect the rates at which the pheromone components are released from the lures, as well as the responses of males to a pheromone source (McNeil 1991). As the experiments were

carried out over a large geographic range, it is inevitable that there were intra- and inter-site variability in climatic conditions during the trapping periods. Thirdly, due to the fact that we used mitochondrial markers for strain diagnosis, which are maternally inherited, we were not able to differentiate between homozygote and heterozygote corn- and rice-strain males in our experiments. Therefore, it is possible that also hybrid males were attracted to our lures. Hybridization in the field has been found to be around 16% (Prowell et al. 2004), and several studies have found a bias of inter-strain matings between rice-strain females and corn-strain males, while corn-strain females hybridize less frequently (Nagoshi and Meagher 2003; Prowell et al. 2004; Nagoshi et al. 2006b; Nagoshi 2010). Consequently, the majority of hybrids occurring in nature have a rice-strain mitochondrial COI gene. If our lures attracted hybrid males, they were thus probably typed as rice-strain males. However, despite these limitations we believe our findings support the idea of geographic variability in the sexual communication system of the fall armyworm.

#### ***Geographic variation in corn-strain male responses***

Testing two different 4-component blends revealed that corn-strain males were equally attracted to both blends in Canada and North Carolina, but preferred Blend 1 over Blend 2 in South America, i.e. in Argentina and Peru. Interestingly, Blend 1 mimics the pheromone composition previously reported for corn-strain females in Florida (Groot et al. 2008). However, corn-strain males in Florida and Puerto Rico showed a preference for Blend 1 in one of the two fields tested at each site, but were equally attracted to Blend 1 and 2 in the other field (Figure 1). This differential male attraction between fields could be caused by habitat-specific volatile differences. For example, the two corn fields in Puerto Rico, which were only 4 km apart, were planted with different corn varieties, were in different phenological states during the trapping period, and were treated with different insecticides. This could result in different background odor profiles, which in turn may have influenced the attraction of corn-strain males to the two different 4-component blends used in the first experiment. Previously, it was shown that corn-strain males varied in their attraction to sex pheromone blends in different fields with different host plants in north and south Florida (Meagher and Nagoshi 2013; Unbehend et al. 2013). Furthermore, both strains have some host plant preferences (Pashley 1986; Nagoshi et al. 2006a; Machado et al. 2008), and *S. frugiperda* males show EAG responses to at least 16 different plant volatiles (Malo et al. 2004). Thus, it seems likely that corn-strain males exhibit different responses to female sex pheromones in different habitats, emitting host or non-host volatiles.

The differential attraction of corn-strain males could also be explained by genetic differences between *S. frugiperda* populations from North America, the Caribbean and South America. Population genetic analyses of *S. frugiperda* samples collected throughout the Western Hemisphere generally found no isolation by

distance between populations from different regions (Clark et al. 2007; Martinelli et al. 2007; Belay et al. 2012), which indicates no geographically restricted gene flow probably due to the high migratory ability of *S. frugiperda* (Luginbill 1928; Sparks 1979). However, these analyses did not take into account strain-specific differences and in several cases the strain-type of captured individuals was unknown (Clark et al. 2007; Belay et al. 2012). Genetic studies on populations from Arkansas and Florida showed significant genetic variation among populations, both within and between the two strains (Lewter et al. 2006). Furthermore, corn-strain individuals exhibit different mitochondrial haplotype profiles between populations from a) Florida, Puerto Rico, Georgia and b) Texas, Brazil, Mississippi, Alabama, Louisiana (Nagoshi et al. 2007, 2008a,b, 2010). Thus, genetic differences could play a role in differential attraction of corn-strain males to synthetic blends in different regions.

Besides different host plant volatiles or genetic differences, the responses of corn-strain males could have been influenced by geographically varying environmental factors like temperature or humidity, which are known to influence sexual communication in insects (McNeil 1991). In a previous wind tunnel study, we observed that males of both strains were highly sensitive to changes in temperature or humidity and stopped their response to calling females whenever humidity or temperature was low (Unbehend et al. 2013). Thus, it is possible that males may respond differently to pheromone blends in regions with dry and cold climate compared to (sub) tropical climate zones.

In addition to variation in response to two synthetic 4-component blends, corn-strain males also exhibited significant geographic differences in their attraction to different doses of Z7-12:OAc. In Florida and Puerto Rico, corn-strain males were more attracted to the 2% dose than to other doses tested, but were equally attracted to 2 and 4% Z7-12:OAc in Peru (Figure 2). Furthermore, some corn-strain males from Puerto Rico were attracted to Z9-14:OAc alone, even though Z7-12:OAc has been considered an essential secondary sex pheromone component, without which males are not attracted (Tumlinson et al. 1986; Unbehend et al. 2013). The fact that the response of corn-strain males to Z7-12:OAc varied between regions suggests that females may also vary in their relative amount of Z7-12:OAc across different regions. Although previous data indicated that the production of Z7-12:OAc is under strong stabilizing selection (Unbehend et al. 2013), in light of the data presented here, selection pressures may be different in different regions.

### ***Geographic variation in rice-strain male responses***

In general, rice-strain males were equally attracted to Blend 1 and Blend 2 in different fields in North America, the Caribbean and South America (Figure 1). Although the response of rice-strain males was significantly different to the one of corn-strain males in a corn field in Puerto Rico and Argentina, the attraction of both

strains looks relatively similar in the first experiment. The results of our statistical analysis should be handled with care, as factors like time-dependent density changes within a population, differences between replicates, as well as seasonal/regional differences caused a considerable variation in trap catches (Figure S1). Overall, the results of our first experiment show that both strains respond similarly to the two different 4-components blends that we tested, but exhibit some geographic variation in their response.

Interestingly, rice-strain males from Florida and Puerto Rico showed a broader response spectrum to different doses of Z7-12:OAc compared to corn-strain males (Figure 2). This small but significant strain-specific difference is likely to be important, because Z7-12:OAc is the critical secondary sex pheromone component which is usually crucial for male attraction (Tumlinson et al. 1986; Unbehend et al. 2013). However, it is possible that the broader response spectrum of rice-strain males is due to attraction of hybrid males, if hybrids prefer more extreme blend ratios than the parental pure strains. This should be tested further, although we do not assume that the response of rice-strain males is masked by the presence of hybrid males, because hybridization frequency in nature is relatively low [70]. In conclusion, the results of our experiments suggest that both strains exhibit rather geographic than strain-specific differences in their response, although the response of rice-strain males seems to be broader than that of corn-strain males.

#### ***Male attraction to the minor compound Z11-16:OAc***

Testing the importance of Z11-16:OAc for male attraction showed that corn-strain males from Peru were equally attracted to blends with and without different doses of Z11-16:OAc, similar to the response of corn- and rice-strain males in Florida (Unbehend et al. 2013). In general, addition of Z11-16:OAc to Z9-14:OAc and Z7-12:OAc did also not decrease male attraction in Florida, Peru, Costa Rica and Pennsylvania (Tumlinson et al. 1986; Andrade et al. 2000; Fleischer et al. 2005; Unbehend et al. 2013). These data suggest that Z11-16:OAc is not an essential component for *S. frugiperda* male attraction, which is supported by the observation that *S. frugiperda* males from Mexico did not respond electrophysiologically to Z11-16:OAc (Malo et al. 2004).

#### ***Geographic variation in male attraction to E7-12:OAc***

So far, the E-isomer of the critical secondary sex pheromone component Z7-12:OAc has only been found in *S. frugiperda* females from Brazil, and males from this region responded electrophysiologically to E7-12:OAc and exhibited a higher attraction to binary blends (Z9-14:OAc and Z7-12:OAc) when E7-12:OAc was added (Batista-Pereira et al. 2006). In our trapping experiments, we found that corn-strain males from Peru were not attracted to traps baited only with E7-12:OAc and Z9-14:OAc, but were similarly attracted to all other blends that contained Z7-

12:OAc (Figure 4). Thus, corn-strain males from Peru appear to distinguish between both isomers and need Z7-12:OAc, but not E7-12:OAc, for attraction. This result contrasts our findings in North Carolina, where males did not differentiate between these two isomers. However, while in Peru males captured were corn-strain individuals, males caught in North Carolina could not be strain-typed due to DNA degradation, but probably belonged to both strains. Hence, we currently cannot exclude the possibility that corn- and rice-strain males show differential strain-specific attraction to E- and Z-7-12:OAc. Different isomers of a pheromone component are usually critical for attraction of males and can even lead to speciation, as shown in the two pheromone strains of *Ostrinia nubilalis* (Hübner) (Lassance 2010). Taken together, geographic variation in response to E7-12:OAc seems to exist, but additional experiments are required to evaluate the importance of E7-12:OAc for both strains in different regions.

### ***Implication for pest management***

As the fall armyworm is a serious agricultural pest, an efficient monitoring system for both strains in different habitats and regions would be helpful to detect infestations early and start pest control. To avoid high costs, monitoring via pheromone baited traps requires an effective ‘minimal’ synthetic lure that equally attracts both strains in all habitats and regions. The results of our study showed that only Z9-14:OAc and Z7-12:OAc are usually required to attract *S. frugiperda* in the field, and although both strain exhibit some strain-specific responses towards different doses of Z7-12:OAc, their response ranges also overlap (Figure 2). More precisely, both strains responded equally well to 2% Z7-12:OAc and therefore, we recommend a monitoring blend consisting of 100% Z9-14:OAc (300 µg) and 2% Z7-12:OAc (6 µg).

### ***Conclusions***

Corn- and rice-strain males exhibited some similarities in their attraction to the different blends that we tested, although corn-strain males showed more differentiation in their response than rice-strain males. We found some geographic variation in attraction of corn- and rice-strain males to two synthetic 4- component blends. In contrast, rice-strain males, but not corn-strain males, showed almost no geographic variation in their attraction to different doses of Z7-12:OAc in different regions. One aspect that merits further attention is the possibility that habitat-specific volatiles influence the male response to pheromone blends in different fields. Furthermore, the minor compound Z11-16:OAc does not seem to affect attraction of *S. frugiperda* males, while region-specific differences in the attraction seem to occur to the compound that has only been identified from female glands in Brazil, E7-12:OAc. Overall, the data show some geographic variation in the response of *S. frugiperda* males to pheromone blends. If this variation coincides

with geographic variation in the female pheromone composition, then geographic differentiation between populations could occur.

### *Acknowledgements*

This research was partly funded by the German Science Foundation (grant number GR3627/2-1), the National Science Foundation (award IOS-1052238 and NSF-DEB-1025217), and the Max-Planck-Gesellschaft. We thank Wilson Rivera González from Monsanto and Jose Santiago from 3<sup>rd</sup>Millenium Genetics for their help with the male trapping experiments conducted in Santa Isabel in Puerto Rico, and Sofia Torres, Carmen Urueña, and Alejandro Vera at EEAOC for their help with the trapping experiment in Tucumán, Argentina. Thanks to Gregg Nuessly from the Everglades Research and Education Center in Belle Glade and to Tommy Toms, manager of the Graham Farms in Moore Haven, for their support with the trapping experiments in Florida. We thank Raul Blas, Jorge Tobaru, German Arellano, Antonio Oblitas, Daniela Gálvez, and Jose Meléndez from Universidad Nacional Agraria La Molina (UNALM), and José Galarza from Servicio Nacional de Sanidad Agraria (SENASA), for their assistance with male trapping experiments and DNA extractions conducted on UNALM campus in Peru. Special thanks to Antje Schmaltz, Susanne Neumann, Katja Müller, Sarah Behr, Simone Sörtl, Maximilian Hemmerling, and Susanne Donnerhacke for their assistance in strain-typing trapped males at the MPICE in Jena, Germany.

### REFERENCES

- ACKERMAN, J. D. 1989. Geographic and seasonal variation in fragrance choices and preferences of male Euglossine bees. *Biotropica* 21:340-347.
- ANDRADE, R., C. RODRIGUEZ, A. C. OEHLISCHLAGER. 2000. Optimization of a pheromone lure for *Spodoptera frugiperda* (Smith) in Central America. *Journal of the Brazilian Chemical Society* 11:609-613.
- BAKER, T. C. 2002. Mechanism for saltational shifts in pheromone communication systems. *Proceedings of the National Academy of Sciences of the United States of America* 99:13368-13370.
- BALABAN, E. 1988. Cultural and genetic variation in swamp sparrows (*Melospiza georgiana*) 1. Song variation, genetic variation, and their relationship. *Behaviour* 105:250-291.
- BATISTA-PEREIRA, L. G., K. STEIN, A. F. DE PAULA, J. A. MOREIRA, I. CRUZ, M. D. FIGUEIREDO, J. PERRI, A. G. CORREA. 2006. Isolation, identification, synthesis, and field evaluation of the sex pheromone of the Brazilian population of *Spodoptera frugiperda*. *Journal of Chemical Ecology* 32:1085-1099.
- BELAY, D. K., P. L. CLARK, S. R. SKODA, D. J. ISENHOUR, J. MOLINA-OCHOA, C. GIANNI, J. E. FOSTER. 2012. Spatial genetic variation among *Spodoptera frugiperda* (Lepidoptera: Noctuidae) sampled from the United States, Puerto Rico, Panama, and Argentina. *Annals of the Entomological Society of America* 105:359-367.
- BERNAL, X. E., C. GUARNIZO, H. LUDDECKE. 2005. Geographic variation in advertisement call and genetic structure of *Colostethus palmatus* (Anura, Dendrobatidae) from the Colombian Andes. *Herpetologica* 61:395-408.

- CARDÉ, R. T. AND A. K. MINKS. 1995. Control of moth pests by mating disruption – successes and constraints Annual Review of Entomology 40:559-585.
- CLARK, P. L., J. MOLINA-OCHOA, S. MARTINELLI, S. R. SKODA, D. J. ISENHOUR, D. J. LEE, J. T. KRUMM, J. E. FOSTER. 2007. Population variation of the fall armyworm, *Spodoptera frugiperda*, in the Western Hemisphere. Journal of Insect Science 7:10.
- DELISLE, J. AND J. N. MCNEIL. 1987. The combined effect of photoperiod and temperature on the calling behavior of the true armyworm, *Pseudaletia unipuncta*. Physiological Entomology 12:157-164.
- DELISLE, J. AND L. ROYER. 1994. Changes in pheromone titer of oblique-banded leafroller, *Choristoneura rosaceana*, virgin females as a function of time of day, age, and temperature. Journal of Chemical Ecology 20:45-69.
- DESCOINS, C., J. F. SILVAIN, B. LALANNECASSOU, H. CHERON. 1988. Monitoring of crop pests by sexual trapping of males in the French West Indies and Guyana. Agriculture Ecosystems & Environment 21:53-65.
- EL-SAYED, A. M., D. M. SUCKLING, J. A. BYERS, E. B. JANG, C. H. WEARING. 2009. Potential of 'lure and kill' in long-term pest management and eradication of invasive species. Journal of Economic Entomology 102:815-835.
- FLEISCHER, S. J., C. L. HARDING, P. E. BLOM, J. WHITE, J. GREHAN. 2005. *Spodoptera frugiperda* pheromone lures to avoid nontarget captures of *Leucania phragmatidicola*. Journal of Economic Entomology 98:66-71.
- GEMENO, C., A. F. LUTFALLAH, K. F. HAYNES. 2000. Pheromone blend variation and cross-attraction among populations of the black cutworm moth (Lepidoptera: Noctuidae). Annals of the Entomological Society of America 93:1322-1328.
- GEMENO, C. AND K. F. HAYNES. 2001. Impact of photoperiod on the sexual behavior of the black cutworm moth (Lepidoptera: Noctuidae). Environmental Entomology 30:189-195.
- GONZALEZ-ZUARTH, C. A., A. VALLARINO, C. MACIAS GARCIA. 2011. Female responsiveness underlies the evolution of geographic variation in male courtship between allopatric populations of the fish *Girardinichthys multiradiatus*. Evolutionary Ecology 25:831-843.
- GREEN, J. P., C. ROSE, J. FIELD. 2012. The role of climatic factors in the expression of an intrasexual signal in the paper wasp *Polistes dominulus*. Ethology 118:766-774.
- GROOT, A. T., R. G. SANTANGELO, E. RICCI, C. BROWNIE, F. GOULD. 2007. Differential attraction of *Heliothis subflexa* males to synthetic pheromone lures in eastern US and western Mexico. Journal of Chemical Ecology 33:353-368.
- GROOT, A. T., M. MARR, G. SCHÖFL, S. LORENZ, A. SVATOS, D. G. HECKEL. 2008. Host strain specific sex pheromone variation in *Spodoptera frugiperda*. Frontiers in Zoology 5.
- GROOT, A. T., O. INGLIS, S. BOWDRIDGE, R. G. SANTANGELO, C. BLANCO, J. D. LOPEZ, A. T. VARGAS, F. GOULD, C. SCHAL. 2009. Geographic and temporal variation in moth chemical communication. Evolution 63:1987-2003.
- GROOT, A. T., C. A. BLANCO, A. CLASSEN, O. INGLIS, R. G. SANTANGELO, J. LOPEZ, D. G. HECKEL, C. SCHAL. 2010a. Variation in Sexual Communication of the Tobacco Budworm, *Heliothis virescens*. Southwestern Entomologist 35:367-372.
- GROOT, A. T., A. CLASSEN, H. STAUDACHER, C. SCHAL, D. G. HECKEL. 2010b. Phenotypic plasticity in sexual communication signal of a noctuid moth. Journal of Evolutionary Biology 23:2731-2738.
- GROOT, A. T., M. MARR, D. G. HECKEL, G. SCHÖFL. 2010c. The roles and interactions of reproductive isolation mechanisms in fall armyworm (Lepidoptera: Noctuidae) host strains. Ecological Entomology 35:105-118.
- JONES, R. L. AND A. N. SPARKS. 1979. (Z)-9-tetradecen-1-ol acetate – A secondary sex pheromone of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera, Noctuidae). Journal of Chemical Ecology 5:721-725.


- JUÁREZ, M. L., M. G. MURUA, M. G. GARCIA, M. ONTIVERO, M. T. VERA, J. C. VILARDI, A. T. GROOT, A. P. CASTAGNARO, G. GASTAMINZA, E. WILLINK. 2012. Host association of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) corn and rice strains in Argentina, Brazil, and Paraguay. *Journal of Economic Entomology* 105:573-582.
- KAEFER, I. L., B. M. TSUJI-NISHIKIDO, A. P. LIMA. 2012. Beyond the river: underlying determinants of population acoustic signal variability in Amazonian direct-developing *Allobates* (Anura: Dendrobatoidea). *Acta Ethologica* 15:187-194.
- KAWAZU, K., Y. SUZUKI, Y. YOSHIYASU, E. B. CASTILLON, H. ONO, P. T. VUONG, F. K. HUANG, T. ADATI, T. FUKUMO, S. TATSUKI. 2005. Attraction of *Chaphalocrocis medinalis* (Lepidoptera: Crambidae) males in Southeast Asia to female sex pheromone traps: Field tests in southernmost China, northern Vietnam and southern Philippines with three synthetic pheromone blends regarding geographic variations. *Applied Entomology and Zoology* 40:483-488.
- KUMAR, H. AND K. N. SAXENA. 1986. Certain environmental factors influencing the acoustic communication in the sexual behavior of the leafhopper *Amrasca devastans* (Distant) (Homoptera: Cicadellidae). *Applied Entomology and Zoology* 21:55-62.
- LAMPERT, K. P., A. S. RAND, U. G. MUELLER, M. J. RYAN. 2003. Fine-scale genetic pattern and evidence for sex-biased dispersal in the tungara frog, *Physalaemus pustulosus*. *Molecular Ecology* 12:3325-3334.
- LANDOLT, P. J. AND T. W. PHILLIPS. 1997. Host plant influences on sex pheromone behavior of phytophagous insects. *Annual Review of Entomology* 42:371-391.
- LASSANCE, J. M. 2010. Journey in the *Ostrinia* world: From pest to model in chemical ecology. *Journal of Chemical Ecology* 36:1155-1169.
- LEBLOIS, R., F. ROUSSET, D. TIKEL, C. MORITZ, A. ESTOUP. 2000. Absence of evidence for isolation by distance in an expanding cane toad (*Bufo marinus*) population: an individual-based analysis of microsatellite genotypes. *Molecular Ecology* 9:1905-1909.
- LEWTER, J. A., A. L. SZALANSKI, R. N. NAGOSHI, R. L. MEAGHER, C. B. OWENS, R. G. LUTTRELL. 2006. Genetic variation within and between strains of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Florida Entomologist* 89:63-68.
- LIMA, E. R. AND J. N. MCNEIL. 2009. Female sex pheromones in the host races and hybrids of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Chemoecology* 19:29-36.
- LU, Y. J. AND M. J. ADANG. 1996. Distinguishing fall armyworm (Lepidoptera: Noctuidae) strains using a diagnostic mitochondrial DNA marker. *Florida Entomologist* 79:48-55.
- LUGINBILL, P. 1928. The fall army worm. U S Dept Agric Tech Bull 34:1-92.
- MACDOUGALL-SHACKLETON, E. A. AND S. A. MACDOUGALL-SHACKLETON. 2001. Cultural and genetic evolution in mountain white-crowned sparrows: Song dialects are associated with population structure. *Evolution* 55:2568-2575.
- MACHADO, V., M. WUNDER, V. D. BALDISSERA, J. V. OLIVEIRA, L. M. FIUZA, R. N. NAGOSHI. 2008. Molecular characterization of host strains of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Southern Brazil. *Annals of the Entomological Society of America* 101:619-626.
- MALO, E. A., V. R. CASTREJON-GOMEZ, L. CRUZ-LOPEZ, J. C. ROJAS. 2004. Antennal sensilla and electrophysiological response of male and female *Spodoptera frugiperda* (Lepidoptera: Noctuidae) to conspecific sex pheromone and plant odors. *Annals of the Entomological Society of America* 97:1273-1284.
- MARTINELLI, S., P. L. CLARK, M. I. ZUCCHI, M. C. SILVA-FILHO, J. E. FOSTER, C. OMOTO. 2007. Genetic structure and molecular variability of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) collected in maize and cotton fields in Brazil. *Bulletin of Entomological Research* 97:225-231.

- MCSELFRESH, J. S. AND J. G. MILLAR. 1999. Geographic variation in sex pheromone blend of *Hemileuca electra* from Southern California. *Journal of Chemical Ecology* 25:2505-2525.
- MCNEIL, J. N. 1991. Behavioral ecology of pheromone-mediated communication in moths and its importance in the use of pheromone traps. *Annual Review of Entomology* 36:407-430.
- MEAGHER, R. L. AND M. GALLO-MEAGHER. 2003. Identifying host strains of fall armyworm (Lepidoptera: Noctuidae) in Florida using mitochondrial markers. *Florida Entomologist* 86:450-455.
- MEAGHER, R. L. AND R. N. NAGOSHI. 2013. Attraction of fall armyworm males (Lepidoptera: Noctuidae) to host strain females. *Environmental Entomology* 42:751-757.
- MILLER, D. R., K. E. GIBSON, K. F. RAFFA, S. J. SEYBOLD, S. A. TEALE, D. L. WOOD. 1997. Geographic variation in response of pine engraver, *Ips pini*, and associated species to pheromone, Lanierone. *Journal of Chemical Ecology* 23:2013-2031.
- MITCHELL, E. R., J. H. TUMLINSON, J. N. MCNEIL. 1985. Field evaluation of commercial pheromone formulations and traps using a more effective sex pheromone blend for the fall armyworm (Lepidoptera, Noctuidae). *Journal of Economic Entomology* 78:1364-1369.
- NAGOSHI, R. N. AND R. MEAGHER. 2003. Fall armyworm FR sequences map to sex chromosomes and their distribution in the wild indicate limitations in interstrain mating. *Insect Molecular Biology* 12:453-458.
- NAGOSHI, R. N., R. L. MEAGHER, J. J. ADAMCZYK, S. K. BRAMAN, R. L. BRANDENBURG, G. NUESSELY. 2006a. New restriction fragment length polymorphisms in the cytochrome oxidase I gene facilitate host strain identification of fall armyworm (Lepidoptera: Noctuidae) populations in the southeastern United States. *Journal of Economic Entomology* 99:671-677.
- NAGOSHI, R. N., R. L. MEAGHER, G. NUESSELY, D. G. HALL. 2006b. Effects of fall armyworm (Lepidoptera: Noctuidae) interstrain mating in wild populations. *Environmental Entomology* 35:561-568.
- NAGOSHI, R. N., P. SILVIE, R. L. MEAGHER, J. LOPEZ, V. MACHADOS. 2007. Identification and comparison of fall armyworm (Lepidoptera: Noctuidae) host strains in Brazil, Texas, and Florida. *Annals of the Entomological Society of America* 100:394-402.
- NAGOSHI, R. N., J. S. ARMSTRONG, P. SILVIE, R. L. MEAGHER. 2008a. Structure and distribution of a strain-biased tandem repeat element in fall armyworm (Lepidoptera: Noctuidae) populations in Florida, Texas, and Brazil. *Annals of the Entomological Society of America* 101:1112-1120.
- NAGOSHI, R. N., R. L. MEAGHER, K. FLANDERS, J. GORE, R. JACKSON, J. LOPEZ, J. S. ARMSTRONG, G. D. BUNTIN, C. SANSONE, B. R. LEONARD. 2008b. Using haplotypes to monitor the migration of fall armyworm (Lepidoptera: Noctuidae) corn-strain populations from Texas and Florida. *Journal of Economic Entomology* 101:742-749.
- NAGOSHI, R. N. 2010. The fall armyworm triose phosphate isomerase (*Tpi*) gene as a marker of strain identity and interstrain mating. *Annals of the Entomological Society of America* 103:283-292.
- NAGOSHI, R. N., R. L. MEAGHER, D. A. JENKINS. 2010. Puerto Rico fall armyworm has only limited interactions with those from Brazil or Texas but could have substantial exchanges with Florida populations. *Journal of Economic Entomology* 103:360-367.
- OLVIDO, A. E., P. R. FERNANDES, T. A. MOUSSEAU. 2010. Relative effects of juvenile and adult environmental factors on mate attraction and recognition in the cricket, *Allonemobius socius*. *Journal of Insect Science* 10:1-17.

- PASHLEY, D. P. 1986. Host-associated genetic differentiation in fall armyworm (Lepidoptera, Noctuidae) – a sibling species complex. *Annals of the Entomological Society of America* 79:898-904.
- PASHLEY, D. P. 1988. Current status of fall armyworm host strains. *Florida Entomologist* 71:227-234.
- PASHLEY, D. P. 1989. Host-associated differentiation in armyworms (Lepidoptera: Noctuidae): An allozymic and mitochondrial DNA perspective. Pp. 12. *Electrophoretic studies on agricultural pests*, Systematics Association. Clarendon, Oxford.
- PASHLEY, D. P., A. M. HAMMOND, T. N. HARDY. 1992. Reproductive isolating mechanisms in fall armyworm host strains (Lepidoptera, Noctuidae). *Annals of the Entomological Society of America* 85:400-405.
- PHELAN, P. L. 1992. *Evolution of sex pheromones and the role of asymmetric tracking*. Chapman & Hall, New York & London.
- PODOS, J. AND P. S. WARREN. 2007. The evolution of geographic variation in bird song. Pp. 403-458 in H. J. Brockmann, T. J. Roper, M. Naguib, K. E. Wynne-Edwards, C. Barnard, and J. Mitani, eds. *Advances in the Study of Behavior*, Vol 37.
- PRÖHL, H., R. A. KOSHY, U. MUELLER, A. S. RAND, M. J. RYAN. 2006. Geographic variation of genetic and behavioral traits in northern and southern tungara frogs. *Evolution* 60:1669-1679.
- PROWELL, D. P., M. MCMICHAEL, J. F. SILVAIN. 2004. Multilocus genetic analysis of host use, introgression, and speciation in host strains of fall armyworm (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America* 97:1034-1044.
- R DEVELOPMENT CORE TEAM. 2007. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria
- REDDY, G. V. P. AND A. GUERRERO. 2004. Interactions of insect pheromones and plant semiochemicals. *Trends in Plant Science* 9:253-261.
- ROELOFS, W. L. AND R. T. CARDÉ. 1974. Sex pheromones in the reproductive isolation of lepidopterous species.
- ROESER-MUELLER, K., E. STROHM, M. KALTENPOTH. 2010. Larval rearing temperature influences amount and composition of the marking pheromone of the male beewolf, *Philanthus triangulum*. *Journal of Insect Science* 10.
- ROYER, L. AND J. N. MCNEIL. 1991. Changes in calling behavior and mating success in the European corn borer (*Ostrinia nubilalis*), caused by relative humidity. *Entomologia Experimentalis et Applicata* 61:131-138.
- ROYER, L. AND J. N. MCNEIL. 1993. Effect of relative humidity conditions on responsiveness of European corn borer (*Ostrinia nubilalis*) males to female sex pheromone in a wind tunnel. *Journal of Chemical Ecology* 19:61-69.
- RYAN, M. J., A. S. RAND, L. A. WEIGT. 1996. Allozyme and advertisement call variation in the tungara frog, *Physalaemus pustulosus*. *Evolution* 50:2435-2453.
- SCHÖFL, G., D. G. HECKEL, A. T. GROOT. 2009. Time-shifted reproductive behaviours among fall armyworm (Noctuidae: *Spodoptera frugiperda*) host strains: evidence for differing modes of inheritance. *Journal of Evolutionary Biology* 22:1447-1459.
- SEPPÄ, P. AND A. LAURILA. 1999. Genetic structure of island populations of the anurans *Rana temporaria* and *Bufo bufo*. *Heredity* 82:309-317.
- SLABBEKOORN, H. AND T. B. SMITH. 2002. Bird song, ecology and speciation. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 357:493-503.
- SMADJA, C. AND R. K. BUTLIN. 2009. On the scent of speciation: the chemosensory system and its role in premating isolation. *Heredity* 102:77-97.
- SPARKS, A. N. 1979. Review of the biology of the fall armyworm (Lepidoptera, Noctuidae). *Florida Entomologist* 62:82-87.

- SPARKS, A. N. 1986. Fall armyworm (Lepidoptera, Noctuidae) – potential for area-wide management. *Florida Entomologist* 69:603-614.
- TACK, E. J., D. A. PUTLAND, T. E. ROBSON, A. W. GOLDIZEN. 2005. Geographic variation in vocalisations of Satin Bowerbirds, *Ptilonorhynchus violaceus*, in south-eastern Queensland. *Emu* 105:27-31.
- TOTH, M., C. LÖFSTEDT, B. W. BLAIR, T. CABELLO, A. I. FARAG, B. S. HANSSON, B. G. KOVALEV, S. MAINI, E. A. NESTEROV, I. PAJOR, A. P. SAZONOV, I. V. SHAMSHEV, M. SUBCHEV, G. SZOCS. 1992. Attraction of male turnip moths *Agrotis segetum* (Lepidoptera, Noctuidae) to sex pheromone components and their mixtures at 11 sites in Europe, Asia, and Africa. *Journal of Chemical Ecology* 18:1337-1347.
- TUMLINSON, J. H., E. R. MITCHELL, P. E. A. TEAL, R. R. HEATH, L. J. MENGELKOCH. 1986. Sex pheromone of fall armyworm, *Spodoptera frugiperda* (J.E. Smith) – identification of components critical to attraction in the field. *Journal of Chemical Ecology* 12:1909-1926.
- UNBEHEND, M., S. HÄNNIGER, R. L. MEAGHER, D. G. HECKEL, A. T. GROOT. 2013. Pheromonal divergence between two strains of *Spodoptera frugiperda*. *Journal of Chemical Ecology* 39:364-376.
- WITZGALL, P., P. KIRSCH, A. CORK. 2010. Sex pheromones and their impact on pest management. *Journal of Chemical Ecology* 36:80-100.
- WU, W. Q., C. B. COTTRELL, B. S. HANSSON, C. LÖFSTEDT. 1999. Comparative study of pheromone production and response in Swedish and Zimbabwean populations of turnip moth, *Agrotis segetum*. *Journal of Chemical Ecology* 25:177-196.
- ZHU, J., S. POLAVARAPU, K.-C. PARK, C. GARVEY, D. MAHR, S. NOJIMA, W. ROELOFS, T. BAKER. 2009. Reidentification of pheromone composition of *Sparganothis sulfureana* (Clemens) and evidence of geographic variation in male responses from two US states. *Journal of Asia-Pacific Entomology* 12:247-252.

## SUPPLEMENTARY MATERIAL



**FIGURE S1. Variation in trap catches of *Spodoptera frugiperda* males in different fields and regions.** Graphs show the total number of corn- and rice-strain males that were caught with Blend 1 (100% Z9-14:OAc + 13% Z11-16:OAc + 2% Z7-12:OAc + 1% Z9-12:OAc) in different regions in the first experiment (Figure 1). All traps were rotated three times and all replicates were conducted in the same field, except for North Carolina, where each replicate was conducted in a different field (replicate 1: soybean field, replicate 2: cotton and grass field, replicate 3: soybean and corn field).