



UvA-DARE (Digital Academic Repository)

Genetic mapping of male pheromone response in the European corn borer identifies candidate genes regulating neurogenesis

Koutroumpa, F.A.; Groot, A.T.; Dekker, T.; Heckel, D.G.

DOI

[10.1073/pnas.1610515113](https://doi.org/10.1073/pnas.1610515113)

Publication date

2016

Document Version

Final published version

Published in

Proceedings of the National Academy of Sciences of the United States of America

License

CC BY-NC-ND

[Link to publication](#)

Citation for published version (APA):

Koutroumpa, F. A., Groot, A. T., Dekker, T., & Heckel, D. G. (2016). Genetic mapping of male pheromone response in the European corn borer identifies candidate genes regulating neurogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, 113(42), E6401-E6408. <https://doi.org/10.1073/pnas.1610515113>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Genetic mapping of male pheromone response in the European corn borer identifies candidate genes regulating neurogenesis

Fotini A. Koutroumpa^{a,b,1}, Astrid T. Groot^{a,b}, Teun Dekker^c, and David G. Heckel^b

^aInstitute for Biodiversity and Ecosystems Dynamics, University of Amsterdam, 1098 XH Amsterdam, The Netherlands; ^bDepartment of Entomology, Max Planck Institute for Chemical Ecology, D-07745 Jena, Germany; and ^cDivision of Chemical Ecology, Department of Plant Protection Biology, Swedish University of Agricultural Sciences, 23053 Alnarp, Sweden

Edited by John G. Hildebrand, University of Arizona, Tucson, AZ, and approved August 26, 2016 (received for review July 8, 2016)

The sexual pheromone communication system of moths is a model system for studies of the evolution of reproductive isolation. Females emit a blend of volatile components that males detect at a distance. Species differences in female pheromone composition and male response directly reinforce reproductive isolation in nature, because even slight variations in the species-specific pheromone blend are usually rejected by the male. The mechanisms by which a new pheromone signal–response system could evolve are enigmatic, because any deviation from the optimally attractive blend should be selected against. Here we investigate the genetic mechanisms enabling a switch in male response. We used a quantitative trait locus-mapping approach to identify the genetic basis of male response in the two pheromone races of the European corn borer, *Ostrinia nubilalis*. Male response to a 99:1 vs. a 3:97 ratio of the E and Z isomers of the female pheromone is governed by a single, sex-linked locus. We found that the chromosomal region most tightly linked to this locus contains genes involved in neurogenesis but, in accordance with an earlier study, does not contain the odorant receptors expressed in the male antenna that detect the pheromone. This finding implies that differences in the development of neuronal pathways conveying information from the antenna, not differences in pheromone detection by the odorant receptors, are primarily responsible for the behavioral response differences among the males in this system. Comparison with other moth species reveals a previously unexplored mechanism by which male pheromone response can change in evolution.

Ostrinia nubilalis | Z chromosome | pheromone response | sexual communication | QTL analysis

Few communication systems in the natural world can rival the sensitivity and elegance of moth sexual pheromones, which reliably bring together males and females of the same species for reproduction (1). The signal is a volatile blend of fatty-acid-derived compounds with various modifications, synthesized by the specialized pheromone gland of the female and released into the air. The detection system is the male antenna, bearing sensillar hairs with specialized neurons that express pheromone receptors that are activated by the binding of individual pheromone components. Species-specific blends activate a behavioral response in which the male flies upwind, following the odor plume, and eventually finds the female (2). Subtle differences in the pheromone compositions of closely related species are sufficient to block this attraction and to maintain reproductive isolation in nature (3, 4). The manifold variety of pheromone systems has been suggested as one of the factors promoting the evolution of the high species diversity of Lepidoptera (4, 5).

However, the origin of such variety poses an evolutionary dilemma. In such a finely tuned communication system, any deviation by the female from the optimally attractive pheromone blend or any preference by the male for an atypical blend would decrease mating success and should be selected against, maintaining the status quo (6–9). There is empirical evidence for such selection against deviance

in moths (4, 10, 11). A coordinated change of both signal and response would seem necessary to overcome this problem but is hard to imagine because genes affecting female pheromone production and male response are different and reside on separate chromosomes (12–17). Moreover, despite significant advances in the biochemistry of pheromone synthesis and the physiology of pheromone detection, the identity and mode of action of most genes that shape these traits are unknown. Knowledge of the underlying genetic basis is a requirement for any theory invoking genetic mechanisms for evolutionary change.

The European corn borer (ECB), *Ostrinia nubilalis* Hübner (Lepidoptera: Crambidae), is a well-studied model system of sexual communication in moths. This species consists of two strains, denoted “E” and “Z,” which use different ratios of the same two pheromone compounds, (E)-11-tetradecenyl acetate and (Z)-11-tetradecenyl acetate (henceforth “E11-14:OAc” and “Z11-14:OAc”). E-strain females produce E11-14:OAc and Z11-14:OAc in a ratio of 99:1, and Z-strain females produce these compounds in a nearly opposite ratio of 3:97 (18). E- and Z-strain males prefer the ratios produced by their respective females (19, 20). This preference leads to reproductive isolation (21), although hybrids are occasionally found in nature (22–25). In the laboratory, hybrids are readily formed by matings; hybrid females produce an intermediate compound ratio of 65:35 (22), whereas hybrid males show a response centered around a 50:50 ratio (18, 26).

Significance

Most male moths find their mates by following species-specific pheromones released by females. Despite the importance of pheromone communication in reproductive isolation, much is still unknown about its genetic basis. We investigated male responses in the two pheromone races of the European corn borer. A reasonable hypothesis, that males of the two races differ in the genes encoding the receptor proteins that respond to the two pheromone components, was previously rejected without a convincing alternative. We found that instead male choice was correlated with genes affecting growth and differentiation of the nerve cells that may contain these receptors. This unexpected finding resolves the dilemma and points to another layer of complexity in the evolution of sexual pheromone communication systems.

Author contributions: A.T.G., T.D., and D.G.H. designed research; F.A.K., A.T.G., T.D., and D.G.H. performed research; F.A.K., A.T.G., and D.G.H. analyzed data; and F.A.K. and D.G.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. Email: fotini.koutroumpa@gmail.com.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1610515113/-DCSupplemental.

The genes controlling these strain differences have been mapped to different chromosomes by interstrain crosses (13). The gene causing variation in female pheromone production is autosomal and encodes a fatty-acyl reductase (27, 28). The main gene controlling the male behavioral response difference, *Resp*, is sex-linked on the Z chromosome (13, 18) (males are ZZ and females ZW in the standard chromosomal nomenclature). The identities of the gene(s) controlling strain variation in male behavioral response in *Ostrinia* are still unknown.

So far, the most promising candidates for the control of male preference in moths are the genes encoding olfactory receptor proteins (ORs) that bind to specific pheromone components (29–33). In strong support of this idea, Gould et al. (14) showed that a cluster of OR genes expressed in the antenna is very tightly linked to an autosomal gene controlling the interspecific difference in male behavioral response among *Heliothis virescens* and *Heliothis subflexa*. Moreover, single-cell recordings showed that species-specific spike-amplitude responses to individual pheromone components were also linked to the OR cluster (14). In *O. nubilalis*, five of the seven pheromone ORs are located within a large cluster on the Z chromosome (34). The pheromone ORs show similar spatial antennal expression patterns in males of both strains (35), but *OnubOR6*, which responds to Z11-14:OAc (36), is expressed at a higher level in Z-strain males (35), and *OnubOR4*, which binds to E11-14:OAc (32), is expressed at a higher level in E-strain males; expression levels of both *OnubOR6* and *OnubOR4* are intermediate in hybrids (35). Three Z-linked genes—*OnubOR4*, *OnubOR6*, and *OnubOR1*—show more sequence differences among the E and Z strains than do the two autosomal ORs; however, linkage mapping of the OR cluster on the Z chromosome placed it ~20 cM distant from the *Resp* gene (31). Although mapping of *Resp* was based on phenotype data from 78 males (13), it is difficult to judge the significance of this result, because the accuracy of the behavioral scoring is unknown. Furthermore, independent replicate backcrosses using the E and Z strains by another group showed a high variability in recombination rates among Z-linked genetic markers (37).

To identify the position of the *Resp* locus more precisely, we conducted a new quantitative trait locus (QTL) analysis, using 470 male progeny from a series of backcrosses. We confirmed that *Resp* is separated from the OR cluster (31) with a more precise estimate of 15 cM. Moreover, we identified several genes that are much more tightly linked to *Resp* and which point to a site of action in establishing the connections between olfactory sensory neurons (OSNs) and the brain or deeper within the brain itself. This finding opens up a set of candidate genes acting at a deeper mechanistic level that could be responsible for variations in male response.

Results

Crossing Design. Hybrid crosses were set up between the Z and E strains, and resulting male progeny were backcrossed to Z-strain females for seven consecutive generations. In each generation we scored behavioral responses to Z, E, and hybrid pheromone blends using artificial lures for 50–150 males (fewer for the third and fourth backcrossed generations). In each generation, one of the scored males with a hybrid phenotype was chosen for backcrossing to a Z-strain female to produce the next generation. In total, 649 males were phenotyped, using the wind tunnel setup described in *Methods*. After the last generation of backcrossing, DNA was isolated from all 649 phenotyped males, and genotypes were scored at nine sex-linked markers (*kettin*, five ORs, *ldh*, *bgi012356*, and *bgi03892*). This preliminary scoring showed a loss of genetic polymorphism correlated with behavioral response in the last two backcross generations, probably because of the inadvertent selection of a homozygous male for continuation of the backcrossing. Therefore, only 470 males from the first five generations could be used for final mapping of 21 sex-linked genes and the

QTL analysis. We also genotyped 143 females (~20 per generation) to aid in map construction.

Behavioral Measurements. The relative intensity of individual male behavioral responses to each of the pheromone blends—Z, hybrid (H), and E—was summarized using a quantitative score that assigned numerical weights to two key behaviors previously shown to be correlated with directed flight toward the pheromone source (38). These behaviors are wing-fanning, a rapid vibration of the wings while remaining in a stationary position, and extrusion of the hair pencil scales at the tip of the abdomen, which release the male pheromone during courtship at close range (Fig. 1). Wing-fanning could be strong and continuous or weak and intermittent (on-off), and hair pencil extrusion could be present or absent. The latency time, i.e., the number of seconds between presentation of the stimulus and the initiation of wing-fanning, was also factored in. An initial set of numerical weights for these behaviors was first chosen arbitrarily, such that the scored response to a given pheromone blend increased with the occurrence and intensity of wing-fanning, increased more with a shorter latency time, and increased even more with the extrusion of hair pencil scales. Results from the three lures were combined by subtracting the H and E lure responses from the Z lure response, so that Z-strain males would have the highest score, because these are male-informative backcross progeny from a pure Z-strain mother. Phenotypic scores then were calculated for each male (see Fig. S1A for details). After an initial QTL analysis, the parameter set of numerical weights was optimized using a discriminant function approach, to discriminate more precisely between the responses of *Resp* locus homozygotes vs. *Resp* locus heterozygotes (Fig. S1B).

Linkage Maps. To examine the contribution of each chromosome to male behavioral response, a preliminary map using amplified fragment length polymorphism (AFLP) markers covering all 31 chromosomes was constructed, using data from the 120 progeny of the first backcross generation (Fig. S2). We then mapped an additional 21 genes on the Z chromosome using the progeny of all backcrosses. These genes were chosen based on their Z-chromosomal location in *Bombyx mori*, and their orthologs in *O. nubilalis* were PCR-amplified using degenerate primers designed from multispecies sequence alignments (Table S1). Overall, the gene order in *O. nubilalis* between *kettin* at one end and *paraplegin* near the other end was the same as in *B. mori*. However, we did not map any genes surrounding *Tpi*, where Wadsworth et al. (39) have detected a region of low recombination that probably corresponds to an inversion. Additionally, the order of the two end markers *pdp1* and *bgi03892* was inverted (Fig. 2). An inversion between *bgi03892* and another marker was also detected in *O. nubilalis* by Kroemer et al. (Fig. 2) (37). Finally, we found that the Z-linked marker




Fig. 1. Quantitative scoring of male behavioral response indicators.

bgi002071 in *B. mori* was autosomal in *O. nubilalis*, as was also shown for marker *G23* just above *kettin* by Kroemer et al. (37). Thus, the segment of the Z chromosome above *kettin* in *B. mori* appears to be autosomal in *O. nubilalis* and therefore segregates independently from the behavioral response trait.

QTL Analyses. A QTL analysis using 198 AFLP markers that were mapped to the 31 chromosomes spread over 45 linkage groups showed that the Z chromosome was the only one with a significant effect on the male behavioral response, (Fig. S2). This result is in agreement with the findings of Dopman et al. (13), who used a qualitative flight assay to map the major *Resp* gene to the Z chromosome in a backcross to the E strain. Our quantitative measurement of male behavior additionally permits an estimate of the contribution of minor genes. In comparison with the Z chromosome with a significant logarithm of odds (LOD) score of 5.0, explaining 25% of the variance in behavioral score, the chromosome with the next largest effect is AFLP linkage group 31, explaining 11% of the variance, but this effect was below the $P < 0.05$ significance threshold (Fig. S2). No other linkage group has an LOD score higher than 1.5 (Fig. S2).

We then conducted a more extensive QTL analysis using 12 genes on the Z chromosome, using data from all 470 phenotyped and genotyped males. Interval mapping produced a single broad peak of the log-likelihood function at position 84.2 cM with an LOD score of 8.7 covering an interval of 36 cM, containing one or more QTLs for the behavioral response. The OR cluster occurred near one edge of this broad interval (Fig. S3A). We used this large interval to optimize the behavioral scoring function (*SI Materials*

and Methods), which increased the LOD score at the peak to 27.8 cM and narrowed its width to 12 cM (Fig. S3B). Finally, for the final QTL analysis we added genotype scores for an additional 12 markers, 6 of which were located within the 12-cM interval (Fig. 3).

The resulting log-likelihood function showed a single narrow peak at position 85.3 cM near the marker *kon-tiki* with an LOD score of 29.6. The 2-LOD confidence interval in which the LOD score exceeds 27.6 is 8 cM wide and covers positions 83 cM to 91 cM. The likelihood that *Resp* occurs at the peak is more than 100 times greater than the likelihood that it occurs outside the 2-LOD confidence interval. The log-likelihood function also showed a second peak at 108 cM with a maximum LOD score of 20.5, flanked by two markers (*paraplegin* and *pdp1*). This peak is approximately 10^7 times less likely to be the *Resp* locus than any position within the 2-LOD confidence interval, but it could represent a second QTL with a smaller effect on the behavioral score. To examine this possibility, we conducted fixed-effects two-way ANOVA using genotypes at the markers nearest the major and minor peaks as factors (Fig. S4). The major peak effect was highly significant ($P < 0.0001$); the minor peak effect and interaction effects were not ($P = 0.46$ and $P = 0.37$, respectively for the marker to the left of the minor peak and $P = 0.49$ and $P = 0.18$ for the marker to the right of the minor peak), indicating that any possible contribution of the minor peak effect and interaction effect to the overall variance of the behavioral score is less than 10% of the contribution of the major QTL.

Evaluation of Candidate Genes for the Behavioral Response. We next evaluated the OR cluster as a candidate for the *Resp* locus. The cluster maps to position 72 cM; in scoring *OnubOR5*, *OnubOR4*,




Fig. 2. Comparison of linkage maps of *O. nubilalis* and linkage and physical maps for *B. mori*. Genes common to all maps are shown in dark gray, and gene inversions are shown in light gray. (A–C) Compare our results with previously published results on *O. nubilalis*. (A) Adapted from Dopman et al. (13). (B) Adapted from Kroemer et al. (37). (C) Our results. (D) *B. mori* sex chromosome gene linkage. Z is the sex chromosome. Thick lines represent the chromosomes, and the dashed line shows the direction in which the chromosome continues.




Fig. 3. Final QTL analysis with 21 sex-linked markers and optimized behavioral scores. The male response locus (*Resp*) is predicted to be inside the gray zone. Every locus outside this zone is at least 100 times less likely to be the *Resp* locus. The distance between the different loci is measured in centimorgans, and the LOD scores obtained for the ORs and the second peak are shown in boldface type.

and *OnubOR6*, we did not detect any crossovers within the cluster (Figs. 2 and 3). The OR genes occur at 85.3 cM, 13.3 cM away from the larger QTL peak, somewhat closer than the distance of 20 cM found by Lassance et al. (31) between the ORs and the *Resp* locus. The OR cluster is even further away (36 cM) from the smaller peak at 108 cM. The LOD score of the OR cluster is 20.5, making it very unlikely to be the *Resp* locus. The peak at 85.3 cM is 10^9 times more likely, and any location within the 2-LOD confidence interval is at least 10^7 times more likely to be the *Resp* locus than is the OR cluster. The slight hump in the LOD score at the OR cluster (Fig. 3) might indicate a smaller contribution to the behavioral response. To test the possibility that the ORs represent a third, minor QTL, we conducted a two-way ANOVA using the marker genotypes at the QTL peak and the ORs as factors (Fig. S5). As before, the main effect of the peak was highly significant ($P < 0.0001$), but neither the main effect of the OR genotype ($P = 0.84$) nor the interaction effect ($P = 0.13$) was significant. Moreover, the nonsignificant effect of the OR genotype depended on the genotype of the markers near the peak. Thus, sequence variation at the OR cluster makes no consistent contribution to male pheromone preference.

The 2-LOD confidence interval around the major peak contains two of the marker genes, *kon-tiki* and *bric-a-brac* (*bab*). Comparing our *O. nubilalis* map with the *B. mori* and *Danaus plexippus* genomes, a total of seven genes have been mapped between *ldh* and *bgi012356* (Fig. 3). Interestingly, six of these genes—*highwire*, *pentaxin*, *tol*, *kon-tiki*, *bab*, and *CCR4-not*—seem to be involved in neurogenesis (Table 1). These genes seem to control neuronal axonal growth and orientation processes specifically in insects and other animals (see references in Table 1). Furthermore, the genome comparisons to identify other candidates for *Resp* showed a total of 12 genes (including *kon-tiki* and *bab*) located within the *Resp* region. Eight have one or more identifiable protein domains, six have homologs in *Drosophila* with some functional information,

and three are not similar to any protein of known function. Although none has an obvious connection to pheromone sensory physiology, several are implicated in neurogenesis; we discuss these possible roles below.

Discussion

By fine-scale mapping of the male response in *O. nubilalis*, we found candidate genes that had not been implicated in previous studies of male pheromone perception in moths. Moreover, confirming previous work (31), we ruled out variation in other candidate genes, i.e., the ORs that detect pheromone components and have been implicated in studies of other species. We will explain the justification for our approach in analyzing behavior, point out some aspects of our mapping methods, reconsider the role of the pheromone receptors, and discuss the possible roles of genes in the chromosomal region we have identified.

In mapping the genes underlying the male behavioral response, we deliberately chose to measure traits different from but correlated with the final, evolutionarily relevant outcome: a mating with a female of the same strain. This outcome has traditionally been measured by observing male behavior and flight toward a female or an artificial pheromone dispenser at the upwind end of a wind tunnel. The ultimate criterion of response is considered to be touching and attempting to copulate with the pheromone source, and this criterion has been used as a qualitative, dichotomous trait in previous mapping studies (13, 18). This approach has been justified by the observation that E-strain males may exhibit earlier signs of attention to the Z lure and the opposite applies to Z-strain males; such signs are unreliable predictors of the final outcome (38). However, recent studies have shown that early responses by Z-strain males can lock them in to subsequent behaviors that are maintained even if the pheromone lure is experimentally switched from Z to E while the moth is in midflight (40). Thus, variation in the early responses also may be a relevant indicator of genetic differences between the Z and E strains. Therefore, we scored male response as a quantitative trait by observing the preflight responses of males to different pheromone blends. This reduction in individual assay time greatly increased the number of males that could be scored against all three pheromone blends over the generations. Even though 68% of the variance in this quantitative trait was nongenetic or was caused by minor genetic factors, QTL analysis could define an interval on the Z chromosome that accounted for 32% of the variance. Our methods enabled the quantitative assessment of previously suspected candidate genes and identified candidates that had not previously been considered; this assessment would have not been possible if the response had been scored as a dichotomous trait. Moreover, our method avoids inaccuracies in fine-scale genetic mapping caused by errors in scoring male response as a dichotomous trait, which are inevitably confounded with recombination (see the discussion of mapping strategies in *SI Materials and Methods*).

The exclusion of the cluster of ORs from the QTL region means that sequence variation in those genes, whether in the coding sequence or in noncoding regions that affect their expression patterns, cannot be directly responsible for the difference in male response. However the possibility remains that differences in OR expression or function may be controlled by a *trans*-acting transcription factor located within the QTL region. We previously found no qualitative difference in OR expression: Both strains have the same number of sensilla and neurons housing them (35). However, we did find quantitative differences in the expression of *OnubOR4* and *OnubOR6* (35). These differences could be caused by a *trans*-acting transcription factor or could be affected by changes in neuron size and number. Nevertheless, the genetic factor underlying male response in the *Ostrinia* pheromone strains must be different from that of *H. virescens* and *H. subflexa*, in which the male behavioral

Table 1. Description of genes near the *resp* locus region

Resp region	Full name (abbreviation)	Homologs/synonyms	BLASTp best hit		Function	Expression	Ref. or source
			Accession number	Organism			
Out	<i>Highwire (hiw)*</i>	PHR protein family: Myc (PAM) [†] Rpm-1 gene [‡] Esrom [§] Lectins	AF262977	*	Ubiquitin ligase, negative regulator of synaptic growth Presynaptically: synaptic growth and axon guidance Postsynaptically: endocytosis of glutamate receptors	Muscle neurons; CNS; retinal cone	(75–78)
Out	<i>Pentraxin (pent)*</i>		U18772	§	Innate immunity Synapse formation and remodeling, neural crest cell migration	Sensory neurons Brain	(79)
Out	<i>terribly reduced optic lobes(trol)*</i>	Heparan sulfate Proteoglycan [¶] Perlecan ^{†,‡}	XP_014364776.1	¶	Activates neuroblasts' adhesion, growth migration and differentiation Signal transduction Regulates neuromuscular junction	Larval brain, imaginal discs, fat body, muscles; adult gonads	(43, 80, 81)
In	<i>Aldose-1 epimerase[#]</i>	Galactose mutarotase [†]	XP_004933180.1	#		Cytoplasm; testis	(82)
In	<i>LOC101738019[#]</i>	KGM_08728	XP_004933181.1	#	Uncharacterized protein	unknown	GenBank
In	<i>Cralbp*</i>	KGM_088417 α -Tocopherol transfer protein	XP_013184856.1	**	Cellular retinaldehyde binding protein α -Tocopherol transfer protein	Fat body, gut: hemocytes	(83)
In	<i>kon-tiki (kon)*</i>	NG2 proteoglycans: Chondroitin sulfate proteoglycan ^{**} Perdido* NG2/MCSP ^{†,§}	XP_013184902.1	**	Muscle development, i.e., filopodium assembly and orientation targeting skeletal positions, directed myotube migration Target-derived signal initiating stable connection Neurogenesis i.e., cell migration and differentiation	Myotube tips: Embryonic founder; Myoblasts: CNS	(51, 52)
In	<i>Tudor domain*</i>	RING finger protein 17	XP_013140855.1	††	Germline development		(53)
In	<i>BAP18[¶]</i>	SANT domain	XP_004933184.1	#	Chromatin modification DNA binding		(54)
In	LIM homeobox	Transcription factor	XP_013184855.1	**	Tissue patterning and differentiation Neuronal patterning		(55)
In	<i>bgi12353A</i>	LOC101738771 [#]	EHJ68261.1		Uncharacterized protein	Testis	GenBank
In	<i>bgi12353B</i>	LOC101738910 [#] KGM_07611 [¶]	XP_013184860.1	**	Uncharacterized protein	Testis	GenBank
In	<i>Archipelago (ago)</i>	F-box protein	XP_013184917.1	**	Cyclin binding Ubiquitin–protein transferase activity: negative regulator of cell growth, including axon guidance	Imaginal tissues: eye; imaginal disk: photoreceptor cell	(56, 57)
In	<i>Bric à brac (bab)*</i>	BTB POZ domain	XP_014364829.1	¶	Leg and antenna segmentation Eye–antennal disk morphogenesis and imaginal disk-derived leg morphogenesis Sex differentiation, i.e., abdominal color sexual dimorphism in <i>D. melanogaster</i>	Leg and antenna imaginal disks; ovaries	(58–60)
Out	<i>CCR4-NOT*</i>		XP_013184897.1	**	Female gonad development Regulation of synaptic growth of the neuromuscular junction Inhibition of miRNA degradation		(61, 62)

Drosophila melanogaster*.†*Homo sapiens*.‡*Caenorhabditis elegans*.§*Mus musculus*.¶*Papilio machaon*.#*Bombyx mori*.||*Danaus plexippus*.*Amyelois transitella*.††*Papilio polytes*.

responses and most of the neurophysiological responses to pheromone components cosegregated with the autosomal OR cluster (14).

The candidate genes in the *Resp* QTL region (Table 1) suggest that other factors affect the structure, function, or connectivity of the OSNs or other neurons in the antennal lobe or elsewhere in the brain. Three candidates are similar to genes of unknown function in the sequenced genomes of other lepidopteran species (Table 1). Two other candidates could potentially play a role in gene regulation: One has a Tudor domain also found in proteins that bind to methylated histones (41), and the other has a SANT (switching-defective protein 3, adaptor 2, nuclear receptor co-repressor, and transcription factor IIIB) domain involved in histone modification (42). Two candidates, aldose epimerase and a lipid-binding α -tocopherol transfer protein, have domains not necessarily specific to the nervous system. Four genes within the QTL region and two just outside it are similar to genes with roles in neurogenesis. Of the latter, *trol* (*terribly reduced optic lobes*, also called *perlecan*) is a large multidomain heparan sulfate proteoglycan in the extracellular matrix with Laminin G domains that binds to and stores other signaling molecules controlling neuroblast proliferation (43). On the other side of the QTL region is *CCR4-Not*, coding for the catalytic subunit of the deadenylase complex (44, 45), which interacts with the conserved P body component HPat/Pat1 to regulate synaptic terminal growth in the *Drosophila* neuromuscular junction (46).

We consider that four genes mapping within the QTL region are most likely to play a relevant role in neurogenesis. In *Drosophila melanogaster*, *kon-tiki* encodes a transmembrane protein that promotes cell migration in muscle development (47, 48). Similar mechanisms have been proposed to generate directed migration and target recognition of myotubes as well as neuronal axon and dendrite growth toward their synaptic partners (49). The second likely candidate is a gene with LIM and homeobox domains characteristic of many transcription factors that has also been described as a neural-patterning gene in several organisms [i.e., *Drosophila*, mice, and *Caenorhabditis elegans* (50)]. The third likely candidate is *archipelago* (*ago*), which encodes an F-box protein and part of a ubiquitin ligase complex that interacts with the Notch signaling pathway and suppresses tissue growth in flies and tumor development in vertebrates (51, 52). Mutations in the fourth likely candidate, *bab*, have many consequences, including disordered arrangements of eggs in the ovaries in *D. melanogaster* (53). In both antennae and tarsi, *bab* causes distal segment fusion (54). The protein encoded by *bab* possesses a BTB domain also found in the *Drosophila* genes *tramtrack* and *Broad-Complex* (55). Moreover, *bab* was recently found to interact with other transcription factors in a network that patterns the developing olfactory tissue in *D. melanogaster* (56).

In addition to differing in their behavioral response to female pheromones, E- and Z-strain males also differ in the connections between the antennal OSNs and the antennal lobe of the brain. In the Z strain, the OSNs that respond to Z11-14:OAc project their axons onto the large, medial glomerulus of the antennal lobe, and the OSNs responding to E11-14:OAc project onto the smaller, lateral glomerulus (57). In the E strain, these connections are reversed (57). In F₁ hybrids, the connections are similar to those in the E strain, but the lateral and medial glomeruli are more similar in size, and glomerular size appears to be Z-linked (58). A Z-linked mutant in *B. mori* shows a similar rewiring of neuronal connections. In wild-type *B. mori* males, OSNs expressing the bombykol receptor BmOR1 target their axons to the larger toroid glomerulus, whereas OSNs expressing the bombykal receptor BmOR3 send their axons to the smaller cumulus (59, 60). In null mutants of the Z-linked *acj6* gene, BmOR1-expressing OSNs are rare or absent, and BmOR3-expressing OSNs project instead to the toroid (61). Mutant males are attracted not by the main pheromone component bombykol but instead by the minor component bombykal. We have con-

firmed that *acj6* is also Z-linked in *O. nubilalis*, but it segregates independently from the QTL interval. Furthermore, *acj6* maps distal to *kettin* on the Z chromosome of *B. mori*, far from the QTL interval (Fig. 2). Thus, despite intriguing similarities in the phenotype caused by mutation of *acj6* in *Bombyx*, *acj6* does not seem to be responsible for male response in *O. nubilalis*.

Nevertheless, it is possible that one of the genes in the QTL interval could have a similar effect on axonal targeting. In *Drosophila* (unlike mammals) the expression of a given OR in a sensory neuron is not directly involved in glomerular targeting (62). Instead, upstream-acting transcription factors involved in neuronal differentiation separately specify both the downstream expression of a particular OR (63) and the axonal guidance of that neuron to its cognate glomerulus in the brain. In *O. nubilalis*, both strains possess the same cluster of ORs for pheromone detection. One of the newly identified candidate genes could control the expression of specific ORs in different OSNs [as inferred by Koutroumpa et al. (35)] or the targeting of these OSNs to the antennal lobe, or both processes. A third possibility is a change in the antennal lobe itself, governing which OSNs will connect to which glomeruli. Such a change is suggested by the results of transplanting antennae of E-strain males onto Z-strain males: The chimeric males responded only to the Z blend (64). In this example, the Z genotype of the recipient bearing the antennal lobe had a greater influence than the E genotype of the OSNs in the transplanted antennae.

One other extensively studied trait is the spike amplitude of OSNs responding to the E or Z component in the two strains. In E-strain males, the E-responding OSN has bigger spike amplitude than the Z-responding OSN. In Z-strain males, the magnitudes of the spike amplitudes are reversed, and in hybrids they are more nearly equal. Spike amplitude is correlated with the diameter of the dendrite projecting into the sensillar lymph (65). Early studies found that spike amplitude was autosomally inherited (66), not sex-linked (18), and was not on the autosome controlling female pheromone production (12). Later studies found that spike amplitude was not correlated with male behavioral response (67) but also may have a minor Z-linked component (15). There is a strain difference in electroantennogram responses that also appears to be Z-linked (58). Further discussion on spike amplitude measurements and the interpretation of these results in the light of sex-linkage can be found in Koutroumpa et al. (35). We did not score spike amplitudes or whole-antennal responses in this study because of the prohibitive workload of neurophysiological preparations of each male after the behavioral assays, but further mapping of these traits would be worthwhile.

In conclusion, our QTL-mapping approach has revealed several genes involved in neurogenesis that may account for the differences in male behavioral response between pheromone strains in *O. nubilalis*. In addition to the previously established importance of changes in the ORs, as supported by phylogenetic analysis, functional expression, and genetics, other changes can shift preference without any change in the ORs themselves. Further advances will result from the application of QTL mapping to other recently diverged species pairs or species with pheromone races, as well as from functional studies of the candidate genes we have identified.

Materials and Methods

Insects. Laboratory colonies of ECB Z and E strains were used. The Z-strain colony derived from cornfield-collected adults in Kéty town, county of Tolna, Hungary in 2004. The E-strain colony was established from larvae extracted from maize stems collected by Magda Rak-Cizej of the Agriculture and Forestry Institute, Novo Mesto, Slovenia. The purity of the strains was monitored by GC analysis of female pheromone production following the protocol of Kárpáti et al. (68), and male response was evaluated with an electroantennogram (58). All insects were reared on a semiartificial diet (69)

until pupation. Adults were fed a 5% honey–water solution throughout their adult lives. All animals were kept at 25 °C, relative humidity 70% under an 18-h/6-h light/dark photoperiod. The day of emergence was considered day 0.

Backcrosses. All crosses were single-pair crosses. Because there is crossing-over in male Lepidoptera (70), we conducted male-informative backcrosses to generate recombination in the Z chromosome. Single-pair matings were set up hybridizing pure Z-strain females with E-strain males (ZE hybrid) and E-strain females with Z-strain males (EZ hybrid). The F₁ hybrid males were backcrossed to pure Z-strain females. We selected one out of 30 hybrid families, namely family EZ2, because it was highly fecund. Sons of the EZ2 family that showed hybrid behavior in the wind tunnel bioassays were backcrossed with pure Z-strain females, giving rise to the first backcross generation (BC1: ZE22). Two BC1 families were continued to second-generation backcrosses, i.e., ZE22-4 and ZE22-7 families. Because only the ZE22-7 family generated enough offspring, we continued the backcrossing procedure with this family for six additional generations (BC2–BC7). One family for each of the subsequent BC2–BC7 families was phenotyped and genotyped.

Wind Tunnel Phenotyping. All 649 backcross males were tested separately in the wind tunnel for response to each of the Z, E, and H blends. Our wind-tunnel assays were based on a series of behavioral male responses to each of the pheromone blends, i.e., resting, wing fanning (on/off), strong wing fanning, and hair pencil extrusion. Each category was given a score (Fig. 1 and Fig. S1), so that we had a weighted and quantitative measurement for each male.

The males were 1–2 d old and were sexually and olfactorily virgin. All males were behaviorally tested at the second hour of their scotophase, with 70% humidity and optimal temperature of 19 °C. Each male was kept in a cylinder 2 cm in diameter with gauze on both ends so that the air plume with the pheromone could pass through the tube. With this setup we were able to keep the males captured during the assay, avoiding the need to recapture them (and possibly damage them) at the end of the experiment. We used the same males for subsequent matings.

The three pheromone blends were ordered from Pherobank and were separately presented to each of the males for 1 min maximally. During one experimental day, all males were tested first with the Z blend, followed successively by the H and E blends, with an interval of 30 min to ventilate the wind tunnel between the tests. After a male showed the hair pencil extrusion response or after 1 min, the male was removed from the wind tunnel. The

latency of each reaction to the blend was recorded. If the behavioral response was ambiguous, the male was retested on the following day.

Mapping and Genotyping. We mapped the chromosomes using AFLP markers, using a protocol adapted from ref. 71. On polyacrylamide gels, the markers scored were present in the pure E-strain female, the F₁ hybrid male, and the heterozygous backcross males and were absent in the pure Z-strain male, the F₁ Z-strain female, and in the homozygous backcross offspring. The opposite pattern was scored as well. We scored 198 AFLP markers from 47 primer combinations in 123 samples (parents and F₁ and BC1 males and females). Using MapMaker 3.0 (72) we grouped 180 of these markers into 45 linkage groups, with at least two markers from different primer pairs. The number of markers for the 35 linkage groups varied from 2 to 11; the average number was 5.14. For fine-scale mapping on the Z chromosome, we used intron-size polymorphisms or SNP variation in 21 Z-linked genes that were scored in all backcross males; then recombination rates were determined in MapMaker as well. We verified that all polymorphic differences used in the mapping also differed among wild-type Z and E males in the laboratory strains.

QTL Analysis. With the phenotypic scores for all male behaviors, we first determined which linkage groups explained a significant portion of the variation, using R-studio using *t* test-based marker regression as implemented in R/qtl (version 0.98.490). We established significance thresholds for LOD scores empirically by permutation tests using 10,000 permutations. After finding a significant LOD score for only the Z chromosome, we conducted QTL analysis with the 21 Z-linked genes using MAPMAKER/QTL 1.1 (73, 74). We used the gene order determined by MapMaker 3.0, with a backcross design. We extracted the numerical values of the LOD scores from the program output and plotted them in Excel.

ACKNOWLEDGMENTS. The behavioral investigations were supported by Grant Vetenskapsrådet (the Swedish Research Council), Project 621-2014-4816 (to T.D.) and by Linnaeus Chemical Ecology, Ethology, and Evolution Initiative (ICE3) grant [The Swedish Research Council for Sustainable Development (FORMAS)/Swedish University of Agricultural Sciences] (to F.A.K. and T.D.). The molecular work and mapping analysis were supported by the Max Planck Institute for Chemical Ecology (D.G.H., A.T.G., and F.A.K.). F.A.K. received further funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement 661322.

- Cardé RT, Baker TC (1984) Sexual communication with pheromones. *Chemical Ecology of Insects*, eds Bell WJ, Cardé RT (Chapman and Hall, New York), pp 355–383.
- Cardé RT, Willis MA (2008) Navigational strategies used by insects to find distant, wind-borne sources of odor. *J Chem Ecol* 34(7):854–866.
- Groot AT, Dekker T, Heckel DG (2016) The genetic basis of pheromone evolution in moths. *Annu Rev Entomol* 61:99–117.
- Löfstedt C (1993) Moth pheromone genetics and evolution. *Philos Trans R Soc Lond B Biol Sci* 340(1292):167–177.
- Johansson BG, Jones TM (2007) The role of chemical communication in mate choice. *Biol Rev Camb Philos Soc* 82(2):265–289.
- Brooks R, et al. (2005) Experimental evidence for multivariate stabilizing sexual selection. *Evolution* 59(4):871–880.
- Kirkpatrick M, Nuismer SL (2004) Sexual selection can constrain sympatric speciation. *Proc Biol Sci* 271(1540):687–693.
- Coyne JA, Orr HA (2004) *Speciation* (Sinauer Associates, Inc., Sunderland, MA), pp 1–545.
- Ritche MG (2007) Sexual selection and speciation. *Annu Rev Ecol Syst* 38(7):79–102.
- Butlin RK, Trickett AJ (1997) *Can Population Genetic Simulations Help to Interpret Pheromone Evolution?* (Chapman & Hall, New York), pp 548–562.
- Groot AT, et al. (2006) Experimental evidence for interspecific directional selection on moth pheromone communication. *Proc Natl Acad Sci USA* 103(15):5858–5863.
- Löfstedt C, Hansson BS, Roelofs W, Bengtsson BO (1989) No linkage between genes controlling female pheromone production and male pheromone response in the European corn borer, *Ostrinia nubilalis* Hübner (Lepidoptera; Pyralidae). *Genetics* 123(3):553–556.
- Dopman EB, Bogdanowicz SM, Harrison RG (2004) Genetic mapping of sexual isolation between E and Z pheromone strains of the European corn Borer (*Ostrinia nubilalis*). *Genetics* 167(1):301–309.
- Gould F, et al. (2010) Sexual isolation of male moths explained by a single pheromone response QTL containing four receptor genes. *Proc Natl Acad Sci USA* 107(19):8660–8665.
- Olsson SB, et al. (2010) *Ostrinia* revisited: Evidence for sex linkage in European Corn Borer *Ostrinia nubilalis* (Hubner) pheromone reception. *BMC Evol Biol* 10:285.
- Lassance JM (2010) Journey in the *Ostrinia* world: From pest to model in chemical ecology. *J Chem Ecol* 36(10):1155–1169.
- Heckel DG (2010) Smells like a new species: Gene duplication at the periphery. *Proc Natl Acad Sci USA* 107(21):9481–9482.
- Roelofs W, et al. (1987) Sex pheromone production and perception in European corn borer moths is determined by both autosomal and sex-linked genes. *Proc Natl Acad Sci USA* 84(21):7585–7589.
- Cardé RT, Kochansky J, Stimmel JF, Wheeler AG, Roelofs WL (1975) Sex-pheromone of European corn-borer (*Ostrinia nubilalis*) - cis-responding and trans-responding males in Pennsylvania. *Environ Entomol* 4(3):413–414.
- Klun JA (1968) Isolation of a sex pheromone of the European corn borer. *J Econ Entomol* 61(2):484–487.
- Malausa T, et al. (2005) Assortative mating in sympatric host races of the European corn borer. *Science* 308(5719):258–260.
- Klun JA, Maini S (1979) Genetic basis of an insect chemical communication system: The European corn borer Lepidoptera, Pyralidae. *Environ Entomol* 8(3):423–426.
- Buechi R, Priesner E, Brunetti R (1982) Das sympatrische vorkommen von zwei pheromonstammen des Maiszünslers, *Ostrinia nubilalis* Hbn., in der Sudschweiz. *Mitt Schweiz Entomol Ges* 55(1-2):33–53.
- Roelofs WL, Du JW, Tang XH, Robbins PS, Eckenrode CJ (1985) Three European corn borer populations in New York based on sex pheromones and voltinism. *J Chem Ecol* 11(7):829–836.
- Bengtsson BO, Löfstedt C (1990) No evidence for selection in a pheromonly polymorphic moth population. *Am Nat* 136(5):722–726.
- Glover TJ, Campbell MG, Linn CE, Roelofs WL (1991) Unique sex-chromosome mediated behavioral-response specificity of hybrid male European corn-borer moths. *Experientia* 47(9):980–984.
- Lassance JM, et al. (2010) Allelic variation in a fatty-acyl reductase gene causes divergence in moth sex pheromones. *Nature* 466(7305):486–489.
- Lassance JM, et al. (2013) Functional consequences of sequence variation in the pheromone biosynthetic gene *pgFAR* for *Ostrinia* moths. *Proc Natl Acad Sci USA* 110(10):3967–3972.
- Carraher C, Authier A, Steinwender B, Newcomb RD (2012) Sequence comparisons of odorant receptors among toxic moth moths reveal different rates of molecular evolution among family members. *PLoS One* 7(6):e38391.
- Krieger J, et al. (2004) Genes encoding candidate pheromone receptors in a moth (*Heliothis virescens*). *Proc Natl Acad Sci USA* 101(32):11845–11850.

31. Lassance J-M, Bogdanowicz SM, Wanner KW, Löfstedt C, Harrison RG (2011) Gene genealogies reveal differentiation at sex pheromone olfactory receptor loci in pheromone strains of the European corn borer, *Ostrinia nubilalis*. *Evolution* 65(6):1583–1593.
32. Leary GP, et al. (2012) Single mutation to a sex pheromone receptor provides adaptive specificity between closely related moth species. *Proc Natl Acad Sci USA* 109(35):14081–14086.
33. Steinwender B, Thrimawithana AH, Crowhurst RN, Newcomb RD (2015) Pheromone receptor evolution in the cryptic leafroller species, *Ctenopseustis obliquana* and *C. herana*. *J Mol Evol* 80(1):42–56.
34. Yasukochi Y, Miura N, Nakano R, Sahara K, Ishikawa Y (2011) Sex-linked pheromone receptor genes of the European corn borer, *Ostrinia nubilalis*, are in tandem arrays. *PLoS One* 6(4):e18843.
35. Koutroumpa FA, et al. (2014) Shifts in sensory neuron identity parallel differences in pheromone preference in the European corn borer. *Front Ecol Evol* 2(65):1–12.
36. Wanner KW, et al. (2010) Sex pheromone receptor specificity in the European corn borer moth, *Ostrinia nubilalis*. *PLoS One* 5(1):e8685.
37. Kroemer JA, et al. (2011) A rearrangement of the Z chromosome topology influences the sex-linked gene display in the European corn borer, *Ostrinia nubilalis*. *Mol Genet Genomics* 286(1):37–56.
38. Glover TJ, Tang XH, Roelofs WL (1987) Sex pheromone blend discrimination by male moths from E and Z strains of European corn borer. *J Chem Ecol* 13(1):143–151.
39. Wadsworth CB, Li X, Dopman EB (2015) A recombination suppressor contributes to ecological speciation in *OSTRINIA* moths. *Heredity (Edinb)* 114(6):593–600.
40. Kárpáti Z, Tasin M, Cardé RT, Dekker T (2013) Early quality assessment lessens pheromone specificity in a moth. *Proc Natl Acad Sci USA* 110(18):7377–7382.
41. Huang Y, Fang J, Bedford MT, Zhang Y, Xu RM (2006) Recognition of histone H3 lysine-4 methylation by the double tudor domain of JMJD2A. *Science* 312(5774):748–751.
42. Badri KR, Zhou Y, Dhru U, Aramgam S, Schuger L (2008) Effects of the SANT domain of tension-induced/inhibited proteins (TIPs), novel partners of the histone acetyltransferase p300, on p300 activity and TIP-6-induced adipogenesis. *Mol Cell Biol* 28(20):6358–6372.
43. Voigt A, Pflanz R, Schäfer U, Jäckle H (2002) Perlecan participates in proliferation activation of quiescent *Drosophila* neuroblasts. *Dev Dyn* 224(4):403–412.
44. Temme C, Zaessinger S, Meyer S, Simonelig M, Wahle E (2004) A complex containing the CCR4 and CAF1 proteins is involved in mRNA deadenylation in *Drosophila*. *EMBO J* 23(14):2862–2871.
45. Villanyi Z, Collart MA (2015) Ccr4-Not is at the core of the eukaryotic gene expression circuitry. *Biochem Soc Trans* 43(6):1253–1258.
46. Pradhan SJ, et al. (2012) The conserved P body component HPat/Pat1 negatively regulates synaptic terminal growth at the larval *Drosophila* neuromuscular junction. *J Cell Sci* 125(Pt 24):6105–6116.
47. Estrada B, Gisselbrecht SS, Michelson AM (2007) The transmembrane protein Perdido interacts with Grip and integrins to mediate myotube projection and attachment in the *Drosophila* embryo. *Development* 134(24):4469–4478.
48. Schnorrer F, Kalchauer I, Dickson BJ (2007) The transmembrane protein Kon-tiki couples to Dgrip to mediate myotube targeting in *Drosophila*. *Dev Cell* 12(5):751–766.
49. Schnorrer F, Dickson BJ (2004) Muscle building; mechanisms of myotube guidance and attachment site selection. *Dev Cell* 7(1):9–20.
50. Hobert O, Westphal H (2000) Functions of LIM-homeobox genes. *Trends Genet* 16(2):75–83.
51. Moberg KH, Mukherjee A, Veraksa A, Artavanis-Tsakonas S, Hariharan IK (2004) The *Drosophila* F box protein archipelago regulates dMyc protein levels in vivo. *Curr Biol* 14(11):965–974.
52. Nicholson SC, Nicolay BN, Frolov MV, Moberg KH (2011) Notch-dependent expression of the archipelago ubiquitin ligase subunit in the *Drosophila* eye. *Development* 138(2):251–260.
53. Godt D, Laski FA (1995) Mechanisms of cell rearrangement and cell recruitment in *Drosophila* ovary morphogenesis and the requirement of *bric à brac*. *Development* 121(1):173–187.
54. Chu J, Dong PDS, Panganiban G (2002) Limb type-specific regulation of *bric à brac* contributes to morphological diversity. *Development* 129(3):695–704.
55. Godt D, Couderc JL, Cramton SE, Laski FA (1993) Pattern formation in the limbs of *Drosophila*: *bric à brac* is expressed in both a gradient and a wave-like pattern and is required for specification and proper segmentation of the tarsus. *Development* 119(3):799–812.
56. Li Q, et al. (2016) A functionally conserved gene regulatory network module governing olfactory neuron diversity. *PLoS Genet* 12(1):e1005780.
57. Kárpáti Z, Dekker T, Hansson BS (2008) Reversed functional topology in the antennal lobe of the male European corn borer. *J Exp Biol* 211(Pt 17):2841–2848.
58. Kárpáti Z, Olsson S, Hansson BS, Dekker T (2010) Inheritance of central neuroanatomy and physiology related to pheromone preference in the male European corn borer. *BMC Evol Biol* 10:286.
59. Nakagawa T, Sakurai T, Nishioka T, Touhara K (2005) Insect sex-pheromone signals mediated by specific combinations of olfactory receptors. *Science* 307(5715):1638–1642.
60. Sakurai T, et al. (2011) A single sex pheromone receptor determines chemical response specificity of sexual behavior in the silkworm *Bombyx mori*. *PLoS Genet* 7(6):e1002115–e1002115.
61. Fujii T, et al. (2011) Sex-linked transcription factor involved in a shift of sex-pheromone preference in the silkworm *Bombyx mori*. *Proc Natl Acad Sci USA* 108(44):18038–18043.
62. Dobritsa AA, van der Goes van Naters W, Warr CG, Steinbrecht RA, Carlson JR (2003) Integrating the molecular and cellular basis of odor coding in the *Drosophila* antenna. *Neuron* 37(5):827–841.
63. Jafari S, et al. (2012) Combinatorial activation and repression by seven transcription factors specify *Drosophila* odorant receptor expression. *PLoS Biol* 10(3):e1001280–e1001280.
64. Linn C, Poole K, Zhang A, Roelofs W (1999) Pheromone-blend discrimination by European corn borer moths with inter-race and inter-sex antennal transplants. *J Comp Physiol A* 184(3):273–278.
65. Hansson BS, Hallberg E, Löfstedt C, Steinbrecht RA (1994) Correlation between dendrite diameter and action potential amplitude in sex pheromone specific receptor neurons in male *Ostrinia nubilalis* (Lepidoptera: Pyralidae). *Tissue Cell* 26(4):503–512.
66. Hansson BS, Löfstedt C, Roelofs WL (1987) Inheritance of olfactory response to sex pheromone components in *Ostrinia nubilalis*. *Naturwissenschaften* 74(10):497–499.
67. Cosse AA, et al. (1995) Pheromone behavioral-responses in unusual male European corn-borer hybrid progeny not correlated to electrophysiological phenotypes of their pheromone-specific antennal neurons. *Experientia* 51(8):809–816.
68. Kárpáti Z, Molnar B, Szoecs G (2007) Pheromone titer and mating frequency of E- and Z-strains of the European corn borer, *Ostrinia nubilalis*: Fluctuation during scotophase and age dependence. *Acta Phytopathol Entomol Hung* 42(2):331–341.
69. Mani E, Riggenbach W, Mendik M (1978) Rearing of the codling moth *Laspeyresia pomonella* on artificial diet 1968–1978. *Mitt Schweiz Entomol Ges* 51(4):315–326.
70. Heckel DG (1993) Comparative genetic linkage mapping in insects. *Annu Rev Entomol* 38:381–408.
71. Sheck AL, et al. (2006) Genetics of sex pheromone blend differences between *Heliothis virescens* and *Heliothis subflexa*: A chromosome mapping approach. *J Evol Biol* 19(2):600–617.
72. Lander ES, et al. (1987) MAPMAKER: An interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1(2):174–181.
73. Lincoln S, Daly M, Lander E (1993) Mapping genes controlling quantitative traits using MAPMAKER/QTL 1.1: A tutorial and reference manual, 2nd edition. *Whitehead Institute Technical Report* (Whitehead Institute for Biomedical Research, Cambridge, MA).
74. Paterson AH, et al. (1991) Mendelian factors underlying quantitative traits in tomato: Comparison across species, generations, and environments. *Genetics* 127(1):181–197.
75. Han S, et al. (2012) The E3 ubiquitin ligase protein associated with Myc (Pam) regulates mammalian/mechanistic target of rapamycin complex 1 (mTORC1) signaling in vivo through N- and C-terminal domains. *J Biol Chem* 287(36):30063–30072.
76. Le Guyader S, Maier J, Jesuthasan S (2005) Esrom, an ortholog of PAM (protein associated with c-myc), regulates pteridine synthesis in the zebrafish. *Dev Biol* 277(2):378–386.
77. Murthy V, et al. (2004) Pam and its ortholog highwire interact with and may negatively regulate the TSC1 center dot TSC2 complex. *J Biol Chem* 279(2):1351–1358.
78. Wan HI, et al. (2000) Highwire regulates synaptic growth in *Drosophila*. *Neuron* 26(2):313–329.
79. Sia GM, et al. (2007) Interaction of the N-terminal domain of the AMPA receptor GluR4 subunit with the neuronal pentraxin NP1 mediates GluR4 synaptic recruitment. *Neuron* 55(1):87–102.
80. Datta S, Kankel DR (1992) L(1) trol and L(1)Devl, loci affecting the development of the adult central-nervous system in *Drosophila melanogaster*. *Genetics* 130(3):523–537.
81. Park Y, et al. (2003) *Drosophila* Perlecan modulates FGF and Hedgehog signals to activate neural stem cell division. *Dev Biol* 253(2):247–257.
82. Thoden JB, Timson DJ, Reece RJ, Holden HM (2004) Molecular structure of human galactose mutarotase. *J Biol Chem* 279(22):23431–23437.
83. Werner T, et al. (2000) A family of peptidoglycan recognition proteins in the fruit fly *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 97(25):13772–13777.
84. Fisher RA (1936) The use of multiple measurements in taxonomic problems. *Ann Eugen* 7:179–188.