UvA-DARE (Digital Academic Repository)

One-Dimensional Quasicrystals from Incommensurate Charge Order

Flicker, F.; van Wezel, J.

DOI

[10.1103/PhysRevLett.115.236401](https://doi.org/10.1103/PhysRevLett.115.236401)

Publication date

2015

Document Version

Final published version

Published in

Physical Review Letters

[Link to publication](#)

Citation for published version (APA):

Flicker, F., & van Wezel, J. (2015). One-Dimensional Quasicrystals from Incommensurate Charge Order. *Physical Review Letters*, 115(23), [236401].
<https://doi.org/10.1103/PhysRevLett.115.236401>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

One-Dimensional Quasicrystals from Incommensurate Charge Order

Felix Flicker*

Department of Physics, University of California, Berkeley, California 94720, USA

Jasper van Wezel†

Institute for Theoretical Physics, University of Amsterdam, 1090 GL Amsterdam, Netherlands

(Received 26 June 2015; published 3 December 2015)

Artificial quasicrystals are nowadays routinely manufactured, yet only two naturally occurring examples are known. We present a class of systems with the potential to be realized both artificially and in nature, in which the lowest energy state is a one-dimensional quasicrystal. These systems are based on incommensurately charge-ordered materials, in which the quasicrystalline phase competes with the formation of a regular array of discommensurations as a way of interpolating between incommensurate charge order at high temperatures and commensurate order at low temperatures. The nonlocal correlations characteristic of the quasicrystalline state emerge from a free-energy contribution localized in reciprocal space. We present a theoretical phase diagram showing that the required material properties for the appearance of such a ground state allow for one-dimensional quasicrystals to form in real materials. The result is a potentially wide class of one-dimensional quasicrystals.

DOI: 10.1103/PhysRevLett.115.236401

PACS numbers: 71.23.Ft, 61.44.Br, 61.44.Fw, 71.45.Lr

While many quasicrystals have been artificially created since their first synthesis in 1984, only two naturally occurring quasicrystals have been identified to date—both found in the same Siberian meteorite [1–3]. In the quasiperiodic, never-repeating, pattern of two or more unit cells defining a quasicrystal (QC) [4,5], locally changing a single cell may require adjustment of an infinite number of other cells in order to maintain the quasiperiodicity [6]. This global property makes it nontrivial to grow a QC cell by cell. Proposed mechanisms in two or higher dimensions include recognizing vertex matching rules that imply “forced tiles,” random tiling models, and relaxation processes from nonquasiperiodic systems [7].

In (quasi)one-dimensional (1D) systems, QCs consist of a quasiperiodic sequence of long and short unit cells. The global nature of 1D QCs is apparent from their correspondence to projections of regular 2D crystals [4,8]. One way to experimentally access properties of 1D QCs is to consider instead a setup containing two periodic subsystems that are incommensurate with respect to one another. The sequence of subsystem species encountered as one traverses the system is aperiodic, and can be mapped onto a similar aperiodic or quasicrystalline sequence of unit cells [see Fig. 1(a)]. Such incommensurate systems have recently been employed in a number of experimental studies of QCs, including optical waveguide arrays simulating 1D crystals with incommensurate periodic on-site potentials, cold-atom condensates in optical lattices, moiré superlattices in graphene derivatives, and in theoretical descriptions of incommensurate charge-density waves [8–12]. While these are of much interest in their own right, they only constitute quasicrystals by way of mathematical

analogy. One key difference with true quasicrystals for example, is that incommensurate systems have no lower bound on the spatial separation between sites of the two subsystems; it is only the sequence which is quasiperiodic (see also the Supplemental Material [13]).

In this Letter we describe a mechanism for generating 1D quasicrystals based on coupling two incommensurate subsystems and allowing them to adjust to minimize their overall free energy. We take as a specific example the case of incommensurate charge-density waves, and we show that, under specific conditions, it is energetically favorable for the atomic lattice to adopt a quasiperiodic structure. The resulting state is thus a true quasicrystal, transcending the mere mathematical equivalence between 1D QCs and incommensurate charge-density modulations on top of a periodic atomic background. We show that the required material properties are physically realistic. Finally, we outline the criteria needed for this mechanism to apply more generally.

Low-dimensional materials are prone to develop charge order. In the ideal 1D case, a Peierls instability always yields a spontaneous periodic modulation of the electron density, accompanied by periodic displacements in the atomic lattice with the same wave vector [16]. In real quasi-1D (or higher-dimensional) materials, charge order typically occurs only if the electron-phonon coupling is sufficiently strong [14,17]. In that case, the order arises when the electronic susceptibility overcomes the cost of populating phonon modes, so that

$$\int d\mathbf{k} |g(\mathbf{k}, \mathbf{k} + \mathbf{q})|^2 \frac{f(\epsilon_{\mathbf{k}}) - f(\epsilon_{\mathbf{k}+\mathbf{q}})}{\epsilon_{\mathbf{k}+\mathbf{q}} - \epsilon_{\mathbf{k}}} \geq \hbar\omega_{\mathbf{q}}, \quad (1)$$

where $g(\mathbf{k}, \mathbf{k} + \mathbf{q})$ is the electron-phonon coupling, which is generically momentum and orbital dependent. $f(\epsilon_{\mathbf{k}})$ is the Fermi-Dirac distribution for an electron with momentum $\mathbf{k}$ and energy $\epsilon_{\mathbf{k}}$, and $\hbar\omega_{\mathbf{q}}$ is the energy of the phonon mode that softens in the charge ordering transition. The full susceptibility $\chi(\mathbf{q})$, defined by the left-hand side of Eq. (1), has its maximum at $\tilde{\mathbf{q}}$, which determines the wave vector $\mathbf{Q}(T_{\text{CDW}}) = \tilde{\mathbf{q}}$ at which the charge order and atomic displacements first form. In the presence of strong nesting, guaranteed in 1D, $\tilde{\mathbf{q}}$ equals the nesting vector. In higher dimensions it is determined by the momentum dependences of both the electron-phonon coupling and the electronic dispersion [14].

The evolution of the charge density wave vector $\mathbf{Q}(T)$, as temperature is lowered beyond the onset temperature T_{CDW} , can be modeled by a Ginzburg-Landau expansion of the free energy [18,19],

$$F = - \int d\mathbf{q} \chi(\mathbf{q}) \rho^2(\mathbf{q}) + \int d\mathbf{x} b \rho^4(\mathbf{x}) - \sum_{n, \mathbf{K}} c_n \cos(\mathbf{K} \cdot \mathbf{x}) \rho^n(\mathbf{x}). \quad (2)$$

The modulation on top of the average electron density is written as $\rho(\mathbf{x}) = \psi \cos[\phi(\mathbf{x})]$, where for a density wave without defects $\phi(\mathbf{x}) = \mathbf{Q} \cdot \mathbf{x}$. For notational convenience the quadratic part of the free energy has been written in reciprocal space, where it is proportional to the full electronic susceptibility χ . This term favors the formation of charge order at wave vector $\tilde{\mathbf{q}}$. The other terms are more conveniently written in real space. Those proportional to c_n , with $n > 2$, represent the local coupling between the atomic lattice and the charge modulations. These favor charge modulations locally commensurate with the lattice by giving a nonzero contribution to F whenever $n\phi(\mathbf{x}) = \mathbf{K} \cdot \mathbf{x}$, with $\mathbf{K}$ a reciprocal lattice vector of the atomic lattice [15,18].

At the transition temperature T_{CDW} , the order parameter amplitude ψ is small, so that the lower-order terms in Eq. (2) dominate, and the order forms at wave vector $\mathbf{Q}(T_{\text{CDW}}) = \tilde{\mathbf{q}}$. As temperature is lowered, ψ increases, and the terms proportional to c_n begin to compete with the second-order term. Because the coupling to the lattice has an effect only for the parts of the density wave that are locally commensurate, the charge order with $\phi(\mathbf{x}) = \mathbf{Q} \cdot \mathbf{x}$ will not gain energy from the final terms for any incommensurate value of $\mathbf{Q}$. Instead, the charge-ordered state can lower its energy by adopting a locally commensurate structure $\phi(\mathbf{x}) = \mathbf{K} \cdot \mathbf{x} - \delta(\mathbf{x})$, where $\delta(\mathbf{x})$ introduces a regular array of broadened phase slips (discommensurations) [20], which render the average propagation vector incommensurate even if the local structure is predominantly commensurate [15]. The evolution as the temperature decreases further is characterized by a sharpening of the initially broad discommensurations [15,19].

In many materials, $\mathbf{Q}(T)$ starts out incommensurate at the onset of the charge-ordered state, evolves towards a close-by commensurate value, and locks in at a second transition temperature T_{IC} below which the charge order remains commensurate. Well-known examples can be seen in $2H\text{-TaSe}_2$ and tetrathiafulvalene tetracyanoquinodimethane (TTF-TCNQ) [21,22]. In other materials, the evolution of $\mathbf{Q}(T)$ starts out in a similar manner, but remains incommensurate to zero temperature. Examples include $\text{K}_{0.3}\text{MoO}_3$, TbTe_3 , and NbSe_2 [22–24]. A proliferation of discommensurations can describe either type of behavior depending on the value of the ratio $c_n/\chi(\tilde{\mathbf{q}})$ in the free energy of Eq. (2), which determines the strength of the coupling to the lattice relative to the maximum strength of the full susceptibility. The presence of discommensurations in these materials has been experimentally verified, for example, in scanning tunneling microscopy experiments on NbSe_2 [25].

The discommensuration state takes advantage of all the terms in the free energy of Eq. (2). Nevertheless, one can imagine conditions under which it may not be the optimal configuration of $\phi(\mathbf{x})$. Consider, for example, incommensurate charge order forming at T_{CDW} with an incommensurate wavelength $\lambda = 2\pi/\tilde{q}$ lying close to halfway between three and four lattice spacings [Fig. 1(a)]. In that case, it is not *a priori* obvious whether the coupling to the lattice will prefer the charge order to lock-in at period 3 or 4. Choosing one, and introducing discommensurations such that the local electronic density modulations are commensurate while the average propagation vector remains incommensurate, necessarily results in a high density of rapid phase changes [shown in Fig. 1(b)].

FIG. 1 (color online). Possible charge orderings when the electron density (blue) develops an instability with wave vector Q incommensurate with the ion lattice (pink). Here we used a wavelength of $2\pi/Q = \sqrt{11.7} \approx 3.42$ lattice spacings. (a) Perfectly incommensurate charge order. Labeling the blue peaks A and the pink ions B , the sequence of A 's and B 's is a quasiperiodic tiling of the units $ABBB$ and $ABBBB$ [14]. (b) Discommensuration state, which is locally commensurate with period 3 but maintains the average wave vector Q [15]. (c) A quasicrystal composed of a quasiperiodic arrangement of period-3 and period-4 unit cells, maintaining the average wave vector Q . This state is reached by shifting each peak A onto the nearest ion B . Dotted lines show the incommensurate state for comparison. In all cases, the ion positions will adjust to the altered electronic charge distribution (not shown). An alternative comparison between these states is provided in the Supplemental Material [13].

These sharp discommensurations involve high-momentum components of $\rho(\mathbf{q})$ (higher harmonics of $\tilde{\mathbf{q}}$), and are, therefore, energetically costly with respect to the quadratic term in F . An alternative way of distorting the incommensurate charge order and gaining local lock-in energy, while retaining the overall incommensurability, is depicted in Fig. 1(c). This configuration is obtained by moving every maximum in the charge-density wave directly onto the atomic position closest to it. The result is a charge modulation that is locally commensurate everywhere, but with two local periodicities. The locally commensurate patches are smoothly connected (although higher derivatives are not smooth), and, therefore, do not involve the high-momentum components characteristic of the discommensurations. For a more detailed description of the difference between this state and the discommensuration state, see the Supplemental Material [13].

It is clear from the incommensurability that the pattern of period-3 and period-4 modulations in the multiply commensurate structure never repeats itself. In fact, it can be shown, using a cut-and-project construction, that the arrangement is quasiperiodic [14]. In reality, the ions will be displaced towards the charge maxima simultaneously with the displacement of the charge maxima towards the ions. The structure shown in Fig. 1(c) will, therefore, not just contain incommensurate charge modulations or a quasiperiodic electronic structure, but will constitute a true 1D quasicrystal. At the onset temperature of the multiply commensurate order, there will be a crystal-to-quasicrystal phase transition breaking all translational symmetries of the original lattice.

The x-ray diffraction pattern of a quasicrystal is characterized by a dense array of sharp peaks [5,7]. As shown in Fig. 2, the Fourier transform of the multiply commensurate charge order indeed contains such an arrangement, centered around the average incommensurate wave vector $\tilde{\mathbf{q}}$. Real

FIG. 2 (color online). Calculated x-ray diffraction pattern for the discommensuration state (blue) and quasicrystal (pink) shown in Fig. 1 (purple indicates overlapping peaks). The black line depicts the approximation to the full susceptibility χ used to calculate the phase diagram of Fig. 3. Both types of order have their main peak at $\tilde{\mathbf{q}}$, but the quasicrystal has a dense covering of secondary peaks close to the main diffraction spot, which benefit from the nonzero χ in that region. The units are arbitrary, but both diffraction patterns have the same scale.

quasi-1D materials have susceptibilities $\chi(\mathbf{q})$ that decay over a finite range of momenta. The clustered peaks around $\tilde{\mathbf{q}}$ can, therefore, also gain energy from the quadratic term in F . This situation should be contrasted with that of the Fourier transform of the discommensuration state, also shown in Fig. 2. In that case, although the central peak gains energy, the higher harmonics typically lie far away in k -space, and cannot utilize the nonzero width of the susceptibility. The difference in the extent to which these structures profit from the quadratic term in F can render the quasicrystal energetically favorable to the discommensuration state.

To quantify the relative stabilities of the quasicrystal and discommensuration states, we consider again the free energy of Eq. (2). We model the full susceptibility as $\chi(\mathbf{q}) = a/[\sigma^2(|\mathbf{q}| - \tilde{\mathbf{q}})^2 + 1]$, where the width $1/\sigma$ of the peak is taken to be small but nonzero. The ratio b/a of the quartic and quadratic component of F at $\mathbf{q} = \tilde{\mathbf{q}}$ determines the overall magnitude of the order parameter ψ , but does not affect the form of $\phi(\mathbf{x})$. Whether the discommensuration or quasicrystalline state is favorable is thus determined by the ratios $(\psi^4 c_4)/(\psi^3 c_3)$ and $(\psi^3 c_3)/(\psi^2 a)$. In Fig. 3(a), we present the phase diagram as a function of these two ratios. The energy of the quasicrystalline state is compared to that of an array of discommensurations with optimized density and widths. A comparison of the energies of individual states, as a function of $\psi c_3/a$ for fixed $\psi c_4/c_3$, is shown in Fig. 3(b).

The energy of the incommensurate state does not depend on the coupling to the lattice, and is constant in $\psi c_3/a$. The commensurate state, on the other hand, does not gain any energy from the susceptibility term, but takes full advantage of the lock-in term, and decreases linearly in energy as $\psi c_3/a$ increases. At high coupling to the lattice, the commensurate state thus becomes favorable to the incommensurate. The discommensuration state interpolates between these two extremes, having its main Fourier component at the incommensurate value but being locally commensurate. It takes advantage of all terms in the

FIG. 3 (color online). (a) The predicted phase diagram emerging from Eq. (2), for $Q = 2\pi/\sqrt{11.7}$ and $b = 1$. (b) The free energies of the different ordered states as a function of $\psi c_3/a$ for fixed $\psi c_4/c_3 = 1.2$. The abbreviations used are QC: quasicrystal, DCn: discommensuration state with commensurate areas of period n , ICDW: incommensurate charge-density wave, and $Q = 1/n$: commensurate charge order with period n .

free energy, and outperforms both the commensurate and incommensurate states.

The quasicrystal also has its dominant Fourier component at the incommensurate value. At intermediate values of $\psi c_3/a$ it additionally takes advantage of the spread in $\chi(\mathbf{q})$ and only pays a modest price for not being optimally commensurate everywhere. In this regime, the quasicrystal is favored over the discommensuration state, giving rise to the extended area of stability shown in Fig. 3(a). At either extreme of $\psi c_3/a$, however, the discommensurations win out: the spectral weight of the dominant QC Fourier peak is constant, so does not approach the weight of the incommensurate state in the limit of zero lattice coupling. The discommensuration state is, therefore, favorable to the quasicrystal at very low values of $\psi c_3/a$. For strong coupling to the lattice, the locally commensurate nature of the quasicrystal allows it to gain lock-in energy, with the overall energy gain a weighted average of the c_3 and c_4 terms. However, if these two coefficients are not equal, the quasicrystal cannot gain as much energy as the discommensuration state, which is always locally commensurate with the most favorable commensurability. The discommensuration state, therefore, also wins out over the quasicrystal at very high values of the coupling to the lattice.

Based on these considerations, the prerequisites for this type of 1D quasicrystal to emerge in any real (natural or artificial) material are (i) a quasi-1D charge-ordered material with an incommensurate wave vector lying close to halfway between two commensurate values, (ii) a sharp, but not infinitely sharp, susceptibility, and (iii) a strong, momentum-independent electron-phonon coupling. The latter is necessary to ensure that the coupling of the charge order to the lattice does not strongly favor any particular commensurate value of the propagation vector.

Quasi-1D materials contain strongly bonded chains of atoms with relatively weak interchain coupling. Their Fermi surfaces typically contain nearly flat pieces, giving a strongly peaked susceptibility. The charge-ordering wave vector is therefore close to $2k_F$, and is generically incommensurate. The question then arises as to why we are not overwhelmed with naturally occurring quasicrystals in quasi-1D materials. First, the charge modulation ρ is small compared to the average charge density, implying that a dense set of peaks neighboring $\tilde{\mathbf{q}}$ in the diffraction pattern may not be straightforward to distinguish experimentally. Second, in real materials it may be expected that the lock-in energy is different at different commensurate values [19], putting most materials close to the left side of Fig. 3(a). Note, however, that the order parameter amplitude ψ increases with decreasing temperature, and thus enhances the value of $\psi c_4/c_3$. Conversely, at the onset of ordering, ψ is vanishingly small, and the free energy is dominated by the second-order term. This implies that the charge-ordered phase starts off at the origin of the phase diagram, and moves upwards and to the right only as temperature

decreases below T_{CDW} . Third, except in cases of extremely strong electronic nesting, the momentum and orbital dependence of the electron-phonon coupling is generically essential for charge order to develop in real materials [14]. These points aside, the results of Fig. 3 suggest that a targeted search may reveal previously unnoticed quasicrystalline phases at low temperatures in quasi-1D materials with incommensurate charge order. Experimentally, the quasicrystalline charge order could be identified, for example, using microscopy to directly identify arrangements of neighboring period-4 and period-3 cells, or using diffraction probes to find a fractal distribution of sharp peaks.

To conclude, we have shown that for an extended range of material parameters quasi-1D systems minimize their free energy by developing quasiperiodic tilings of two inequivalent unit cells. This provides a wide class of 1D materials with the potential to become quasicrystalline. Candidate materials, both natural and artificial, could be revealed by a directed search for materials exhibiting incommensurate charge order, with a wavelength close to halfway between two commensurate values, whose electronic structures are well nested, and whose electron-phonon couplings are strong but momentum independent. The quasicrystalline state also extends the set of possible low-energy configurations of incommensurately charge-ordered systems, adding to the known incommensurate and commensurate states and their interpolation via discommensurations.

Existing proposals for quasicrystal growth mechanisms have centered on single-cell additions at defect sites, or local adjustments based on propagating defects [7]. The quasicrystalline state discussed here, on the other hand, arises from a global adjustment of two periodic but incommensurate parent structures. The required coordination for this global effect in real space derives from the fact that it is a localized peak of the susceptibility in reciprocal space that determines the dominant wave vector of the quasicrystalline state.

Because the free energy contains two incommensurate periods, it is perhaps unsurprising that a ground state exists that respects the lack of translational symmetry. Closely related states have been found in both the quantum Hall effect in the Tao-Thouless limit [26] and the formation of Wigner lattices in the Hubbard model [27]. In the latter case, the Wigner lattice is invoked to explain the optical spectra of nearly quasicrystalline states in the 1D charge-ordered material TTF-TCNQ [28,29]. Both systems, however, concern high-denominator rational commensurate periods rather than the incommensurate situation considered here, and neither depends on the particular free-energy arguments arising from Eq. (2).

The mechanism for forming quasicrystals considered in this Letter has the potential to arise whenever two incommensurate periodicities are simultaneously present in the same system. If an appropriate coupling between the

subsystems can be introduced in the presence of a global constraint, a free energy similar to that of Eq. (2) may be expected to apply, and a quasicrystalline state will generically emerge over an extensive portion of phase space.

The authors wish to thank Shane Farnsworth for helpful discussions. J. v. W. acknowledges support from a VIDI grant financed by the Netherlands Organisation for Scientific Research (NWO). F. F. acknowledges support from a Lindemann Trust Fellowship of the English Speaking Union.

*flicker@physics.org

†vanwezel@uva.nl

- [1] D. Shechtman, I. Blech, D. Gratias, and J. W. Cahn, *Phys. Rev. Lett.* **53**, 1951 (1984).
- [2] L. Bindi, P. J. Steinhardt, N. Yao, and P. J. Lu, *Science* **324**, 1306 (2009).
- [3] L. Bindi, N. Yao, C. Lin, L. S. Hollister, C. L. Andronikos, V. V. Distler, M. P. Eddy, A. Kostin, V. Kryachko, G. J. MacPherson, W. M. Steinhardt, M. Yudovskaya, and P. J. Steinhardt, *Sci. Rep.* **5**, 9111 (2015).
- [4] J. E. S. Socolar and P. J. Steinhardt, *Phys. Rev. B* **34**, 617 (1986).
- [5] M. Senechal, *Quasicrystals and Geometry* (Cambridge University Press, Cambridge, England, 1995).
- [6] C. V. Achim, M. Schmiedeberg, and H. Löwen, *Phys. Rev. Lett.* **112**, 255501 (2014).
- [7] C. Janot, *Quasicrystals: A Primer* (Clarendon Press, Oxford, 1994).
- [8] F. Flicker and J. van Wezel, *Europhys. Lett.* **111**, 37008 (2015).
- [9] Y. E. Kraus, Y. Lahini, Z. Ringel, M. Verbin, and O. Zilberberg, *Phys. Rev. Lett.* **109**, 106402 (2012).
- [10] Y. E. Kraus and O. Zilberberg, *Phys. Rev. Lett.* **109**, 116404 (2012).
- [11] M. Aidelsburger, M. Lohse, C. Schweizer, M. Atala, J. T. Barreiro, S. Nascimbene, N. R. Cooper, I. Bloch, and N. Goldman, *Nat. Phys.* **11**, 162 (2015).
- [12] C. R. Dean, L. Wang, P. Maher, C. Forsythe, F. Ghahari, Y. Gao, J. Katoch, M. Ishigami, P. Moon, M. Koshino, T. Taniguchi, K. Watanabe, K. L. Shepard, J. Hone, and P. Kim, *Nature* **497**, 598 (2013).
- [13] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.115.236401> for a detailed description of the differences between the various charge-ordered phases discussed in the main text.
- [14] F. Flicker and J. van Wezel, *Nat. Commun.* **6**, 7034 (2015).
- [15] W. L. McMillan, *Phys. Rev. B* **14**, 1496 (1976).
- [16] R. Peierls, *More Surprises in Theoretical Physics* (Princeton University Press, Princeton, NJ, 1991).
- [17] S. K. Chan and V. Heine, *J. Phys. F* **3**, 795 (1973).
- [18] W. L. McMillan, *Phys. Rev. B* **12**, 1187 (1975).
- [19] Y. Feng, J. van Wezel, J. Wang, F. Flicker, D. M. Silevitch, P. B. Littlewood, and T. F. Rosenbaum, *Nat. Phys.* **11**, 865 (2015).
- [20] We employ the standard assumption of the array being periodic. Although strong lattice coupling may in principle alter the distribution, this will not affect the qualitative difference between the discommensuration state and any of the other states discussed here. Only the character of the transition between them may be altered.
- [21] S. Kagoshima, T. Ishiguro, and H. Anzai, *J. Phys. Soc. Jpn.* **41**, 2061 (1976).
- [22] D. E. Moncton, J. D. Axe, and F. J. DiSalvo, *Phys. Rev. B* **16**, 801 (1977).
- [23] A. H. Moudden, J. D. Axe, P. Monceau, and F. Levy, *Phys. Rev. Lett.* **65**, 223 (1990).
- [24] N. Ru, C. L. Condon, G. Y. Margulis, K. Y. Shin, J. Laverock, S. B. Dugdale, M. F. Toney, and I. R. Fisher, *Phys. Rev. B* **77**, 035114 (2008).
- [25] A. Soumyanarayanan, M. M. Yee, Y. He, J. van Wezel, D. J. Rahn, K. Rossnagel, E. W. Hudson, M. R. Norman, and J. E. Hoffman, *Proc. Natl. Acad. Sci. U.S.A.* **110**, 1623 (2013).
- [26] E. J. Bergholtz and A. Karlhede, *Phys. Rev. B* **77**, 155308 (2008).
- [27] J. Hubbard, *Phys. Rev. B* **17**, 494 (1978).
- [28] J. B. Torrance, B. A. Scott, and F. B. Kaufman, *Solid State Commun.* **17**, 1369 (1975).
- [29] J. Tanaka, M. Tanaka, T. Kawai, T. Takabe, and O. Maki, *Bull. Chem. Soc. Jpn.* **49**, 2358 (1976).