



**UvA-DARE (Digital Academic Repository)**

**Searching for Primordial Black Holes in the Radio and X-Ray Sky**

Gaggero, D.; Bertone, G.; Calore, F.; Connors, R.M.T.; Lovell, M.R.; Markoff, S.B.; Storm, E.M.

*Published in:*  
Physical Review Letters

*DOI:*  
[10.1103/PhysRevLett.118.241101](https://doi.org/10.1103/PhysRevLett.118.241101)

[Link to publication](#)

*Citation for published version (APA):*

Gaggero, D., Bertone, G., Calore, F., Connors, R. M. T., Lovell, M., Markoff, S., & Storm, E. (2017). Searching for Primordial Black Holes in the Radio and X-Ray Sky. *Physical Review Letters*, 118(24), [241101]. DOI: 10.1103/PhysRevLett.118.241101

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

## Searching for Primordial Black Holes in the Radio and X-Ray Sky

Daniele Gaggero,<sup>1,\*</sup> Gianfranco Bertone,<sup>1</sup> Francesca Calore,<sup>1,2</sup> Riley M. T. Connors,<sup>1,3</sup> Mark Lovell,<sup>1,4</sup>  
Sera Markoff,<sup>1,3</sup> and Emma Storm<sup>1</sup>

<sup>1</sup>GRAPPA, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, Netherlands

<sup>2</sup>LAPTh, CNRS and Université Savoie Mont Blanc, 9 Chemin de Bellevue, 74941 Annecy-le-Vieux, France

<sup>3</sup>API, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, Netherlands

<sup>4</sup>MPIA, Königstuhl 17, D-69117 Heidelberg, Germany

(Received 9 December 2016; revised manuscript received 12 May 2017; published 12 June 2017)

We model the accretion of gas onto a population of massive primordial black holes in the Milky Way and compare the predicted radio and x-ray emission with observational data. We show that, under conservative assumptions on the accretion process, the possibility that  $\mathcal{O}(10)M_{\odot}$  primordial black holes can account for all of the dark matter in the Milky Way is excluded at  $5\sigma$  by a comparison with a Very Large Array radio catalog at 1.4 GHz and at  $\approx 40\sigma$  by a comparison with a Chandra x-ray catalog (0.5–8 keV). We argue that this method can be used to identify such a population of primordial black holes with more sensitive future radio and x-ray surveys.

DOI: 10.1103/PhysRevLett.118.241101

*Introduction.*—The first direct detection of a gravitational wave signal, announced by the LIGO Collaboration recently [1], demonstrated the existence of  $\sim 30 M_{\odot}$  black holes (BHs), prompting the suggestion [2,3] that these objects are *primordial* black holes (PBHs) that may account for all of the dark matter (DM) [4–6] in the Universe. The connection between PBHs and DM has been extensively studied in the past (see, e.g., [7–12]), and a number of constraints exist on the cosmic abundance of PBHs over a very wide mass range (see the discussion below and, e.g., Ref. [13] for a recent review).

In this Letter, we consider for the first time, in the context of PBH searches, the x-ray and radio emission from the Galactic Ridge region produced by the accretion of interstellar gas onto a population of  $\mathcal{O}(10)M_{\odot}$  PBHs in the Milky Way. Given current estimates of the bulge mass [14], if PBHs constitute all of the DM, there should be  $\mathcal{O}(10^9)$  such objects within 2 kpc from the Galactic center (GC). Since the inner part of the bulge contains high gas densities [15], a significant fraction would inevitably form an accretion disk and emit a broadband spectrum of radiation. We show (Fig. 1) that radio and x-ray data in the Galactic Ridge region rule out, at  $5\sigma$  and  $40\sigma$ , respectively, the possibility that PBHs constitute all of the DM in the Galaxy, even under conservative assumptions on the physics of accretion.

Our limits arise from a realistic modeling of the accretion process, based on the observational evidence for inefficient accretion in the Milky Way today [16,17], and corroborate, with a completely independent approach, the exclusion of massive PBHs as DM candidates.

*Accretion on black holes.*—We should expect the accretion rates  $\dot{M}$  of a Galactic population of PBHs accreting from interstellar gas to be well below the Eddington limit

$\dot{M}_{\text{Edd}}$ . Even under the unrealistic assumption of Bondi-Hoyle-Lyttleton accretion [18,19] and typical velocities as low as  $\sim 10$  km/s, the accretion rate would definitely be sub-Eddington:  $\dot{M} \sim 10^{-5}(n_{\text{gas}}/\text{cm}^{-3})\dot{M}_{\text{Edd}}$ .

BHs accreting at  $\dot{M} < 0.01\dot{M}_{\text{Edd}} \equiv \dot{M}_{\text{crit}}$  are radiatively inefficient, such that the luminosity scales nonlinearly with  $\dot{M}$  [20]. The prevailing physical pictures adopted to explain the weak emission properties are advection-dominated accretion, in which the gas cooling time scales greatly exceed the dynamical time scales [21], and mass loss from




FIG. 1. Upper limits on the fraction of DM in PBHs of a given mass  $M$ , arising from the nonobservation of bright x-ray (blue shaded regions) and radio (red) BH candidates at the GC. We assume a conservative value of  $\lambda$ , regulating the departure from the Bondi accretion rate:  $\lambda = 0.02$ . The dotted gray line corresponds to  $30 M_{\odot}$  PBH, and the hatched gray region is unphysical ( $f_{\text{DM}} > 1$ ).

the disk or internal convective flows, such that the accretion rate itself has decreased once the gas reaches the inner edge of the disk [22,23]. It is likely that both mechanisms are at play, a view supported by both radio and x-ray constraints on the gas density around Sgr A\*, the supermassive BH at the center of the Galaxy, the least luminous accreting BH observed to date (in Eddington units) and thus a well-studied source from the point of view of weak accretion physics [24–26]. We compute the accretion rates and the radiative efficiencies of a Galactic population of PBHs in the low-efficiency limit, following the formalism presented in Refs. [27,28]. We take into account the findings of previous studies regarding the accretion of interstellar gas onto isolated black holes [29–31].

We model the radiative efficiency  $\eta$ , defined by the relation for the bolometric luminosity  $L_B = \eta \dot{M} c^2$ , as  $\eta = 0.1 \dot{M} / \dot{M}_{\text{crit}}$  for  $\dot{M} < \dot{M}_{\text{crit}}$  (if we were to assume instead efficient accretion above the critical rate,  $\dot{M} > \dot{M}_{\text{crit}}$ , then we would have a constant  $\eta = 0.1$ ). As already discussed, all our sources fall below this critical accretion rate, such that they are all inefficient accretors: This means the luminosity scales nonlinearly with the accretion rate,  $L \propto \dot{M}^2$ .

We parameterize the accretion rate as  $\dot{M} = \lambda \dot{M}_{\text{Bondi}}$ , such that

$$\dot{M} = 4\pi\lambda(GM_{\text{BH}})^2\rho(v_{\text{BH}}^2 + c_s^2)^{-3/2}, \quad (1)$$

where  $G$  is the gravitational constant,  $v_{\text{BH}}$  is the velocity of the BH, and  $c_s$  is the sound speed of the accreted gas, which is below 1 km/s in cold, dense environments.

An important element that needs consideration is the temperature of the accreted gas due to radiative preheating [27]. Photoionizing radiation will lead to an ionization bubble surrounding the source, known as the Strömgen sphere [32], with a characteristic radius  $R_{\text{Str}}$ . In the following, we assume that the gas around the BH is fully ionized—and therefore, we set  $c_s = 10$  km/s—if the time scale for the ionization of the Strömgen sphere is shorter than the time scale associated with the incoming flux of fresh, unprocessed material.

Regarding  $\lambda$ , we choose a reference value of 0.02. Given the degeneracy between  $\lambda$  and the angular momentum and temperature of the accreted gas, this value is consistent with isolated neutron star population estimates and studies of active Galactic nuclei accretion [16,17,26].

This prescription is the same as that adopted by Ref. [28]; however, we consider  $M_{\text{BH}} = 30 M_{\odot}$  and rescale the value of  $\dot{M}_{\text{crit}} = 0.01 \dot{M}_{\text{Edd}}$  used in that work across the full 10–100  $M_{\odot}$  mass range.

We convert bolometric luminosity to x-ray luminosity via the approximate factor  $L_X \approx 0.3 L_B$  following Ref. [28].

Motivated by the results presented in Ref. [33] and by the discussion in Refs. [27,28], we assume the presence of a jet—thus requiring a system with a surplus of angular momentum or a dynamically important magnetic field

combined with a spinning black hole—emitting radio waves in the gigahertz domain with an optically thick, almost flat spectrum, while the x-ray emission is non-thermally dominated, originating from optically thin regions closer to the BH. In order to convert the x-ray luminosity into a gigahertz radio flux, we adopt the universal empirical relation discussed, e.g., in Ref. [34], also known as the *fundamental plane* (FP), which applies for a remarkably large class of compact objects of different masses, from x-ray binary systems to active Galactic nuclei. We calculate the x-ray luminosity in the 2–10 keV band in accordance with the FP, assuming a hard power-law x-ray spectrum with photon index  $\alpha$ , and a typical range for hard state x-ray binaries of 1.6–2.0 (see [35]). We extrapolate this power-law spectrum into the 0.5–8 and 10–40 keV bands in order to also make comparisons with the Chandra and NuSTAR catalogs. We then use the FP relation to calculate the 5 GHz radio flux from the 2–10 keV x-ray flux and assume a flat radio spectrum, such that  $F_{5 \text{ GHz}} = F_{1.4 \text{ GHz}}$ , allowing a direct comparison with the 1.4 GHz source catalog from a Very Large Array (VLA) survey of the GC region.

*Primordial black hole population.*—In order to derive a bound from x-ray and radio data, we set up a Monte Carlo simulation for each PBH mass, assuming a delta mass function.

We populate the Galaxy with PBHs following the Navarro-Frenk-White distribution [36] (other more conservative choices are discussed below). We implement the accurate 3D distribution of molecular, atomic, ionized gas in the inner bulge presented in Ref. [15]; that distribution includes a detailed model of the 3D structure of the central molecular zone (CMZ), a 300 pc wide region characterized by large molecular gas density and centered on the GC, i.e., in the region where the largest density of PBHs is expected.

For each PBH, the velocity is drawn randomly from a Maxwell-Boltzmann distribution. The characteristic velocity of the distribution is position dependent. The velocity distribution at a given radius is a crucial ingredient, because the accretion rate scales as  $v^{-3}$  [Eq. (1)]. In order to derive such a distribution, we consider the recent state-of-the-art model for the mass distribution in the Milky Way described in Ref. [37], where six axis-symmetric components are taken into account (bulge, DM halo, thin and thick stellar disks, and HI and molecular gas disks). We then assume that the velocity distribution at a distance  $R$  from the GC is a Maxwell-Boltzmann with  $v_{\text{mean}} = v_{\text{circ}}(R) = \sqrt{[GM(<R)/R]}$ . Under the assumption of isotropic orbits, an exact computation of the phase-space density could be performed by means of the Eddington formalism [38], as done, e.g., in Ref. [39]. (we verified that, in the high-resolution Aquarius  $N$ -body simulations, the anisotropy parameter  $\beta = 1 - \sigma_t/\sigma_r$  is consistent with 0 in the whole range of radii we are interested in; therefore, the assumption of isotropic orbits is solid.)

We checked that our simple approach is equivalent in the low-velocity tail, up to  $v \approx 40$  km/s [40]. Since our results depend only on PBHs with velocities  $\lesssim 10$  km/s (see below), we can safely neglect the high-velocity tail and adopt the simple formalism described above.

Given the mass, position, and velocity of each PBH (and the gas density), we compute the accretion rate, x-ray, and radio emission adopting the prescriptions discussed in the previous section.

*Radio BH candidates.*—The 1.4 GHz source catalog from a VLA survey of the GC region [41] contains 170 sources in a  $1^\circ \times 1^\circ$  region centered on the GC. The minimum detectable flux for this catalog is  $\sim 1$  mJy.

In order to compare our predictions to the observations, we carry out a data analysis on the VLA catalog and check if there can be any BH candidate among the detected sources.

If any of these sources are accreting BHs, their x-ray and radio emissions should be colocated. We therefore compare the radio catalog with the x-ray point source catalog from Ref. [42], which contains 9017 sources detected by Chandra in the 0.5–8 keV band in a  $2^\circ \times 0.8^\circ$  band centered on the GC, and search for all sources in both catalogs that have positions within  $10''$  of each other. (This is a very conservative separation. The positional accuracy of Chandra is  $< 1''$ . For the VLA, the positional accuracy is typically a small fraction of the synthesized beam,  $2''.4 \times 1''.3$  for the survey in Ref. [41], taken in the A configuration. A separation of  $10''$  is chosen in Ref. [41] to search for positional coincidences in other radio catalogs; we therefore also adopt  $10''$  as the maximum allowed separation).

We find 24 sources in both the x-ray and radio catalogs within  $10''$  of each other. If we assume that these sources are accreting BHs, then their x-ray and radio fluxes should lie on the FP, as explained above. So, we use the FP (considering masses from 10 to  $100 M_\odot$ ) to predict the x-ray flux from the radio flux of each of these objects (24 in the very conservative case, nine if we exclude likely foreground sources).

We find that the predicted x-ray fluxes are substantially larger ( $\sim 3$ – $7$  orders of magnitude) than the flux reported in the catalog from Ref. [42] in the whole mass range we consider. We therefore conclude that none of the 24 (or nine likely Galactic) VLA sources with overlapping positions lie on the FP, and therefore, given the assumptions described above regarding the presence of a jet, we have no BH candidate in our sample.

*X-ray BH candidates.*—Hard x-ray emission ( $> 10$  keV) suffers from far less Galactic absorption than soft x-ray emission and is therefore a good band to search for emission from accreting BHs.

We consider sources in the Chandra catalog [42] in the 0.5–8 keV band and those detected by NuSTAR in the 10–40 keV band [43]. For Chandra (NuSTAR), we consider a small region of interest (ROI) including the high-density region of the Galactic Ridge:  $-0.9^\circ < l < 0.7^\circ$ ;  $-0.3^\circ < b < 0.3^\circ$  ( $-0.9^\circ < l < 0.3^\circ$ ;  $-0.1^\circ < b < 0.4^\circ$ ). There are

483 likely Galactic x-ray sources in the Chandra catalog above a flux threshold of  $2 \times 10^{-6}$  ph cm $^{-2}$  s $^{-1}$  and 70 NuSTAR sources. (“Likely Galactic” sources are defined in Ref. [42] based on their hardness ratios. The exposure across the Chandra survey region is variable, and the flux threshold used here is a compromise between maximizing the ROI and the completeness, per Ref. [42].) Since in all cases the corresponding radio flux predicted with the FP would be 3–7 orders of magnitude below the detection threshold of the VLA survey in Ref. [41], we cannot draw any conclusions on the nature of these x-ray sources. Therefore, we consider all of them in our analysis as potential BH candidates (we remove only  $\sim 40\%$  of the detected NuSTAR sources that are thought to be cataclysmic variables [35]).

*Results.*—The main result of the Letter is presented in Fig. 1. We display the  $2\sigma$ ,  $3\sigma$ , and  $5\sigma$  constraints on the DM fraction as a function of the PBH mass.

The upper limits are derived as follows. We perform  $\mathcal{O}(100)$  Monte Carlo simulations for ten reference values of the mass in the 10– $100 M_\odot$  interval, assuming a DM fraction  $f_{\text{DM}} = 1$ . We determine the mean and standard deviation of the distributions of the predicted number of PBHs with radio fluxes above the VLA threshold and with x-ray fluxes exceeding the Chandra (NuSTAR) threshold, in the corresponding ROIs. We verify that the number of bright PBHs is compatible with Poisson statistics and the average predicted number scales linearly with  $f_{\text{DM}}$ . We derive the radio and x-ray bounds by comparing the number of predicted PBHs with the number of BH candidates derived from the analysis of radio and x-ray catalogs described in the previous section. For the x-ray bound, we show the result obtained with the more sensitive Chandra catalog. The NuSTAR bound is slightly weaker: It allows us to exclude at  $2\sigma$  values of  $f_{\text{DM}}$  as low as 0.4 (for  $30 M_\odot$ ).

In Fig. 2, we show the PBHs detectable by VLA at 1.4 GHz assuming a PBH mass of  $30 M_\odot$  and a DM




FIG. 2. Example of the distribution of  $30 M_\odot$  PBHs detectable by VLA in the ROI, for one Monte Carlo realization. The colored background depicts the column gas density. The size of the black points is proportional to the PBH velocity in the range 0.3–3 km/s (for detectable PBHs).

fraction equal to 1, for one specific Monte Carlo realization. This scenario predicts, on average,  $40 \pm 6$  sources above the VLA flux threshold for  $30 M_{\odot}$ , and, thus, it is excluded by more than  $5\sigma$  from radio observations. However, it is important to understand where the constraining power comes from: The PBHs above the detection threshold, and thus the ones with the larger x-ray flux, lie in the very inner region of the Galaxy where the column gas density is the highest and show very small velocities, in the range  $\sim 0.3\text{--}3$  km/s. Therefore, the constraints arise from the very-low-velocity tail of the distribution and from regions correlated with very high column densities, e.g., CMZ, as already mentioned above.

*Discussion and conclusions.*—In this Letter, we derive new, strong constraints on the hypothesis that PBHs comprise all of the DM in the Universe. In particular, we find that PBHs with  $M \approx 30 M_{\odot}$ , that could be responsible for the gravitational waves detected by LIGO, contribute less than 20% to the whole DM density.

In the mass window  $10\text{--}100 M_{\odot}$ , our constraints are competitive with (and even stronger than) those arising from the study of microlensing events with the massive compact halo object project [44] (for  $\gtrsim 15 M_{\odot}$ ) and with those from halo wide binaries [45,46] (for  $\gtrsim 60 M_{\odot}$ ). For  $M \gtrsim 10 M_{\odot}$ , they are also comparable to or stronger than the constraints from the survival of central star clusters in faint dwarf galaxies, in particular, in Eridanus II [47,48]. Even more stringent constraints arise, in principle, from the analysis of the cosmic microwave background (CMB) [49]. However, those arising from the analysis of spectral distortions (based on Far-Infrared Absolute Spectrophotometer data) turned out to be much weaker than originally thought [50], while the ones based on the study of CMB anisotropies (see also the recent results by Ref. [51]) are based on assumptions on the accretion of gas on PBHs in the early Universe that are still under debate, as the modeling of the accretion process is based on theoretical arguments and not directly supported by observations as in our case (see also the discussion in Ref. [50]).

In contrast with Ref. [49], in fact, we adopt a very conservative prescription, compatible with current astronomical observations, for both the accretion rate and the radiative efficiency, setting the ratio of the actual accretion rate to the Bondi rate  $\lambda$  equal to 0.02. We remark that  $\lambda$  probably follows some distribution and is also likely degenerate with  $c_S$  and  $v_{\text{BH}}$ —future studies are required to disentangle these. Moreover, we exploit for the first time in this context the empirical FP relation between radio and x-ray emission, which has been observed on a wide class of sources in a large mass range, from x-ray binaries to active Galactic nuclei. By adopting such a relation, we are able to predict the expected radio and x-ray luminosities of a population of PBHs in the Galaxy compatible with the DM phase-space distribution, as well as to look for BH candidates in radio and x-ray catalogs. We set upper limits

on the DM PBH fraction using both radio (VLA) and x-ray (Chandra and NuSTAR) pointlike source catalogs, by comparing the number of expected PBHs above observational thresholds and the observed number of BH candidates in a very narrow region about the GC.

These bounds are robust with respect to the modeling of the full velocity distribution, since the predicted number of bright PBHs depends only on the very-low-velocity tail ( $<10$  km/s) where we checked the agreement among different numerical or analytical methods. Moreover, our limits are independent of the details of the gas distribution (we checked that the bound is still present even with a naive modeling of the CMZ as a sphere with uniform density compatible with the mass constraints provided in [15]). They are also not affected significantly by a shallower DM profile as proposed, e.g., in Ref. [52]; however, assuming an even flatter profile like the Burkert one (an extremely conservative assumption for our Galaxy), the bound is present only under the assumption of Bondi accretion.

We recall that our limits hold for a narrow mass function; a detailed study of the impact of different mass distributions is beyond the scope of the present Letter and postponed to a future work.

Although our radio and x-ray bounds vanish for  $\lambda \lesssim 10^{-2}$ , future instruments will be able to verify better the accretion model as well as the PBH DM fraction. In particular, given the significant increase in the sensitivity of future radio telescopes, we expect an important part of the yet-allowed parameter space to be probed by upcoming facilities such as MeerKAT and, later, square kilometer array (SKA). Using the radiometer equation [53], the minimum ( $1\sigma$ ) detectable radio flux is  $S_{\nu,\text{rms}} = (T_{\text{sky}} + T_{\text{rx}})/(G\sqrt{2T_{\text{obs}}\Delta\nu})$ . For SKA1-MID (1.4 GHz), we assume gain  $G = 15$  K/Jy, receiver temperature  $T_{\text{rx}} = 25$  K, sky temperature towards the GC  $T_{\text{sky}} = 70$  K, and bandwidth  $\Delta\nu = 770$  MHz [54].

For a one-hour exposure, the instrumental detection sensitivity of SKA1-MID turns out to be  $\sim 2.7 \mu\text{Jy}$  (significantly above the source confusion limit), which would give  $\mathcal{O}(2000)$  detectable PBHs for our reference value  $\lambda = 0.02$  ( $f_{\text{DM}} = 1$  and  $M = 30 M_{\odot}$ ).

SKA will therefore be able to either place a very strong constraint in the absence of BH candidate detection or detect a subdominant population of PBHs (although the expected population of astrophysical BHs becomes comparable with the primordial one for DM fractions lower than  $\sim 10^{-3}$ ). With an even longer exposure ( $\approx 1000$  h, 85 nJy sensitivity), achievable for dedicated deep field continuum observations, such strong constraints can be placed by SKA even under the assumption of extremely low values of  $\lambda$ ,  $\mathcal{O}(10^{-3})$ .

Interestingly, our procedure can also be applied in order to extend work on searches for astrophysical BHs in the Galaxy [27,28,31], adopting the realistic spatial and velocity distributions expected for those objects. Our formalism

has the potential to characterize this guaranteed population of objects in future analyses.

We are indebted to M. Fornasa for verifying of the phase-space velocity at low radii with the Eddington formalism. We thank P. D. Serpico for his useful comments on the manuscript and D. Baumann, P. Crumley, B. Freivogel, A. King, A. Urbano, J. Vink, and C. Weniger for useful discussions. Finally, we thank the anonymous reviewers for their insightful comments and suggestions. R. C. is supported by the Netherlands Research School for Astronomy (NOVA). S. M. acknowledges support from the Netherlands Organization for Scientific Research (NWO) VICI Grant No. 639.043.513. E. S. is supported by the Netherlands Organization for Scientific Research (NWO) through a Vidi grant (PI: C. Weniger).

\*d.gaggero@uva.nl

- [1] B. P. Abbott *et al.* (Virgo, LIGO Scientific Collaboration), *Phys. Rev. Lett.* **116**, 061102 (2016).
- [2] S. Bird, I. Cholis, J. B. Munoz, Y. Ali-Haimoud, M. Kamionkowski, E. D. Kovetz, A. Raccanelli, and A. G. Riess, *Phys. Rev. Lett.* **116**, 201301 (2016).
- [3] S. Clesse and J. García-Bellido, *Phys. Dark Universe* **15**, 142 (2017).
- [4] G. Jungman, M. Kamionkowski, and K. Griest, *Phys. Rep.* **267**, 195 (1996).
- [5] G. Bertone, D. Hooper, and J. Silk, *Phys. Rep.* **405**, 279 (2005).
- [6] G. Bertone, *Nature (London)* **468**, 389 (2010).
- [7] P. Ivanov, P. Naselsky, and I. Novikov, *Phys. Rev. D* **50**, 7173 (1994).
- [8] M. Yu. Khlopov, *Res. Astron. Astrophys.* **10**, 495 (2010).
- [9] B. J. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, *Phys. Rev. D* **81**, 104019 (2010).
- [10] D. Blais, C. Kiefer, and D. Polarski, *Phys. Lett. B* **535**, 11 (2002).
- [11] N. Afshordi, P. McDonald, and D. N. Spergel, *Astrophys. J.* **594**, L71 (2003).
- [12] P. H. Frampton, M. Kawasaki, F. Takahashi, and T. T. Yanagida, *J. Cosmol. Astropart. Phys.* **04** (2010) 023.
- [13] B. Carr, F. Kuhnel, and M. Sandstad, *Phys. Rev. D* **94**, 083504 (2016).
- [14] M. Portail, C. Wegg, and O. Gerhard, *Mon. Not. R. Astron. Soc.* **450**, L66 (2015).
- [15] K. Ferrière, W. Gillard, and P. Jean, *Astron. Astrophys.* **467**, 611 (2007).
- [16] R. Perna, J. McDowell, K. Menou, J. Raymond, and M. V. Medvedev, *Astrophys. J.* **598**, 545 (2003).
- [17] S. Pellegrini, *Astrophys. J.* **624**, 155 (2005).
- [18] F. Hoyle and R. A. Lyttleton, *Proc. Cambridge Philos. Soc.* **35**, 405 (1939).
- [19] H. Bondi and F. Hoyle, *Mon. Not. R. Astron. Soc.* **104**, 273 (1944).
- [20] S. Heinz and R. A. Sunyaev, *Mon. Not. R. Astron. Soc.* **343**, L59 (2003).
- [21] R. Narayan and I. Yi, *Astrophys. J. Lett.* **428**, L13 (1994).
- [22] R. D. Blandford and M. C. Begelman, *Mon. Not. R. Astron. Soc.* **303**, L1 (1999).
- [23] E. Quataert and A. Gruzinov, *Astrophys. J.* **539**, 809 (2000).
- [24] G. C. Bower, M. C. H. Wright, H. Falcke, and D. C. Backer, *Astrophys. J.* **588**, 331 (2003).
- [25] D. P. Marrone, J. M. Moran, J.-H. Zhao, and R. Rao, *Astrophys. J. Lett.* **654**, L57 (2007).
- [26] Q. D. Wang, M. A. Nowak, S. B. Markoff, F. K. Baganoff, S. Nayakshin, F. Yuan, J. Cuadra, J. Davis, J. Dexter, A. C. Fabian, N. Grosso, D. Haggard, J. Houck, L. Ji, Z. Li, J. Neilsen, D. Porquet, F. Ripple, and R. V. Shcherbakov, *Science* **341**, 981 (2013).
- [27] T. J. Maccarone, *Mon. Not. R. Astron. Soc.* **360**, L30 (2005).
- [28] R. P. Fender, T. J. Maccarone, and I. Heywood, *Mon. Not. R. Astron. Soc.* **430**, 1538 (2013).
- [29] Y. Fujita, S. Inoue, T. Nakamura, T. Manmoto, and K. E. Nakamura, *Astrophys. J.* **495**, L85 (1998).
- [30] P. J. Armitage and P. Natarajan, *Astrophys. J. Lett.* **523**, L7 (1999).
- [31] E. Agol and M. Kamionkowski, *Mon. Not. R. Astron. Soc.* **334**, 553 (2002).
- [32] B. Strömberg, *Astrophys. J.* **89**, 526 (1939).
- [33] R. P. Fender, *Mon. Not. R. Astron. Soc.* **322**, 31 (2001).
- [34] R. M. Plotkin, S. Markoff, B. C. Kelly, E. Körding, and S. F. Anderson, *Mon. Not. R. Astron. Soc.* **419**, 267 (2012).
- [35] J. Hong *et al.*, *Astrophys. J.* **825**, 132 (2016).
- [36] J. F. Navarro, C. S. Frenk, and S. D. M. White, *Astrophys. J.* **462**, 563 (1996).
- [37] P. J. McMillan, *Mon. Not. R. Astron. Soc.* **465**, 76 (2017).
- [38] A. S. Eddington, *Mon. Not. R. Astron. Soc.* **76**, 572 (1916).
- [39] M. Fornasa and A. M. Green, *Phys. Rev. D* **89**, 063531 (2014).
- [40] M. Fornasa (private communication).
- [41] T. J. W. Lazio and J. M. Cordes, *Astrophys. J. Suppl. Ser.* **174**, 481 (2008).
- [42] M. P. Muno *et al.*, *Astrophys. J. Suppl. Ser.* **181**, 110 (2009).
- [43] F. A. Harrison *et al.*, *Astrophys. J.* **770**, 103 (2013).
- [44] C. Alcock *et al.*, *Astrophys. J. Lett.* **550**, L169 (2001).
- [45] D. P. Quinn, M. I. Wilkinson, M. J. Irwin, J. Marshall, A. Koch, and V. Belokurov, *Mon. Not. R. Astron. Soc.* **396**, L11 (2009).
- [46] M. A. Monroy-Rodríguez and C. Allen, *Astrophys. J.* **790**, 159 (2014).
- [47] T. D. Brandt, *Astrophys. J.* **824**, L31 (2016).
- [48] T. S. Li *et al.*, *Astrophys. J.* **838**, 8 (2017).
- [49] M. Ricotti, J. P. Ostriker, and K. J. Mack, *Astrophys. J.* **680**, 829 (2008).
- [50] S. Clesse and J. García-Bellido, arXiv:1610.08479.
- [51] L. Chen, Q.-G. Huang, and K. Wang, *J. Cosmol. Astropart. Phys.* **12** (2016) 044.
- [52] F. Calore, N. Bozorgnia, M. Lovell, G. Bertone, M. Schaller, C. S. Frenk, R. A. Crain, J. Schaye, T. Theuns, and J. W. Trayford, *J. Cosmol. Astropart. Phys.* **12** (2015) 053.
- [53] R. Dewey, G. Stokes, D. Segelstein, J. Taylor, and J. Weisberg, in *Birth and Evolution of Neutron Stars: Issues Raised by Millisecond Pulsars*, edited by S. P. Reynolds and D. R. Stinebring (National Radio Astronomy Observatory, Green Bank, 1984), p. 234.
- [54] F. Calore, M. Di Mauro, F. Donato, J. W. T. Hessels, and C. Weniger, *Astrophys. J.* **827**, 143 (2016).