



UvA-DARE (Digital Academic Repository)

The outcome of institutional youth care compared to non-institutional youth care for children of primary school age and early adolescence

A multi-level meta-analysis

Strijbosch, E.L.L.; Huijs, J.A.M.; Stams, G.J.J.M.; Wissink, I.B.; van der Helm, G.H.P.; de Swart, J.J.W.; van der Veen, Z.

Published in:

Children and Youth Services Review

DOI:

[10.1016/j.chidyouth.2015.09.018](https://doi.org/10.1016/j.chidyouth.2015.09.018)

[Link to publication](#)

Citation for published version (APA):

Strijbosch, E. L. L., Huijs, J. A. M., Stams, G. J. J. M., Wissink, I. B., van der Helm, G. H. P., de Swart, J. J. W., & van der Veen, Z. (2015). The outcome of institutional youth care compared to non-institutional youth care for children of primary school age and early adolescence: A multi-level meta-analysis. *Children and Youth Services Review*, 58, 208-218. <https://doi.org/10.1016/j.chidyouth.2015.09.018>


General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<http://dare.uva.nl>)



The outcome of institutional youth care compared to non-institutional youth care for children of primary school age and early adolescence: A multi-level meta-analysis



E.L.L. Strijbosch^{a,*}, J.A.M. Huijs^b, G.J.J.M. Stams^b, I.B. Wissink^b, G.H.P. van der Helm^c, J.J.W. de Swart^d, Z. van der Veen^b

^a Juzt (Quality and Innovation unit), Erasmusweg 34, 4834 AA Breda, The Netherlands

^b University of Amsterdam (Department of Forensic Child and Youth Care Sciences), Nieuwe Achtergracht 127, 1018 WS Amsterdam, The Netherlands

^c Leiden University of Applied Sciences (Youth Expert Centre), Zernikedreef 11, 2333 CK Leiden, The Netherlands

^d Windesheim University of Applied Sciences, Campus 2-6, 8017 CA Zwolle, The Netherlands

ARTICLE INFO

Article history:

Received 1 December 2014

Received in revised form 20 September 2015

Accepted 20 September 2015

Available online 25 September 2015

Keywords:

Institutional youth care

Outcome

Meta-analysis

Primary school age

Early adolescence

Evidence-based treatment

ABSTRACT

Objective: The outcome of institutional youth care for children is heavily debated. This multilevel meta-analysis aims to address the outcome of institutional youth care compared to non-institutional youth care for children of primary school age and early adolescence in economically developed countries. A gain of knowledge in this area may help the decision for referral of children to institutional youth care or other types of care (e.g., foster care or community-based care), and improve outcomes for children in youth care.

Methods: Of 19 controlled studies (15,526 participants), 63 effect sizes of behaviour problems (externalizing, internalizing, and total), skills (social and cognitive) and delinquency were computed based on comparisons between institutional Evidence-Based Treatment (EBT), institutional Care As Usual (CAU), non-institutional EBT, and non-institutional CAU.

Results: Institutional CAU showed a small-to-medium negative significant effect compared to non-institutional CAU ($d = -0.342$). Furthermore, children in institutional care showed slightly more delinquent behaviour compared to children in non-institutional care ($d = -0.329$). Significant moderating effects were also found for study design, year of publication and sex of the child.

Conclusions: Children receiving non-institutional CAU (mostly foster care) had slightly better outcomes than children in institutional CAU (regular group care). No differences were found between institutional and non-institutional care when institutional treatment was evidence-based. More research is needed on the conditions that make established treatment methods work in institutional care for (young) children.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Under the United Nations Convention on the Rights of the Child, children have the right to grow up in a stable and safe environment where they receive the warmth and support they need for their development (Children's Rights Alliance, 2010; Höfte, Van der Helm, & Stams, 2012; United Nations, 1989). Unfortunately, not every parent is able to offer a stable and secure home, and some children have to live in foster care or institutional youth care (Manso, García-Baamonde, Alonso, & Barona, 2011).

There is an ongoing debate about the appropriateness of institutional care for children, (Chance, Dickson, Bennett, & Stone, 2010; Dozier et al., 2014; Souverein, Van der Helm, & Stams, 2013). Mainly since the last decennium, but beginning in the 1980s, there has been a shift from institutional towards community based care, and a change from a deficit-focused to a strength-focused approach, in particular building on family strengths and resources (Knapp, 2006; Kumpfer & Alvarado, 2003; Leichtman, 2008; Lonne, Parton, Thomson, & Harries, 2009; Lösel & Farrington, 2012; Melkman, 2015; Weick, Rapp, Sullivan, & Kisthardt, 1989). From this perspective, a growing number of (evidence-based) treatment alternatives have been developed, such as Multi-Systemic Therapy (MST; Henggeler, Pickrel & Brondino, 1999b; Van der Stouwe, Asscher, Stams, Dekovic, & Van der Laan, 2014) and Multidimensional Treatment Foster Care (MTFC; Chamberlain & Reid, 1998). However, the appropriateness of institutional youth care compared to non-institutional youth care should still be judged taking the type and severity of the problems of children into account

* Corresponding author at: Researcher at the Quality and Innovation unit at Juzt, Erasmusweg 34, 4834 AA Breda, The Netherlands.

E-mail addresses: e.strijbosch@juzt.nl (E.L.L. Strijbosch), g.j.j.m.stams@uva.nl (G.J.J.M. Stams), i.b.wissink@uva.nl (I.B. Wissink), helm.vd.p@hsleiden.nl (G.H.P. van der Helm).

(Andrews & Bonta, 2010; Souverein et al., 2013) as well as the children's age.

Khoo, Skoog, and Dalin (2012) pointed out that, whereas adolescents are often referred because of their own behaviour problems and delinquency, children are often brought to the attention of social services because of parents' shortcomings and problems in the home. These problems are often interwoven with serious emotional and behavioural disorders that interfere with children's development and their ability to function at home, in school and in their communities (Benzies, Harrison, & Magill-Evans, 2004; Linville et al., 2010; Raine, Brennan, Mednick, & Mednick, 1996). Many studies addressing outcomes of institutional youth care versus other care forms have focused on (late) adolescents. Especially within the current timeframe of de-institutionalization, it is important to also pay close attention to the outcomes of institutional youth care for children and young adolescents. In the next paragraphs, we provide an overview of research on institutional versus non-institutional youth care and evidence-based treatment (EBT) versus care as usual (CAU). In this article we consider 'treatment' as particular behavioural interventions targeting problems that hamper adaptive functioning (James, 2011). Evidence-based treatment refers to structured and often manualized interventions based on empirically supported theories about what causes and maintains problems, which have been proven to be effective (to some degree) in (quasi-) experimental research (Chorpita et al., 2013; Wampold, Goodheart, & Levant, 2007; Weisz et al., 2013a).

1.1. Institutional and non-institutional care for children

Since young children are extremely vulnerable and develop rapidly at the physical, emotional and cognitive level, treatment in a family or family-like environment (i.e., non-institutional care) is usually preferred over institutional care (Dozier et al., 2014). Available evidence-based treatment methods (non-institutional EBT), such as Functional Family Therapy (Alexander, Pugh, Parsons, & Sexton, 2000), Multi-Systemic Therapy (MST; Henggeler et al., 1999b) and several kinds of behavioural parent training (for an overview, see Weisz et al., 2013a) focus on care assistance and treatment at the youth's home and community locations, such as the school and contexts involving structured and unstructured free time activities (Dijkstra, Creemers, Asscher, Dekovic, & Stams, 2015; Pennell & Burford, 2000). Weisz et al. (2013a) performed a meta-analysis based on research within the last four decades and found that non-institutional evidence-based treatment for psychopathology in children and adolescents outperformed non-institutional usual care, but the advantages proved to be modest, and moderated by youth, location and assessment characteristics. Non-institutional CAU mostly includes non-structured and non-manualized treatment, (intensive) case management, several forms of foster care with or without the involvement of professional support, and counselling. Non-institutional CAU might also include interventions that are promising from a theoretical perspective, such as Family Group Conferencing, but still lack sufficient empirical support (Asscher, Dijkstra, Stams, Dekovic, & Creemers, 2014; Crampton, 2007; Dijkstra, Creemers, Asscher, & Stams, 2014; Frost, Abram, & Burgess, 2014).

Although a family-environment is preferred for every child, non-institutional community-based treatment and/or living in a foster home can be extremely difficult for children showing persistent aggressive and anti-social behaviour, with a risk for frequent placement disruptions (Dekker, Van Miert, Roest, & Van der Helm, 2012; Jakobsen, 2013; Oosterman, Schuengel, Slot, Bullens, & Doreleijers, 2007; Van Oijen, 2010). The prevalence of placement instability should not be underestimated; it can aggravate emotional and behavioural problems (Barber & Delfabbro, 2003; Hussey & Guo, 2005; James, Landsverk, Slymen, & Leslie, 2004; Rubin, Alessandrini, Feudtner, Localio & Hadley, 2004; Ryan & Testa, 2005).

Compared to children in non-institutional care, children in institutional care show more aggressive behaviour, and have more often been diagnosed with oppositional defiant disorder or conduct disorder (Handwerk, Field, & Friman, 2001; Lee & Thompson, 2007; Vermaes & Nijhof, 2014). Recent studies show that the severe behaviour problems can be associated with abnormal brain development as a result of neglect and traumatization (Fairchild et al., 2013; Raine, 2013). Providing the right treatment for children in institutional care is therefore very complex. Besides, living in an institutional setting can in itself have a negative or positive impact on the development of children (Dunn, Culhane, & Taussig, 2010; Preyde, Adams, Cameron, & Frensch, 2009). For example, as a result of the separation from their parents, children may develop internalizing problems (White & King, 2011), externalizing problems (Van der Helm, Stams, & Van der Laan, 2011) and attachment problems (Johnson, Browne, & Hamilton-Giachritsis, 2006; Van den Bergh, Weterings, & Schoenmakers, 2011; Van den Dries, Juffer, Van IJzendoorn, & Bakermans-Kranenburg, 2009). Also, negative peer influences, such as 'deviancy training' (Dishion, McCord, & Poulin, 1999), can affect the development of children in institutional care. Children's aggressive behaviour can trigger coercive behaviour in professionals, with a detrimental effect on the living group climate. The institutional setting can, on the other hand, also provide the safety and protection children coming from harmful circumstances need. For a discussion on the negative and positive consequences of institutional youth care, see Souverein et al. (2013).

There is little consensus in the literature about the effectiveness and appropriateness of institutional youth care compared to non-institutional youth care, and how the above-mentioned problems are being addressed (Preyde et al., 2011; Souverein et al., 2013). In particular the long-term outcomes for children and adolescents living in institutional youth care have been questioned (Chor, McClelland, Weiner, Jordan, & Lyons, 2012; Dregan & Gulliford, 2012; Frensch & Cameron, 2002). There are some studies that indicate positive outcomes, but they are mostly based on small samples, and control groups are often missing (Bean, White, & Lake, 2005). Some pre-experimental studies showed a reduction of behavioural and emotional problems after treatment in institutional youth care (Larzelere, Daly, Davis, Chmelka, & Handwerk, 2004; Leichtman, Leichtman, Barber, & Neese, 2001). More recently, Dregan, Brown and Armstrong (2011) have investigated the effectiveness of institutional youth care and foster care, and showed that children in both conditions were at increased risk of behavioural and emotional problems in adulthood. Relatively better outcomes were related to the involvement of families during placement, e.g., by offering family therapy (Chance et al., 2010; Schubert, Mulvey, Loughran, & Losoya, 2012). Also a short length of stay, a positive living group climate, aftercare services and minimizing placement instability were important factors associated with better outcomes in institutional youth care (Hoagwood & Cunningham, 1993; Khoo et al., 2012; Schubert et al., 2012).

Many studies are less positive about institutional youth care. Negative peer influences are often mentioned (Aguilar-Vafaie, Roshani, Hassanabadi, Masoudian, & Afruz, 2011; Orobio de Castro, Merk, Koops, Veerman, & Bosch, 2005; Whitehead, Keshet, Lombrowski, Domenico, & Green, 2007). Whitehead et al. (2007) maintained that institutional youth care focuses too much on the child itself instead of on the entire child system (peers, school, and parents). Additionally, Manso et al. (2011) showed that many children in institutional care do not only have problems with their personal and social functioning, but also have educational problems. Dregan and Gulliford (2012) concluded that children in institutional care develop less favourably compared to children in foster care. As a possible explanation for this result, they mentioned that foster care provides more positive care experiences because it is a relatively stable placement with early admission to care and, as opposed to institutional care, a limited number of different caregivers. Their study did not adjust for pre-care characteristics. As some studies indicate (Barth,

Greeson, Green, Hurley, & Sisson, 2007; Berger, Bruch, Johnson, James, & Rubin, 2009; Vermaes & Nijhof, 2014), children who are most disturbed and in need of specialized treatment are often not selected for foster care.

1.2. Institutional care for children: EBT versus CAU

Even though research findings on the effectiveness of institutional versus non-institutional care are not consistent, it can be assumed that institutional care will be needed for a certain group of children, and it is therefore important to address the differences between several kinds of institutional care. Institutional youth care is carried out in 24-hour care facilities for children and youth with emotional and behavioural problems (Boendermaker, Van Rooijen, & Berg, 2012; Lee & Thompson, 2007). These facilities aim to provide a safe and structured environment in which the child receives either regular (long-term) care to be able to grow up in a sound manner (institutional CAU), or regular care combined with (short-term) specific child- and family-centred treatments from multidisciplinary teams that create individual care or treatment plans for each child (Preyde et al., 2009), that is, institutional EBT. Institutional EBT contains group as well as individual treatment, which is mostly based on behavioural, cognitive and solution focused treatment models (Rose, 2014; Van der Helm & Hanrath, 2011; Whittaker, Del Valle, & Holmes, 2015). For a recent overview of evidence-based treatment methods applied in institutional settings, see James, Alemi, and Zepeda (2013). Examples of group based programs are EQUIP (Van Stam et al., 2014) and Re-Art (Hoogsteder et al., 2014). In many economically developed countries the services provided by the institutions are defined, regulated and (governmentally) monitored on the basis of guidelines, principles and standardized assessment of performances (Behar, Friedman, Pinto, Katz-Leavy, & Jones, 2007; Gudbrandsson, 2008).

In their meta-analysis on the effectiveness and implementation of evidence-based practices in institutional care settings, James et al. (2013) demonstrated that EBT can be implemented and tested within the context of institutional care, and notwithstanding the limitations within the underlying studies, they found overall encouraging outcomes that indicated improvements in multiple domains of functioning. However, this meta-analysis mainly concerned (late) adolescents, and may not be representative for children of primary school age and early adolescence.

1.3. Selection factors

As mentioned earlier, the type and severity of the problems, as well as the age of the child may determine whether a child is referred to institutional or non-institutional care. Youths who are placed out of their family home tend to display more problems, fewer strengths, and more risk factors than youth who remain at home (Farmer, Mustillo, Burns, & Holden, 2008). Risk factors may pertain to child maltreatment, persistent juvenile delinquency or both. Notably, Asscher, Van der Put, and Stams (in press) showed that many juvenile delinquents, in particular girls, have a history of child maltreatment. Stith et al. (2009), in their meta-analytic review, found that parent characteristics (parent anger/hyper-reactivity, parent perceiving the child as a problem, parent's level of stress, parent self-esteem) and family factors (high family conflict, low family cohesion, low parent-child relationship quality) were most predictive of physical abuse and neglect. Assink et al. (2015) conducted a meta-analytic review on risk factors for persistent juvenile delinquency, and they found relatively large effects for the criminal history, aggressive behaviour, and alcohol/drug abuse domains, and small effects for the family, neurocognitive and attitude domains. The physical health, background and neighbourhood domains did not yield significant effects. Farmer, Southerland, Mustillo, and Burns (2009) showed that the likelihood of moving back home after institutional care was related to race, age, gender, family income and total

child behaviour problems. Decreased strengths and more child-level risk factors predicted instability of reunification. Generally, the accumulation of risk factors may result in less positive outcomes (Evans, Li, & Whipple, 2013). To summarize, parent and family factors as well as behavioural problems, skills and delinquency may explain referral to institutional or non-institutional care. The extent to which the (dynamic) risk factors are diminished during treatment, as well as certain (static) background variables, may influence the chance of moving back home successfully.

1.4. Previous meta-analyses addressing institutional versus non-institutional care

Although there is a vast body of research examining the outcomes of institutional care for children and adolescents, it is still unclear whether institutional youth care yields better or worse outcomes compared to non-institutional youth care (Preyde et al., 2009). Institutional youth care is very costly (Frensch & Cameron, 2002), and with a shift towards more community and strengths based care, it is important to investigate outcomes of institutional youth care compared to alternatives, by means of meta-analytic reviews (James, 2011; Ziviani, Feeney, Cuskelly, Meredith, & Hunt, 2012).

Recently two meta-analytic studies have been performed comparing institutional youth care to non-institutional youth care. Van den Dries et al. (2009) focused on children under the age of four and found that, compared to children raised in institutions, (early) adopted children had more secure attachment relationships with their caregivers. De Swart et al. (2012) examined outcomes of institutional youth care over the past three decades for youth and young adults up to 23 years of age. In this study, it was concluded that institutional youth care can be equally effective as non-institutional youth care. Furthermore, De Swart et al. (2012) showed that institutional evidence-based treatment yielded better outcomes than institutional care as usual.

The study of De Swart et al. (2012) focused on a broad age range, and did not differentiate between several kinds of outcomes within the studies. For several reasons, which are outlined in the next paragraph, it is important to address outcomes of institutional care compared to non-institutional care for primary school age children and early adolescents in a separate meta-analytic study. This will add to existing knowledge about the effects of institutional youth care, and hopefully assist those who refer children to institutional care or other types of care.

1.5. The present study

The present study examines the outcomes of institutional youth care compared to non-institutional care and the outcomes of institutional evidence-based treatment (EBT) compared to institutional care as usual (CAU) for primary school age children and early adolescents. This meta-analysis is based on the meta-analysis of De Swart et al. (2012), but there are two main differences. First, the present meta-analysis focuses on the younger age group (children mainly in the range from 4–14 years, with a maximum of 17 years), whereas De Swart et al. (2012) primarily focused on adolescents and young adults (mainly in the range from 14–18 years, with a maximum age of 23). These different age groups have different cognitive abilities and developmental tasks and are subject to different socialization practices. Children and young adolescents are thought to be more vulnerable to the separation from their parents, whereas (late) adolescents regularly have formed their own identity and have attained more independence from their parents (De Wit, Slot, & van Aken, 2013). Because of these specific differences, it is important to increase the knowledge on effectiveness of institutional care in the younger age group, and be able to distinguish between different outcome measures. The second main difference from De Swart et al. (2012) is that the present study uses multilevel techniques in order to be able to include more effect sizes of the same study, increasing statistical power and allowing the

examination of more moderators than can be achieved in a standard meta-analysis.

The purpose of this multilevel meta-analysis is to examine the effects of institutional youth care on behaviour problems (externalizing, internalizing and total), skills (social and cognitive) and delinquency in primary school age children and early adolescents. To accomplish this, four comparisons have been made. First, institutional Evidence-Based Treatment (EBT), i.e., structured (individual and group) treatment based on theoretical and empirical evidence (mostly behavioural, cognitive and solution focused therapies), is compared to institutional Care As Usual (CAU), i.e., group care offering daily care and structure, mostly in a (psychiatric) living group setting. Second, institutional EBT is compared to non-institutional EBT, such as Functional Family Therapy. Third, institutional CAU is compared to non-institutional EBT. Fourth, institutional CAU is compared to non-institutional CAU, mostly foster care. In addition, the following moderators were also examined because of their possible relevance for the interpretation of the outcomes: year of publication, journal impact factor, study quality, study design, time of measurement, type of intervention, data source, sex, mean age, percentage of girls, target group, ethnicity and control for pre-test differences in outcomes between the experimental and control group.

2. Methods

2.1. Study selection

Three search methods were used to collect the studies. First, we collected the studies with children and youth between 4 and 17 years old that De Swart et al. (2012) included in their meta-analysis. Second, we searched for studies on a broad domain of institutional youth care in the period from 1970 to 2013 in electronic databases: ScienceDirect, PsychInfo, Picarta, Springerlink, ERIC, Medline and Google Scholar. We used the following keywords in various combinations: residential care, institutional care, group care, foster care, child, youth, comparison and effectiveness. Finally, we searched in the reference lists of all eligible studies. Two researchers applied the search strategy independently of each other.

2.2. Inclusion criteria

Studies were included if (1) the (quasi-) experimental group received institutional EBT or institutional CAU; (2) the control group received institutional CAU, non-institutional EBT, or non-institutional CAU; (3) the average age of the children was under 15, and the total age range fell between 4 and 17 years; (4) studies provided at least post-test scores or follow-up scores in order to be able calculate effect sizes for differences between the experimental and control group; and (5) studies were published.

The search led to a total of 9 new studies that had not been included in the De Swart et al., 2012 meta-analysis. From the meta-analysis of De Swart et al., 15 studies were selected. All 24 studies were read by three researchers. In several meetings the researchers reached consensus on which of the studies would be included. Out of the 9 new studies, 4 could be included in the meta-analysis. Reasons for exclusion were mainly that the maximum age of the total population was too high, or that the study compared different groups but did not have a control group. This strategy finally resulted in a total of 19 studies ($N = 15,526$ children and youth) meeting the inclusion criteria (see Table 1).

2.3. Coding the studies

Within the studies that met the inclusion criteria, three researchers searched for available outcome variables and consensus was reached to categorize them as total behaviour problems, externalizing behaviour

problems, internalizing behaviour problems, social skills, cognitive skills and delinquency.

Two researchers coded the first three studies together. Subsequently, the other sixteen studies were divided between the two researchers and were coded independently. Additionally, a third researcher coded eight of the studies by random selection. Inter-rater agreement was analysed by calculating the percentage of agreement, using Kappa for categorical variables and intraclass correlation for interval and ratio variables. The inter-rater reliability proved to be satisfactory, with Kappa's ranging from .834 to 1.000, and intraclass correlations ranging from .963 to 1.000. In a few cases an initial disagreement occurred about the type of comparison. More specifically, the question was raised whether certain treatment methods could be counted as evidence-based or not. Again this was discussed, and consensus was reached after consulting with an expert in the field of evidence-based treatment. When computed effect sizes were not the same among researchers, the differences were only marginal (in the third decimal place) and resolved after discussion. These effect sizes were recalculated using the right numbers.

When coding the studies, the institutional group was always the experimental group and the non-institutional group was the control group. In cases of two institutional groups, the group receiving evidence-based treatment was regarded as the experimental group. In 9 of the 19 included studies, pre-test effect sizes could be subtracted from the post-test effect sizes. This was done in order to account for initial group differences between the experimental and control group. A moderator was added to distinguish between the two types of studies (accounted for pre-test differences, yes or no).

Each study included in the meta-analysis was coded for methodological and sample characteristics. Methodological characteristics were year of publication, journal impact factor, study quality, study design, time of measurement, type of comparison, type of intervention, and the data source. A study quality index (QI) was used to evaluate the study quality (Downs & Black, 1998). Sample characteristics were mean age, sex, percentage of girls, target group, ethnicity and the outcome variables.

These moderators were divided into continuous moderators and discrete moderators. Continuous moderators were year of publication, journal impact factor, study quality, mean age, percentage of girls and ethnicity (percentage of non-Caucasian participants). Discrete moderators were sex (boys or mixed), time of measurement (post-test or follow-up), control on pre-test (yes or no), type of comparison (institutional EBT vs non-institutional EBT, institutional EBT vs institutional CAU, institutional CAU vs non-institutional EBT and institutional CAU vs non-institutional CAU), study design (matched, non-matched or RCT), type of intervention (behavioural treatment, cognitive behaviour therapy, skills training, system treatment or no treatment), target group (civil, criminal, psychiatric or mixed), data source (official report, parent report, professional report or mixed) and the outcomes (total behaviour problems, externalizing behaviour problems, internalizing behaviour problems, social skills, cognitive skills or delinquency).

2.4. Publication bias

Studies reporting strong significant associations are more likely to be accepted for publication in a journal. Therefore, studies that report less strong significant associations are more difficult to find. Subsequently, conclusions of this meta-analysis may be incomplete, which is called the file drawer problem (Rosenthal & Hershstein, 1979). File drawer bias was examined using a funnel plot of the distribution of effect sizes. Each individual study's effect size is plotted on the horizontal axis against its sample size, standard error or precision (the reciprocal of the standard error) on the vertical axis. The distribution of effect sizes should be shaped as a funnel if no publication bias is present since the more numerous studies with small sample sizes are expected to show a larger variation in the magnitude of effect sizes than the

Table 1
Study characteristics.

Study (publication year)	Impact factor journal	Study quality rating	World part of origin	Comparison	Design	Time of measurement	Intervention institutional	N	Seks	Average age	Percentage non-Caucasian	Target group	Control on pre-test	N of effect sizes
Barth et al. (2007)	2.21	14	North America	Inst CAU – Noninst EBT	Matched control	Follow-up	Behaviour modification	786	Mixed	11,42	39,35	Civil	No	1
Berger et al. (2009)	3.63	16	Western Europe	Inst CAU – Noninst CAU	Matched control	Follow-up	Behaviour modification	2453	Mixed	9,62	52	Civil	Yes	3
Frensch, Cameron, and Preyde (2009)	0.00	15	North America	Inst EBT – Noninst EBT	Non-matched	Post-test and follow-up	CBT	210	Mixed	11,55	2	Civil	Yes	2
Henggeler et al. (1999a)	4.98	13	North America	Inst CAU – Noninst EBT	RCT	Post-test	Behaviour modification	113	Mixed	13,00	66	Psychiatric	Yes	5
Henggeler et al. (2003)	4.98	15	North America	Inst CAU – Noninst EBT	RCT	Follow-up	Behaviour modification	160	Mixed	12,90	67	Psychiatric	No	3
James, Roesch, and Zhang (2012)	1.28	17	North America	Inst CAU – Noninst CAU	Matched control	Post-test	Behaviour modification	1191	Mixed	8,95	58,70	Civil	No	3
Kazdin (1987)	4.46	15	North America	Inst EBT – Noninst EBT	Non-matched	Post-test and follow-up	CBT	37	Mixed	10,90	23,20	Psychiatric	Yes	10
Kolko, Loar, and Sturnick (1990)	4.98	16	Western Europe	Inst EBT – Inst CAU	Matched control	Post-test	CBT	56	Mixed	10,35	50,32	Psychiatric	Yes	1
Leve and Chamberlain (2005)	2.97	17	North America	Inst CAU – Noninst CAU	RCT	Post-test	Behaviour modification	153	Mixed	14,88	21,57	Civil	No	1
Mattejat, Hirt, Wilken, Schmidt, and Remschmidt (2001)	2.38	15	Western Europe	Inst CAU – Noninst EBT	RCT	Post-test and follow-up	Behaviour modification	68	Mixed	11,90	Unknown	Psychiatric	Yes	4
McCrae, Bethany, Barth, and Rauktis (2010)	0.67	18	North America	Inst CAU – Noninst CAU	Matched control	Post-test	Behaviour modification	124	Mixed	12,25	67,70	Civil	Yes	5
Moody (1997)	0.00	11	North America	Inst EBT – Inst CAU	Non-matched	Post-test	Behaviour modification	28	Boys	14,30	89,29	Criminal	Yes	3
Preyde et al. (2009)	1.12	13	North America	Inst CAU – Noninst EBT	Non-matched	Post-test and follow-up	Behaviour modification	210	Mixed	13,91	Unknown	Civil	Yes	5
Preyde et al. (2011)	0.00	12	North America	Inst CAU – Noninst EBT	Non-matched	Post-test and follow-up	Behaviour modification	112	Mixed	11,57	Unknown	Civil	Yes	3
Robst, Armstrong, and Dollard (2011)	1.27	16	North America	Inst CAU – Noninst CAU	Matched control	Post-test	Behaviour modification	842	Mixed	Unknown	52,20	Civil	Yes	1
Ryan, Marshall, Herz, and Hernandez (2008)	1.17	18	North America	Inst CAU – Noninst CAU	Matched control	Post-test	Behaviour modification	8226	Mixed	8,45	83	Civil	No	1
Scholte and Van der Ploeg (2003)	0.00	13	Western Europe	Inst EBT – Noninst EBT	Non-matched	Post-test	Skills	105	Mixed	14,90	35	Mixed	No	3
Thompson et al. (1996)	1.12	15	North America	Inst EBT – Inst CAU	Non-matched	Post-test	Skills	587	Mixed	14,55	29,60	Mixed	Yes	1
Wilmshurst (2002)	0.00	17	North America	Inst EBT – Noninst EBT	Non-matched	Post-test and Follow-up	Skills	65	Mixed	10,74	5	Civil	Yes	8

less numerous studies with large effect sizes. A violation of funnel plot symmetry reflects publication bias, that is, a selective inclusion of studies showing positive or negative outcomes (Sutton, Duval, Tweedie, Abrams, & Jones, 2000). Funnel plot asymmetry was tested by regressing the standard normal deviate, defined as the effect size divided by its standard error, against the estimate's precision (the inverse of the standard error) that largely depends on sample size (see Egger, Smith, Schneider, & Minder, 1997). If there is asymmetry, the regression line does not run through the origin and the intercept significantly deviates from zero.

2.5. Analysis

For each of the studies Cohen's d was calculated for the effectiveness of institutional youth care on the basis of differences between institutional and non-institutional youth care and difference between evidence-based institutional treatment and institutional care as usual. Both the post-test data and the follow-up data were used. Effect sizes were calculated on the basis of means and standard deviations, percentages and t -, F -, χ^2 -, p -values. For this purpose, Wilson's effect size determination program (Wilson, 2001) was used. An effect size of $d = 0.20$ is considered as small, an effect size of $d = 0.50$ is considered as medium and an effect size of $d = 0.80$ is considered as large (Cohen, 1988). Outliers were checked on the basis of z -values larger than 3.29 or smaller than -3.29 ($p < 0.005$; Tabachnick & Fidell, 2001). No outliers were found. After that, categorical variables were turned into dichotomous dummy codes and continuous moderator variables were centred around their mean in order to be able to conduct multilevel meta-analysis.

The homogeneity of the combined, total effect size was tested with a z -test of the between study variance (total study variance divided by its standard error). If this z -test is significant, there is heterogeneity. In case of significant heterogeneity, moderators may account for differences between studies, and it is imperative to conduct categorical and/or continuous moderator analyses.

Finally, multilevel analysis was conducted by using the program MLwiN (Hox, 2002). The multilevel random effects model takes the hierarchical structure of the data into account in which the effect sizes (the lowest level) are nested within studies (the highest level). Iterative maximum likelihood procedures were applied to estimate unknown parameters.

3. Results

This meta-analysis included 19 studies, comprising 63 effect sizes, with the results based on $N = 15,526$ children and youth between the ages of 4 and 17 years. Table 2 shows a representation of the overall mean effect size and the significant discrete moderators. The overall mean effect size was $d = -0.02$. The individual study effect sizes ranged from -1.14 to 1.56 . Possible publication bias was examined by testing funnel plot asymmetry. The standard normal deviate was regressed against the estimate's precision. As the intercept did not significantly deviate from zero ($t = 1.879$; $p = 0.08$), there was no indication of funnel plot asymmetry and therefore no indication of publication bias. These findings suggest that the mean effect size can be considered robust. Finally, the overall mean effect size proved to be heterogeneous ($z = 2.875$), indicating that the effect was not the same in all studies. This is a precondition to be able to expect significant outcomes from moderator analyses.

Table 2 shows that type of comparison was a significant moderator: $\chi^2(3) = 8.932$, $p < .05$. The reference category was the comparison between institutional evidence-based treatment and non-institutional evidence-based treatment ($d = 0.342$, ns). The institutional Care As Usual (CAU) versus non-institutional CAU comparison differed significantly from the reference group comparison ($z = 2.601$, $p < .01$), yielding a small-to-medium negative significant effect ($d = -0.342$; $z = 2.280$, $p < .05$). This means that children in non-institutional CAU, mostly foster care, had somewhat better outcomes than children in institutional CAU, whereas there were no significant differences in the effects of EBT between institutional and non-institutional care.

Another moderator variable with a significant effect was study design: $\chi^2(2) = 9.656$, $p < 0.01$. Non-matched studies differed significantly from matched studies: $z = 3.217$,

$p < 0.01$. Matched studies yielded a negative and significant effect size ($d = -0.309$; $z = 2.255$,

$p < 0.05$), whereas non-matched studies showed a positive and significant effect size.

($d = 0.299$; $z = 2.300$, $p < 0.05$). The non-matched studies showed better outcomes for children in institutional youth care and the matched studies showed better outcomes for children in non-institutional youth care.

Finally, type of outcome was a significant moderator: $\chi^2(5) = 25.115$, $p < 0.001$. Delinquency differed significantly from the reference

Table 2
Results for the overall mean effect size and discrete moderators (bivariate models).

Moderator variables	# studies	# ES	Mean d	z_1	RC	z_2	Heterogeneity	Fit χ^2
Overall	19	63	-0.018	0.176			2.875**	
Type of comparison							2.804**	8.932*
Inst EBT vs Noninst EBT (RG)	3	20	0.342	1.591				
Inst EBT vs Inst CAU	4	8	0.285	1.397+	-0.057	0.193		
Inst CAU vs Noninst EBT	6	21	-0.038	0.252	-0.380	1.445		
Inst CAU vs Noninst CAU	6	14	-0.342	2.280*	-0.684	2.601**		
Study design							2.814**	9.656**
Matched (RG)	7	15	-0.309	2.255*				
Non-matched	8	35	0.299	2.300*	0.608	3.217**		
RCT	4	13	-0.131	0.712	0.177	0.773		
Type of outcome							2.833**	25.115***
Total problems (RG)	11	14	0.035	0.343				
Externalizing behaviour	10	13	0.168	1.680	0.133	2.180*		
Internalizing behaviour	10	13	-0.051	0.510	-0.085	1.393		
Social skills	4	6	0.089	0.605	0.054	0.409		
Cognitive skills	9	12	0.051	0.490	0.016	0.208		
Delinquency	5	5	-0.329	2.179*	-0.364	2.318*		

Note. RG = reference group; # studies = number of studies; # ES = number of effect sizes; Mean d = mean effect size; z_1 = significance of moderator; RC = slope; z_2 = differences in mean d with reference group; Heterogeneity = within class heterogeneity (z); Fit = difference with model without moderators (χ^2); + = one-sided trend = 0.08.

* $p < 0.05$.

** $p < 0.01$.

*** $p < 0.001$.

Table 3
Results for the continuous moderators (bivariate models).

Moderator variables	# studies	# ES	β_0 (SE)	β_1 (SE)	z	Heterogeneity	Fit χ^2
<i>Methodological moderator</i>							
Year of publication	19	63	0.055 (0.089)	−0.035	2.692**	2.800**	6.806**
<i>Sample characteristic</i>							
Girls exp. group	19	45	−0.060 (0.089)	−0.013	2.167*	2.558*	4.966*
Girls cont group	19	45	−0.070 (0.089)	−0.013	2.167*	2.545*	4.586*
Total girls	19	63	0.022 (0.092)	−0.015	2.500*	2.840**	5.696*

Note. # studies = number of studies; # ES = number of effect sizes; β_0 (SE) = intercept; β_1 (SE) = slope; Z = significance of moderator; Heterogeneity = within class heterogeneity (z); Fit = difference with model without moderators (χ^2).

* $p < 0.05$.

** $p < 0.01$.

group, which was total problems: $z = 2.318$, $p = 0.05$, yielding a small-to-medium negative significant effect ($d = -0.329$; $z = 2.179$, $p < 0.05$). This means that children in institutional youth care showed more delinquent behaviour compared to children in non-institutional youth care. The other outcome categories did not show a significant effect, and were therefore not mentioned in Table 2.

Table 3 shows the results for the continuous moderators. Year of publication was a significant moderator ($z = 2.692$, $p < 0.01$), indicating that earlier published studies were associated with larger effect sizes ($\beta_1 = -0.035$). Also with regard to sex of the child a significant moderating effect was found ($z = 2.500$, $p < 0.05$). Studies with a high percentage of females were associated with smaller effect sizes ($\beta_1 = -0.015$).

4. Discussion

The purpose of the present meta-analysis was to examine the outcome of institutional youth care compared to non-institutional care and the outcome of institutional evidence-based treatment (EBT) compared to institutional care as usual (CAU) for primary school age children and early adolescents. This differs from the meta-analysis of De Swart et al. (2012), which had a broader age range and focused primarily on middle and late adolescents and young adults. In addition, the present study uses multilevel techniques in order to be able to include more effect sizes of the same study, preserving all relevant information, which increases statistical power and enables the examination of more moderators than can be achieved in a standard meta-analysis.

The main conclusion is that institutional CAU showed a small-to-medium negative significant effect compared to non-institutional CAU ($d = -0.342$). This indicates that children in non-institutional CAU, mostly foster care, had somewhat better outcomes than children in institutional CAU, which is regular group care offering daily care and structure in a (psychiatric) living group setting. This conclusion is in line with the retrospective study by Dregan and Gulliford (2012). They found that children in institutional youth care develop less favourably compared to children in foster care, and institutional care is associated with increased risk of adult criminal convictions and depression. Our findings differ from those in the meta-analytic study of the De Swart et al. (2012), which focused on a broader age range, with an overrepresentation of late adolescents and young adults. Although De Swart et al. (2012) found that non-institutional CAU had slightly better outcomes than institutional CAU, this difference just failed to reach statistical significance. Notably, selection effects should be taken into account when interpreting this result, because children with more complex problems generally require more intensive and specialized treatment (Evans et al., 2013; Farmer et al., 2008; Stith et al., 2009). This specialized treatment can be provided in non-institutional care, but may increase the risk for placement disruptions (Oosterman et al., 2007; Van Oijen, 2010; Vanschoonlandt, Vanderfaillie, Van Hoken, De Maeyer, & Andries, 2012). As a result, these children may be referred less often to foster care. Sometimes children start in foster

care and are later referred to institutional care because of the severity of their problems (Hussey & Guo, 2005).

No differences were found between institutional and non-institutional youth care when institutional treatment was evidence-based, which is in line with findings from the meta-analysis of De Swart et al. (2012). However, we did not find a significant advantage in providing youth with institutional EBT instead of institutional CAU. De Swart et al. (2012) found a significant difference between institutional EBT and institutional CAU for middle and late adolescents as well as young adults, with better outcomes for institutional EBT. In the present study the effect size was small-to-medium in favour of institutional EBT for children and young adolescents, but did not reach significance. A possible explanation for this result is a lack of treatment integrity, which is a shortcoming in the application of evidence-based treatment methods in institutional care (James et al., 2013). Another possible explanation is that group workers may not be responsive enough to the individual needs of the children in their group, which would be a basic requirement to be able to generate positive results of institutional EBT (Van der Helm, 2011). Variations in the way children adapt to institutional care do not only depend on importation factors (Gover, MacKenzie, & Armstrong, 2000; Hussey & Guo, 2005), i.e. characteristics of individuals before entering institutional care shaping their adjustment, but also on environmental factors, such as the living group climate. Recent studies on this topic, mainly performed in forensic settings, suggest that group climate factors influence treatment outcomes. For example, Van der Helm, Klapwijk, Stams, and Van der Laan (2009) and Schubert et al. (2012) showed that a positive or 'open' living group climate characterized by warmth and responsiveness from group workers, opportunities for development, and a safe and structured environment, had a positive effect on the treatment of juvenile delinquents. A 'closed' or repressive climate, on the other hand, has been found to be associated with negative developmental outcomes and aggression (Van der Helm, Stams, Van Genabeek, & Van der Laan, 2012). Therapeutic alliance has also been identified in the literature as a predictor of positive treatment outcomes and can be influenced (Harder, Knorth, & Kalverboer, 2012; Hurley, Lambert, Van Ryzin, Sullivan, & Stevens, 2013; Karver, Handelsman, Fields, & Bickman, 2006). In addition, although there is abundant evidence indicating the prevalence of trauma exposure among youth in institutional care, few models exist for creating trauma-informed milieu treatment (Brown, McCauley, Navalta, & Saxe, 2013), which could also hamper better results for institutional EBT.

Moderator analyses indicated that the design of the studies influenced the conclusion about the effectiveness of institutional youth care. Institutional youth care showed better outcomes in non-matched studies, whereas matched studies showed less favourable outcomes for institutional youth care. A possible explanation is that a relatively greater therapeutic gain can be achieved when problem rates are higher at the start, which is often the case in institutional care (Barth et al., 2007; Berger et al., 2009). The main benefit of matching studies is high internal conclusion validity, because the participants in the experimental group and control group are supposed to be comparable. For

this reason, more value is attached to the outcomes of the matched studies, which yield less positive results for institutional youth care. Notably, a possible drawback of matching is that children with the most severe problems are often left out of the analyses because they are less likely to match with juveniles in the control group. This may violate the external validity or, in other words, clinical representativeness of studies using a matched control design (Goodman et al., 1997).

In contrast to the study of De Swart et al. (2012), the present study showed a significant moderator effect for delinquency. Children in institutional youth care showed more delinquent behaviour compared to children in non-institutional youth care. Differences between institutional and non-institutional care with regard to delinquent problem behaviour may be more extreme in the younger age group, because presently a (young) child is only referred to institutional care when no other options are available. Other types of outcomes did not show significant differences between institutional and non-institutional care in this meta-analysis, which is in line with results reported by De Swart et al. (2012).

Finally, significant moderating effects were found for year of publication and sex of the child. Studies that were published more recently and studies with a higher percentage of females yielded smaller effect sizes. This first finding may be explained by the fact that statistical techniques, such as propensity score matching, have become more advanced over time, which enables better control for multiple confounders that are responsible for differences between institutional and non-institutional care. In addition, and probably even more important, growing insights into offering qualitatively good institutional care may have diminished the differences between institutional and non-institutional youth care (Farrington & Welsh, 2006). The association between sex of the child and magnitude of the effect sizes could be explained by the fact that girls in institutional care seem to suffer more from neglect (Gunther Moor, 2011) and more often have a history of trauma and sexual abuse, resulting in relatively more internalizing problems (Zurbriggen, Gobin, & Freyd, 2010). Research indicates that children with externalizing behavioural problems seem to make more progress in institutional care than youth with internalizing behavioural problems, which are mostly girls (Knorth, Harder, Zandberg, & Kendrick, 2007). This could be an explanation for the smaller differences between institutional and non-institutional youth care when the percentage of girls increases.

4.1. Implications for future research

In the first place, given the non-significant and small-to-medium effect size of the comparison between institutional EBT and institutional CAU, more research is necessary to examine the effectiveness of evidence-based treatment in institutional youth care for children. Future research should focus on the conditions that make established treatment methods work in institutional youth care for (young) children: what works for whom and under what circumstances? An interesting discussion on this topic, introducing a new model on translational research for the field of residential child care, is found in Nunno, Sellers, and Holden (2014).

Furthermore, in general, future effect studies focussing on institutional care versus alternatives should report more information on (cumulative) risk factors (Evans et al., 2013; Farmer et al., 2008; Stith et al., 2009), because changes in dynamic risk factors are critical to establish the effects of the care provided. In many effect studies hardly any information is reported on parent and family factors. Tools that assist clinicians in identifying and matching the level of risk to the intensity of treatment are available; see, for example Augimeri, Walsh, Woods, and Jiang (2012). In addition, protective factors should be examined in future studies, which is in line with the shift towards strengths-based care (Kumpfer & Alvarado, 2003). Results on changes in risk and protective factors would enable refinements of future meta-analytic studies.

Another recommendation for future effect studies is to use child self-report measures. Self-report measures can be a valuable supplement to the official, parental and professional reports (Dunn et al., 2010). Yet another important factor to be taken into account in future studies is IQ. Many children in institutional youth care have a mild intellectual disability, which can have a great impact on treatment results (Van Nieuwenhuijzen, Vriens, Scheepmaker, Smit, & Porton, 2011).

4.2. Implications for policy and practice

The main conclusion of this study is that non-institutional CAU had slightly better outcomes than institutional CAU, which supports the idea that institutional care should be considered as a last resort, to be used only when non-residential alternatives are less appropriate. When problems are too severe and institutional care seems to be the best choice given the situation, it is recommended that attention be paid to the living group climate, therapeutic alliances and trauma-sensitivity as preconditions to be able to provide adequate (evidence-based) institutional care for children. These factors should be regularly monitored, and group workers should receive proper training and coaching in order to make necessary improvements and enable children to express their thoughts and feelings about living in a residential setting more easily (Hunt, 2010; Leichtman, 2008). For example, promoting positive outcomes for children with a special emphasis on developing safer environments and healthy relationships is described in the Children and Residential Experiences (CARE) programme (Holden, 2009).

4.3. Limitations of the study

A general limitation of meta-analysis is that it cannot account for all underlying differences between studies by means of moderator analyses, such as the impact of culture and care systems among countries (Weisz, Ugueto, Cheron, & Herren, 2013b). There are also some technical limitations of this meta-analysis that should be noted. First, in some moderator categories the number of effect sizes was very small, which may have resulted in less reliable outcomes. Second, not every study reported the range of the age of the children. When a mean age lower than 15 years was reported without the exact age range, the study was included. As a result, we do not know if the children included in these studies were mainly primary school age children or early adolescents, which hampers more specific generalizability of our study findings. Another limitation is that evidence-based treatment (EBT) and care as usual (CAU) are broadly defined categories. The limited number of studies and effect sizes did not allow making more refined comparisons between different kinds of treatments.

5. Conclusion

Despite its limitations, this meta-analysis provides new insights into the outcome of institutional youth care compared to non-institutional youth care and the outcome of institutional evidence-based treatment (EBT) compared to institutional care as usual (CAU) for children of primary school age and early adolescence. Also, the application of multi-level analysis made it possible to include more effect sizes than can be achieved with regular meta-analysis and to examine more moderators. This study indicates that a particular group of children seems to have better outcomes in non-institutional care as usual, such as foster care, compared to institutional care. An explanation is that a more stable (family) environment provides better conditions for development and treatment, such as the development of secure child-caregiver attachment relationships (Dozier et al., 2014). Additionally, within such an environment, children may be less affected by negative peer influences (Aguilar-Vafaie et al., 2011; Whitehead et al., 2007). When problems are too severe to treat at home or in foster care, specialized institutional treatment may still be required. In order to offer optimal (evidence-

based) care within institutions, one should besides child and family related characteristics (risk and protective factors) also take environmental factors (e.g., group climate) and therapeutic alliance into account, and provide a trauma-sensitive treatment milieu. Making careful decisions in referring children to specific forms of care prevents problems from getting worse later in life and diminishes recidivism and re-entry in care, thereby also diminishing costs for society in the long term.

Acknowledgements

We thank Martha J. Holden, MS, Senior Extension Associate with the Bronfenbrenner Center for Translational Research at the College of Human Ecology, Cornell University, for her help with reviewing the English in this article.

References¹

- Aguilar-Vafaei, M.E., Roshani, M., Hassanabadi, H., Masoudian, Z., & Afruz, G.A. (2011). Risk and protective factors for residential foster care adolescents. *Children and Youth Services Review*, 33, 1–15. <http://dx.doi.org/10.1016/j.childyouth.2010.08.005>.
- Alexander, J., Pugh, D., Parsons, B., & Sexton, T. (2000). Functional family therapy. In D.S. Elliot (Ed.), *Blueprints for violence prevention*. 2. Boulder, CO: Venture Publishing.
- Andrews, D.A., & Bonta, J. (2010). Rehabilitating criminal justice policy and practice. *Psychology, Public Policy, and Law*, 16, 39–55. <http://dx.doi.org/10.1037/a0018362>.
- Asscher, J.J., Dijkstra, S., Stams, G.J.J.M., Dekovic, M., & Creemers, H.E. (2014). Family Group Conferencing in youth care: Characteristics of the decision making model, implementation and effectiveness of the family group (FG) plans. *BMC Public Health*, 14, 154. <http://dx.doi.org/10.1186/1471-2458-14-154>.
- Asscher, J.J., Van der Put, C.E., & Stams, G.J.J.M. (2015). *Gender differences in the association between child maltreatment and adolescent offending*. Journal of Family Violence (in press).
- Assink, M., Van der Put, C.E., Hoeve, M., De Vries, L.A., Stams, G.J.J.M., & Oort, F.J. (2015). *Risk factors for persistent delinquent behavior among juveniles: A meta-analytic review*. (Manuscript submitted for publication).
- Augimeri, L., Walsh, M., Woods, S., & Jiang, D. (2012). Risk assessment and clinical risk management for young antisocial children: The forgotten group. *Universitas Psychologica*, 11, 1147–1156.
- Barber, J.G., & Delfabbro, P.H. (2003). Placement stability and the psychosocial well-being of children in foster care. *Research on Social Work Practice*, 13, 415–431. <http://dx.doi.org/10.1177/1049731503013004001>.
- Barth, R.P., Greeson, J.K., Green, R.L., Hurley, S., & Sisson, J. (2007*). Outcomes for youth receiving intensive in-home therapy or residential care: A comparison using propensity scores. *American Journal of Orthopsychiatry*, 77, 497–505. <http://dx.doi.org/10.1037/0002-9432.77.4.497>.
- Bean, P., White, L., & Lake, P. (2005). A prototype residential model for adolescents with substance abuse and mental health disorders. *Journal of Addictive Diseases*, 24 (132–132).
- Behar, L., Friedman, R.M., Pinto, A., Katz-Leavy, J., & Jones, W.G. (2007). Protecting youth placed in unlicensed, unregulated residential “treatment” facilities. Special issue: The American Bar Association’s youth at risk initiative. *Family Court Review*, 45, 399–413. <http://dx.doi.org/10.1111/j.1744-1617.2007.00155.x>.
- Benzies, K.M., Harrison, M.J., & Magill-Evans, J. (2004). Parenting stress, marital quality, and child behaviour problems at age 7 years. *Public Health Nursing*, 21(2), 111–121. <http://dx.doi.org/10.1111/j.0737-1209.2004.021204.x>.
- Berger, L.M., Bruch, S.K., Johnson, E.L., James, S., & Rubin, D. (2009*). Estimating the “impact” of out-of-home placement on child well-being: Approaching the problem of selection bias. *Child Development*, 80, 1856–1876. <http://dx.doi.org/10.1111/j.1467-8624.2009.01372.x>.
- Boendermaker, L., Van Rooijen, K., & Berg, T. (2012). *Residentieële Jeugdzorg: Wat werkt? [Residential youth care: What works?]*. Utrecht: Nederlands Jeugdinstituut.
- Brown, A.D., McCauley, K., Navalta, C.P., & Saxe, G.N. (2013). Trauma Systems Therapy in residential settings: Improving emotion regulation and the social environment of traumatized children and youth in congregate care. *Journal of Family Violence*, 28, 693–703. <http://dx.doi.org/10.1007/s10896-013-9542-9>.
- Chamberlain, P., & Reid, J.B. (1998). Comparison of two community alternatives to incarceration for chronic juvenile offenders. *Journal of Consulting and Clinical Psychology*, 66, 624–633. <http://dx.doi.org/10.1037/0022-006X.66.4.624>.
- Chance, S., Dickson, D., Bennett, P.M., & Stone, S. (2010). How fundamental changes in residential care can improve the ways we help children and families. *Residential Treatment for Children & Youth*, 27, 127–148. <http://dx.doi.org/10.1080/08865711003738522>.
- Children’s Rights Alliance (2010). *The united nations convention on the rights of the child*. Dublin: Children’s Rights Alliance.
- Chor, K.H.B., McClelland, G.M., Weiner, D.A., Jordan, N., & Lyons, J.S. (2012). Predicting outcomes of children in residential treatment: A comparison of a decision support algorithm and a multidisciplinary team decision model. *Children and Youth Services Review*, 34, 2345–2352. <http://dx.doi.org/10.1016/j.childyouth.2012.08.016>.
- Chorpita, B.F., Weisz, J.R., Daleiden, E.L., Schoenwald, S.K., Palinkas, L.A., Miranda, J., ... Gibbons, R.D. (2013). Long-term outcomes for the child STEPs randomized effectiveness trial: A comparison of modular and standard treatment designs with usual care. *Journal of Consulting and Clinical Psychology*, 81, 999–1009. <http://dx.doi.org/10.1037/a0034200>.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: Lawrence Erlbaum.
- Crampton, D. (2007). Research review: Family group decision-making: A promising practice in need of more programme theory and research. *Child and Family Social Work*, 12, 202–209. <http://dx.doi.org/10.1111/j.1365-2206.2006.00442.x>.
- De Swart, J.J.W., Van den Broek, H., Stams, G.J.J.M., Asscher, J.J., Van der Laan, P.H., Holsbrink-Engels, G.A., & Van der Helm, G.H.P. (2012). The effectiveness of institutional youth care over the past three decades: A meta-analysis. *Children and Youth Services Review*, 34, 1818–1824. <http://dx.doi.org/10.1016/j.childyouth.2012.05.015>.
- De Wit, J., Slot, W., & van Aken, M. (2013). *Psychologie van de adolescentie. Basisboek*. Baarn: HB Uitgevers.
- Dekker, A. L., Van Miert, V. S. L., Roest, J. J., & Van der Helm, G. H. P. (2012). Van onhandelbaar naar behandelbaar [from intractable to treatable]. Unpublished manuscript, Leiden University of Applied Sciences.
- Dijkstra, S., Creemers, H.E., Asscher, J.J., Dekovic, M., & Stams, G.J.J.M. (2015). *The effectiveness of family group conferencing (FGC) in youth care*. (Manuscript submitted for publication).
- Dijkstra, S., Creemers, H.E., Asscher, J.J., & Stams, G.J.J.M. (2014). *De inzet van familienetwerkberaden in de jeugdzorg: Een systematische review en meta-analyse van de uitkomsten van studies naar de effectiviteit van familienetwerkberaden [Family Group Conferencing in child welfare: A systematic review and meta-analytic study on the effectiveness of Family Group Conferencing in child welfare]*. Amsterdam: University of Amsterdam.
- Dishion, T.J., McCord, J., & Poulin, F. (1999). When interventions harm. Peer groups and problem behaviour. *American Psychologist*, 54, 755–764. <http://dx.doi.org/10.1037/0003-066X.54.9.755>.
- Downs, S.H., & Black, N. (1998). The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *Journal of Epidemiology and Community Health*, 52, 377–384. <http://dx.doi.org/10.1136/jech.52.6.377>.
- Dozier, M., Kaufman, J., Kobak, R., O’Connor, T.G., Sagi-Schwartz, A., Scott, S., ... Zeanah, C.H. (2014). Consensus statement on group care for children and adolescents: A statement of policy of the American orthopsychiatric association. *American Journal of Orthopsychiatry*, 84, 219–225. <http://dx.doi.org/10.1037/ort0000005>.
- Dregan, A., & Gulliford, M.C. (2012). Foster care, residential care and public care placement patterns are associated with adult life trajectories: Population-based cohort study. *Social Psychiatry and Psychiatric Epidemiology*, 47, 1517–1526. <http://dx.doi.org/10.1007/s00127-011-0458-5>.
- Dregan, A., Brown, J., & Armstrong, D. (2011). Do adult emotional and behavioural outcomes vary as a function of diverse childhood experiences of the public care system? *Psychological Medicine*, 41, 2213–2220. <http://dx.doi.org/10.1017/S0033291711000274>.
- Dunn, D.M., Culhane, S.E., & Taussig, H.N. (2010). Children’s appraisals of their experiences in out-of-home care. *Children and Youth Services Review*, 32, 1324–1330. <http://dx.doi.org/10.1016/j.childyouth.2010.05.001>.
- Egger, M., Smith, D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315, 629–634. <http://dx.doi.org/10.1136/bmj.315.7109.629>.
- Evans, G.W., Li, D., & Whipple, S.S. (2013). Cumulative risk and child development. *Psychological Bulletin*, 139, 1342–1396. <http://dx.doi.org/10.1037/a0031808>.
- Fairchild, G., Hagan, C.C., Walsh, N.D., Passamonti, L., Calder, A.J., & Goodyer, I.M. (2013). Brain structure abnormalities in adolescent girls with conduct disorder. *Journal of Child Psychology and Psychiatry*, 54, 86–95. <http://dx.doi.org/10.1111/j.1469-7610.2012.02617.x>.
- Farmer, E.M.Z., Mustillo, S., Burns, B.J., & Holden, E.W. (2008). Use and predictors of out-of-home placements within systems of care. *Journal of Emotional and Behavioral Disorders*, 16, 5–14. <http://dx.doi.org/10.1177/1063426607310845>.
- Farmer, E.M.Z., Southerland, D., Mustillo, S.A., & Burns, B.J. (2009). Returning home in systems of care: Rates, predictors, and stability. *Journal of Emotional and Behavioral Disorders*, 17, 133–146. <http://dx.doi.org/10.1177/1063426608327002>.
- Farrington, D.P., & Welsh, B.C. (2006). A half-century of randomized experiments on crime and justice. In M. Tonry (Ed.), *Crime and Justice*. Vol. 34. (pp. 55–132). Chicago: University of Chicago Press.
- Frensch, K., & Cameron, G. (2002). Treatment of choice or a last resort? A review of residential mental health placements for children and youth. *Child & Youth Care Forum*, 31, 307–339. <http://dx.doi.org/10.1023/A:1016826627406>.
- Frensch, K., Cameron, G., & Preyde, M. (2009*). Community adaptation of youth accessing residential programs or a home-based alternative: School attendance and academic functioning. *Child & Youth Care Forum*, 38, 287–303. <http://dx.doi.org/10.1007/s10566-009-9083-9>.
- Frost, N., Abram, F., & Burgess, H. (2014). Family group conferences: Evidence, outcomes and future research. *Child & Family Social Work*, 19, 501–507. <http://dx.doi.org/10.1111/cfs.12049>.
- Goodman, S.H., Lahey, B.B., Fielding, B., Dulcan, M., Narrow, W., & Regier, D. (1997). Representativeness of clinical samples of youths with mental disorders: A preliminary population-based study. *Journal of Abnormal Psychology*, 106, 3–14. <http://dx.doi.org/10.1037/0021-843X.106.1.3>.
- Gover, A.R., MacKenzie, K.L., & Armstrong, C.S. (2000). Importational and deprivation explanations of juveniles’ adjustment to correctional facilities. *International Journal of*

¹ References marked with an asterisk (*) are included in the meta-analysis.

- Offender Therapy and Comparative Criminology, 44, 450–467. <http://dx.doi.org/10.2277/0306624X00444004>.
- Gudbrandsson, B. (2008). *Rights of Children in Institutions: Report on the implementation of the Council of Europe Recommendation Rec(2005)5 on the rights of children living in residential institutions*. Iceland: Council of Europe.
- Gunther Moor, B. (2011). *Being left out: Neurobiological correlates of social rejection and their developmental trajectory*. Amsterdam: Ipskamp.
- Handwerk, M.L., Field, C.E., & Friman, P.C. (2001). The iatrogenic effects of group intervention for antisocial youth: Premature extrapolations? *Journal of Behavioral Education*, 10, 223–238. <http://dx.doi.org/10.1023/A:1012299716053>.
- Harder, A.T., Knorth, E.J., & Kalverboer, M.E. (2012). Securing the downside up: Client and care factors associated with outcomes of secure residential youth care. *Child & Youth Care Forum*, 41, 259–276. <http://dx.doi.org/10.1007/s10566-011-9159-1>.
- Henggeler, S.W., Rowland, M.D., Halliday-Boykins, C., Sheidow, A.J., Ward, D.M., & Randall, J. (2003*). One-year follow-up of multisystemic therapy as an alternative to the hospitalization of youths in psychiatric crisis. *Journal of the American Academy of Child and Adolescent Psychiatry*, 42, 543–551. <http://dx.doi.org/10.1097/01.CHI.0000046834.09750.5F>.
- Henggeler, S.W., Rowland, M.D., Randall, J., Ward, D.M., Pickrel, S.G., Cunningham, P.B., ... Santos, A.B. (1999a). Home-based multisystemic therapy as an alternative to the hospitalization of youths in psychiatric crisis: Clinical outcomes. *Journal of the American Academy of Child and Adolescent Psychiatry*, 38, 1331–1339. <http://dx.doi.org/10.1097/00004583-199911000-00006>.
- Henggeler, S., Pickrel, S. G., & Brondino, M. J. (1999b). Multisystemic treatment of substance abusing and dependent delinquents: Outcomes, treatment fidelity, and transportability. *Mental Health Services Research*, 1, 171–184. <http://dx.doi.org/10.1023/A:1022373813261>.
- Hoagwood, K., & Cunningham, M. (1993). Outcomes of children with emotional disturbance in residential treatment for educational purposes. *Journal of Child and Family Studies*, 1, 129–140. <http://dx.doi.org/10.1007/BF01321281>.
- Höfte, S.J., Van der Helm, G.H.P., & Stams, G.J.J.M. (2012). *Het internationaal recht en de gesloten jeugdzorg: Adviezen voor de praktijk [International justice and secure residential care: Practice advice]*. *Justitiële Verkenningen*, 6, 84–99.
- Holden, M.J. (2009). *Children and residential experiences: Creating conditions for change*. Arlington, VA: Child Welfare League of America.
- Hoogsteder, L.M., Kuijpers, N., Stams, G.J.J.M., Van Horn, J.E., Hendriks, J., & Wissink, I.B. (2014). Study on the effectiveness of Responsive Aggression Regulation Therapy (Re-ART). *International Journal of Forensic Mental Health*, 13, 25–35. <http://dx.doi.org/10.1080/14999013.2014.893711>.
- Hox, J. (2002). *Multilevel analysis: Techniques and applications*. Mahway: Lawrence Erlbaum Associates.
- Hunt, K.F. (2010). The impact of brief play therapy training on the emotional awareness of care workers in a young children's residential care setting in Australia. *British Journal of Guidance and Counselling*, 38, 287–299. <http://dx.doi.org/10.1080/03069885.2010.492208>.
- Hurley, K.D., Lambert, M.C., Van Ryzin, M., Sullivan, J., & Stevens, A. (2013). Therapeutic alliance between youth and staff in residential group care: Psychometrics of the therapeutic alliance quality scale. *Children and Youth Services Review*, 35, 56–64. <http://dx.doi.org/10.1016/j.chilyouth.2012.10.009>.
- Hussey, D.O., & Guo, S. (2005). Characteristics and trajectories of treatment foster care youth. *Child Welfare*, 84(4), 485–506.
- Jakobsen, T.B. (2013). Anti-social youth? Disruptions in care and the role of 'behavioral problems'. *Children and Youth Services Review*, 35, 1455–1462. <http://dx.doi.org/10.1016/j.chilyouth.2013.05.012>.
- James, S. (2011). What works in group care? A structured review of treatment models for group homes and residential care. *Children and Youth Services Review*, 33, 308–321. <http://dx.doi.org/10.1016/j.chilyouth.2010.09.014>.
- James, S., Alemi, Q., & Zepeda, V. (2013). Effectiveness and implementation of evidence-based practices in residential care settings. *Children and Youth Services Review*, 35, 642–656. <http://dx.doi.org/10.1016/j.chilyouth.2013.01.007>.
- James, S., Landsverk, J., Slymen, D., & Leslie, L. (2004). Predictors of outpatient mental health service use: the role of foster care placement change. *Mental Health Services Research*, 6(3), 127–141. <http://dx.doi.org/10.1023/B:MHSR.0000036487.39001.51>.
- James, S., Roesch, S., & Zhang, J.J. (2012*). Characteristics and behavioral outcomes for youth in group care and family-based care: A propensity score matching approach using national data. *Journal of Emotional and Behavioral Disorders*, 20(3), 144–156. <http://dx.doi.org/10.1177/1063426611409041>.
- Johnson, R., Browne, K.D., & Hamilton-Giachritsis, C.E. (2006). Young children in institutional care at risk of harm. *Trauma, Violence & Abuse*, 7(1), 34–60. <http://dx.doi.org/10.1177/1524838005283696>.
- Karver, M.S., Handelsman, J., Fields, S., & Bickman, L. (2006). Meta-analysis of therapeutic relationship variables in youth and family therapy: The evidence for different relationship variables in the child and adolescent treatment outcome literature. *Clinical Psychology Review*, 26, 50–65. <http://dx.doi.org/10.1016/j.cpr.2005.09.001>.
- Kazdin, A.E. (1987*). Treatment of antisocial behavior in children: Current status and future directions. *Psychological Bulletin*, 10, 187–203. <http://dx.doi.org/10.1037/0033-2909.102.2.187>.
- Khoo, E., Skoog, V., & Dalin, R. (2012). In and out of care. A profile and analysis of children in the out-of-home care system in Sweden. *Children and Youth Services Review*, 34, 900–907. <http://dx.doi.org/10.1016/j.chilyouth.2012.01.019>.
- Knapp, M. (2006). The economics of group care practice. *Children and Youth Services*, 28, 259–284 Chapter 11. (doi:10.1300/J024v28n01_05).
- Knorth, E.J., Harder, A.T., Zandberg, T., & Kendrick, A.J. (2007). Under one roof: A review and selective meta-analysis on the outcomes of institutional child and youth care. *Children and Youth Services Review*, 30, 123–140. <http://dx.doi.org/10.1016/j.chilyouth.2007.09.001>.
- Kolko, D.J., Loar, L.L., & Sturnick, D. (1990*). Inpatient social-cognitive skills training groups with conduct disordered and attention deficit disordered children. *Journal of Child Psychology and Psychiatry*, 31, 737–748.
- Kumpfer, K.L., & Alvarado, R. (2003). Family-strengthening approaches for the prevention of youth problem behaviors. *American Psychologist*, 58, 457–465. <http://dx.doi.org/10.1037/0003-066X.58.6-7.457>.
- Larzelere, R.E., Daly, E.L., Davis, J.L., Chmelka, M.B., & Handwerk, M.L. (2004). Outcome evaluation of Girls and Boys Town's Family Home Program. *Education and Treatment of Children*, 27(2), 130–149.
- Lee, B.R., & Thompson, R. (2007). Comparing outcomes for youth in treatment foster care and family-style group care. *Children and Youth Services Review*, 30, 746–757. <http://dx.doi.org/10.1016/j.chilyouth.2007.12.002>.
- Leichtman, M. (2008). The essence of residential treatment: III. Change and adaptation. *Residential Treatment for Children & Youth*, 25, 189–207. <http://dx.doi.org/10.1080/08865710802429663>.
- Leichtman, M., Leichtman, M.L., Barber, C., & Neese, D.T. (2001). Effectiveness of intensive short-term treatment with severely disturbed adolescents. *American Journal of Orthopsychiatry*, 71, 227–235. <http://dx.doi.org/10.1037/0002-9432.71.2.227>.
- Leve, L.D., & Chamberlain, P. (2005*). Association with delinquent peers: Intervention effects for youth in the juvenile justice system. *Journal of Abnormal Child Psychology*, 33, 339–347.
- Linville, D., Chronister, K., Dishion, T., Todahl, J., Miller, J., Shaw, D., ... Wilson, M. (2010). A longitudinal analysis of parenting practices, couple satisfaction, and child behavior problems. *Journal of Marital and Family Therapy*, 36, 244–255. <http://dx.doi.org/10.1111/j.1752-0606.2009.00168.x>.
- Lonne, B., Parton, N., Thomson, J., & Harries, M. (2009). *Reforming child protection*. London: Routledge.
- Lösel, F., & Farrington, D.P. (2012). Direct protective and buffering protective factors in the development of youth violence. *American Journal of Preventive Medicine*, 43, 8–23. <http://dx.doi.org/10.1016/j.amepre.2012.04.029>.
- Manso, J.M.M., García-Baamonde, M.E., Alonso, M.B., & Barona, E.G. (2011). An analysis of how children adapt to residential care. *Children and Youth Services Review*, 33, 1981–1988. <http://dx.doi.org/10.1016/j.chilyouth.2011.05.024>.
- Mattejat, F., Hirt, B.R., Wilken, J., Schmidt, M.H., & Remschmidt, H. (2001*). Efficacy of inpatient and home treatment in psychiatrically disturbed children and adolescents. *European Child and Adolescent Psychiatry*, 10, 71–79. <http://dx.doi.org/10.1007/s007870170008>.
- McCrae, J.S., Bethany, R.L., Barth, R.P., & Rautkis, M.E. (2010*). Comparing three years of well-being outcomes for youth in group and nonkinship foster care. *Child Welfare*, 89, 229–249.
- Melkman, E. (2015). Risk and protective factors for problem behaviors among youth in residential care. *Children and Youth Services Review*, 51, 117–124. <http://dx.doi.org/10.1016/j.chilyouth.2015.02.004>.
- Moody, E.E. (1997*). Lessons from pair counseling with incarcerated juvenile delinquents. *Journal of Addictions and Offender Counseling*, 18, 10–25. <http://dx.doi.org/10.1002/j.2161-1874.1997.tb00122.x>.
- Nunno, M.A., Sellers, D.E., & Holden, M.J. (2014). Implications of translational research for the field of residential child care. *Scottish Journal of Residential Child Care*, 13(3).
- Oosterman, M., Schuengel, C., Slot, N.W., Bullens, R.A.R., & Doreleijers, T.A.H. (2007). Disruptions in foster care: A review and meta-analysis. *Children and Youth Services Review*, 29, 53–76. <http://dx.doi.org/10.1016/j.chilyouth.2006.07.003>.
- Orobio de Castro, B., Merk, W., Koops, W., Veerman, J.W., & Bosch, J.D. (2005). Emotions in social information processing and their relations with reactive and proactive aggression in referred aggressive boys. *Journal of Clinical Child & Adolescent Psychology*, 34, 105–116. http://dx.doi.org/10.1207/s15374424ccp3401_10.
- Pennell, J., & Burford, G. (2000). Family group decision making: protecting children and women. *Child Welfare*, 79(2), 131–158.
- Preyde, M., Adams, G., Cameron, G., & Frensch, K. (2009*). Outcomes of children participating in mental health residential and intensive family services: Preliminary findings. *Residential Treatment for Children & Youth*, 26(1), 1–20. <http://dx.doi.org/10.1080/08865710802689555>.
- Preyde, M., Frensch, K., Cameron, G., White, S., Penny, S., & Lazure, K. (2011*). Long-term outcomes of children and youth accessing residential or intensive home-based treatment: Three year follow up. *Journal of Child and Family Studies*, 20, 660–668. <http://dx.doi.org/10.1007/s10826-010-9442-z>.
- Raine, A. (2013). *The anatomy of violence: The biological roots of crime*. New York: Vintage Books.
- Raine, A., Brennan, P., Mednick, B., & Mednick, S.A. (1996). High rates of violence, crime, academic problems, and behavioural problems in males with both early neuromotor deficits and unstable family environments. *Archives of General Psychiatry*, 53, 544–549. <http://dx.doi.org/10.1001/archpsyc.1996.01830060090012>.
- Robst, J., Armstrong, M., & Dollard, N. (2011*). Comparing outcomes for youth served in treatment foster care and treatment group care. *Journal of Child and Family Studies*, 20, 696–705. <http://dx.doi.org/10.1007/s10826-011-9447-2>.
- Rose, J. (2014). *Working with young people in secure accommodation*. New York: Routledge.
- Rosenthal, R., & Hershstein, R.J. (1979). The file drawer problem and tolerance for null results. *Psychological Bulletin*, 86, 638–641. <http://dx.doi.org/10.1037/0033-2909.86.3.638>.
- Rubin, D.M., Alessandrini, E.A., Feudtner, C., Mandell, D.S., & Localio, A.R. (2004). Placement stability and mental health costs for children in foster care. *Pediatrics*, 113, 1336–1341.
- Ryan, J.P., & Testa, M.F. (2005). Child maltreatment and juvenile delinquency: Investigating the role of placement and placement instability. *Children and Youth Services Review*, 27, 227–249. <http://dx.doi.org/10.1016/j.chilyouth.2004.05.007>.

- Ryan, J.P., Marshall, J.M., Herz, D., & Hernandez, P.M. (2008*). Juvenile delinquency in child welfare: Investigating group home effects. *Children and Youth Services Review*, 30, 1088–1099. <http://dx.doi.org/10.1016/j.chidyouth.2008.02.004>.
- Scholte, E.M., & Van der Ploeg, J.D. (2003*). Effectiveness of residential treatment methods for youngsters with severe behavioural problems: Findings from a one year follow-up study. *International Journal of Child and Family Welfare*, 6, 185–197.
- Schubert, C.A., Mulvey, E.P., Loughran, T.A., & Losoya, S.H. (2012). Perceptions of institutional experience and community outcomes for serious adolescent offenders. *Criminal Justice and Behavior*, 39, 71–94. <http://dx.doi.org/10.1177/0093854811426710>.
- Souverein, F.A., Van der Helm, G.H.P., & Stams, G.J.J.M. (2013). Nothing works in secure residential youth care? *Children and Youth Services Review*, 35, 1941–1945. <http://dx.doi.org/10.1016/j.chidyouth.2013.09.010>.
- Stith, S.M., Liu, T., Davies, L.C., Boykin, E.L., Alder, M.C., Harris, J.M., ... Dees, J.E.M.E.G. (2009). Risk factors in child maltreatment: A meta-analytic review of the literature. *Aggression and Violent Behavior*, 14, 13–29. <http://dx.doi.org/10.1016/j.avb.2006.03.006>.
- Sutton, A.J., Duval, S.J., Tweedie, R.L., Abrams, K.R., & Jones, D.R. (2000). Empirical assessment of effect of publication bias on meta-analyses. *British Medical Journal*, 320(7249), 1574–1577. <http://dx.doi.org/10.1136/bmj.320.7249.1574>.
- Tabachnick, B.G., & Fidell, L.S. (2001). *Using multivariate statistics*. New York: Harper & Row.
- Thompson, R.W., Smith, G.L., Osgood, D.W., Dowd, T.P., Friman, P.C., & Daly, D.L. (1996*). Residential care: A study of short- and long-term educational effects. *Children and Youth Services Review*, 18, 221–242. [http://dx.doi.org/10.1016/0190-7409\(96\)00002-3](http://dx.doi.org/10.1016/0190-7409(96)00002-3).
- United Nations (1989). Convention on the rights of the child. Retrieved from <http://www.ohchr.org/en/professionalinterest/pages/crc.aspx>
- Van den Bergh, P.M., Weterings, A.M., & Schoenmakers, M. (2011). Gehechtheid en loyaliteit bij pleegkinderen. Een analyse vanuit de theorie en de praktijk [Attachment and loyalty in foster children. An analysis from theory and practice]. *Tijdschrift voor Orthopedagogiek, Kinderpsychiatrie en Klinische Kinderpsychologie*, 36(3), 128–143.
- Van den Dries, L., Juffer, F., Van Ijzendoorn, M.H., & Bakermans-Kranenburg, M.J. (2009). Fostering security? A meta-analysis of attachment in adopted children. *Children and Youth Services Review*, 31, 410–421. <http://dx.doi.org/10.1016/j.chidyouth.2008.09.008>.
- Van der Helm, G.H.P. (2011). *First do no harm: Living group climate in secure juvenile correctional institutions*. Amsterdam: Uitgeverij SPW.
- Van der Helm, G.H.P., & Hanrath, J. (Eds.). (2011). *Wat werkt in de gesloten jeugdzorg? What works in secure residential care?*. Amsterdam: SWP.
- Van der Helm, G.H.P., Klapwijk, M., Stams, G.J.J.M., & Van der Laan, P.H. (2009). 'What works' for juvenile prisoners: The role of group climate in a youth prison. *Journal of Children's Services*, 4(2), 36–48. <http://dx.doi.org/10.1108/17466660200900011>.
- Van der Helm, G.H.P., Stams, G.J.J.M., & Van der Laan, P.H. (2011). Measuring group climate in prison. *The Prison Journal*, 91, 158–177. <http://dx.doi.org/10.1177/0032885511403595>.
- Van der Helm, G.H.P., Stams, G.J.J.M., Van Genabeek, M., & Van der Laan, P.H. (2012). Group climate, personality, and self-reported aggression in incarcerated male youth. *The Journal of Forensic Psychiatry & Psychology*, 23(1), 23–39. <http://dx.doi.org/10.1080/14789949.2011.633615>.
- Van der Stouwe, T., Asscher, J.J., Stams, G.J.J.M., Dekovic, M., & Van der Laan, P.H. (2014). The effectiveness of multisystemic therapy (MST): A meta-analysis. *Clinical Psychology Review*, 34, 468–481. <http://dx.doi.org/10.1016/j.cpr.2014.06.006>.
- Van Nieuwenhuijzen, M., Vriens, A., Scheepmaker, M., Smit, M., & Porton, E. (2011). The development of a diagnostic instrument to measure social information processing in children with mild to borderline intelligence. *Research in Developmental Disabilities*, 32, 358–370. <http://dx.doi.org/10.1016/j.ridd.2010.10.012>.
- Van Oijen, S.V. (2010). *Resultaat van pleegzorgplaatsingen: Een onderzoek naar breakdown en de ontwikkeling van adolescente pleegkinderen bij langdurige pleegzorgplaatsingen [Results of foster care placements: A study about breakdown and the development of adolescent foster children in long term foster care]*. Groningen: RUG.
- Van Stam, M.A., Van der Schuur, W.A., Tserkezis, S., Van Vugt, E.S., Asscher, J.J., Gibbs, J.C., & Stams, G.J.J.M. (2014). The effectiveness of EQUIP on sociomoral development and recidivism reduction: A meta-analytic study. *Children and Youth Services Review*, 38, 44–51. <http://dx.doi.org/10.1016/j.chidyouth.2014.01.002>.
- Vanschoonlandt, F., Vanderfaellie, J., Van Holen, F., De Maeyer, S., & Andries, C. (2012). Kinship- and non-kinship foster care: Differences in contact with parents and foster child's mental health problems. *Children and Youth Services Review*, 34, 1533–1539. <http://dx.doi.org/10.1016/j.chidyouth.2012.04.010>.
- Vermaes, I.P.R., & Nijhof, K.S. (2014). Zijn jongeren in JeugdzorgPlus Anders dan jongeren in de open residentiële jeugdzorg? [Are juveniles in secure youth care different from juveniles in open residential youth care?]. *Orthopedagogiek: Onderzoek & Praktijk*, 53, 33–46.
- Wampold, B.E., Goodheart, C.D., & Levant, R.F. (2007). Clarification and elaboration on evidence-based practice in psychology. *American Psychologist*, 62, 616–618. <http://dx.doi.org/10.1037/0003-066X62.6.616>.
- Weick, A., Rapp, C., Sullivan, W.P., & Kisthardt, W. (1989). A strengths perspective for social work practice. *Social Work*, 34, 350–354.
- Weisz, J.R., Kuppens, S., Eckshtain, D., Ugueto, A.M., Hawley, K.M., & Jensen-Doss, A. (2013a). Performance of evidence-based youth psychotherapies compared with usual clinical care: A multilevel meta-analysis. *JAMA Psychiatry*, 70, 750–761. <http://dx.doi.org/10.1001/jamapsychiatry.2013.1176>.
- Weisz, J.R., Ugueto, A.M., Cheron, D.M., & Herren, J. (2013b). Evidence-based youth psychotherapy in the mental health ecosystem. *Journal of Clinical and Adolescent Psychology*, 42, 274–286. <http://dx.doi.org/10.1080/15374416.2013.764824>.
- White, C.P., & King, K. (2011). Is maternal fatigue mediating the relationship between maternal depression and child outcomes? *Journal of Child and Family Studies*, 20, 844–853. <http://dx.doi.org/10.1007/s10826-011-9452-5>.
- Whitehead, K., Keshet, M., Lombrowski, B., Domenico, A., & Green, D. (2007). Definition and accountability: A youth perspective. *American Journal of Orthopsychiatry*, 77, 348–349. <http://dx.doi.org/10.1037/0002-9432.77.3.34>.
- Whittaker, J.K., Del Valle, J.F., & Holmes, L. (Eds.). (2015). *Therapeutic residential care with children and youth: Developing evidence-based international practice*. London/Philadelphia: Jessica Kingsley Publishers.
- Wilmschurst, L.A. (2002*). Treatment programs for youth with emotional and behavioral disorders: An outcome study of two alternate approaches. *Mental Health Services Research*, 4, 85–96. <http://dx.doi.org/10.1023/A:1015200200316>.
- Wilson, D.B. (2001). Effect size determination program. [Mason.gmu.edu/~dwilson/downloads/ES_Calculator.xls](http://mason.gmu.edu/~dwilson/downloads/ES_Calculator.xls)
- Ziviani, J., Feeney, R., Cuskelly, M., Meredith, P., & Hunt, K. (2012). Effectiveness of support services for children and young people with challenging behaviours related to or secondary to disability, who are in out-of-home care: A systematic review. *Children and Youth Services Review*, 34, 758–770. <http://dx.doi.org/10.1016/j.chidyouth.2012.01.002>.
- Zurbriggen, E.L., Gobin, R.L., & Freyd, J.J. (2010). Childhood emotional abuse predicts late adolescent sexual aggression perpetration and victimization. *Journal of Aggression, Maltreatment & Trauma*, 19(2), 204–223. <http://dx.doi.org/10.1080/10926770903539631>.