UvA-DARE (Digital Academic Repository)

Tunable, nonlinear Hong-Ou-Mandel interferometer

Oehri, D.; Pletyukhov, M.; Gritsev, V.; Blatter, G.; Schmidt, S.

DOI

[10.1103/PhysRevA.91.033816](https://doi.org/10.1103/PhysRevA.91.033816)

Publication date

2015

Document Version

Final published version

Published in

Physical Review A

[Link to publication](#)

Citation for published version (APA):

Oehri, D., Pletyukhov, M., Gritsev, V., Blatter, G., & Schmidt, S. (2015). Tunable, nonlinear Hong-Ou-Mandel interferometer. *Physical Review A*, 91(3), [033816]. <https://doi.org/10.1103/PhysRevA.91.033816>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Tunable, nonlinear Hong-Ou-Mandel interferometer

D. Oehri,¹ M. Pletyukhov,² V. Gritsev,³ G. Blatter,¹ and S. Schmidt¹¹*Theoretische Physik, ETH Zurich, CH-8093 Zurich, Switzerland*²*Institute for Theory of Statistical Physics and JARA – Fundamentals of Future Information Technology, RWTH Aachen, 52056 Aachen, Germany*³*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, Postbus 94485, 1098 XH Amsterdam, The Netherlands*

(Received 23 December 2014; published 13 March 2015)

We investigate the two-photon scattering properties of a Jaynes-Cummings (JC) nonlinearity consisting of a two-level system (qubit) interacting with a single-mode cavity, which is coupled to two waveguides, each containing a single incident photon wave packet initially. In this scattering setup, we study the interplay between the Hong-Ou-Mandel (HOM) effect arising due to quantum interference and effective photon-photon interactions induced by the presence of the qubit. We calculate the two-photon scattering matrix of this system analytically and identify signatures of interference and interaction in the second-order auto- and cross-correlation functions of the scattered photons. In the dispersive regime, when qubit and cavity are far detuned from each other, we find that the JC nonlinearity can be used as an almost linear, *in situ* tunable beam splitter giving rise to ideal Hong-Ou-Mandel interference, generating a highly path-entangled two-photon NOON state of the scattered photons. The latter manifests itself in strongly suppressed waveguide cross-correlations and Poissonian photon number statistics in each waveguide. If the two-level system and the cavity are on resonance, the JC nonlinearity strongly modifies the ideal HOM conditions leading to a smaller degree of path entanglement and sub-Poissonian photon number statistics. In the latter regime, we find that photon blockade is associated with bunched autocorrelations in both waveguides, while a two-polariton resonance can lead to bunched as well as antibunched correlations.

DOI: [10.1103/PhysRevA.91.033816](https://doi.org/10.1103/PhysRevA.91.033816)

PACS number(s): 42.50.Pq, 42.79.Fm, 42.79.Gn, 11.55.Ds

I. INTRODUCTION

The Hong-Ou-Mandel (HOM) effect [1] lies at the heart of linear optical quantum computing [2,3] and is utilized to test the degree of indistinguishability of photons as well as the quality of single-photon sources. Conventionally, the HOM effect is demonstrated as a two-photon interference effect at a linear 50:50 beam splitter: when two indistinguishable photon wave packets impinge on the beam splitter from two different waveguides, they form an entangled two-photon NOON state with both photons in the same output waveguide, which produces a characteristic dip in the coincidence probability of finding photons in both waveguides simultaneously. This destructive interference is complete only for indistinguishable photons with equal energy and zero time delay. Any deviation from the ideal conditions leads to a diminishing of the dip, which thus constitutes a measure for indistinguishability.

The HOM effect has been demonstrated experimentally using parametric down-conversion [1] and pulsed single-photon sources operating at optical frequencies [4–8]. Recently, the HOM effect has attracted new attention in the context of waveguide and circuit QED [9–13], where it was demonstrated with unprecedented precision for microwave photons as well [14,15], using recently developed microwave single-photon sources and beam splitters. This development paves the way for all-integrated linear optical quantum computing at microwave frequencies.

In all experiments so far, the central element of the HOM effect, i.e., a 50:50 beam splitter, was based on a linear and static device, whose single-photon transmission and reflection probabilities (which are constant over a broad range of frequencies) are fixed by the manufacturing process. From an experimental point of view it would be desirable to also utilize

beam splitters with *in situ* tunable reflection and transmission probabilities.

In this paper we study theoretically two-photon scattering at a Jaynes-Cummings nonlinearity [16,17], i.e., a coupled qubit-cavity system, connected directly to two transmission lines. Such a scattering setup is readily realizable using state-of-the-art circuit QED technology. Hereby, the transition frequency of the two-level system (qubit) is a flux-tunable parameter, which allows the effective single-photon resonances of the qubit-cavity system to be energetically shifted. Due to the finite width of the resonances induced by the coupling to the transmission lines, one can fine-tune the single-photon transmission and reflection probabilities of the scattering target. The Jaynes-Cummings nonlinearity may thus act as an *in situ* tunable beam splitter in a waveguide QED setup. This tunable beam splitter works best for energies of the incoming photons with a bandwidth ξ smaller than the cavity decay rate κ . In circuit QED, this condition can be satisfied by using a driven qubit-cavity system as a single-photon source [18,19] which couples more weakly to the waveguide (leading to sharp wave packets) than the beam splitter cavity.

In addition, the two-level system introduces a nonlinearity into the system, which may modify the ideal HOM conditions. Here, we investigate in detail the interplay between the two-photon interference and effective photon-photon interactions in a Jaynes-Cummings system. For this purpose, we utilize an analytic scattering matrix approach recently developed for a two-level system embedded in a chiral photonic waveguide [20]. Based on this analytic approach, we derive the exact two-photon scattering matrix of the proposed setup and identify experimentally measurable signatures of the HOM effect and effective photon-photon interactions in the second-order cross- and autocorrelation functions of the two output modes. In the

FIG. 1. Schematic of the scattering setup with a qubit-cavity system interacting via the light-matter interaction strength g and directly connected to two transmission lines giving rise to a cavity decay rate $\kappa = 2\pi g_w^2$. We consider two single-photon wave packets with spatial width v/ξ impinging on the cavity from two different directions, e.g., emitted by two independent single-photon sources [18,19].

dispersive regime, i.e., when the two-level system and the cavity are far detuned from each other, the Jaynes-Cummings nonlinearity gives rise to a qubitlike and a cavitylike scattering resonance. In the vicinity of these resonances, the two-photon scattering matrix of a qubit and a Kerr-like nonlinearity, previously studied in the context of the HOM effect [11], are derived as limiting cases.

II. MODEL

Figure 1 shows the scattering setup considered in this paper. The total Hamiltonian of this system is given by

$$H = H_{JC} + H_w + V \quad (1)$$

with the Jaynes-Cummings (JC) Hamiltonian

$$H_{JC} = \omega_c b^\dagger b + \omega_q \sigma^+ \sigma^- + g(b^\dagger \sigma^- + b \sigma^+) \quad (2)$$

describing the coupled qubit-cavity system; we set $\hbar = 1$. The Hamiltonian H_w of the transmission lines describes photons in chiral states (see Fig. 1) with linear dispersion $v = vk$ with wave vector $k > 0$ and velocity v , i.e.,

$$H_w = \sum_{i=1,2} \int dv v a_{iv}^\dagger a_{iv}, \quad (3)$$

which linearly couple to the scatterer via

$$V = g_w \sum_{i=1,2} \int dv (b^\dagger a_{iv} + \text{H.c.}). \quad (4)$$

Here, we have introduced the cavity photon operator b and the operators a_{iv} describing photons with energy v in waveguide i . The two-level system, described by the raising and lowering operators for a two-level system $\sigma^\pm$, introduces a nonlinearity into the system due to the light-matter coupling $\sim g$. In our model, we neglect a coupling of the two-level system to other modes of the system and environment, because the associated spontaneous emission and dephasing rates can be engineered to be several orders of magnitude smaller than the qubit-cavity coupling g and cavity decay rate $\kappa = 2\pi g_w^2$ to the leads (for typical parameter values in circuit QED systems, see Ref. [21]).

The scattering resonances are determined by the eigenstates of the Jaynes-Cummings Hamiltonian: The ground state $|\psi_0\rangle = |0, \downarrow\rangle$ consists of zero photons in the cavity and the two-level system in its ground state with energy $\varepsilon_0 = 0$. The

excited states ($n > 0$) are

$$|\psi_{n+}\rangle = \cos \frac{\theta_n}{2} |n-1, \uparrow\rangle + \sin \frac{\theta_n}{2} |n, \downarrow\rangle, \quad (5)$$

$$|\psi_{n-}\rangle = -\sin \frac{\theta_n}{2} |n-1, \uparrow\rangle + \cos \frac{\theta_n}{2} |n, \downarrow\rangle, \quad (6)$$

with the angle $\tan \theta_n = -2g\sqrt{n}/\delta$ and the detuning $\delta = \omega_c - \omega_q$. They correspond to n excitations, i.e., polariton quasiparticles, which form a superposition of the state $|n, \downarrow\rangle$ with n photons in the cavity and the two-level system in the ground state and the state $|n-1, \uparrow\rangle$ with $(n-1)$ photons in the cavity and the two-level system in the excited state. The corresponding energies are given by

$$\varepsilon_{n\pm} = n\omega_c - \delta/2 \pm \sqrt{(\delta/2)^2 + ng^2}. \quad (7)$$

The JC eigenenergies give rise to single-photon scattering resonances with Lorentzian peaks at $\varepsilon_{1\pm}$ as a function of incoming photon energy. Due to the coupling to the waveguide, the cavity states also attain a finite lifetime $\sim 1/\kappa$ with $\kappa = 2\pi g_w^2$. In the next section, we will derive explicit expressions for the single-photon and two-photon scattering matrix of this system.

III. SCATTERING FORMALISM

We consider the generic situation, where an initial one- or two-photon state $|\psi_{\text{in}}\rangle$ is prepared inside the waveguides at $t_{\text{in}} = -\infty$, interacts (scatters) at the Jaynes-Cummings nonlinearity, and is observed as the outgoing state $|\psi_{\text{out}}\rangle$ in a photon detector at $t_{\text{out}} = +\infty$. The relation between the initial and the scattered state is given by the S matrix, i.e.,

$$|\psi_{\text{out}}\rangle = S|\psi_{\text{in}}\rangle \quad \text{with} \quad S = \left[-i \int_{-\infty}^{\infty} V(t) dt \right]. \quad (8)$$

Here, we have written the coupling Hamiltonian in the interaction representation, i.e., $V(t) = \exp(iH_0 t) V \exp(-iH_0 t)$ with $H_0 = H_{JC} + H_w$.

In the following, we will need the one-photon S matrix (for the scattering at an empty cavity or qubit, i.e., void of excitations) written in a Fock-state basis as

$$P_0 S^{(1)} P_0 = P_0 \sum_{j,v} S_{v'v}^{ij} a_{iv'}^\dagger a_{jv} \quad (9)$$

with matrix elements $S_{v'v}^{ij} = \langle 0 | a_{iv'} S^{(1)} a_{jv}^\dagger | 0 \rangle$. Here $|0\rangle \equiv |\psi_0\rangle \otimes |0\rangle_w$ is the state void of excitations, i.e., no photons in the cavity and waveguide, and the two-level system residing in its ground state; $P_0 = |\psi_0\rangle \langle \psi_0|$ is the projector onto the dark state of the cavity, i.e., $|\psi_0\rangle = |0, \downarrow\rangle$. The symbol $\sum = \sum_{ij} \int_{v'v}$ denotes summation (integration) over all arabic (greek) indices. Similarly, the two-photon scattering matrix is defined as

$$P_0 S^{(2)} P_0 = \frac{P_0}{4} \sum_{j_1, j_2} \sum_{v'_1, v'_2, v_1, v_2} S_{v'_1 v'_2 v_1 v_2}^{i_1 i_2 j_1 j_2} a_{i_1 v'_1}^\dagger a_{i_2 v'_2}^\dagger a_{j_1 v_1} a_{j_2 v_2}, \quad (10)$$

with $S_{v'_1 v'_2 v_1 v_2}^{i_1 i_2 j_1 j_2} = \langle 0 | a_{i_2 v'_2} a_{i_1 v'_1} S^{(2)} a_{j_1 v_1}^\dagger a_{j_2 v_2}^\dagger | 0 \rangle$.

Both single- and two-photon scattering matrix elements are calculated in Appendix A,¹ by making use of the formalism developed in Ref. [20]. For the single-photon matrix elements in (9) we obtain

$$S_{\nu'\nu}^{11} = S_{\nu'\nu}^{22} = r_\nu \delta(\nu' - \nu), \quad (11)$$

$$S_{\nu'\nu}^{12} = S_{\nu'\nu}^{21} = t_\nu \delta(\nu' - \nu), \quad (12)$$

with the reflection (scattering to the same waveguide) amplitude

$$r_\nu = \frac{(\nu - \omega_q)(\nu - \omega_c) - g^2}{(\nu - \omega_q)(\nu - \omega_c + i\kappa) - g^2} \quad (13)$$

and the transmission (scattering to the other waveguide) amplitude

$$t_\nu = \frac{-i\kappa(\nu - \omega_q)}{(\nu - \omega_q)(\nu - \omega_c + i\kappa) - g^2}. \quad (14)$$

The reflection amplitude vanishes for $\nu = \varepsilon_{1,\pm}$, giving rise to resonances in transmission; see Fig. 2(b). The one-photon resonance energies and widths are determined by $\tilde{\varepsilon}_{1,\pm}$ which is obtained from $\varepsilon_{1,\pm}$ [cf. Eq. (7)], by replacing the cavity frequency ω_c with $\tilde{\omega}_c = \omega_c - i\kappa$ (and correspondingly replacing the detuning δ by $\tilde{\delta} = \delta - i\kappa$; the n -excitation resonances $\tilde{\varepsilon}_{n,\pm}$ are defined in the same way). For large positive detuning $\delta \gg g$, the upper polariton resonance at $\varepsilon_{1,+}$ is cavitylike with a width κ , while the lower polariton resonance at $\varepsilon_{1,-}$ is qubitlike with an effective width $\kappa g^2/\delta^2$.

In the context of the Hong-Ou-Mandel effect it is also useful to define the single-photon energies

$$\Omega_\sigma^{(1,2)} = \frac{1}{2}[\omega_c + \omega_q \pm \kappa + \sigma\sqrt{(\delta \pm \kappa)^2 + 4g^2}], \quad \sigma = \pm, \quad (15)$$

where the single-photon transmission and reflection probabilities are equal to 1/2, corresponding to a 50:50 beam splitter operation; see Fig. 2(b).

The two-photon scattering matrix in (10) is of the form

$$S_{\nu'_1 \nu'_2 \nu_1 \nu_2}^{i_1 i_2 j_1 j_2} = S_{\nu'_1 \nu_1}^{i_1 j_1} S_{\nu'_2 \nu_2}^{i_2 j_2} + S_{\nu'_2 \nu_1}^{i_2 j_1} S_{\nu'_1 \nu_2}^{i_1 j_2} + i\mathcal{T}_{\nu'_1 \nu'_2 \nu_1 \nu_2}^{(2)}. \quad (16)$$

The first two terms describe two uncorrelated single-photon scattering events as obtained from Eqs. (11) and (12). The third term on the right-hand side (RHS) of Eq. (16) describes the nontrivial two-photon scattering process described by the T -matrix element

$$\begin{aligned} \mathcal{T}_{\nu'_1 \nu'_2 \nu_1 \nu_2}^{(2)} &= \delta(\nu'_1 + \nu'_2 - \nu_1 - \nu_2) \frac{\kappa^2 g^4}{\pi} \frac{(\nu_1 + \nu_2 - \tilde{\varepsilon}_{1,+} - \tilde{\varepsilon}_{1,-})}{\prod_{\alpha=\pm} (\nu_1 + \nu_2 - \tilde{\varepsilon}_{2,\alpha})} \\ &\times \frac{\prod_{\alpha=\pm} (\nu_1 + \nu_2 - 2\tilde{\varepsilon}_{1,\alpha})}{\prod_{i=1,2} \prod_{\alpha=\pm} (\nu'_i - \tilde{\varepsilon}_{1,\alpha})(\nu_i - \tilde{\varepsilon}_{1,\alpha})}. \end{aligned} \quad (17)$$

Note that the T -matrix elements do not depend on the waveguide indices due to the symmetric coupling of the two

¹We extend the integration range in all energy integrals to $\nu \in (-\infty, \infty)$ which is a good approximation as long as the frequency of the incoming photons $\nu_0 > 0$ (comparable to the resonator and two-level system frequencies) sets the largest energy scale in the problem.

FIG. 2. (a) Energy level diagram of the Jaynes-Cummings Hamiltonian showing the upper and lower one- and two-polariton resonances $\varepsilon_{1\sigma}$ and $\varepsilon_{2\sigma}$ defined in Eq. (7) as functions of the detuning δ . Dashed (gray) lines correspond to the two-photon energies $2\Omega_\sigma^{(1,2)}$ defined in Eq. (15) leading to the ideal HOM effect for sharp wave packets with $\xi \rightarrow 0$. Dotted lines correspond to the two-polariton energies $\varepsilon_{2\sigma}$ and $2\varepsilon_{1\sigma}$ as functions of detuning δ . Their difference measures the nonlinearity in the Jaynes-Cummings spectrum $U = \varepsilon_{2\sigma} - 2\varepsilon_{1\sigma}$. (b) The single-photon transmission probability (i.e., transmission to the other waveguide) $T_\nu = |t_\nu|^2$ exhibits two Lorentzian resonances located approximately around the two-level system frequency ω_q and the cavity frequency ω_c in the dispersive regime with $|\delta| = |\omega_c - \omega_q| \gg g$. The energies $\Omega_\sigma^{(1,2)}$ correspond to ideal HOM conditions with $T_\nu = R_\nu = 1/2$ (where $R_\nu = |r_\nu|^2$ denotes the reflection probability and we have chosen $\delta/g = 3$ and $\kappa/g = 0.1$).

transmission lines. The results in (9)–(17) are consistent with those derived in Ref. [16,17] (using a different approach) and enable us to calculate the wave function as well as photon statistics in the output modes for arbitrary incoming states containing up to two photons. In the next section we apply these results to the specific situation depicted in Fig. 1 with two single-photon wave packets approaching the cavity in different waveguides.

IV. HONG-OU-MANDEL EFFECT

The Hong-Ou-Mandel effect [1] is a two-particle interference effect describing the spatial bunching of two indistinguishable photons which arrive at the same time at an ideal 50:50 beam splitter from two different incoming arms and always end up in the same outgoing arm. The effect is fundamentally related to the bosonic exchange properties of

the photons. In the following, we will discuss such a Hong-Ou-Mandel (HOM) effect for two photon scattering at a JC nonlinearity. As discussed above, the single-photon scattering characteristics of a JC nonlinearity provide equal transmission and reflection probability at energies $\Omega_{\pm}^{(1,2)}$ (around the two polariton energies $\varepsilon_{1,\pm}$) such that the cavity-qubit system may act as an ideal 50:50 beam splitter. However, due to the finite width of the photon wave packets, the ideal beam splitter conditions can be satisfied for only one spectral component of the wave packets. Furthermore, the nonlinearity of the JC system induced by the coupling of the photons to the two-level system modifies the two-photon scattering properties compared to a linear 50:50 beam splitter. Both finite wave-packet width and nonlinearity may wash out the quantum interference leading to the HOM effect and thus will be investigated in detail in the following.

In order to investigate the HOM effect, we consider two photons incoming in different waveguides described by the (normalized) incoming state

$$|\psi_{\text{in}}\rangle = \iint d\nu_1 d\nu_2 f_{\nu_1}^{(1)} f_{\nu_2}^{(2)} a_{1\nu_1}^\dagger a_{2\nu_2}^\dagger |0\rangle \quad (18)$$

with the function $f_v^{(i)}$ describing the incoming photon wave packet in lead $i = 1, 2$, satisfying $\int d\nu |f_v^{(i)}|^2 = 1$. The outgoing state after scattering is related to the incoming state via $|\psi_{\text{out}}\rangle = S|\psi_{\text{in}}\rangle$ which leads to

$$\begin{aligned} |\psi_{\text{out}}\rangle &= |\psi_{\text{out}}^{11}\rangle + |\psi_{\text{out}}^{22}\rangle + |\psi_{\text{out}}^{12}\rangle \\ &= \iint d\nu_1 d\nu_2 f_{\nu_1}^{(1)} f_{\nu_2}^{(2)} \\ &\quad \times \left(\frac{1}{2} \iint d\nu'_1 d\nu'_2 S_{\nu'_1 \nu'_2, \nu_1 \nu_2}^{11,12} a_{1\nu'_1}^\dagger a_{1\nu'_2}^\dagger |0\rangle \right. \\ &\quad + \frac{1}{2} \iint d\nu'_1 d\nu'_2 S_{\nu'_1 \nu'_2, \nu_1 \nu_2}^{22,12} a_{2\nu'_1}^\dagger a_{2\nu'_2}^\dagger |0\rangle \\ &\quad \left. + \iint d\nu'_1 d\nu'_2 S_{\nu'_1 \nu'_2, \nu_1 \nu_2}^{12,12} a_{1\nu'_1}^\dagger a_{2\nu'_2}^\dagger |0\rangle \right) \end{aligned} \quad (19)$$

with three contributions; the first two contributions describe two photons scattered into the same waveguide and the third contribution describes the scattering into different waveguides. The HOM effect arises if the last contribution $|\psi_{\text{out}}^{12}\rangle$ vanishes. We thus define a HOM parameter

$$\gamma = \langle \psi_{\text{out}} | n_1 n_2 | \psi_{\text{out}} \rangle = \langle \psi_{\text{out}}^{12} | \psi_{\text{out}}^{12} \rangle, \quad (20)$$

with $n_i = \int d\nu a_{i\nu}^\dagger a_{i\nu}$ which yields the coincidence probability of finding one photon in the left and one photon in the right waveguide. Thus, $\gamma = 0$ corresponds to a superposition of two states, one with both photons in the left and one with both photons in the right waveguide, i.e., a two-photon entangled NOON state with respect to the waveguide degrees of freedom. Using Eq. (19), we obtain from (20)

$$\gamma = \iint d\nu'_1 d\nu'_2 \left| \iint d\nu_1 d\nu_2 S_{\nu'_1 \nu'_2, \nu_1 \nu_2}^{12,12} f_{\nu_1}^{(1)} f_{\nu_2}^{(2)} \right|^2. \quad (21)$$

In the following, we will consider Lorentzian wave packets described by

$$f_v^{(i)} = \sqrt{\frac{\xi}{2\pi}} \frac{e^{-i\nu(t-t_i)}}{\nu - \nu_{0i} + i\xi/2} \quad (22)$$

around energies $\nu_{0i} > 0$ with width ξ . The times t_i correspond to the instants when the fronts of the wave packets reach the mirrors of the cavity, such that the time delay between them is given by $\Delta t = t_1 - t_2$. The integrals in Eq. (21) can be calculated analytically which results in explicit algebraic expressions for the HOM parameter, which, however, are rather lengthy and thus have been omitted here for brevity.

Figure 3 shows the behavior of the HOM parameter γ as obtained from the exact calculation of Eq. (21) for $\Delta t = 0$ and equal energies of the incoming photons $\nu_{01} = \nu_{02}$. Before discussing these results in detail, it is instructive to separate uncorrelated and correlated contributions in Eq. (21). From the result for the two-photon scattering matrix in (16), we can

FIG. 3. (a) HOM parameter γ [defined in Eq. (20)] as a function of the detuning δ and the two-photon energy $e_0 = \nu_{01} + \nu_{02} - 2\omega_c$ (defined relative to the cavity frequency) for $\kappa/g = 0.1$ and $\xi/\kappa = 0.1$. (b) Inset in (a) (black box) together with the two photon energies depicted in Fig. 2(a) [dashed, $2\Omega_{\pm}^{(1,2)}$, and dotted, $2\varepsilon_{1,+}$ and $\varepsilon_{2,+}$, lines with the same notation as in Fig. 2(a)]. Thin horizontal and vertical lines correspond to the cuts in (c) and (d); see below. (c) Cuts through (b) at constant two-photon energy $e_0/g = 1$ for wave packets of width $\xi/\kappa = 0.1$ (black lines) and $\xi/\kappa = 1$ (light gray lines). The dotted lines correspond to the linear approximation (for $\xi/\kappa = 0.1$), where correlation contributions due to the T -matrix elements in Eq. (26) are neglected. Vertical arrows correspond to the detuning for which $\nu_{01} + \nu_{02} = 2\Omega_{\pm}^{(1,2)}$ (ideal HOM conditions). (d) Cuts through (b) at constant detuning $\delta = 0.5g$ (upper panel) and $\delta = 2g$ (lower panel).

rewrite

$$\gamma = \iint d\nu'_1 d\nu'_2 |\mathcal{A}_{\nu'_1, \nu'_2}^{12}|^2 \quad (23)$$

with

$$\mathcal{A}_{\nu'_1, \nu'_2}^{12} = \mathcal{A}_{\nu'_1, \nu'_2}^{12, \text{lin}} + \mathcal{C}_{\nu'_1, \nu'_2}, \quad (24)$$

where

$$\mathcal{A}_{\nu'_1, \nu'_2}^{12, \text{lin}} = t_{\nu'_1} t_{\nu'_2} f_{\nu'_1}^{(1)} f_{\nu'_2}^{(2)} + r_{\nu'_1} r_{\nu'_2} f_{\nu'_1}^{(1)} f_{\nu'_2}^{(2)} \quad (25)$$

and

$$\mathcal{C}_{\nu'_1, \nu'_2} = i \iint d\nu_1 d\nu_2 \mathcal{T}_{\nu'_1 \nu'_2, \nu_1 \nu_2}^{(2)} f_{\nu_1}^{(1)} f_{\nu_2}^{(2)}. \quad (26)$$

In the linear approximation, we neglect the contributions in (23) arising from the $T^{(2)}$ matrix. Assuming furthermore infinitely sharp wave packets $\xi \ll \kappa$ as well as zero initial energy detuning, i.e., $\nu_{01} = \nu_{02} = \nu_0$, and zero time delay $\Delta t = 0$, we can write the HOM parameter in a simplified form as

$$\begin{aligned} \gamma_{\text{lin}} &= \iint d\nu'_1 d\nu'_2 |\mathcal{A}_{\nu'_1, \nu'_2}^{12, \text{lin}}|^2 \\ &= \prod_{\sigma=\pm} \frac{(2\nu_0 - 2\Omega_{\sigma}^{(1,2)})^2 (2\nu_0 - 2\Omega_{\sigma}^{(2)})^2}{|2\nu_0 - 2\tilde{\varepsilon}_{1,\sigma}|^4} \end{aligned} \quad (27)$$

with $\Omega_{\sigma}^{(1,2)}$ defined in Eq. (15). As expected, the result of the linear approximation in (27) predicts perfect HOM-like interference with $\gamma_{\text{lin}} = 0$ for infinitely sharp wave packets at the single-photon energies $\nu_{01} = \nu_{02} = \Omega_{\sigma}^{(1,2)}$ where ideal 50:50 beam splitter conditions prevail; see Fig. 2.

Figure 3(a) shows γ for nonideal but favorable conditions $\kappa/g = 0.1$ and $\xi/\kappa = 0.1$. The energies $\Omega_{\sigma}^{(1,2)}$, located around the one-polariton energy $\varepsilon_{1,\sigma}$, give rise to a double-dip feature in Fig. 3(a) [see the zoom in Fig. 3(b) and cuts in Fig. 3(d)]. For large detuning $\delta \gg g$, we find two well-separated (approximately by κ) dips for positive (negative) detuning around the upper (lower) polariton resonance. For negative (positive) detuning the two HOM features around the upper (lower) polariton resonance are very sharp and separated by $\kappa g^2/\delta^2$ only. The two dips merge into one dip due to the finite wave-packet width for $\xi \lesssim \kappa g^2/\delta^2$ and are completely washed out for $\xi > \kappa g^2/\delta^2$. Figure 3(b) shows a good agreement between the exact result as obtained from (21) and the dip location predicted from the linear approximation as discussed above. In fact, the outgoing state with the highest degree of path entanglement (smallest value of γ) is always found for the ideal HOM conditions [dashed lines in Fig. 3(b)] independent of the detuning. The overall degree of entanglement, however, becomes maximal ($\gamma \approx 0$) only in the dispersive regime. Additionally, one observes a dipole-induced-transparency-like

(DIT-like) effect [22] with $|t_{\nu}| \approx 1$ if the photons are tuned into resonance with the one-polariton state $2\nu_0 \approx 2\varepsilon_{1+}$ [upper dotted line in Fig. 3(b)] leading to an almost nonentangled outgoing state with one photon in each waveguide, similar to the incoming state ($\gamma \approx 1$).

Figures 3(c) and 3(d) represent cuts through the inset in Fig. 3(b) at fixed energy (c) and detuning (d) of the incoming photons. Both parameters can be used to tune in or out of the HOM-like interference. Correlation effects attributed to the difference between solid and dotted lines in Figs. 3(c) and 3(d) come into play for small detuning and wash out the HOM effect, but are irrelevant in the dispersive regime with moderately large detuning. We also note that for broad wave packets with $\xi \sim \kappa$ (gray curve) the HOM interference is washed out independent of the detuning. This is due to the fact that ideal beam splitter conditions with 50:50 transmission and reflection probability can be achieved only for infinitely sharp wave packets with $\xi \rightarrow 0$, otherwise not all parts of the wave packet are scattered with equal probabilities as mentioned at the beginning of this section.

The reason for the different behavior at positive (negative) detuning in Fig. 3 is due to the change of the qubit or photon nature of the one-polariton resonance: with increasing positive (negative) detuning, the lower (upper) polariton resonance becomes more qubit (photon)-like and is thus strongly decoupled from (coupled to) the photon scattering and interference process, which leads to the HOM effect. More specifically, in the strongly dispersive limit corresponding to large positive detuning $\delta \gg g$ with $\theta_n \rightarrow \pi^-$ in (5), the polariton state $|\psi_{1+}\rangle$ becomes photonlike, i.e., $|\psi_{1+}\rangle \rightarrow |1, \downarrow\rangle$, while the state $|\psi_{1-}\rangle$ becomes qubitlike, i.e., $|\psi_{1-}\rangle \rightarrow |0, \uparrow\rangle$. For large negative detuning $-\delta \gg g$ with $\theta_n \rightarrow 0^+$, the situation is reversed: $|\psi_{1+}\rangle \rightarrow |0, \uparrow\rangle$, $|\psi_{1-}\rangle \rightarrow |1, \downarrow\rangle$.

In the limit $\delta \gg g$ ($-\delta \gg g$), photons with energies around ε_{1+} (ε_{1-}) effectively scatter at a Kerr nonlinearity described by the Hamiltonian $H_{\text{Kerr}} = \bar{\omega}_c b^\dagger b + (U/2) b^\dagger b^\dagger b b$ with energy $\bar{\omega}_c \approx \omega_c + g^2/\delta - g^4/\delta^3$ and a weak nonlinearity $U \approx -2g^4/\delta^3$, where the polaronic shift in the energy as well as the nonlinearity are induced by the presence of the two-level system. Note that $\text{sgn}(U)$ is opposite to $\text{sgn}(\delta)$. On the other hand, photons with energies around ε_{1-} (ε_{1+}) scatter mostly at the weakly coupled two-level system with transition frequency $\bar{\omega}_q \approx \omega_q - g^2/\delta + g^4/\delta^3$, where the weak coupling gives rise to a small width $\bar{\kappa} \approx \kappa g^2/\delta^2$.

The single-photon and two-photon scattering matrices of the JC nonlinearity given by Eqs. (13), (14), and (17) simplify to the scattering matrices of a Kerr nonlinearity [23] a two-level system [24] (TLS) in the corresponding energy ranges $2\nu_0 \approx 2\omega_q$ ($2\nu_0 \approx 2\omega_c$) at large detuning, as shown in Appendix B. Making use of these scattering matrices, we may calculate the HOM coefficient γ from Eq. (21) in the Kerr regime, yielding

$$\begin{aligned} \gamma &= 1 - \frac{(2\kappa)^2 U [2e_{0c} \xi + U(2\kappa + 3\xi)]}{\xi [e_{0c}^2 + (2\kappa + 3\xi)^2] (U^2 + 4\xi^2)} + \frac{2(2\kappa)^2 e_{0c} \xi - [2\xi^2 + (2\kappa)\xi - (2\kappa)^2] (2\kappa) U}{\xi U \{e_{0c}^2 + [\xi + (2\kappa)]^2\}} \\ &\quad + \frac{4\xi (2\kappa)^2 \{[\xi - (2\kappa)]U - 2e_{0c} \xi\}}{U [U^2 + 4\xi^2] [(e_{0c} - U)^2 + (\xi + 2\kappa)^2]} \end{aligned} \quad (28)$$

with $e_{0c} = \nu_{01} + \nu_{02} - 2\bar{\omega}_c$. In the TLS regime, we obtain

$$\gamma = 1 + \frac{-2i\bar{\kappa}}{e_{0q} + i(2\bar{\kappa} + \xi)} + \frac{(-2i\bar{\kappa})^2}{[e_{0q} + i(2\bar{\kappa} + \xi)][e_{0q} + i(2\bar{\kappa} + 3\xi)]} + \frac{2i\bar{\kappa}}{e_{0q} - i(2\bar{\kappa} + \xi)} + \frac{(2i\bar{\kappa})^2}{[e_{0q} - i(2\bar{\kappa} + \xi)][e_{0q} - i(2\bar{\kappa} + 3\xi)]} \quad (29)$$

with $e_{0q} = \nu_{01} + \nu_{02} - 2\bar{\omega}_q$. These simple expressions show good agreement with the HOM features in the dispersive limit in the corresponding energy regimes.

V. CORRELATIONS

The HOM parameter γ discussed in the previous section is rather difficult to measure directly. Instead, it is more convenient to study the second-order correlation function

$$G_{ij}^{(2)}(\tau) = \int dt \langle a_{ix}^\dagger(t) a_{jx}^\dagger(t + \tau) a_{jx}(t + \tau) a_{ix}(t) \rangle, \quad (30)$$

where

$$a_{ix}(t) = \frac{1}{\sqrt{2\pi}} \int dv e^{-iv(t-x/v)} a_{iv} \quad (31)$$

represents the photon annihilation operator at a particular position $x \rightarrow +\infty$ in waveguide i [such that $G_{ij}^{(2)}(\tau)$ is independent of position] and v is the photon group velocity. Here and in the following, expectation values are calculated with respect to the outgoing state $\langle \cdot \rangle = \langle \psi_{\text{out}} | \cdot | \psi_{\text{out}} \rangle$.

It is straightforward to show that the cross-correlation function $G_{12}^{(2)}(\tau)$ is related to the Hong-Ou-Mandel parameter by simple integration, i.e.,

$$\int d\tau G_{12}^{(2)}(\tau) = \langle n_1 n_2 \rangle = \gamma. \quad (32)$$

For a perfect HOM interference, both photons end up in the same waveguide and thus the cross-correlation function vanishes altogether.

By integrating the autocorrelation function $G_{11}^{(2)}(\tau)$ over time, one obtains the difference between the second and the first moments of the photon number distribution,² i.e.,

$$\int d\tau G_{11}^{(2)}(\tau) = \langle n_1(n_1 - 1) \rangle = \langle n_1^2 \rangle - \langle n_1 \rangle. \quad (33)$$

In the special case of two indistinguishable photons with zero time delay and zero energy detuning, there is a perfect symmetry between waveguides 1 and 2 which leads to $\langle n_1 \rangle = \langle n_2 \rangle = 1$ and $\langle n_1^2 \rangle = \langle n_2^2 \rangle$. Furthermore, making use of $\langle (n_1 + n_2)^2 \rangle = 4$, we find $\langle n_1^2 \rangle = 2 - \gamma$ and correspondingly

$$\langle \Delta n_1^2 \rangle = \langle n_1^2 \rangle - \langle n_1 \rangle^2 = 1 - \gamma. \quad (34)$$

Note that the expressions in (33) and (34) are identical in this special case. Thus, in this case, perfect HOM interference ($\gamma = 0$) is associated with a Poissonian photon number distribution characterized by $\langle \Delta n_1^2 \rangle = \langle n_1 \rangle$, while any deviations from the

ideal HOM conditions lead to sub-Poissonian statistics with $\langle \Delta n_1^2 \rangle < \langle n_1 \rangle$ (no super-Poissonian statistics is possible for the two-photon scattering considered here). The time-integrated autocorrelation function thus also serves as a useful measure for the degree of path entanglement generated by the HOM effect.

Making use of the outgoing state Eq. (19), we find

$$G_{ij}^{(2)}(\tau) = \frac{1}{2\pi} \int dE \left| \int d\Delta e^{i\Delta\tau} \mathcal{A}_{E/2+\Delta, E/2-\Delta}^{ij} \right|^2 \quad (35)$$

with $\mathcal{A}_{v'_1, v'_2}^{ij} = \mathcal{A}_{v'_1, v'_2}^{ij, \text{lin}} + \mathcal{C}_{v'_1, v'_2}$, where

$$\mathcal{A}_{v'_1, v'_2}^{11, \text{lin}} = r_{v'_1} t_{v'_2} f_{v'_1}^{(1)} f_{v'_2}^{(2)} + t_{v'_1} r_{v'_2} f_{v'_2}^{(1)} f_{v'_1}^{(2)} \quad (36)$$

and $\mathcal{A}_{v'_1, v'_2}^{12, \text{lin}}$ and $\mathcal{C}_{v'_1, v'_2}$ are defined in Eqs. (25) and (26). The expression (35) can again be evaluated analytically. In Figs. 4 and 5 we show the normalized second-order correlation functions

$$g_{ij}^{(2)}(\tau) = \frac{G_{ij}^{(2)}(\tau)}{G_{ij, \infty}^{(2)}}, \quad (37)$$

where

$$G_{ij, \infty}^{(2)} = \lim_{\Delta t \rightarrow \infty} [G_{ij}^{(2)}(\tau = \Delta t) + G_{ij}^{(2)}(\tau = -\Delta t)] \quad (38)$$

denotes the uncorrelated contribution to $G_{ij}^{(2)}$ obtained from an incoming state with a large time delay $\Delta t \gg 1/\xi$ between the two wave packets describing the case of independent scattering of two distinguishable (classical) particles.

The correlation functions $g_{ij}^{(2)}(\tau)$ are affected by both the statistical nature of the photons and the correlation effects induced by the nonlinearity. In the following, we will first describe the signatures of the HOM effect in the cross-correlation function and then discuss correlation effects due to the Jaynes-Cummings nonlinearity.

A. Signatures of HOM interference

Figures 4(a) and 4(b) show that the HOM effect at $2\nu_0 \approx 2\Omega_\sigma^{(1,2)}$ (dashed lines) manifests itself as a suppression of the cross-correlations at zero time delay with $g_{12}^{(2)}(0) \approx 0$ only in the dispersive regime ($|\delta| \gg g$) for the upper polaritons at positive detuning ($\delta > 0$) and the lower polaritons at negative detuning ($\delta < 0$), where nonlinear effects are weak and the JC nonlinearity acts as an almost ideal beam splitter. The different behavior at small and large detuning is analyzed in more detail in Fig. 4(c) showing cuts through the inset in Fig. 4(b) (indicated by vertical lines): strong deviations from the linear result (dotted lines) are observed for small detuning (upper panel), but can be neglected for $|\delta| \gg g$ (lower

²Note that the lowest two moments fully characterize the full counting statistics in the case of two-photon scattering.

FIG. 4. (a) Normalized cross-correlation at zero time delay $g_{12}^{(2)}(0)$ as defined in Eq. (37) as a function of detuning δ and two-photon energy $e_0 = 2\nu_0 - 2\omega_c$ as defined in the caption of Fig. 3. (b) Inset in (a) (see black box) together with the two-photon energies depicted in Fig. 2(a) [dashed and dotted lines with the same notation as in Fig. 2(a)]. (c) Cuts through (b) at constant detuning [indicated with vertical lines in (b)] for wave packets of width $\xi/\kappa = 0.1$. The dotted lines correspond to the linear approximation, where correlation contributions due to the T -matrix elements in Eq. (26) are neglected. The arrows correspond to $2\Omega_{+}^{(1,2)}$. (d) Time dependence of the normalized second-order cross-correlation function for the parameter values indicated by the star in the lower panel of (c), i.e., for almost ideal HOM conditions, but for different time delay Δt of the initial single-photon wave packets. The cross-correlation function displays characteristic side peaks at time delay Δt , whose height, i.e. $g_{12}^{(2)}(\tau = \Delta t)$, converges at large time delays ($\Delta t \gg 1/\xi \gg 1/\kappa$), i.e., for completely distinguishable photon wave packets, towards $1/2$ (see text for explanation).

panel). The zero-time-delay cross-correlation function $g_{12}^{(2)}(0)$ thus provides a useful signature for path entanglement only in the dispersive regime. Note that for an ideal HOM effect in the dispersive regime with $\gamma = 0$ the wave-packet width should also tend to zero. The nonlinear effects at small detuning will be discussed further in the next section.

The quantum interference process leading to the HOM effect relies on the indistinguishability of the two photons at zero time delay and equal energies. Thus, a time delay between the incoming photons leads to distinguishability and suppresses the HOM interference. Figure 4(d) shows the cross-correlation function with a time delay Δt of the two-photon wave packets in the dispersive regime for parameter settings corresponding to the star in the lower panel of Fig. 4(c) (HOM condition). We observe broad peaks centered at the time delay $\tau \approx \Delta t$, where the first photon has maximal overlap with the second. The overall width $\sim 1/\xi$ of the side peak is

FIG. 5. (a) Normalized autocorrelation function $g_{11}^{(2)}(0)$ at zero time delay as defined in Eq. (37) as a function of detuning δ and two-photon energy $e_0 = 2\nu_0 - 2\omega_c$ for two photons incoming from different waveguides ($\xi/\kappa = 0.1$). (b) Time dependence of the autocorrelation function $g_{11}^{(2)}(\tau)$ for the case of bunching [upper panel, indicated by star in (a)] and antibunching [lower panel, indicated by circle in (a)]. (c) Normalized correlation function of transmitted (left) and reflected (right) photons $\tilde{g}_{22}^{(2)}(0)$ and $\tilde{g}_{11}^{(2)}(0)$ at zero time delay for two photons incoming from the same waveguide 1. Here, the dotted lines indicate the resonances at $2\varepsilon_{1,+}$ and $\varepsilon_{2,+}$, with the same notation as in Fig. 2(a).

given by the width of the wave packet while the cusplike feature on top arises due to the coupling to the waveguides $\sim 1/\kappa$ and is suppressed in the limit $\xi/\kappa \rightarrow 0$. For large time delay and sharp wave packets such that $\Delta t \gg 1/\xi \gg 1/\kappa$, the cross-correlation function is given by

$$G_{12}^{(2)}(\tau) \approx \frac{\xi}{8} (e^{-\xi|\tau-\Delta t|} + e^{-\xi|\tau+\Delta t|}), \quad (39)$$

which yields the asymptotic value $g_{12}^{(2)}(\tau = \Delta t) \approx 1/2$ in agreement with Fig. 4(d). The asymptotic value of $1/2$ originates from the fact that only one of the two classical processes contributes to $G_{12}^{(2)}(\tau = \Delta t)$ while the normalization in Eq. (38) is obtained by summing over both classical paths.

The appearance of a maximum in the cross-correlation function at finite times thus serves as a measure for the distinguishability of the two photons due to an initial time delay and can be used to benchmark the two single-photon sources attached to both transmission lines. These findings, valid in the dispersive regime, are consistent with recent experimental circuit QED results for a static, linear 50:50 beam splitter in Refs. [14,15].

B. Signatures of photon nonlinearities

In the previous section, we mostly discussed signatures of the HOM effect in the dispersive regime, where nonlinear effects can be neglected. We will now focus on a discussion of the resonant regime, where photon nonlinearities are strong.

When both photons are resonant with the one-polariton state ($2\nu_0 = 2\varepsilon_{1,\pm}$), dipole-induced transparency with reflection amplitude $r_{\nu=\varepsilon_{1,\pm}} = 0$ [cf. Fig. 2(b)] would lead to complete and independent transmission of the two particles in the absence of any correlations, i.e., if one neglects the contributions of the T matrix in Eq. (35). However, when qubit and cavity are on resonance ($\delta \approx 0$), those contributions dominate and lead to photon blockade. In that case, a single photon already present in the cavity blocks the transmission of a second photon, which is thus reflected and preferentially drags the cavity photon into the same waveguide leading to bunched correlations with $g_{11}^{(2)}(0) \gg 1$ [see Fig. 5(a)]. Interestingly, when the total energy of both photons matches the two-polariton state energy ($2\nu_0 = \varepsilon_{2,\pm}$), bunched as well as antibunched correlations can be observed, depending on the value of the detuning parameter.

Note that this behavior should be contrasted with the case where both photons impinge on the qubit-cavity system from the same waveguide, i.e., for an incoming state $|\psi_{\text{in}}\rangle \sim \int \int d\nu_1 d\nu_2 f_{\nu_1}^{(1)} f_{\nu_2}^{(2)} a_{1\nu_1}^\dagger a_{1\nu_2}^\dagger |0\rangle$. This case is shown in Fig. 5(c), where photon blockade at $2\nu_0 = 2\varepsilon_{1,\pm}$ leads to bunching in reflection (right panel), but antibunching in transmission (left panel). In the latter case, bunching correlations are found for energies lying between the two antibunched regions. These findings are in agreement with previous theoretical [17] as well as experimental studies [25].

VI. CONCLUSION

In this work, we have studied in detail the interplay of quantum interference leading to the HOM effect and effective photon-photon interactions in a waveguide QED system, where two single-photon wave packets are incident on a Jaynes-Cummings nonlinearity from two separate waveguides. For this purpose, we have calculated the cross- and autocorrelation functions of the two waveguides analytically based on a scattering matrix approach. A central result of our study is that the proposed setup can be used as an *in situ* tunable HOM interferometer in the dispersive regime for already moderate detuning between the two-level system and the cavity and rather sharp wave packets of width ξ smaller than the cavity decay width κ . In the opposite regime, where the two-level system and cavity are on resonance, HOM interference is washed out and signatures of photon blockade due to effective photon-photon interactions induced by strong coupling of photons and the two-level system, manifest as bunched correlations in the second-order autocorrelation function. Interestingly, for the case considered here of two photons impinging on the system from two different waveguides, the two-photon resonance can lead to bunching as well as antibunching correlations depending on the sign of the qubit-cavity detuning. This result is different from the standard case of two photons impinging on the system from the same waveguide. It would thus be worthwhile to compare this case with a cw-pump scheme where two coherent microwave sources drive the system continuously from two waveguides. Such an analysis as well

as the generalization of our work to the case of N -particle scattering are interesting topics for future work.

ACKNOWLEDGMENTS

We acknowledge support from the Swiss NSF through an Ambizione Fellowship (SS) under Grant No. PP00P2-123519/1.

APPENDIX A: DERIVATION OF THE SCATTERING MATRIX

1. Even and odd modes

To calculate the one and two-photon scattering matrices in (9) and (10), it is convenient to first decompose the modes of waveguides 1 and 2 into even and odd modes [24,26] by introducing

$$a_v^{(e,o)} = (a_{1v} \pm a_{2v})/\sqrt{2} \quad (\text{A1})$$

and rewriting the waveguide Hamiltonian as $H_w = H_w^{(e)} + H_w^{(o)}$ with $H_w^{(e,o)} = \int d\nu \nu a_v^{(e,o)\dagger} a_v^{(e,o)}$. With this transformation only the even modes couple to the cavity via

$$V = \sqrt{2}g_w \int d\nu (b^\dagger a_v^{(e)} + b a_v^{(e)\dagger}), \quad (\text{A2})$$

and thus the Hamiltonian for the odd modes becomes trivial. The scattering matrix associated with the odd modes is given simply by the identity matrix. The Hamiltonian in the even modes,

$$H^{(e)} = H_w^{(e)} + H_{\text{JC}} + V, \quad (\text{A3})$$

then describes one chiral mode linearly coupled to the cavity. The scattering matrix for the even modes is nontrivial and can be calculated using the formalism in Ref. [20] (see next section).

Once we have calculated the one-photon scattering matrices in the even subspace, i.e., $S_{\nu\nu'}^{(e)} = \langle 0 | a_{\nu'}^{(e)} S a_{\nu}^{(e)\dagger} | 0 \rangle$, we transform back to physical space via the relations

$$S_{\nu'\nu}^{11} = S_{\nu'\nu}^{22} = \frac{1}{2} (S_{\nu'\nu}^{(e)} + \delta_{\nu'\nu}), \quad (\text{A4})$$

$$S_{\nu'\nu}^{12} = S_{\nu'\nu}^{21} = \frac{1}{2} (S_{\nu'\nu}^{(e)} - \delta_{\nu'\nu}), \quad (\text{A5})$$

where the δ function on the right-hand side stems from the trivial contribution of the odd modes. From the two-photon scattering matrix in the even space $S_{\nu'_1\nu'_2\nu_1\nu_2}^{(e)} = \langle 0 | a_{\nu'_2}^{(e)} a_{\nu'_1}^{(e)} S a_{\nu_1}^{(e)\dagger} a_{\nu_2}^{(e)\dagger} | 0 \rangle = S_{\nu'_1\nu_1}^{(e)} S_{\nu'_2\nu_2}^{(e)} + S_{\nu'_1\nu_2}^{(e)} S_{\nu'_2\nu_1}^{(e)} + i\mathcal{T}_{\nu'_1\nu'_2,\nu_1\nu_2}^{(2e)}$ we obtain the two-photon scattering matrix in Eq. (10) from the results above together with the simple relation

$$\mathcal{T}_{\nu'_1\nu'_2,\nu_1\nu_2}^{(2)} = \mathcal{T}_{\nu'_1\nu'_2,\nu_1\nu_2}^{(2e)} / 4. \quad (\text{A6})$$

2. T matrix

To calculate the single- and two-photon scattering matrix of the even modes we make use of the formalism introduced in Ref. [20]. The scattering matrix is conveniently expressed through the T matrix which contains the nontrivial part of the scattering, i.e.,

$$S = 1 - 2\pi i \delta(E_{\text{in}} - E_{\text{out}}) T(E_{\text{in}}) \quad (\text{A7})$$

with the energies E_{in} and E_{out} of the incoming and outgoing states. The T matrix can be expressed through the full

Green's function $\hat{G}(\omega) = (\omega - H + i0^+)^{-1}$ with $H^{(e)}$ defined in Eq. (A3) as

$$T(\omega) = V + V\hat{G}(\omega)V. \quad (\text{A8})$$

The series representation of the full Green's function $\hat{G}(\omega) = \sum_{n=0}^{\infty} G_0(\omega) [VG_0(\omega)]^n$ in terms of the free Green's function $G_0(\omega) = (\omega - H_0 + i0^+)^{-1}$ and the interaction V given in Eq. (A2) yields the corresponding series representation of the T matrix. For photon-number-conserving scattering processes we obtain $T(\omega) = V \sum_{n=1}^{\infty} [G_0(\omega)V]^n$. The main result of Ref. [20] is to show that the T matrix for an incoming N -photon state can be expressed as

$$T^{(N)}(\omega) = \dot{:} G_0^{-1} \tilde{G} (V \tilde{G})^{2N} G_0^{-1} \dot{:}, \quad (\text{A9})$$

with the dressed Green's function

$$\tilde{G}(\omega) = (\omega - H_0 - \Sigma)^{-1}, \quad (\text{A10})$$

where the operator Σ (of the form of a self-energy) accounts for the coupling of the atomic system to the waveguide and for the linear coupling in Eq. (A2) attains the remarkably simple form

$$\Sigma = -2i\pi g_w^2 b^\dagger b. \quad (\text{A11})$$

The operation $\dot{:}(\cdot)\dot{:}$ is a version of the normal ordering, which removes contractions in pairs of V 's, but does not account for contractions between V and H_0 . The latter can be effectively accounted for by shifts in the ω arguments of the dressed Green's functions $\tilde{G}$ occurring in the expansion (A9).

3. Scattering matrix for even modes

We are now in the position to calculate, e.g., the one-photon scattering matrix elements

$$S_{\nu'\nu}^{(e)} = \langle 0 | a_{\nu'}^{(e)} S a_{\nu}^{(e)\dagger} | 0 \rangle = \delta(\nu' - \nu) - 2\pi i \delta(\nu' - \nu) \langle 0 | a_{\nu'}^{(e)} T^{(1)}(\nu) a_{\nu}^{(e)\dagger} | 0 \rangle \quad (\text{A12})$$

from the result in (A9), i.e.,

$$T^{(1)}(\omega) = \dot{:} G_0^{-1}(\omega) \tilde{G}(\omega) V \tilde{G}(\omega) V \tilde{G}(\omega) G_0^{-1}(\omega) \dot{:}. \quad (\text{A13})$$

Here, the combination $\tilde{G}(E_{\text{in}})G_0^{-1}(E_{\text{in}})$ at the end of the operator chain on the RHS of Eq. (A13) acts as a projector on the dark states (nonbroadened states). This can be seen from the expression $\tilde{G}(E_{\text{in}})G_0^{-1}(E_{\text{in}})|\psi_{\text{in}}\rangle$, which is zero for any incoming eigenstate of H_0 since $G_0^{-1}(E_{\text{in}}) = (E_{\text{in}} - H_0)$ except when $\tilde{G}(E_{\text{in}}) = G_0(E_{\text{in}})$, i.e., for the nonbroadened states with $\Sigma|\psi_{\text{in}}\rangle = 0$. In the case of the JC nonlinearity, the only nonbroadened state is the ground state of the combined cavity-qubit system $|\psi_0\rangle = |0, \downarrow\rangle$, such that we can replace $\tilde{G}(E_{\text{in}})G_0^{-1}(E_{\text{in}})$ by the projector $P_0 = |\psi_0\rangle\langle\psi_0|$. The same is true for $G_0^{-1}(E_{\text{in}})\tilde{G}(E_{\text{in}})$ on the left of the expression (as $E_{\text{in}} = E_{\text{out}}$). By expressing the dressed Green's function in (A10) through the projector $\tilde{P}_{n\sigma} = |\tilde{\psi}_{n\sigma}\rangle\langle\tilde{\psi}_{n\sigma}|$ on the Jaynes-Cummings resonances $|\tilde{\psi}_{n\sigma}\rangle$ [given by the expressions for $|\psi_{n\sigma}\rangle$ in (5) with the angle θ_n replaced by $\tilde{\theta}_n$ with $\omega_c \rightarrow \tilde{\omega}_c = \omega_c - i\kappa$] as $\tilde{G}(\omega) = \sum_{n\alpha} \tilde{P}_{n\alpha}(\omega - H_w - \tilde{\epsilon}_{n\alpha})^{-1}$ and calculating the corresponding matrix elements in (A13),

one arrives at the final result $S_{\nu'\nu}^{(e)} = s_{\nu'}^{(e)}\delta(\nu' - \nu)$ with

$$s_{\nu}^{(e)} = \frac{(\nu - \tilde{\epsilon}_{1+}^*)(\nu - \tilde{\epsilon}_{1-}^*)}{(\nu - \tilde{\epsilon}_{1+})(\nu - \tilde{\epsilon}_{1-})}. \quad (\text{A14})$$

A similar calculation for the two-photon T matrix

$$T^{(2)}(\omega) = \dot{:} P_0 V \tilde{G}(\omega) V \tilde{G}(\omega) V \tilde{G}(\omega) V P_0 \dot{:} \quad (\text{A15})$$

yields after some lengthy algebra the result stated in Eq. (17).

APPENDIX B: STRONGLY DISPERSIVE REGIME: LIMIT OF KERR NONLINEARITY AND TLS SCATTERING MATRIX

We show here, that in the dispersive limit $\delta \gg g$ the JC scattering matrix given by Eqs. (13), (14), and (17) can be simplified to the scattering matrix of the Kerr nonlinearity for incoming photon energies $\nu_{0i} \approx \omega_c$ and to the scattering matrix of a TLS for energies $\nu_{0i} \approx \omega_q$. (The limit $-\delta \gg g$ can be treated analogously.)

Let us start with the Kerr regime: We approximate the energies (7) as $\epsilon_{n+} = \omega_c + ng^2/\delta - n^2g^4/\delta^3$, such that the two lowest levels effectively form a Kerr nonlinearity with $\epsilon_{1+} = \tilde{\omega}_c$ and $\epsilon_{2+} = 2\tilde{\omega}_c + U$ with $\tilde{\omega}_c \approx \omega_c + g^2/\delta - g^4/\delta^3$ and $U \approx -2g^4/\delta^3$. The energies of the off-resonant states $|\psi_{n-}\rangle$ are approximated as $\epsilon_{n-} \approx n\omega_c - \delta$. To approximate the single-photon scattering matrix coefficients t_ν and r_ν given by Eqs. (13) and (14), it is useful to note that they can be written as $r_\nu = (s_\nu^{(e)} + 1)/2$ and $t_\nu = (s_\nu^{(e)} - 1)/2$. As $\tilde{\epsilon}_{1-}$ is off resonant, we approximate $\nu - \tilde{\epsilon}_{1-} \approx \epsilon_{1+} - \tilde{\epsilon}_{1-} \approx \delta$, which leads to $s_\nu^{(e)} = (\nu - \tilde{\omega}_c - i\kappa)/(\nu - \tilde{\omega}_c + i\kappa)$ in agreement with Ref. [23] and to $r_\nu = (\nu - \tilde{\omega}_c)/(\nu - \tilde{\omega}_c + i\kappa)$ and $t_\nu = -i\kappa/(\nu - \tilde{\omega}_c + i\kappa)$. In the same way, we approximate the T matrix given by Eq. (17): Only the single-photon resonances at energy ϵ_{1+} , the two-photon resonances at energies ϵ_{2+} , as well as the energy dependence in $\nu_1 + \nu_2 - 2\epsilon_{1+}$ are relevant. In all the other factors which do not contain ϵ_{n+} we approximate $\nu_i \approx \epsilon_{1+}$ and obtain to leading order in g/δ

$$T_{\nu'_1\nu'_2,\nu_1\nu_2}^{(2)} \approx -\frac{\kappa^2 U}{\pi} \frac{\nu_1 + \nu_2 - 2\tilde{\omega}_c + 2i\kappa}{\nu_1 + \nu_2 - 2\tilde{\omega}_c - U + 2i\kappa} \times \frac{\delta_{\nu'_1+\nu'_2,\nu_1+\nu_2}}{\prod_{i=1,2} (\nu'_i - \tilde{\omega}_c + i\kappa)(\nu_i - \tilde{\omega}_c + i\kappa)}, \quad (\text{B1})$$

with $\tilde{\omega}_c$ and U as introduced above.

For incoming photon energies ν_{0i} close to ϵ_{1-} , we proceed similarly. In this regime, only the resonance at $\tilde{\epsilon}_{1-}$ is relevant, which we approximate to leading order in g/δ by $\tilde{\epsilon}_{1-} = \tilde{\omega}_q - i\bar{\kappa}$ with $\tilde{\omega}_q = \omega_q - g^2/\delta$ and $\bar{\kappa} = \kappa g^2/\delta^2$. All the other factors which do not contain $\tilde{\epsilon}_{1-}$ are off resonant and approximated by $\tilde{\epsilon}_{n+} \approx \epsilon_{n+} \approx n\omega_c$ and $\tilde{\epsilon}_{n-} \approx \epsilon_{n-} \approx n\omega_c - \delta$ for $n > 1$. Approximating $\nu - \tilde{\epsilon}_{1+} \approx -\delta$ in Eq. (A14), we obtain $s_\nu^{(e)} = (\nu - \tilde{\omega}_q - i\bar{\kappa})/(\nu - \tilde{\omega}_q + i\bar{\kappa})$, in agreement with Ref. [24]. As above, we approximate ν_i in all terms by ω_q , except for the single-photon resonance at $\tilde{\epsilon}_{1-}$ and in the factor $\nu_1 + \nu_2 - 2\tilde{\epsilon}_{1-}$, and obtain to leading order in g/δ

$$T_{\nu'_1\nu'_2,\nu_1\nu_2}^{(2)} \approx \frac{\bar{\kappa}^2}{\pi} \frac{\delta_{\nu'_1+\nu'_2,\nu_1+\nu_2}(\nu_1 + \nu_2 - 2\tilde{\omega}_q + 2i\bar{\kappa})}{\prod_{i=1,2} (\nu'_i - \tilde{\omega}_q + i\bar{\kappa})(\nu_i - \tilde{\omega}_q + i\bar{\kappa})}, \quad (\text{B2})$$

in agreement with the result for the TLS in Ref. [24].

- [1] C. K. Hong, Z. Y. Ou, and L. Mandel, *Phys. Rev. Lett.* **59**, 2044 (1987).
- [2] H. J. Kimble, *Nature (London)* **453**, 1023 (2008).
- [3] E. Knill, R. Laflamme, and G. J. Milburn, *Nature (London)* **409**, 46 (2001).
- [4] C. Santori, D. Fattal, J. Vučković, G. S. Solomon, and Y. Yamamoto, *Nature (London)* **419**, 594 (2002).
- [5] H. de Riedmatten, I. Marcikic, W. Tittel, H. Zbinden, and N. Gisin, *Phys. Rev. A* **67**, 022301 (2003).
- [6] T. Legero, T. Wilk, M. Hennrich, G. Rempe, and A. Kuhn, *Phys. Rev. Lett.* **93**, 070503 (2004).
- [7] J. Beugnon, M. P. A. Jones, J. Dingjan, B. Darquié, G. Messin, A. Browaeys, and P. Grangier, *Nature (London)* **440**, 779 (2006).
- [8] P. Maunz, D. L. Moehring, S. Olmschenk, K. C. Younge, D. N. Matsukevich, and C. Monroe, *Nat. Phys.* **3**, 538 (2007).
- [9] A. Blais, R.-S. Huang, A. Wallraff, S. M. Girvin, and R. J. Schoelkopf, *Phys. Rev. A* **69**, 062320 (2004).
- [10] A. Wallraff, D. I. Schuster, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. Kumar, S. M. Girvin, and R. J. Schoelkopf, *Nature (London)* **431**, 162 (2004).
- [11] P. Longo, J. H. Cole, and K. Busch, *Opt. Express* **20**, 12326 (2011).
- [12] M. Laakso and M. Pletyukhov, *Phys. Rev. Lett.* **113**, 183601 (2014).
- [13] A. Roulet, H. N. Le, and V. Scarani, [arXiv:1411.6411](https://arxiv.org/abs/1411.6411).
- [14] C. Lang, C. Eichler, L. Steffen, J. M. Fink, M. J. Woolley, A. Blais, and A. Wallraff, *Nat. Phys.* **9**, 345 (2013).
- [15] M. J. Woolley, C. Lang, C. Eichler, A. Wallraff, and A. Blais, *New J. Phys.* **15**, 105025 (2013).
- [16] J. T. Shen and S. Fan, *Phys. Rev. A* **79**, 023837 (2009).
- [17] T. Shi, S. Fan, and C. P. Sun, *Phys. Rev. A* **84**, 063803 (2011).
- [18] A. A. Houck, D. I. Schuster, J. M. Gambetta, J. A. Schreier, B. R. Johnson, J. M. Chow, L. Frunzio, J. Majer, M. H. Devoret, S. M. Girvin, and R. J. Schoelkopf, *Nature (London)* **449**, 328 (2007).
- [19] C. Lang, D. Bozyigit, C. Eichler, L. Steffen, J. M. Fink, A. A. Abdumalikov, Jr., M. Baur, S. Filipp, M. P. da Silva, A. Blais, and A. Wallraff, *Phys. Rev. Lett.* **106**, 243601 (2011).
- [20] M. Pletyukhov and V. Gritsev, *New J. Phys.* **14**, 095028 (2012).
- [21] S. Schmidt and J. Koch, *Ann. Phys. (Leipzig)* **525**, 395 (2013).
- [22] E. Waks and J. Vuckovic, *Phys. Rev. Lett.* **96**, 153601 (2006).
- [23] J.-Q. Liao and C. K. Law, *Phys. Rev. A* **82**, 053836 (2010).
- [24] J. T. Shen and S. Fan, *Phys. Rev. A* **76**, 062709 (2007).
- [25] A. Reinhard, T. Volz, M. Winger, A. Badolato, K. J. Hennessy, E. Hu, and A. Imamoglu, *Nat. Photonics* **6**, 93 (2012).
- [26] T. Shi and C. P. Sun, *Phys. Rev. B* **79**, 205111 (2009).