

UvA-DARE (Digital Academic Repository)

High-resolution IR absorption spectroscopy of polycyclic aromatic hydrocarbons: the realm of anharmonicity

Maltseva, E.; Petrigani, A.; Candian, A.; Mackie, C.J.; Huang, X.; Lee, T.J.; Tielens, A.G.G.M.; Oomens, J.; Buma, W.J.

DOI

[10.1088/0004-637X/814/1/23](https://doi.org/10.1088/0004-637X/814/1/23)

Publication date

2015

Document Version

Final published version

Published in

Astrophysical Journal

[Link to publication](#)

Citation for published version (APA):

Maltseva, E., Petrigani, A., Candian, A., Mackie, C. J., Huang, X., Lee, T. J., Tielens, A. G. G. M., Oomens, J., & Buma, W. J. (2015). High-resolution IR absorption spectroscopy of polycyclic aromatic hydrocarbons: the realm of anharmonicity. *Astrophysical Journal*, 814(1), Article 23. <https://doi.org/10.1088/0004-637X/814/1/23>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

HIGH-RESOLUTION IR ABSORPTION SPECTROSCOPY OF POLYCYCLIC AROMATIC HYDROCARBONS: THE REALM OF ANHARMONICITY

ELENA MALTSEVA¹, ANNEMIEKE PETRIGNANI^{2,3}, ALESSANDRA CANDIAN², CAMERON J. MACKIE², XINCHUAN HUANG⁴,
TIMOTHY J. LEE⁴, ALEXANDER G. G. M. TIELENS², JOS OOMENS³, AND WYBREN JAN BUMA¹

¹University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands; w.j.buma@uva.nl

²Leiden Observatory, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands

³Radboud University, Toernooiveld 7, 6525 ED Nijmegen, The Netherlands; petrignani@strw.leidenuniv.nl

⁴SETI Institute, 189 Bernardo Avenue, Suite 100, Mountain View, CA 94043, USA

Received 2015 June 25; accepted 2015 October 12; published 2015 November 12

ABSTRACT

We report on an experimental and theoretical investigation of the importance of anharmonicity in the 3- μm CH stretching region of polycyclic aromatic hydrocarbon (PAH) molecules. We present mass-resolved, high-resolution spectra of the gas-phase cold (~ 4 K) linear PAH molecules naphthalene, anthracene, and tetracene. The measured IR spectra show a surprisingly high number of strong vibrational bands. For naphthalene, the observed bands are well separated and limited by the rotational contour, revealing the band symmetries. Comparisons are made to the harmonic and anharmonic approaches of the widely used Gaussian software. We also present calculated spectra of these acenes using the computational program SPECTRO, providing anharmonic predictions with a Fermi-resonance treatment that utilizes intensity redistribution. We demonstrate that the anharmonicity of the investigated acenes is strong, dominated by Fermi resonances between the fundamental and double combination modes, with triple combination bands as possible candidates to resolve remaining discrepancies. The anharmonic spectra as calculated with SPECTRO lead to predictions of the main bands that fall within 0.5% of the experimental frequencies. The implications for the aromatic infrared bands, specifically the 3- μm band, are discussed.

Key words: astrochemistry – ISM: molecules – line: identification – methods: laboratory: molecular – techniques: spectroscopic

1. INTRODUCTION

Polycyclic aromatic hydrocarbon (PAH) species are considered to be responsible for the family of infrared (IR) emission features, the so-called aromatic infrared bands (AIBs), that dominate the spectrum of objects ranging from protoplanetary disks to entire galaxies (see, Joblin & Tielens 2011, and reference therein). PAH molecules absorb visible to ultraviolet (UV) photons and then relax through IR emission (Sellgren 1984; Allamandola et al. 1989; Puget & Léger 1989). These spectral signatures are due to vibrational modes that are typical of a family of molecules showing the same chemical bond types.

Over the years, a wealth of theoretical (see, e.g., the NASA Ames and Cagliari databases; Mallocci et al. 2007; Boersma et al. 2011, 2014a) and experimental (see, e.g., Oomens 2011, and reference therein) studies of the vibrational spectrum of different types of PAH molecules (neutral, charged, heteroatom substituted, etc.) have been reported with the aim of aiding the identification of subclasses of PAHs when compared of the global characteristics of the AIBs in well-known astronomical sources (Boersma et al. 2012, 2013, 2014b; Candian et al. 2012).

Both theoretical and experimental studies have known caveats. Apart from rare exceptions (Cané et al. 2007; Pirali et al. 2009; Maris et al. 2015), density functional theory (DFT) calculations have been performed exclusively with the double harmonic approximation, thus neglecting the effects of anharmonicity. The calculations by Pirali et al. (2009) have been the most extensive in incorporating the effects of anharmonicity, but have been restricted to naphthalene. Moreover, Pirali et al. (2009) focused on addressing the detailed profiles of the lower frequency, IR active modes, and in

particular, assessing the relative contribution of anharmonicity effects (e.g., hot bands) and rotational structure to measured emission profiles. More specifically Pirali et al. (2009) considered Fermi (and Darling–Dennison) resonances. What was not included was derivatives beyond the first order of the dipole moment. In contrast, here we present the first results of an extensive study on the effects of anharmonicity on the IR spectrum of PAHs, combining high-resolution absorption spectra with theoretical studies which allow for anharmonicity as well as Fermi resonances. As the effects of anharmonicity and Fermi resonances are expected to be the largest for the CH stretching modes, we focus here on the 3000 cm^{-1} region. In addition, unlike almost all previous gas-phase studies—with the notable exception of Huneycutt et al. (2004)—, the spectra are measured at very low temperature in order to limit the effects of rotational broadening that may obstruct observation of individual bands and to enable an easier and unambiguous assignment of bands. Under such conditions we also avoid (mode-specific) perturbations by the matrix environment that are difficult to predict but inevitably present in matrix-isolation spectroscopy (MIS) studies. As will be shown in the present study, these aspects (resolution and isolated molecules) are key to assessing the quality of the calculations and fine-tuning the computational approach.

Commonly, correction factors of about 0.96 (Langhoff 1996) are applied to calculated harmonic frequencies to bring them in line with peak positions measured using MIS techniques. As the latter are affected by ill-understood matrix effects, proper validation of the theoretical spectra against laboratory spectra of cold, gas-phase PAHs are imperative before they can be used in analysis studies of observed interstellar spectra. To properly account for their astrophysical properties, supersonic molecular

beam methods in combination with laser spectroscopy provide an attractive way to study the vibrational spectrum of very cold PAH molecules under isolated conditions at a resolution that is only determined by the laser bandwidth. Recent advances in computational chemistry (Bloino & Barone 2012) enable efficient anharmonic vibrational analyses of medium sized molecules, providing theoretical tools to properly interpret the experimental results.

In the present study we apply IR–UV double resonance spectroscopy to record mass-selected IR absorption spectra in the $3\ \mu\text{m}$ region of three small linear PAH molecules: naphthalene, anthracene, and tetracene. We show that these spectra are at odds with harmonic and anharmonic calculations as employed so far, but are in good agreement with our new anharmonic spectra calculated with the program SPECTRO that properly incorporates multiple resonances.

2. METHODS

2.1. Experimental

IR absorption spectra were recorded using an IR–UV depletion scheme. The molecular beam setup employed for these measurements has been described in detail before (Smolarek et al. 2011). Briefly, a heated supersonic pulsed source with a typical pulse duration of $190\ \mu\text{s}$ and argon at a backing pressure of 2.2 bar as a buffer gas were used to create a cold molecular beam. A constant mass-selected ion signal was created via a two-color Resonant Enhanced MultiPhoton Ionization (REMPI) scheme using a Nd:YAG pumped frequency-doubled dye laser to excite the molecules to the first excited state and then an ArF excimer laser to ionize the excited molecules. The resonant excitation in combination with the supersonic cooling conditions ensures that all of the bands start from the vibrationless ground state. The IR light in the $3\text{-}\mu\text{m}$ region produced by a Nd:YAG pumped frequency-mixed dye laser with a line width of $0.07\ \text{cm}^{-1}$ preceded these two laser beams by 200 ns, leading to dips in the ion signal upon IR absorption. The mass-resolved nature of our experiments and the use of supersonically cooled molecules ensures that in our experiments we can unambiguously conclude that all of the measured absorption features concern excitations from the vibrational ground state of each PAH, and that they do not derive from isotopically substituted species.

2.2. Computational Details

We employed three computational methods; the standard harmonic and anharmonic vibrational approaches, both using Gaussian 09 (Frisch et al. 2009), and an anharmonic method utilizing a modified version (Martin et al. 1995) of the SPECTRO program (Willetts et al. 1990; Gaw et al. 1996) that incorporates intensity sharing due to Fermi resonances (Mackie et al. 2015b), referred to in this work as G09-h, G09-anh, and SP15, respectively. All calculations apply DFT using a similar integration grid as in Boese & Martin (2004), the B9-71 functional (Hamprrecht et al. 1998), and the TZ2P basis set (Dunning 1971), which provide the best performance on organic molecules (Boese & Martin 2004; Cané et al. 2007). SP15 takes the G09-h intensities and G09-anh force constants and implements its own vibrational second-order perturbation method (Mackie et al. 2015a, 2015b). It treats couplings between multiple resonances (states falling within $200\ \text{cm}^{-1}$ of each other) simultaneously and redistributes intensity among




Figure 1. Absorption spectra of naphthalene as predicted with (a) SP15, (b) G09-anh, and (c) G09-h together with the measured spectrum (this exp.).

the modes. Traditionally, two vibrational states of the same symmetry with energies close to each other may interact. The resonance types included in our second-order perturbation theory treatment are beyond the traditional Fermi I and II type, and also include 1-1 and 2-2 type resonances. The coupling terms involve both cubic and quartic anharmonic force constants, and the explicit formula for the 1-1 and 2-2 type resonances are given in Lehmann (1988, 1993). The implementation of this polyad approach into SPECTRO is described in Martin et al. (1995). No intensity resonances are considered in this study, as we use the Gaussian double-harmonic intensities for intensity distributions.

3. RESULTS AND DISCUSSION

All IR absorption spectra were recorded between 3.17 and $3.4\ \mu\text{m}$ (2950 and $3150\ \text{cm}^{-1}$). With the present signal-to-noise ratios (S/Ns), no other IR bands were observed outside the displayed range. The respective ion signals were created by fixing the frequency of the first UV photon to the 0-0 band of the $S_1 \leftarrow S_0$ electronic transition of the respective PAHs. The exact frequencies used are $32028.18\ \text{cm}^{-1}$, $27697.0\ \text{cm}^{-1}$, and $22402.43\ \text{cm}^{-1}$ for naphthalene, anthracene, and tetracene, respectively. These were determined by scanning a small range around the previously reported values from Hiraya et al. (1985), Lambert et al. (1984), Zhang et al. (2008) for the $S_1(^1B_{3u}) \leftarrow S_0(^1A_g)$, $S_1(^1B_{1u}) \leftarrow S_0(^1A_g)$, and $S_1(^1B_{2u}) \leftarrow S_0(^1A_g)$ transitions, respectively.

3.1. Naphthalene ($C_{10}H_8$)

Figure 1 displays the experimental and theoretical IR absorption spectra for naphthalene. The spectrum shows narrow bands with widths varying between 1 and $3\ \text{cm}^{-1}$ and more

Table 1The Measured Bands of Naphthalene, Anthracene, and Tetracene with Frequencies in cm^{-1} and Intensities Normalized to the Respective Strongest Transitions

Naphthalene (C_{10}H_8)			Anthracene ($\text{C}_{14}\text{H}_{10}$)			Tetracene ($\text{C}_{18}\text{H}_{12}$)		
This Work			CRDS	This Work		CRDS	This Work	
Freq.	Rel. Int.	Symm.	Freq.	Freq.	Rel. Int.	Freq.	Freq.	Rel. Int.
2963.8	0.2	...	2965.3	2973.2	0.10	...	3008.9	0.39
2972.4	0.3	b _{1u}	2973.1	2979.6	0.08	...	3015.3	0.2
2981.3	0.2	...	2980.9	2992.5	0.07	...	3023.7	0.44
2989.0	0.28	b _{1u}	2989.1	3011.8	0.15	...	3032.6	0.4
3014.0	0.3	b _{2u}	3013.7	3022.0	0.30	3021.7	3038.5	0.45
3029.0	0.31	b _{1u}	3029.1	3030.0	0.25	...	3039.7	0.46
3034.5 [*]	0.15	b _{1u}	...	3033.7	0.20	3032.1	3046.2	0.34
3039.5 [*]	0.19	b _{2u}	...	3046.7	0.30	3047.5	3050.6	0.18
3042.3 ^(*)	0.29	b _{1u}	...	3055.4	0.54	...	3054.8	0.38
3043.8 ^(*)	0.35	b _{2u})3043.7	3062.3	0.45	...	3056.8	0.51
3048.2	0.19	b _{2u}	3048.9	3065.3	0.72	3064.3	3061.1	1
3052.2 [*]	0.17	3066.9	0.64	3067.5	3066.6	0.2
3058.1	0.65	b _{2u}	3057.9	3071.9	1.00	3072.5	3069.5	0.23
3060.5	0.54	b _{1u}	3061.1	3077.8	0.40	3077.9	3077.6	0.8
3065.2	0.99	b _{1u}	3065.1	3081.8	0.24	...	3080.4	0.27
3071.4	0.1	b _{2u}	3069.9	3095.9	0.24	...	3087.9	0.23
3076.2	0.37	b _{2u}	3074.1	3109.6	0.32	3109.9	3094.1	0.31
3079.2	1	b _{2u}	3079.1	3098.8	0.38
3083.9 [*]	0.16	b _{2u}	3101.5	0.19
3092.6 [*]	0.21
3100.2	0.45	b _{2u}	3100.1
3102.6	0.21	...	3102.2
3109.4	0.45	b _{2u}	3109.3

Note. Comparison is made to previous gas-phase frequencies (cm^{-1}) recorded with cavity ring down spectroscopy (CRDS; Huneycutt et al. 2004).

^{*} These are the newly measured lines. The brackets denote that these two lines were previously measured as one.

than 16 well-separated bands of which a considerable fraction have intensities exceeding 20% of the strongest band. In total, 23 experimental lines are identified and listed in Table 1. Based on group theory, four IR-active transitions are present in the CH stretch region, with either b_{1u} or b_{2u} symmetries.⁵ Indeed, G09-h predicts two intense bands that may be associated with the strongest and the two less intense bands that are of ambiguous assignment. A scaling factor of 0.966 was found to give the best agreement with experiments, in close agreement with Cané et al. (2007), considering that only the CH stretch region is considered here.

Table 1 also shows the line positions (in cm^{-1}) reported in a previous study using cavity ring-down spectroscopy (CRDS) (Huneycutt et al. 2004). Most experimental bands are within 0.2 cm^{-1} . Some of the weaker bands deviate up to 1 cm^{-1} , which may be attributed to the inaccuracy with which the center can be determined. Our error in the positions is most likely smaller due to superior S/N. We observe six additional bands not reported before (asterisks in Table 1 and Figure 1). Our spectra do not give evidence of the presence of the very weak bands reported before in Huneycutt et al. (2004) (3064.1 , 3068.5 , and 3098.9 cm^{-1}). The present study also shows narrower line widths, indicating lower internal temperatures. The improved resolution significantly changes the intensity distribution of the majority of the bands; e.g., the bands around 3100 cm^{-1} show the smallest line widths (1.0 and 1.1 cm^{-1} , respectively) and have intensities of about 45% of the intensity of the strongest band, whereas in the CRDS study the intensity

⁵ b_{3u} transitions are IR-active, however, none with this symmetry fall in the CH stretch region.

of these bands was less than 20% and displayed a broader structure.

Figure 1 also shows the spectrum of naphthalene as predicted with G09-anh, convolved with 1 cm^{-1} to resemble experimental resolution. A total of 11 double excitation bands are predicted, i.e., bands arising from $\Delta\nu_i = 2$ or $\Delta\nu_i = 1$; $\Delta\nu_j = 1$ transitions. The G09-anh frequencies are in very good agreement with experiments, the typical deviation of the strongest bands being only 0.6%. This remaining deviation has been corrected for in the figure by a small scaling factor which also serves to facilitate easy comparison between observed and predicted spectra. Interestingly, none of the anharmonic bands have intensities comparable to those observed in experiments. This lack of intensity can arise from two possible scenarios. First, intensity is obtained from the anharmonicity of the potential energy and dipole moment surfaces along the relevant coordinates, making the $\Delta\nu = 1$ selection rule no longer strictly valid. Second, intensity is acquired through Fermi coupling to the CH-stretch fundamentals, leading to intensity borrowing.

To distinguish between these two, experiments on deuterated naphthalene were performed as the fundamental CD-stretch modes of naphthalene-d₈ are displaced to $\sim 4 \mu\text{m}$ while the overtones and combination bands originate from normal modes that are much less affected by deuteration. In the first scenario only slight shifts are expected, and in the second case deuteration will lead to a dramatic reduction of intensity. We scanned a range of $3.12\text{--}3.85 \mu\text{m}$ ($2600\text{--}3200 \text{ cm}^{-1}$) taking into account the possible shifts of the combination bands due to deuteration, and found no signal. We thus conclude that the additional bands in naphthalene-h₈ derive their intensity from

Fermi coupling to fundamental CH-stretch transitions. This means that all bands are either of b_{1u} or b_{2u} symmetry with corresponding rotational contours, the envelope of b_{2u} bands being narrower than for b_{1u} bands. A rotational-contour analysis of the two strongest bands gives the best agreement using a rotational temperature of 4 K and a homogeneous line width of 0.5 cm^{-1} . Indeed, two sets of band widths were measured; relatively large widths between 2.3 and 2.8 cm^{-1} attributed to b_{1u} transitions and narrower widths between 1.1 and 1.9 cm^{-1} attributed to b_{2u} transitions.

To properly include Fermi resonances, we calculated the spectrum of naphthalene using SP15. Like G09-anh, in total 11 IR-active combination bands are within the experimental range. The slight overestimation of the anharmonicity is even smaller, within 0.4%. Unlike G09-anh, SP15 does give appropriate intensities and the agreement with experiments is improved although discrepancies remain. The major discrepancy concerns the number of observed bands. The calculations predict 15 bands while 23 bands are observed. We therefore performed a combinatorial analysis based on experimental data. We determined the frequencies of all possible double combination bands using the sums of measured fundamental frequencies of naphthalene in the $6 \mu\text{m}$ region (see Hewett et al. 1994 and references therein), including both IR and Raman active modes. On the basis of these modes (all CC stretches except for one CH bend) and the symmetry restrictions, 11 IR-active combination bands fall within a broad range of 3.15 – $3.5 \mu\text{m}$ (2900 – 3150 cm^{-1}). Both calculation and combinatorial analysis are thus not able to account for all the observed bands. A similar exercise can be performed for triple combination bands, which provides a multitude of additional candidates with the correct symmetry that fall within the experimental range. Interestingly, we find that these dominantly involve CH bending modes, which are prone to coupling with CH stretching modes. We conclude that the $3 \mu\text{m}$ region of the absorption spectrum of naphthalene is dominated by Fermi resonances and that triple combination bands need to be taken into account as well. The latter would require incorporation of even higher-order couplings than currently included in the anharmonic analyses. A detailed assignment of the double combination bands in naphthalene as well as anthracene and tetracene (*vide infra*) will be presented in a separate study Mackie et al. (2015b). Also, methods that incorporate higher-order terms are presently under investigation and will be presented in future work.

3.2. Anthracene ($C_{14}H_{10}$)

The experimental and theoretical absorption spectra of anthracene are shown in Figure 2. The positions of the observed bands are listed in Table 1. Again, the experimental spectrum reveals more bands with appreciable intensities than predicted by harmonic calculations, which gives the best agreement, with a scaling factor of 0.966. The G09-anh spectrum again slightly overestimates the anharmonicity, requiring a scaling factor less than 0.5% to overlap the strongest bands. As before, G09-anh predicts many additional bands with low to zero intensity. Interestingly, the band at $3.2 \mu\text{m}$ is much higher than observed in experiment and might be caused by a missing resonance. Indeed, incorporating Fermi coupling using SP15 permits us to redistribute intensity over multiple transitions, which not only leads to a lower intensity




Figure 2. Absorption spectra of anthracene as predicted with (a) SP15, (b) G09-anh, and (c) G09-h together with the measured spectrum (this exp.).

band at $3.22 \mu\text{m}$ but also to better overall agreement with experiment.

Table 1 shows the comparison to the previous CRDS study (Huneycutt et al. 2004). We measure seven more bands, which can be attributed to improved cooling conditions. All other band positions are within experimental error, with the exception of the previously reported band at 3032.1 cm^{-1} . This band consists of two close-lying bands at 3030.0 and 3033.7 cm^{-1} in our experiment (arrows in Figure 2). Bands with similar widths as in naphthalene are observed, indicating similar cooling conditions ($\sim 2.5 \text{ cm}^{-1}$). Simulation of the rotational contours that can be expected for sensible rotational temperatures under the present experimental conditions show—in contrast to what was observed for naphthalene—that the widths of the observed bands are not determined by their rotational contours. Instead, one has to conclude that these bands consist of several overlapping bands, and that a combinatorial analysis of the entire spectrum as was done for naphthalene is not possible. Moreover, the state density in anthracene at these energies is already so large that this approach would likely not lead to a conclusive assignment.

3.3. Tetracene ($C_{18}H_{12}$)

Figure 3 shows the experimental and theoretical absorption spectra of tetracene. We observe more than 19 well-resolved bands with line widths between 2.3 and 4.4 cm^{-1} (Table 1). As for anthracene, it is most likely that the width of these bands is not determined by the rotational contour of one specific transition, but results from the overlap of several unresolved transitions. The G09-h calculation predicts that only four of the twelve symmetry-allowed CH stretch bands have an appreciable intensity. For these four bands, a rather poor agreement




Figure 3. Absorption spectra of tetracene as predicted with (a) SP15, (b) G09-anh, and (c) G09-h together with the measured spectrum (this exp.).

with the experiments is observed, in contrast to naphthalene and anthracene, where a reasonably good agreement could be obtained for the fundamental bands after scaling. The G09-anh calculation, on the other hand, performs much better in this case; its frequencies are in close agreement with the experimentally observed strong bands. When including the Fermi coupling with SP15, even better agreement can be achieved.

Since for tetracene no gas-phase IR absorption spectra have been reported before, we compare the present data with data obtained in Ar matrix isolation studies (MIS) (Hudgins & Sandford 1998a). Figure 4 shows all measured spectra compared to their MIS counterparts. The agreement in all three cases is very high, with superior S/N and cooling in the present study, leading to narrower and more resolved lines. As expected from matrix-induced effects (Joblin et al. 1994), relatively small spectral shifts are observed for most bands. The larger shifts are probably due to a redistribution of relative intensities; e.g., for tetracene, an apparent large shift (of 6.1 cm^{-1}) seems to occur for the most intense band; however, it is more likely that the relative intensities of the double structure around $3.27 \mu\text{m}$ are different for the MIS spectrum, leading to another intense band.

4. ASTROPHYSICAL IMPLICATIONS

The present results show that harmonic DFT calculations fail dramatically at predicting high-resolution experimental absorption spectra of PAHs in the $3\text{-}\mu\text{m}$ region. This is important since such calculations are routinely used to help in the interpretation of astronomical observations, and in particular, in efforts to shed light on the evolution of the interstellar carbon inventory. In the past, the discrepancies between frequencies




Figure 4. Absorption spectra of (a) naphthalene, (b) anthracene, and (c) tetracene (this exp.) compared to matrix isolation spectroscopy measurements (MIS) (Hudgins & Sandford 1998a).

calculated in the harmonic approximation and MIS spectra have commonly been overcome by introducing a scaling factor of 0.966, corresponding to a shift in the order of 100 cm^{-1} . In this context, five key observations are made in the present study.

First, MIS spectra agree well with the gas phase spectra measured here in terms of the position of the bands. However, there are differences in relative intensities, reflecting the importance of Fermi resonances (which act differently in a matrix than in the gas phase), and this may lead to subtle shifts in the intensity averaged peak position (Huneycutt et al. 2004).

Second, for the strongest bands in the spectrum, the anharmonic predictions obtained using SP15 all fall within 0.5% from the experimental value, significantly reducing the error introduced by the use of empirical scaling factors.

Third, in order to “translate” spectroscopic data obtained from calculations or low-temperature experiments into emission spectra useful for comparison with observational data, a typical empirical redshift of 15 cm^{-1} is commonly applied (Bauschlicher et al. 2010). As demonstrated by the pioneering study on the low-frequency modes of naphthalene (Pirali et al. 2009), mode-specific shifts in emission spectra are more accurately predictable as our knowledge of the anharmonic potential expands. We are presently working on such modeling efforts.

Fourth, Fermi resonances may contribute significantly to the profile and structure of the $3\text{-}\mu\text{m}$ band. The frequencies of the major CH-stretch fundamental bands are within a spread of around 50 cm^{-1} , about the same for all types of PAHs (Hudgins & Sandford 1998a, 1998b). The frequencies of the modes that are potentially involved in combination bands, on

the other hand, are much more sensitive to the finer details of the structure. These observations match astronomical studies of IR emission that show a very prominent emission band at $3.29\ \mu\text{m}$, corresponding to the 1-0 transition of the fundamental CH-stretch bands, and a broad plateau in the $3.1\text{--}3.7\ \mu\text{m}$ region indicated as the $3\text{-}\mu\text{m}$ plateau or vibrational quasi-continuum (Allamandola et al. 1989; Geballe et al. 1989). DFT calculations so far have not been able to explain and characterize this plateau. Our studies demonstrate that anharmonic couplings can be responsible for some of the structure observed on the main band as well as the wings on both the blue and the red side of the main astronomical band.

Fifth, as our study demonstrates anharmonicity and Fermi resonances are particularly important in the $3\text{-}\mu\text{m}$ region of the spectrum. An assessment of their influence for the other bands remains to be determined.

It may be noticed that the spectral range over which features are observed when comparing the three experimental spectra seems to become narrower with increasing PAH size. This aspect is further discussed in separate studies on the size and structure dependence of the $3\text{-}\mu\text{m}$ region (Mackie et al. 2015b; C. J. Mackie et al. 2015, in preparation; E. Maltseva et al. 2015, in preparation).

5. CONCLUSIONS

We report on the absorption spectra of linear PAHs in the $3\text{-}\mu\text{m}$ region using UV–IR double resonance laser spectroscopy under cold and isolated conditions. Efficient cooling and excellent S/N lead to well-resolved spectra that show a plethora of vibrational transitions. Comparison with harmonic predictions shows that the fraction of intensity that is not associated with fundamental transitions may easily exceed 50%, and therefore cannot be neglected. A detailed analysis of the absorption spectrum of naphthalene has demonstrated that the additional activity originates from vibrational coupling of the bath of “dark” states with the intensity-carrying “bright” states, and not purely from anharmonic effects. The present studies emphasize the necessity of properly incorporating Fermi resonances and higher-order vibrational couplings. First results incorporating Fermi resonances with SPECTRO are rather promising, as they indeed seem to lead to a qualitatively correct description of the intensity distribution. However, the measured spectra show more bands than the number of possible double combination bands. This indicates that incorporation of higher-order couplings, i.e., triple combination bands involving CH bending modes, may be necessary to obtain a proper quantitative description.

Detailed studies of the spectra of PAHs are very timely. The launch of the *James Webb Space Telescope* will open up a revolutionary window on the PAH spectrum, with unprecedented spatial and spectral resolution. Studies on larger PAHs are needed to determine in more detail how the size of a PAH affects the intensity distribution over the $3\text{-}\mu\text{m}$ band. Analogous studies on molecules with the same number of rings but with a different structure—leading among others to the presence of bay and non-bay hydrogen atoms—are also of considerable interest: if a relation can be derived between the structure of the molecule and the shape and position of the $3\text{-}\mu\text{m}$ band, progress could be made toward determining the chemical composition of interstellar objects (Candian et al. 2012). Such IR absorption studies on larger PAHs and

on PAHs with different structures are presently being performed in our lab (E. Maltseva et al. 2015, in preparation).

The experimental work was supported by The Netherlands Organization for Scientific Research (NWO). Studies of interstellar PAHs at Leiden Observatory have been supported through the advanced European Research Council Grant 246976 and a Spinoza award. Computing time has been made available by NWO Exacte Wetenschappen (project MP-270-13 and MP-264-14) and calculations were performed at the LISA Linux cluster of the SurfSARA supercomputer center in Almere, The Netherlands. X.H. and T.J.L. gratefully acknowledge support from the NASA 12-APRA12-0107 grant. X.H. acknowledges the support from NASA/SETI Co-op Agreement NNX15AF45A.

REFERENCES

- Allamandola, L. J., Tielens, A. G. G. M., & Barker, J. R. 1989, *ApJS*, **71**, 733
 Bauschlicher, C. W., Jr., Boersma, C., Ricca, A., et al. 2010, *ApJS*, **189**, 341
 Bloino, J., & Barone, V. 2012, *JChPh*, **136**, 124108
 Boersma, C., Allamandola, L. J., Bauschlicher, C. W., Jr., et al. 2011, *EAS Publications Series*, **46**, 109
 Boersma, C., Bauschlicher, C. W., Jr., & Ricca, A. 2014a, *ApJS*, **211**, 8
 Boersma, C., Bregman, J. D., & Allamandola, L. J. 2013, *ApJ*, **769**, 117
 Boersma, C., Bregman, J. D., & Allamandola, L. J. 2014b, *ApJ*, **795**, 110
 Boersma, C., Rubin, R. H., & Allamandola, L. J. 2012, *ApJ*, **753**, 168
 Boese, A. D., & Martin, J. M. L. J. 2004, *Phys. Chem. A*, **108**, 3085
 Candian, A., Kerr, T. H., Song, I.-O., McCombie, J., & Sarre, P. J. 2012, *MNRAS*, **426**, 389
 Cané, E., Miani, A., & Trombetti, A. 2007, *JPCA*, **111**, 8111
 Dunning, T. H. J. 1971, *CP*, **55**, 716
 Frisch, M. J., Trucks, G. W., Schlegel, H. B., et al. 2009, Gaussian 09.D01 (Pittsburgh, PA: Gaussian, Inc.)
 Gaw, J. F., Willets, A., Green, W. H., & Handy, N. C. 1996, SPECTRO Program, Version 3.0
 Geballe, T. R., Tielens, A. G. G. M., Allamandola, L. J., Moorhouse, A., & Brand, P. W. J. L. 1989, *ApJ*, **341**, 278
 Hamprecht, F. A., Cohen, A. J., Tozer, D. J., & Handy, N. C. J. 1998, *JChPh*, **109**, 6264
 Hewett, K. B., Shen, M., Brummel, C. L., & Philips, L. A. 1994, *JChPh*, **100**, 4077
 Hiraya, A., Achiba, Y., Mikami, N., & Kimura, K. 1985, *JChPh*, **82**, 1810
 Hudgins, D. M., & Sandford, S. A. 1998a, *JPCA*, **102**, 329
 Hudgins, D. M., & Sandford, S. A. 1998b, *JPCA*, **102**, 344
 Huneycutt, A. J., Casaes, R. N., McCall, B. J., et al. 2004, *ChemPhysChem*, **5**, 321
 Joblin, C., d’Hendecourt, L., Léger, A., & Défourneau, D. 1994, *ApJ*, **281**, 923
 Joblin, C., & Tielens, A. G. G. M. (ed.) 2011, in *EAS Publications Series 46, PAHs and the Universe (Les Ulis: EDP Sciences)*, 191
 Lambert, W. R., Felker, P. M., Syage, J. A., & Zewail, A. H. 1984, *JChPh*, **81**, 2195
 Langhoff, S. R. 1996, *JPhCh*, **100**, 8
 Lehmann, K. K. 1988, *MolPh*, **66**, 1129
 Lehmann, K. K. 1993, *MolPh*, **75**, 739
 Mackie, C. J., Candian, A., Huang, X., et al. 2015a, *JChPh*, **142**, 244107
 Mackie, C. J., Candian, A., Huang, X., et al. 2015b, *JChPh*, submitted
 Mallocci, G., Joblin, C., & Mulas, G. 2007, *CP*, **332**, 353
 Maris, A., Calabrese, C., Melandri, S., & Blanco, S. 2015, *JChPh*, **142**, 024317
 Martin, J. M., Lee, T. J., Taylor, P. R., & François, J. P. 1995, *JChPh*, **103**, 2589
 Oomens, J. 2011, *EAS Publications Series*, **46**, 61
 Pathak, A., & Rastogi, S. 2005, *CP*, **313**, 133
 Pirali, O., Vervloet, M., Mulas, G., Mallocci, G., & Joblin, C. 2009, *PCCP*, **11**, 3443
 Puget, J. L., & Léger, A. 1989, *ARA&A*, **27**, 161
 Sellgren, K. 1984, *ApJ*, **277**, 623
 Smolarek, S., Vdovin, A., Rijs, A., et al. 2011, *JPCA*, **115**, 9399
 Willets, A., Gaw, J. F., Green, W. H., & Handy, N. C. 1990, in *Advances in Molecular Vibrations and Collision Dynamics*, ed. J. M. Bowman (Greenwich, CT: JAI), 169
 Zhang, J., Pei, L., & Kong, W. 2008, *JChPh*, **128**, 104301



ERRATUM: “HIGH-RESOLUTION IR ABSORPTION SPECTROSCOPY OF POLYCYCLIC AROMATIC HYDROCARBONS: THE REALM OF ANHARMONICITY” (2015, ApJ, 814, 23)

ELENA MALTSEVA¹, ANNEMIEKE PETRIGNANI^{2,3}, ALESSANDRA CANDIAN², CAMERON J. MACKIE², XINCHUAN HUANG⁴,
TIMOTHY J. LEE⁵, ALEXANDER G. G. M. TIELENS², JOS OOMENS³, AND WYBREN JAN BUMA¹

¹University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands; w.j.buma@uva.nl

²Leiden Observatory, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands; petrignani@strw.leidenuniv.nl

³Radboud University, Toernooiveld 7, 6525 ED Nijmegen, The Netherlands

⁴SETI Institute, 189 Bernardo Avenue, Suite 100, Mountain View, CA 94043, USA

⁵NASA Ames Research Center, Moffett Field, CA 94035-1000, USA

Received 2016 February 4; accepted 2016 February 16; published 2016 March 21

In the original manuscript, the middle panel of Figure 3 is accidentally identical to the middle panel of Figure 2. This erratum shows Figure 3 with the correct middle panel. Table 1 is unaffected, since it reports only experimental data. This change does not affect the scientific results of the paper.




Figure 3. Absorption spectra of tetracene as predicted with (a) SP15, (b) G09-anh, and (c) G09-h together with the measured spectrum.