



## UvA-DARE (Digital Academic Repository)

### Of moths, mites and microbes - The role of bacteria in the life history of two arthropod herbivores

Staudacher, H.

**Publication date**

2016

**Document Version**

Final published version

[Link to publication](#)

**Citation for published version (APA):**


Staudacher, H. (2016). *Of moths, mites and microbes - The role of bacteria in the life history of two arthropod herbivores*. [Thesis, fully internal, Universiteit van Amsterdam].

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).



**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.




# Of Moths, Mites and Microbes

The role of bacteria in the life history of two arthropod herbivores



**Heike Staudacher**





# **OF MOTHS, MITES AND MICROBES**

**THE ROLE OF BACTERIA IN THE LIFE HISTORY  
OF TWO ARTHROPOD HERBIVORES**

Staudacher H., 2016. Of moths, mites and microbes – The role of bacteria in the life history of two arthropod herbivores

PhD thesis, University of Amsterdam, The Netherlands

ISBN: 978 94 91407 29 1

Cover design: Heike Staudacher & Jan van Arkel

Thesis lay-out: Heike Staudacher, with assistance of Jan Bruin

# **OF MOTHS, MITES AND MICROBES**

## **THE ROLE OF BACTERIA IN THE LIFE HISTORY OF TWO ARTHROPOD HERBIVORES**

ACADEMISCH PROEFSCHRIFT

ter verkrijging van de graad van doctor  
aan de Universiteit van Amsterdam  
op gezag van de Rector Magnificus  
prof. dr. D.C. van den Boom  
ten overstaan van een door het college voor promoties  
ingestelde commissie, in het openbaar te verdedigen  
in de Aula der Universiteit

op vrijdag 19 februari 2016, te 11.00 uur

door  
HEIKE STAUDACHER  
geboren te Schramberg, Duitsland

***Promotiecommissie***

Promotor	prof. dr. S.B.J. Menken <sup>1</sup>
Co-promotor	dr. A.T. Groot <sup>1,2</sup>
Overige leden	prof. dr. J.C. Biesmeijer <sup>1</sup> dr. J.A.J. Breeuwer <sup>1</sup> prof. dr. M. Kaltenpoth <sup>2,3</sup> prof. dr. G. Muijzer <sup>1</sup> dr. J.C. de Roode <sup>4</sup> prof. dr. M. Schilthuisen <sup>5</sup> dr. I.M. Smallegange <sup>1</sup>

***Faculteit der Natuurwetenschappen, Wiskunde en Informatica***

<sup>1</sup>University of Amsterdam, The Netherlands

<sup>2</sup>Max Planck Institute for Chemical Ecology, Jena, Germany

<sup>3</sup>Johannes Gutenberg University, Mainz, Germany

<sup>4</sup>Emory University, Georgia, USA

<sup>5</sup>Leiden University, The Netherlands

# Contents

Chapter 1	General introduction	7
Chapter 2	Variability of bacterial communities in the moth <i>Heliothis virescens</i> indicates transient association with the host	27
Chapter 3	Sex-specific consequences of an induced immune response on reproduction in a moth	53
Chapter 4	Effects of immune challenge on the oviposition strategy of a noctuid moth	77
Chapter 5	The bacterial symbionts <i>Wolbachia</i> , <i>Cardinium</i> and <i>Spiroplasma</i> affect gene expression and survival of their spider mite host and impact distinct induced responses in plants	97
Chapter 6	General discussion	159
	Summary	174
	Samenvatting	177
	Author contributions	180
	Author addresses	181
	Biography	182
	Acknowledgements	183



1

## General introduction

## General introduction

Microbes are omnipresent in the environment of macro-organisms and innumerable examples of interactions between micro and macro-organisms have been described. Microbes may colonize different organs or body cavities of macro-organisms and are then referred to as ‘symbionts’ that reside in a ‘host’. Symbiosis was originally defined as the ‘living together of unlike organisms’ by Anton de Bary (1879). Bacterial symbionts can have beneficial, neutral (commensal symbiont), harmful (parasites and pathogens) or mixed effects on the host (Dillon & Dillon, 2004; Baumann, 2005; Werren *et al.*, 2008; Himler *et al.*, 2011; Engel & Moran, 2013). The role of symbionts in hosts may also change during evolution and may be context dependent and thus differ under different circumstances (Werren *et al.*, 2008; Zug & Hammerstein, 2015). First, I will describe and discuss non-pathogenic microbe-host interactions. Thereafter, I will focus on pathogens and the effects of immune challenge on host life history.

### ***1. Non-pathogenic associations of bacteria with arthropod hosts***

Symbioses can be categorized according to the co-dependence between symbionts and their hosts. Long-term, stable associations between symbionts and hosts are more likely to lead to mutualistic symbiosis than short-term, loose associations, because host and symbionts together form a unit of selection during evolution (Zilber-Rosenberg & Rosenberg, 2008; Rosenberg *et al.*, 2009; Salem *et al.*, 2015).

#### *Obligate endosymbionts*

Obligate bacterial endosymbionts are strictly necessary for host survival, growth and/or reproduction. Obligate symbioses are evolutionarily ancient and the phylogeny of these so-called primary endosymbionts is strictly congruent with the phylogeny of their host (Munson *et al.*, 1991; Chen *et al.*, 1999; Baumann, 2005; Dale & Moran, 2006). The co-speciation of the symbiotic partners can result in specializations of the host and the bacterium (Moran & Baumann, 2000; Baumann, 2005; Dale & Moran, 2006). For instance, endosymbionts live inside specialized host cells (bacteriocytes), which can form special structures in the host, the bacteriomes and are maternally transmitted inside the eggs (Buchner, 1965; Baumann, 2005; Moran *et al.*, 2008). This tight coevolutionary living together has resulted in genomes of obligate endosymbionts that can be very small, and that are characterized by the loss of functional genes that are necessary for, e.g., recombination, replication or transcriptional regulation, while genes for the synthesis of nutrients that the host diet lacks, such as essential amino acids, are retained (Moran & Wernegreen, 2000; Baumann, 2005; Dale & Moran, 2006; Moran *et al.*, 2008).

#### *Facultative endosymbionts*

Facultative (so-called secondary) endosymbionts are not strictly necessary for survival or reproduction of their host. Secondary endosymbionts are mostly maternally transmitted inside the eggs, although horizontal transmission can also occur

(Huigens *et al.*, 2004; Caspi-Fluger *et al.*, 2012). Consequently, there is only a partial congruence between symbiont and host phylogeny (Dale & Moran, 2006; Werren *et al.*, 2008). The prevalence of facultative, maternally transmitted bacteria in a host population depends on the efficiency of vertical transmission of the bacteria, but also on host fitness (Werren, 1997; Oliver *et al.*, 2010). Many facultative endosymbionts were shown to benefit the host e.g., by protecting them against parasitoids (Oliver *et al.*, 2003), or by supporting host plant use (Tsuchida *et al.*, 2011; Su *et al.*, 2015).

Endosymbiotic reproductive parasites, such as *Wolbachia*, ‘*Candidatus Cardinium*’, *Spiroplasma* and *Rickettsia* are known to increase their prevalence in a host population by manipulating the reproduction of their host. Since these bacteria are maternally transmitted, reproductive manipulation increases the proportion of infected females in a population via mechanisms such as feminization, parthenogenesis, male-killing or cytoplasmic incompatibility (Werren, 1997; Stouthamer *et al.*, 1999; Weeks *et al.*, 2001; Werren *et al.* 2008; Duron *et al.*, 2008).

Besides reproductive manipulation, reproductive parasites may increase and strengthen their persistence in a host population by benefitting the host (Hoffmann *et al.*, 1998; Fry *et al.*, 2004; Werren *et al.*, 2008; Himler *et al.*, 2011; Zug & Hammerstein, 2015). Accordingly, reproductive parasites have recently been found to enhance host fitness and to increase host resistance to bacterial and viral pathogens (Hedges *et al.*, 2008; Teixeira *et al.*, 2008; Moreira *et al.*, 2009) or parasitoids (Xie *et al.*, 2010, 2014), act as nutritional mutualist (Hosokawa *et al.*, 2010; Nikoh *et al.*, 2014), or alter host physiology to their host’s benefit (Kaiser *et al.*, 2010).

### *Gut bacteria*

Gut bacteria do not live intracellularly, but inhabit different regions or compartments of the gut. Associations between gut bacteria and a host range from permanent (core community) to transient (Shade & Handelsman, 2012). Gut bacteria are in some cases necessary for host survival, development or reproduction (Hosokawa *et al.*, 2007; Salem *et al.*, 2013). When gut bacteria are permanently associated with a host, they are often vertically transmitted, may reside in special compartments or pouches in the guts and provide diverse beneficial services to their hosts. If gut bacteria are vertically transmitted, transmission occurs mostly extracellularly via various mechanisms. For instance, gut bacteria may be smeared on the egg surface by a female after oviposition or, as is the case of the stinkbug *Megacopta punctatissima*, provided in capsules together with the egg mass, which ensures high transmission rates (Fukatsu & Hosokawa, 2002; Engel & Moran, 2013; Salem *et al.*, 2015). Other insects, such as the stinkbug *Riptortus pedestris*, take up bacterial symbionts each generation again from the environment (Kikuchi *et al.*, 2011). Another route of transmission for gut bacteria is coprophagy, in which bacteria are excreted together with feces and can be taken up by conspecifics (Beard *et al.*, 2002). In social insects, gut bacteria may be taken up during contact with nest mates or hive contents and may

be transferred together with food from adults to larvae (Koch & Schmid-Hempel, 2011; Martinson *et al.*, 2012; Marsh *et al.*, 2014; Powell *et al.*, 2014).

Gut bacteria may also only be transiently present in the gut if random uptake from the environment is not followed by stable colonization (Shade & Handelsman, 2012; Engel & Moran, 2013). High variability of bacterial communities between individuals of a host population or variability due to diet or developmental stage may indicate transient association of gut bacteria with a host (Shade & Handelsman, 2012). Nevertheless, the metabolic capabilities that bacteria have evolved independently of the host may suffice to enhance the fitness of an animal that takes these bacteria up into its gut, even if transiently (Mason *et al.*, 2014). For instance, some plant-associated bacteria are able to use methanol as carbon source, fix nitrogen or break down secondary plant metabolites (Sy *et al.*, 2005; Fürnkranz *et al.*, 2008; Vorholt, 2012). The latter was shown to benefit larvae of the moth *Lymantria dispar* as described below (Mason *et al.*, 2014).

### ***1.1 Host plant utilization of herbivores in relation to bacterial symbionts***

Many herbivores have colonized and feed on plants and plant parts that are poor or unbalanced in nutrients or are hard to digest (Schoonhoven *et al.*, 2005). Moreover, herbivores have to deal with physical and chemical defences that plants have evolved to reduce the damage inflicted by herbivores (Schoonhoven *et al.*, 2005; Karban & Baldwin, 2007). Symbiotic bacteria can play an important role in the ability of herbivores to use plant species or plant tissues as food and thus determine their host's ecological niche (Douglas, 2009; Feldhaar, 2011; Hansen & Moran, 2014).

#### *Direct facilitation of host plant use*

Symbiotic bacteria may directly facilitate host plant use by providing amino acids, vitamins or co-factors that are absent from the host diet or by producing enzymes that enhance the digestion of plant tissues (Buchner, 1965; Douglas, 2009; Feldhaar, 2011; Engel & Moran, 2013; Hansen & Moran, 2014). Obligate endosymbionts are well known for synthesizing nutrients for their hosts. For instance, plant sap is poor in amino acids. Hence, phloem feeders such as aphids are provided with essential amino acids by their primary endosymbionts, *Buchnera aphidicola* (Shigenobu *et al.*, 2000; van Ham *et al.*, 2003; Baumann, 2005). Nutrient provisioning is not restricted to obligate endosymbionts, although gut bacteria can fulfil this function as well. For instance, the gut symbiont of the firebug *Pyrhocoris apterus* was shown to provide its host with vitamin B, which proved essential for its ability to reproduce (Salem *et al.*, 2013, 2014). The plataspid stinkbug *Megacopta punctatissima* relies on its gut symbiont *Ishikawaella capsulata* for normal growth, development and reproduction (Fukatsu & Hosokawa, 2002; Hosokawa *et al.*, 2007). In ants, there is strong indication that gut bacteria from the order Rhizobiales may be involved in nitrogen fixation (Russell *et al.*, 2009). Even bacteria that only transiently inhabit the

gut may affect host plant use of herbivores. For instance, larval growth of the gypsy moth, *Lymantria dispar*, was enhanced by bacteria that inhabit the foliage of its host plant aspen, when phenolic glycosides, the major defence metabolites of aspen, were added to the larval (artificial) diet (Mason *et al.*, 2014). The authors further showed that these bacteria can reduce the concentration of phenolic glycosides *in vitro* (Mason *et al.*, 2014).

#### *Indirect facilitation of host plant use*

Besides the direct nutritional effects mentioned above, herbivore-associated bacteria may manipulate plant physiology to their host's benefit (Kaiser *et al.*, 2010; Chung *et al.*, 2013; Hansen & Moran, 2014; Zhu *et al.*, 2014; Su *et al.*, 2015). *Wolbachia* was shown to enhance the fitness of the lepidopteran leaf-miner *Phyllonorycter blancardella*, by preserving photosynthetically active and nutrient rich (green) islands in senescent leaves, thereby increasing host fitness (Kaiser *et al.*, 2010).

Few examples show that symbionts can also interfere with herbivore-induced plant defences. Phytohormones such as jasmonic acid (JA) or salicylic acid (SA) are central to the coordination of plant defences. The induction of phytohormone biosynthesis is specific to the type of attacker: in general, feeding by herbivorous insects induces the JA-related plant responses (Karban & Baldwin, 2007; Erb *et al.*, 2012). JA-responses were shown to cause reduced larval growth, amount of feeding and survival of herbivores and may attract their natural enemies (Howe *et al.*, 1996; Li *et al.*, 2002; Thaler *et al.*, 2002). Biotrophic phytopathogens generally induce SA-related responses in plants (Glazebrook, 2005). Importantly, JA- and SA-responses are interrelated and JA-responses can be suppressed via the induction of SA-responses and vice versa (JA-SA crosstalk) (Pieterse *et al.*, 2012; Thaler *et al.*, 2012). *Hamiltonella defensa*, the secondary symbiont of the whitefly *Bemisia tabaci*, was found to suppress JA-responses in tomato, via small non-proteinaceous molecules in the whitefly's saliva (Su *et al.*, 2015). Also, bacteria associated with the Colorado potato beetle, *Leptinotarsa decemlineata*, were shown to reduce herbivore-induced JA-responses and to increase the larval growth of the beetle (Chung *et al.*, 2013). In both examples, JA suppression did not occur in SA-deficient plants, indicating that induction of SA-related responses caused the suppression of JA-responses (Chung *et al.*, 2013)

#### *The role of bacterial symbionts in agriculture*

The ability of bacterial symbionts to determine the host plant range of herbivores can have implications for agriculture, in particular pest management. Many herbivores are destructive agricultural and forest pests. Microbes that are harboured by arthropod pests may affect their pest status by influencing the dietary range of the pest or by affecting pest management. For instance, the pest status of the stinkbug *Megacopta punctatissima* on legume crop was found to be determined by the genotype of its obligate gut symbiont *Ishikawaella capsulata* (Hosokawa *et al.*,

2007). In this study, egg hatch rate of the pest species *M. punctatissima* decreased significantly when its symbionts were replaced for symbionts of a closely related non-pest species (*M. cribraria*). On the other hand, egg hatch rate of *M. cribraria* was significantly increased after receiving symbionts of *M. punctatissima* (Hosokawa *et al.*, 2007). It was demonstrated in aphids that symbiotic bacteria are able to expand host plant range of its insect herbivore host: transfection of the facultative endosymbiont *Regiella insecticola* from the pea aphid (*Acyrtosiphon pisum*) to the vetch aphid (*Megoura crassicaula*) enabled the latter to survive and reproduce on clover (Tsuchida *et al.*, 2011).

Bacteria can also interfere with chemical pest control measures. For example, bacteria of the genus *Burkholderia* have been found to confer insecticide resistance to the bean bug, *Riptortus pedestris*, which is a pest on leguminous crops (Kikuchi *et al.*, 2012). Further, larval mortality in response to the insecticidal toxins of *Bacillus thuringiensis* was shown to be reduced upon removal of the bacterial midgut community in the gypsy moth, *Lymantria dispar*, and other Lepidoptera (Broderick *et al.*, 2006, 2009). The authors suggest that benign gut bacteria of these Lepidoptera may breach the midgut barrier after it was damaged by *B. thuringiensis* toxin and become opportunistic pathogens (Broderick *et al.*, 2006, 2009). The presence of bacterial symbionts may also affect biological control measures, e.g., *H. defensa* rendered its aphid host more resistant to parasitoid wasps (Oliver *et al.*, 2003).

### **2. Pathogenic effects of bacteria on insect hosts**

Besides commensal and beneficial bacteria, insects may be exposed to a suite of pathogenic bacteria during their lifetime. Pathogens cause disease in organisms and pathogenicity of bacteria depends on virulence factors, such as the ability to attach to and invade host cells, the production of toxins and the secretion of proteins that modulate host cellular functions (Farthing, 2004; Galán, 2009). However, whether bacteria cause disease also strongly depends on the host or the host tissue (Ruby *et al.*, 2004).

#### *The immune system of insects*

As a first line of defence against pathogens, insects possess physical and chemical barriers, such as the cuticle or the peritrophic matrix (Dunn, 1990; Lehane, 1997; Kuraishi *et al.*, 2011; Davis & Engström, 2012). Nevertheless, pathogens might gain access to the insect hemocoel via wounds or may be ingested and be able to cross the midgut epithelium (Ashida *et al.*, 2012). As a consequence, insects have evolved an efficient innate immune system to fight pathogenic intruders. Recognition of self and non-self is the first step of immune defence (Janeway, 1992; Royet, 2004). Insects have a wide range of proteins that function as pattern recognition receptors (PRRs) that recognize pathogen-associated molecular patterns (PAMPs) (Medzhitov & Janeway, 1997). PAMPs include surface molecules of bacteria such as lipopolysaccharides, peptidoglycans or lipoteichoic acid.

The insect immune system is a combination of cellular and humoral immune responses. Cellular immune responses include processes like phagocytosis and encapsulation that are mediated by different types of blood cells (hemocytes) (Lavine & Strand, 2002; Strand, 2008). During phagocytosis, biotic or abiotic entities are enclosed by individual hemocytes. Encapsulation response targets large invaders, such as nematodes or parasitoid larvae, that cannot be enclosed by a single cell and are therefore bound by multiple hemocytes (Strand, 2008). Phagocytized or encapsulated organisms are killed and degraded (Reeves *et al.*, 2002; Stuart & Ezekowitz, 2005; Stuart *et al.*, 2007).

Humoral defences include a broad range of antimicrobial peptides (AMPs) that are produced by the fat body or hemocytes and that are released into the hemolymph (Bulet *et al.*, 1999; Zasloff, 2002). AMPs vary in their antimicrobial spectrum. Most AMPs show activity against a wide range of pathogens, with particular efficiency against certain groups, depending on their mode of action. For example, lysozymes, which destroy bonds between peptidoglycan forming molecules (Rupley, 1967) can efficiently destroy gram-positive bacteria but are also active against gram-negative bacteria. Other AMPs have a more narrow antimicrobial spectrum. For instance, drosomycin only shows activity against filamentous fungi (Bulet *et al.*, 1999; Zhang & Zhu, 2009).

The melanin-synthesis cascade is another important part of the insect immune system. Phenoloxidase (PO) is the key enzyme in this cascade and catalyzes the reactions of phenols to toxic quinones which are polymerized to melanin (Nappi & Vass, 1993; Cerenius & Söderhäll, 2004). Intermediates of melanogenesis as well as melanin itself are cytotoxic and have been reported to be lethal for pathogens (Nappi & Vass, 1993, 1998; Nappi *et al.*, 1995; Söderhäll & Cerenius, 1998; Nappi & Christensen, 2005). Further, melanin and its intermediates have functions in the encapsulation and phagocytosis of pathogens as well as in wound healing processes (Cerenius *et al.*, 2008).

#### *The cost of immune defence and trade-offs with other life history traits*

In life history theory, the concept of trade-offs is based on the assumption that organisms have limited internal resources available that need to be partitioned among the different life history traits (Stearns, 1989; Roff & Fairbairn, 2007). Life history traits are therefore negatively linked to each other: a resource investment into one trait which increases fitness may reduce resources that are available for another fitness-relevant trait (Stearns, 1989; Roff & Fairbairn, 2007).

The immune system may prevent an organism from being colonized by pathogens. However, an immune system implicates different types of costs which may negatively affect other life history traits (Sheldon & Verhulst, 1996; Zuk & Stoehr, 2002; Schmid-Hempel, 2005). Physiological costs include costs of nutrients and energy which are necessary for the maintenance and deployment of the immune system (Schmid-Hempel, 2011). Maintenance costs are incurred to keep the immune

system at a level of readiness, even when the organism is not infected (Schmid-Hempel, 2005, 2011; Kraaijeveld & Wertheim, 2009). Deployment costs are incurred when the immune system is activated (Kraaijeveld & Wertheim, 2009; Schmid-Hempel, 2011). Immune system activation has been shown to trade off with various other life history traits such as life span (Moret & Schmid-Hempel, 2000; Krams *et al.*, 2014), growth (Soler *et al.*, 2003; Brommer, 2004) and reproduction (Gustafsson *et al.*, 1994; Sheldon & Verhulst, 1996; Adamo *et al.*, 2001; Rolff & Siva-Jothy, 2002; Hanssen *et al.*, 2005; French *et al.*, 2007).

Evolutionary costs of immunity arise through the negative genetic covariance between immunity and other fitness-related traits in uninfected organisms (Schmid-Hempel, 2005, 2011). For instance, several studies showed that highly pathogen-resistant insect strains perform worse than less resistant strains, e.g., have longer developmental time, lower larval competitive ability or lower fecundity in the absence of pathogens (Boots & Begon, 1993; Kraaijeveld & Godfray, 1997; McKean *et al.*, 2008).

### *Behavioural changes in the face of infection*

As an alternative (or in addition) to a costly immune system (activation), organisms may have behavioural strategies that reduce the frequency of infection and minimize adverse fitness effects (Schmid-Hempel, 2011; de Roode & Lefèvre, 2012). An important behavioural adaptation to possible infections is to avoid contracting them. This behaviour can include spatial or temporal avoidance of parasites (Orr, 1992; Christe *et al.*, 1994), avoiding feeding on contaminated food (Hutchings *et al.*, 2001; Alma *et al.*, 2010) or avoiding infected conspecifics, such as infected mates (Kavaliers & Colwell, 1995; Penn & Potts, 1998; Penn *et al.*, 1998).

Behavioural adaptations to reduce pathogen loads once infection has been contracted include self-medication behaviour, in which organisms consume medicinal substances to reduce pathogen loads and increase their fitness while infected (Huffman *et al.*, 1996; Singer *et al.*, 2009; Manson *et al.*, 2010). In this context, infected females of the butterfly *Danaus plexippus* were found to engage in trans-generational medication behaviour and oviposit more on plants with high levels of toxic secondary metabolites than on plants with lower levels than uninfected females (Lefèvre *et al.*, 2010).

Another adaptive behaviour of infected animals, which does not directly aim to reduce pathogen loads but may enhance life time fecundity of infected organisms, is terminal investment. If life expectancy decreases due to age or infection, organisms may make a 'terminal investment', i.e., invest their remaining resources into current reproduction in order to maximize their fitness in the short time that remains (Williams, 1966; Clutton-Brock, 1984). Shortening of the lifespan can be caused by the pathogen that may kill the organism or by the costs of immune defence which can trade off with lifespan (Sheldon & Verhulst, 1996). A shift of investment from future to current reproduction may imply an increase in courtship activity, parental

care, oviposition rate or acceptance of less preferred host plants for oviposition (Minchella & Loverde, 1981; Part *et al.*, 1992; Adamo, 1999; Bonneaud *et al.*, 2004; Javošič & Tammaru, 2004; Creighton *et al.*, 2009).

### 3. Study organisms

In this thesis, I investigated possible effects of bacteria on the life histories of two generalist herbivorous arthropods. In the noctuid moth *Heliothis virescens* (Fabricius, 1777) (Lepidoptera: Noctuidae), I determined negative correlations between immunity and reproduction as well as adaptive behavioural changes in the oviposition strategy of this moth in the face of infection. Moreover, I identified bacterial communities that are associated with *H. virescens*.

In the two-spotted spider mite *Tetranychus urticae* Koch, 1836 (Acari: Tetranychidae), which is well known to harbour various (endo)symbiotic bacteria such as *Wolbachia*, ‘*Candidatus* Cardinium’ and *Spiroplasma* (Breeuwer & Jacobs, 1996; Gotoh *et al.*, 2003, 2007a; Enigl & Schausberger, 2007), I investigated effects of these bacteria on mite fitness parameters, mite transcriptome and induced plant responses of tomato leaflets after mite feeding.

#### *Heliothis virescens*

The tobacco budworm, *H. virescens* (FIGURE 1.1), is distributed throughout the American continent (Fitt, 1989). Its larvae are polyphagous and have been reported to feed on at least 37 plant species in 14 families (Barber, 1937; Stadelbacher, 1981; Fitt, 1989; Waldvogel & Gould, 1990; Sheck & Gould, 1993; Blanco *et al.*, 2007). Among the host plants of *H. virescens* are important economic crops like tobacco, cotton and chickpea (Morgan & Chamberlin, 1927; Barber, 1937; Neunzig, 1969; Blanco *et al.*, 2007), which makes this species a major agricultural pest.

*Heliothis virescens* females can oviposit up to 1500 eggs in their lifetime of about 30 days in the laboratory at 25 °C (Proshold *et al.*, 1982; Willers *et al.*, 1987; Fitt, 1989). Eggs are laid singly on plants throughout the night, starting at dusk (Fitt, 1989;




FIGURE 1.1 *Heliothis virescens* (a) larva, (b) pupa, and (c) mating adults. © Jan van Arkel.

## General introduction

Ramaswamy, 1990). *Heliothis virescens* has five larval stages and developmental time from eggs to pupae takes about 17 days at 25 °C (Fye & McAda, 1972). Larvae pupate in the soil for 13 days at 25 °C after which the adults eclose.

*Heliothis virescens* was reported to have an efficient immune system (Barthel *et al.*, 2014). However, possible effects of immune response on life history traits have not been investigated in this species. Furthermore, not much is known about its microbiome, i.e., bacteria that are associated with *H. virescens*, particularly in the field.

### *Tetranychus urticae*

The two-spotted spider mite, *T. urticae* (FIGURE 1.2) is an extremely polyphagous herbivore that occurs worldwide and has been reported to feed on over 1100 plant species, among them are economically important crops like tomato, cucumber, strawberry, bean and cotton (Bolland *et al.*, 1998; Grbic *et al.*, 2011; Migeon *et al.*, 2011) which makes this mite an important agricultural pest.

The life cycle of *T. urticae* includes egg, one larval stage, which is followed by two nymphal stages (protonymph and deutonymph) and the adult stage. The generation time of *T. urticae* takes about two weeks at 27 °C (Chain-ing *et al.*, 1976). Females have been reported to lay over 100 eggs during their lifetime (Carey & Bradley, 1982). Thus this mite can reach large population sizes in a short amount of time.

Spider mites are stylet feeders that pierce plant cells and suck out the cell content. Like other stylet feeders, *T. urticae* has been shown to induce jasmonic and salicylic acid-related plant defences during feeding (Walling, 2000; Kant *et al.*, 2004, 2008; Alba *et al.*, 2015).

*Tetranychus urticae* was found to harbour different endosymbionts, including *Wolbachia*, ‘*Candidatus* Cardinium’ and *Spiroplasma* (Breeuwer & Jacobs, 1996; Gotoh *et al.*, 2003, 2007a; Enigl & Schausberger, 2007). *Wolbachia* and ‘*Candida-*



**FIGURE 1.2** *Tetranychus urticae* adult female. © Jan van Arkel.

*tus Cardinium*' were shown to cause cytoplasmic incompatibility in some populations of *T. urticae* (Breeuwer, 1997; Gotoh *et al.*, 2007a,b). In other *T. urticae* populations, *Wolbachia* did not cause cytoplasmic incompatibility (Gotoh *et al.*, 2003). Further, the effects of *Wolbachia* on mite fitness have been assessed with all possible outcomes, and *Wolbachia* had a negative effect, no effect or a positive effect on female fecundity (Breeuwer, 1997; Vala *et al.*, 2000; Perrot-Minnot *et al.*, 2002; Xie *et al.*, 2011). The presence of *Wolbachia* on mite gene expression has been investigated (Zhang *et al.*, 2015). Less is known about effects of *Spiroplasma* and '*Candidatus Cardinium*' or any combination of endosymbionts on *T. urticae*. Furthermore, the effect of endosymbionts on host-plant use of this mite has not yet been investigated.

## THESIS OUTLINE

**CHAPTER 2.** As outlined above, bacteria can have diverse effects on their hosts. On the one hand, herbivore-associated bacteria may benefit the host, for instance by facilitating plant utilization of the host (Engel & Moran, 2013; Hansen & Moran, 2014). On the other hand, bacterial pathogens may negatively affect life history traits and thus the fitness of an organism (Grenfell & Dobson, 1995; Poulin, 2007; Schmid-Hempel, 2011). Against this background, we investigated the bacterial community that is associated with field and laboratory populations of *H. virescens* and assessed the spatial and temporal variability of this bacterial community as it depends on diet and life stage.

**CHAPTER 3.** The concept of 'Bateman's principle and immunity' predicts that females, who maximize their fitness by increasing longevity, invest more in immunity than males, who maximize fitness by increasing their number of matings (Rolff, 2002). Most moths, including *H. virescens*, do not correspond to the 'classical' sex roles in which females invest more into offspring than males. In this species, females and males invest substantially into offspring: females produce up to 1500 eggs whereas males produce a spermatophore that can comprise up to 5% of their bodymass (Fye & McAda, 1972; Proshold *et al.*, 1982; Blanco *et al.*, 2009). Both sexes produce a sex pheromone, but until now only males have been shown to differentiate between female sex pheromones of different quality, indicating that males are the choosy sex in this species (Vetter & Baker, 1983; Groot *et al.*, 2009, 2014). Because of the deviations in the life history of *H. virescens* from the classical Bateman case, it can be expected that differences in immunity between the sexes do not follow the predictions for 'Bateman's principle and immunity'. To assess the concept of 'Bateman's principle and immunity' in *H. virescens*, we first investigated immune system activation of males and females via immune gene expression experiments using dead cells of the entomopathogen *Serratia entomophila*. Secondly, we assessed the influence of immune activation on mating success of both sexes in mate choice experiments. Thirdly, we tested if immune defence activation

would affect the composition of the female sex pheromone as well as female calling behaviour.

**CHAPTER 4.** Infection may shorten the life span of an organism. One strategy for organisms to maximize fitness in the face of infection is therefore terminal investment, in which investment into future reproduction is shifted towards current reproduction (Williams, 1966; Clutton-Brock, 1984). In this chapter, we investigated whether terminal investment is part of the life history strategy of *H. virescens* and whether females of this moth adaptively change their oviposition strategy in response to an immune challenge with dead cells of the entomopathogen *S. entomophila*. Specifically, we assessed whether *H. virescens* females would increase their egg output in response to an immune challenge, and whether oviposition site choice of *H. virescens* females would be influenced by an immune challenge.

**CHAPTER 5.** Endosymbionts, such as *Wolbachia*, ‘*Candidatus Cardinium*’ and *Spiroplasma*, are well-known reproductive manipulators which may increase the proportion of infected females to increase their prevalence in a host population (Werren, 1997; Duron *et al.*, 2008; Werren *et al.*, 2008; Engelstädter & Hurst, 2009). However, several studies indicate that these bacteria may also positively affect host fitness (Teixeira *et al.*, 2008; Kaiser *et al.*, 2010; Xie *et al.*, 2010; Zug & Hammerstein, 2015). We studied the role of bacterial (endo)symbionts in two strains of the two-spotted spider mite *T. urticae*, one strain is a plant defence inducer and the other strain a plant defence suppressor. We investigated the bacterial community associated with these two strains and tested if the presence of (endo)symbionts was correlated with performance and gene expression profiles of the mites. Further, we assessed if the symbionts were associated with induction or suppression of plant responses in tomato (*Solanum lycopersicum*).

## REFERENCES

- Adamo, S.A. 1999. Evidence for adaptive changes in egg laying in crickets exposed to bacteria and parasites. *Anim Behav* **57**: 117-124.
- Adamo, S.A., Jensen, M. & Younger, M. 2001. Changes in lifetime immunocompetence in male and female *Gryllus texensis* (formerly *G. integer*): trade-offs between immunity and reproduction. *Anim Behav* **62**: 417-425.
- Alba, J.M., Schimmel, B.C.J., Glas, J.J., Ataide, L.M.S., Pappas, M.L., Villarroel, C.A., Schuurink, R.C., Sabelis, M.W. & Kant, M.R. 2015. Spider mites suppress tomato defenses downstream of jasmonate and salicylate independently of hormonal crosstalk. *New Phytol* **205**: 828-840.
- Alma, C.R., Gillespie, D.R., Roitberg, B.D. & Goettel, M.S. 2010. Threat of infection and threat-avoidance behavior in the predator *Dicyphus hesperus* feeding on whitefly nymphs infected with an entomopathogen. *J Insect Behav* **23**: 90-99.
- Ashida, H., Ogawa, M., Kim, M., Mimuro, H. & Sasakawa, C. 2012. Bacteria and host interactions in the gut epithelial barrier. *Nat Chem Biol* **8**: 36-45.
- Barber, G.W. 1937. Seasonal availability of food plants of two species of *Heliothis* in eastern Georgia. *J Econ Entomol* **30**: 150-158.

- Barthel, A., Kopka, I., Vogel, H., Zipfel, P., Heckel, D.G. & Groot, A.T. 2014. Immune defence strategies of generalist and specialist insect herbivores. *Proc R Soc B Biol Sci* **281**: 20140897.
- Baumann, P. 2005. Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Annu Rev Microbiol* **59**: 155-189.
- Beard, C.B., Cordon-Rosales, C. & Durvasula, R.V. 2002. Bacterial symbionts of the triatominae and their potential use in control of Chagas disease transmission. *Annu Rev Entomol* **47**: 123-141.
- Blanco, C.A., Terán-Vargas, A.P., López Jr, J.D., Kauffman, J.V. & Wei, X. 2007. Densities of *Heliothis virescens* and *Helicoverpa zea* (Lepidoptera: Noctuidae) in three plant hosts. *Fla Entomol* **90**: 742-750.
- Blanco, C.A., Guadalupe Rojas, M., Groot, A.T., Morales-Ramos, J. & Abel, C.A. 2009. Size and chemical composition of *Heliothis virescens* (Lepidoptera: Noctuidae) spermatophores. *Ann Entomol Soc Am* **102**: 629-637.
- Bolland, H.R., Gutierrez, J. & Flechtman, C.H. 1998. *World catalogue of the spider mite family (Acari: Tetranychidae)*. Brill, Leiden, The Netherlands.
- Bonneaud, C., Mazuc, J., Chastel, O., Wester Dahl, H. & Sorci, G. 2004. Terminal investment induced by immune challenge and fitness traits associated with major histocompatibility complex in the house sparrow. *Evolution* **58**: 2823-2830.
- Boots, M. & Begon, M. 1993. Trade-offs with resistance to a granulosis virus in the Indian meal moth, examined by a laboratory evolution experiment. *Funct Ecol*: 528-534.
- Breeuwer, J.A.J. & Jacobs, G. 1996. *Wolbachia*: intracellular manipulators of mite reproduction. *Exp Appl Acarol* **20**: 421-434.
- Breeuwer, J.A. 1997. *Wolbachia* and cytoplasmic incompatibility in the spider mites *Tetranychus urticae* and *T. turkestanii*. *Heredity* **79**: 41-47.
- Broderick, N.A., Raffa, K.F. & Handelsman, J. 2006. Midgut bacteria required for *Bacillus thuringiensis* insecticidal activity. *Proc Natl Acad Sci USA* **103**: 15196-15199.
- Broderick, N.A., Robinson, C.J., McMahon, M.D., Holt, J., Handelsman, J. & Raffa, K.F. 2009. Contributions of gut bacteria to *Bacillus thuringiensis*-induced mortality vary across a range of Lepidoptera. *BMC Biol* **7**: 11.
- Brommer, J.E. 2004. Immunocompetence and its costs during development: an experimental study in blue tit nestlings. *Proc R Soc B Biol Sci* **271**: S110-S113.
- Buchner, P. 1965. *Endosymbiosis of animals with plant microorganisms*. Interscience, New York, USA.
- Bulet, P., Hetru, C., Dimarcq, J.-L. & Hoffmann, D. 1999. Antimicrobial peptides in insects; structure and function. *Dev Comp Immunol* **23**: 329-344.
- Carey, J. & Bradley, J. 1982. Developmental rates, vital schedules, sex ratios and life tables for *Tetranychus urticae*, *T. turkestanii* and *T. pacificus* (Acarina: Tetranychidae) on cotton. *Acarologia* **23**: 333-345.
- Caspi-Fluger, A., Inbar, M., Mozes-Daube, N., Katzir, N., Portnoy, V., Belausov, E., Hunter, M.S. & Zchori-Fein, E. 2012. Horizontal transmission of the insect symbiont *Rickettsia* is plant-mediated. *Proc R Soc B Biol Sci* **279**: 1791-1796.
- Cerenius, L. & Söderhäll, K. 2004. The prophenoloxidase-activating system in invertebrates. *Immunol Rev* **198**: 116-126.
- Cerenius, L., Lee, B.L. & Söderhäll, K. 2008. The proPO-system: pros and cons for its role in invertebrate immunity. *Trends Immunol* **29**: 263-271.
- Chain-ing, T.S., Poe, S.L. & Cromroy, H.L. 1976. Biology, life table, and intrinsic rate of increase of *Tetranychus urticae*. *Ann Entomol Soc Am* **69**: 362-364.
- Chen, X., Li, S. & Aksoy, S. 1999. Concordant evolution of a symbiont with its host insect species: molecular phylogeny of genus *Glossina* and its bacteriome-associated endosymbiont, *Wigglesworthia glossinidia*. *J Mol Evol* **48**: 49-58.
- Christe, P., Oppliger, A. & Richner, H. 1994. Ectoparasite affects choice and use of roost sites in the great tit, *Parus major*. *Anim Behav* **47**: 895-898.

## General introduction

- Chung, S.H., Rosa, C., Scully, E.D., Peiffer, M., Tooker, J.F., Hoover, K., Luthe, D.S. & Felton, G.W. 2013. Herbivore exploits orally secreted bacteria to suppress plant defenses. *Proc Natl Acad Sci USA* **110**: 15728-15733.
- Clutton-Brock, T.H. 1984. Reproductive effort and terminal investment in iteroparous animals. *Am Nat*: 212-229.
- Creighton, J.C., Heflin, N.D. & Belk, M.C. 2009. Cost of reproduction, resource quality, and terminal investment in a burying beetle. *Am Nat* **174**: 673-684.
- Dale, C. & Moran, N.A. 2006. Molecular interactions between bacterial symbionts and their hosts. *Cell* **126**: 453-465.
- Davis, M.M. & Engström, Y. 2012. Immune response in the barrier epithelia: lessons from the fruit fly *Drosophila melanogaster*. *J Innate Immun* **4**: 273-283.
- Dillon, R. & Dillon, V. 2004. The gut bacteria of insects: nonpathogenic interactions. *Annu Rev Entomol* **49**: 71-92.
- Douglas, A. 2009. The microbial dimension in insect nutritional ecology. *Funct Ecol* **23**: 38-47.
- Dunn, P.E. 1990. Humoral immunity in insects. *BioScience*: 738-744.
- Duron, O., Bouchon, D., Boutin, S., Bellamy, L., Zhou, L., Engelstädter, J. & Hurst, G.D. 2008. The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol* **6**: 27.
- Engel, P. & Moran, N.A. 2013. The gut microbiota of insects—diversity in structure and function. *FEMS Microbiol Rev* **37**: 699-735.
- Engelstädter, J. & Hurst, G.D. 2009. The ecology and evolution of microbes that manipulate host reproduction. *Annu Rev Ecol Evol Syst* **40**: 127-149.
- Enigl, M. & Schausberger, P. 2007. Incidence of the endosymbionts *Wolbachia*, *Cardinium* and *Spiroplasma* in phytoseiid mites and associated prey. *Exp Appl Acarol* **42**: 75-85.
- Erb, M., Meldau, S. & Howe, G.A. 2012. Role of phytohormones in insect-specific plant reactions. *Trends Plant Sci* **17**: 250-259.
- Farthing, M.J. 2004. Bugs and the gut: an unstable marriage. *Best Pract Res Clin Gastroenterol* **18**: 233-239.
- Feldhaar, H. 2011. Bacterial symbionts as mediators of ecologically important traits of insect hosts. *Ecol Entomol* **36**: 533-543.
- Fitt, G.P. 1989. The ecology of *Heliothis* species in relation to agroecosystems. *Annu Rev Entomol* **34**: 17-53.
- French, S.S., DeNardo, D.F. & Moore, M.C. 2007. Trade-offs between the reproductive and immune systems: facultative responses to resources or obligate responses to reproduction? *Am Nat* **170**: 79-89.
- Fry, A., Palmer, M. & Rand, D. 2004. Variable fitness effects of *Wolbachia* infection in *Drosophila melanogaster*. *Heredity* **93**: 379-389.
- Fukatsu, T. & Hosokawa, T. 2002. Capsule-transmitted gut symbiotic bacterium of the Japanese common plataspid stinkbug, *Megacopta punctatissima*. *Appl Environ Microbiol* **68**: 389-396.
- Fürnkranz, M., Wanek, W., Richter, A., Abell, G., Rasche, F. & Sessitsch, A. 2008. Nitrogen fixation by phyllosphere bacteria associated with higher plants and their colonizing epiphytes of a tropical lowland rainforest of Costa Rica. *ISME J* **2**: 561-570.
- Fye, R.E. & McAda, W. 1972. *Laboratory studies on the development, longevity, and fecundity of six lepidopterous pests of cotton in Arizona*. Agricultural Research Service, US Dept. of Agriculture.
- Galán, J.E. 2009. Common themes in the design and function of bacterial effectors. *Cell Host Microbe* **5**: 571-579.
- Glazebrook, J. 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol* **43**: 205-227.

- Gotoh, T., Noda, H. & Hong, X. 2003. *Wolbachia* distribution and cytoplasmic incompatibility based on a survey of 42 spider mite species (Acari: Tetranychidae) in Japan. *Heredity* **91**: 208-216.
- Gotoh, T., Noda, H. & Ito, S. 2007a. *Cardinium* symbionts cause cytoplasmic incompatibility in spider mites. *Heredity* **98**: 13-20.
- Gotoh, T., Sugasawa, J., Noda, H. & Kitashima, Y. 2007b. *Wolbachia*-induced cytoplasmic incompatibility in Japanese populations of *Tetranychus urticae* (Acari: Tetranychidae). *Experimental and Applied Acarology* **42**: 1-16.
- Grbic, M., Van Leeuwen, T., Clark, R.M., Rombauts, S., Rouze, P., Grbic, V., Osborne, E.J., Dermauw, W., Thi Ngoc, P.C., Ortego, F., Hernandez-Crespo, P., Diaz, I., Martinez, M., Navajas, M., Sucena, E., Magalhaes, S., Nagy, L., Pace, R.M., Djuranovic, S., Smagghe, G., Iga, M., Christiaens, O., Veenstra, J.A., Ewer, J., Villalobos, R.M., Hutter, J.L., Hudson, S.D., Velez, M., Yi, S.V., Zeng, J., Pires-daSilva, A., Roch, F., Cazaux, M., Navarro, M., Zhurov, V., Acevedo, G., Bjelica, A., Fawcett, J.A., Bonnet, E., Martens, C., Baele, G., Wissler, L., Sanchez-Rodriguez, A., Tirry, L., Blais, C., Demeestere, K., Henz, S.R., Gregory, T.R., Mathieu, J., Verdon, L., Farinelli, L., Schmutz, J., Lindquist, E., Feyerisen, R. & Van de Peer, Y. 2011. The genome of *Tetranychus urticae* reveals herbivorous pest adaptations. *Nature* **479**: 487-492.
- Grenfell, B.T. & Dobson, A.P. 1995. *Ecology of infectious diseases in natural populations*. Cambridge University Press, Cambridge, UK.
- Groot, A.T., Inglis, O., Bowdridge, S., Santangelo, R.G., Blanco, C., Lopez Jr, J.D., Vargas, A.T., Gould, F. & Schal, C. 2009. Geographic and temporal variation in moth chemical communication. *Evolution* **63**: 1987-2003.
- Groot, A.T., Schöfl, G., Inglis, O., Donnerhacke, S., Classen, A., Schmalz, A., Santangelo, R.G., Emerson, J., Gould, F. & Schal, C. 2014. Within-population variability in a moth sex pheromone blend: genetic basis and behavioural consequences. *Proc R Soc B Biol Sci* **281**: 20133054.
- Gustafsson, L., Nordling, D., Andersson, M., Sheldon, B. & Qvarnstrom, A. 1994. Infectious diseases, reproductive effort and the cost of reproduction in birds. *Philos Trans R Soc B Biol Sci* **346**: 323-331.
- Hansen, A.K. & Moran, N.A. 2014. The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol Ecol* **23**: 1473-1496.
- Hanssen, S.A., Hasselquist, D., Folstad, I. & Erikstad, K.E. 2005. Cost of reproduction in a long-lived bird: incubation effort reduces immune function and future reproduction. *Proc R Soc B Biol Sci* **272**: 1039-1046.
- Hedges, L.M., Brownlie, J.C., O'Neill, S.L. & Johnson, K.N. 2008. *Wolbachia* and virus protection in insects. *Science* **322**: 702-702.
- Himler, A.G., Adachi-Hagimori, T., Bergen, J.E., Kozuch, A., Kelly, S.E., Tabashnik, B.E., Chiel, E., Duckworth, V.E., Dennehy, T.J. & Zchori-Fein, E. 2011. Rapid spread of a bacterial symbiont in an invasive whitefly is driven by fitness benefits and female bias. *Science* **332**: 254-256.
- Hoffmann, A.A., Hercus, M. & Dagher, H. 1998. Population dynamics of the *Wolbachia* infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* **148**: 221-231.
- Hosokawa, T., Kikuchi, Y., Shimada, M. & Fukatsu, T. 2007. Obligate symbiont involved in pest status of host insect. *Proc R Soc B Biol Sci* **274**: 1979-1984.
- Hosokawa, T., Koga, R., Kikuchi, Y., Meng, X.-Y. & Fukatsu, T. 2010. *Wolbachia* as a bacteriocyte-associated nutritional mutualist. *Proc Natl Acad Sci USA* **107**: 769-774.
- Howe, G.A., Lightner, J. & Ryan, C. 1996. An octadecanoid pathway mutant (JL5) of tomato is compromised in signaling for defense against insect attack. *Plant Cell* **8**: 2067-2077.
- Huffman, M.A., Page, J.E., Sukhdeo, M.V., Gotoh, S., Kalunde, M.S., Chandrasiri, T. & Towers, G.N. 1996. Leaf-swallowing by chimpanzees: a behavioral adaptation for the control of strongyle nematode infections. *Int J Primatol* **17**: 475-503.

## General introduction

- Huigens, M., De Almeida, R., Boons, P., Luck, R. & Stouthamer, R. 2004. Natural interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia* in *Trichogramma* wasps. *Proc R Soc B Biol Sci* **271**: 509-515.
- Hutchings, M.R., Kyriazakis, I. & Gordon, I.J. 2001. Herbivore physiological state affects foraging trade-off decisions between nutrient intake and parasite avoidance. *Ecology* **82**: 1138-1150.
- Janeway, C.A. 1992. The immune system evolved to discriminate infectious nonself from noninfectious self. *Immunol Today* **13**: 11-16.
- Javoiš, J. & Tammaru, T. 2004. Reproductive decisions are sensitive to cues of life expectancy: the case of a moth. *Anim Behav* **68**: 249-255.
- Kaiser, W., Huguet, E., Casas, J., Commin, C. & Giron, D. 2010. Plant green-island phenotype induced by leaf-miners is mediated by bacterial symbionts. *Proc R Soc B Biol Sci*: rspb20100214.
- Kant, M.R., Ament, K., Sabelis, M.W., Haring, M.A. & Schuurink, R.C. 2004. Differential timing of spider mite-induced direct and indirect defenses in tomato plants. *Plant Physiol* **135**: 483-495.
- Kant, M.R., Sabelis, M.W., Haring, M.A. & Schuurink, R.C. 2008. Intraspecific variation in a generalist herbivore accounts for differential induction and impact of host plant defences. *Proc R Soc B Biol Sci* **275**: 443-452.
- Karban, R. & Baldwin, I.T. 2007. *Induced responses to herbivory*. University of Chicago Press.
- Kavaliers, M. & Colwell, D.D. 1995. Discrimination by female mice between the odours of parasitized and non-parasitized males. *Proc R Soc B Biol Sci* **261**: 31-35.
- Kikuchi, Y., Hosokawa, T. & Fukatsu, T. 2011. Specific developmental window for establishment of an insect-microbe gut symbiosis. *Appl Environ Microbiol* **77**: 4075-4081.
- Kikuchi, Y., Hayatsu, M., Hosokawa, T., Nagayama, A., Tago, K. & Fukatsu, T. 2012. Symbiont-mediated insecticide resistance. *Proc Natl Acad Sci USA* **109**: 8618-8622.
- Koch, H. & Schmid-Hempel, P. 2011. Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proc Natl Acad Sci USA* **108**: 19288-19292.
- Kraaijeveld, A. & Godfray, H. 1997. Trade-off between parasitoid resistance and larval competitive ability in *Drosophila melanogaster*. *Nature* **389**: 278-280.
- Kraaijeveld, A.R. & Wertheim, B. 2009. Costs and genomic aspects of *Drosophila* immunity to parasites and pathogens. In: *Insect Infection and Immunity* (J. Rolff & C. Reynolds, eds), pp. 187-205. Oxford University Press, Oxford.
- Krams, I., Kecko, S., Kangassalo, K., Moore, F.R., Jankevics, E., Inashkina, I., Krama, T., Lietuviētis, V., Meija, L. & Rantala, M.J. 2014. Effects of food quality on trade-offs among growth, immunity and survival in the greater wax moth *Galleria mellonella*. *Insect Sci* **22**: 431-439.
- Kuraishi, T., Binggeli, O., Oputa, O., Buchon, N. & Lemaître, B. 2011. Genetic evidence for a protective role of the peritrophic matrix against intestinal bacterial infection in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **108**: 15966-15971.
- Lavine, M.D. & Strand, M.R. 2002. Insect hemocytes and their role in immunity. *Insect Biochem Mol Biol* **32**: 1295-1309.
- Lefèvre, T., Oliver, L., Hunter, M.D. & De Roode, J.C. 2010. Evidence for trans-generational medication in nature. *Ecol Lett* **13**: 1485-1493.
- Lehane, M. 1997. Peritrophic matrix structure and function. *Annu Rev Entomol* **42**: 525-550.
- Li, C., Williams, M.M., Loh, Y.-T., Lee, G.I. & Howe, G.A. 2002. Resistance of cultivated tomato to cell content-feeding herbivores is regulated by the octadecanoid-signaling pathway. *Plant Physiol* **130**: 494-503.
- Manson, J.S., Otterstatter, M.C. & Thomson, J.D. 2010. Consumption of a nectar alkaloid reduces pathogen load in bumble bees. *Oecol* **162**: 81-89.

- Marsh, S.E., Poulsen, M., Pinto-Tomás, A. & Currie, C.R. 2014. Interaction between workers during a short time window is required for bacterial symbiont transmission in *Acromyrmex* leaf-cutting ants. *PLoS ONE* **9**: e103269.
- Martinson, V.G., Moy, J. & Moran, N.A. 2012. Establishment of characteristic gut bacteria during development of the honeybee worker. *Appl Environ Microbiol* **78**: 2830-2840.
- Mason, C.J., Couture, J.J. & Raffa, K.F. 2014. Plant-associated bacteria degrade defense chemicals and reduce their adverse effects on an insect defoliator. *Oecol* **175**: 901-910.
- McKean, K.A., Yourth, C.P., Lazzaro, B.P. & Clark, A.G. 2008. The evolutionary costs of immunological maintenance and deployment. *BMC Evol Biol* **8**: 76.
- Medzhitov, R. & Janeway, C.A. 1997. Innate immunity: the virtues of a nonclonal system of recognition. *Cell* **91**: 295-298.
- Migeon, A., Nouguié, E. & Dorkeld, F. 2011. Spider Mites Web: a comprehensive database for the Tetranychidae. In: *Trends in Acarology* (M.W. Sabelis & J. Bruin, eds), pp. 557-560. Springer, The Netherlands.
- Minchella, D.J. & Loverde, P.T. 1981. A cost of increased early reproductive effort in the snail *Biomphalaria glabrata*. *Am Nat* **118**: 876-881.
- Moran, N.A. & Baumann, P. 2000. Bacterial endosymbionts in animals. *Curr Opin Microbiol* **3**: 270-275.
- Moran, N.A. & Wernegreen, J.J. 2000. Lifestyle evolution in symbiotic bacteria: insights from genomics. *Trends Ecol Evol* **15**: 321-326.
- Moran, N.A., McCutcheon, J.P. & Nakabachi, A. 2008. Genomics and evolution of heritable bacterial symbionts. *Annu Rev Genet* **42**: 165-190.
- Moreira, L.A., Iturbe-Ormaetxe, I., Jeffery, J.A., Lu, G., Pyke, A.T., Hedges, L.M., Rocha, B.C., Hall-Mendelin, S., Day, A. & Riegler, M. 2009. A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and *Plasmodium*. *Cell* **139**: 1268-1278.
- Moret, Y. & Schmid-Hempel, P. 2000. Survival for immunity: the price of immune system activation for bumblebee workers. *Science* **290**: 1166-1168.
- Morgan, A. & Chamberlin, F.S. 1927. tobacco budworm and its control in the Georgia and Florida tobacco-growing region. *USDA Farmer's Bull* **1531**: 1-91.
- Munson, M., Baumann, P., Clark, M., Baumann, L., Moran, N., Voegtlin, D. & Campbell, B. 1991. Evidence for the establishment of aphid-eubacterium endosymbiosis in an ancestor of four aphid families. *J Bacteriol* **173**: 6321-6324.
- Nappi, A. & Vass, E. 1993. Melanogenesis and the generation of cytotoxic molecules during insect cellular immune reactions. *Pigm Cell Res* **6**: 117-126.
- Nappi, A. & Vass, E. 1998. Hydrogen peroxide production in immune-reactive *Drosophila melanogaster*. *J Parasitol* **84**: 1150-1157.
- Nappi, A. & Christensen, B. 2005. Melanogenesis and associated cytotoxic reactions: applications to insect innate immunity. *Insect Biochem Mol Biol* **35**: 443-459.
- Nappi, A.J., Vass, E., Frey, F. & Carton, Y. 1995. Superoxide anion generation in *Drosophila* during melanotic encapsulation of parasites. *Eur J Cell Biol* **68**: 450-456.
- Neunzig, H. 1969. Biology of the tobacco budworm and the corn earworm in North Carolina. *North Carolina Agr. Exp. Sta. Tech. Bull.* **196**: 1-76.
- Nikoh, N., Hosokawa, T., Moriyama, M., Oshima, K., Hattori, M. & Fukatsu, T. 2014. Evolutionary origin of insect-*Wolbachia* nutritional mutualism. *Proc Natl Acad Sci USA* **111**: 10257-10262.
- Oliver, K.M., Russell, J.A., Moran, N.A. & Hunter, M.S. 2003. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. *Proc Natl Acad Sci USA* **100**: 1803-1807.
- Oliver, K.M., Degan, P.H., Burke, G.R. & Moran, N.A. 2010. Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. *Annu Rev Entomol* **55**: 247-266.

## General introduction

- Orr, M.R. 1992. Parasitic flies (Diptera: Phoridae) influence foraging rhythms and caste division of labor in the leaf-cutter ant, *Atta cephalotes* (Hymenoptera: Formicidae). *Behav Ecol Sociobiol* **30**: 395-402.
- Part, T., Gustafsson, L. & Moreno, J. 1992. 'Terminal investment' and a sexual conflict in the collared flycatcher (*Ficedula albicollis*). *Am Nat* **140**: 868-882.
- Penn, D. & Potts, W.K. 1998. Chemical signals and parasite-mediated sexual selection. *Trends Ecol Evol* **13**: 391-396.
- Penn, D., Schneider, G., White, K., Slev, P. & Potts, W. 1998. Influenza infection neutralizes the attractiveness of male odour to female mice (*Mus musculus*). *Ethol* **104**: 685-694.
- Perrot-Minnot, M.J., Cheval, B., Migeon, A. & Navajas, M. 2002. Contrasting effects of *Wolbachia* on cytoplasmic incompatibility and fecundity in the haplodiploid mite *Tetranychus urticae*. *J Evol Biol* **15**: 808-817.
- Pieterse, C.M., Van der Does, D., Zamioudis, C., Leon-Reyes, A. & Van Wees, S.C. 2012. Hormonal modulation of plant immunity. *Annu Rev Cell Dev Biol* **28**: 489-521.
- Poulin, R. 2007. *Evolutionary ecology of parasites*. Princeton University Press, New Jersey.
- Powell, J.E., Martinson, V.G., Urban-Mead, K. & Moran, N.A. 2014. Routes of acquisition of the gut microbiota of the honey bee *Apis mellifera*. *Appl Environ Microbiol* **80**: 7378-7387.
- Proshold, F., Karpenko, C. & Graham, C. 1982. Egg production and oviposition in the tobacco budworm: effect of age at mating. *Ann Entomol Soc Am* **75**: 51-55.
- Ramaswamy, S.B. 1990. Periodicity of oviposition, feeding, and calling by mated female *Heliothis virescens* in a field cage. *J Insect Behav* **3**: 417-427.
- Reeves, E.P., Lu, H., Jacobs, H.L., Messina, C.G., Bolsover, S., Gabella, G., Potma, E.O., Warley, A., Roes, J. & Segal, A.W. 2002. Killing activity of neutrophils is mediated through activation of proteases by K<sup>+</sup> flux. *Nature* **416**: 291-297.
- Roff, D. & Fairbairn, D. 2007. The evolution of trade-offs: where are we? *J Evol Biol* **20**: 433-447.
- Rolff, J. 2002. Bateman's principle and immunity. *Proc R Soc B Biol Sci* **269**: 867-872.
- Rolff, J. & Siva-Jothy, M.T. 2002. Copulation corrupts immunity: a mechanism for a cost of mating in insects. *Proc Natl Acad Sci USA* **99**: 9916-9918.
- Rosenberg, E., Sharon, G. & Zilber-Rosenberg, I. 2009. The hologenome theory of evolution contains Lamarckian aspects within a Darwinian framework. *Environ Microbiol* **11**: 2959-2962.
- Royet, J. 2004. Infectious non-self recognition in invertebrates: lessons from *Drosophila* and other insect models. *Mol Immunol* **41**: 1063-1075.
- Ruby, E., Henderson, B. & McFall-Ngai, M. 2004. We get by with a little help from our (little) friends. *Science* **303**: 1305-1307.
- Rupley, J. 1967. The binding and cleavage by lysozyme of N-acetylglucosamine oligosaccharides. *Proc R Soc B Biol Sci* **167**: 416-428.
- Russell, J.A., Moreau, C.S., Goldman-Huertas, B., Fujiwara, M., Lohman, D.J. & Pierce, N.E. 2009. Bacterial gut symbionts are tightly linked with the evolution of herbivory in ants. *Proc Natl Acad Sci USA* **106**: 21236-21241.
- Salem, H., Kreutzer, E., Sudakaran, S. & Kaltenpoth, M. 2013. Actinobacteria as essential symbionts in firebugs and cotton stainers (Hemiptera, Pyrrhocoridae). *Environ Microbiol* **15**: 1956-1968.
- Salem, H., Bauer, E., Strauss, A.S., Vogel, H., Marz, M. & Kaltenpoth, M. 2014. Vitamin supplementation by gut symbionts ensures metabolic homeostasis in an insect host. *Proc R Soc B Biol Sci* **281**: 20141838.
- Salem, H., Florez, L., Gerardo, N. & Kaltenpoth, M. 2015. An out-of-body experience: the extracellular dimension for the transmission of mutualistic bacteria in insects. *Proc R Soc B Biol Sci* **282**: 20142957.
- Schmid-Hempel, P. 2005. Evolutionary ecology of insect immune defenses. *Annu Rev Entomol* **50**: 529-551.

- Schmid-Hempel, P. 2011. *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford University Press, New York.
- Schoonhoven, L.M., van Loon, J.J.A. & Dicke, M. 2005. *Insect-Plant Biology, 2nd edn*. Oxford University Press, Oxford.
- Shade, A. & Handelsman, J. 2012. Beyond the Venn diagram: the hunt for a core microbiome. *Environ Microbiol* **14**: 4-12.
- Sheck, A. & Gould, F. 1993. The genetic basis of host range in *Heliothis virescens*: larval survival and growth. *Entomol Exp Appl* **69**: 157-172.
- Sheldon, B.C. & Verhulst, S. 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol Evol* **11**: 317-321.
- Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y. & Ishikawa, H. 2000. Genome sequence of the endocellular bacterial symbiont of aphids *Buchnera* sp. APS. *Nature* **407**: 81-86.
- Singer, M.S., Mace, K.C. & Bernays, E.A. 2009. Self-medication as adaptive plasticity: increased ingestion of plant toxins by parasitized caterpillars. *PLoS ONE* **4**: e4796-e4796.
- Söderhäll, K. & Cerenius, L. 1998. Role of the prophenoloxidase-activating system in invertebrate immunity. *Curr Opin Immunol* **10**: 23-28.
- Soler, J.J., de Neve, L., Pérez-Contreras, T., Soler, M. & Sorci, G. 2003. Trade-off between immunocompetence and growth in magpies: an experimental study. *Proc R Soc B Biol Sci* **270**: 241-248.
- Stadelbacher, E. 1981. Role of early-season wild and naturalized host plants in the buildup of the F1 generation of *Heliothis zea* and *H. virescens* in the Delta of Mississippi. *Environ Entomol* **10**: 766-770.
- Stearns, S.C. 1989. Trade-offs in life-history evolution. *Funct Ecol* **3**: 259-268.
- Stouthamer, R., Breeuwer, J.A. & Hurst, G.D. 1999. *Wolbachia pipientis*: microbial manipulator of arthropod reproduction. *Annu Rev Microbiol* **53**: 71-102.
- Strand, M.R. 2008. The insect cellular immune response. *Insect Sci* **15**: 1-14.
- Stuart, L., Boulais, J., Charriere, G., Hennessy, E., Brunet, S., Jutras, I., Goyette, G., Rondeau, C., Letarte, S. & Huang, H. 2007. A systems biology analysis of the *Drosophila* phagosome. *Nature* **445**: 95-101.
- Stuart, L.M. & Ezekowitz, R.A.B. 2005. Phagocytosis: elegant complexity. *Immunity* **22**: 539-550.
- Su, Q., Oliver, K.M., Xie, W., Wu, Q., Wang, S. & Zhang, Y. 2015. The whitefly-associated facultative symbiont *Hamiltonella defensa* suppresses induced plant defences in tomato. *Funct Ecol* **29**: 1007-1018.
- Sy, A., Timmers, A.C., Knief, C. & Vorholt, J.A. 2005. Methylothetic metabolism is advantageous for *Methylobacterium extorquens* during colonization of *Medicago truncatula* under competitive conditions. *Appl Environ Microbiol* **71**: 7245-7252.
- Teixeira, L., Ferreira, Á. & Ashburner, M. 2008. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol*: e1000002
- Thaler, J.S., Farag, M.A., Paré, P.W. & Dicke, M. 2002. Jasmonate-deficient plants have reduced direct and indirect defences against herbivores. *Ecol Lett* **5**: 764-774.
- Thaler, J.S., Humphrey, P.T. & Whiteman, N.K. 2012. Evolution of jasmonate and salicylate signal crosstalk. *Trends Plant Sci* **17**: 260-270.
- Tsuchida, T., Koga, R., Matsumoto, S. & Fukatsu, T. 2011. Interspecific symbiont transfection confers a novel ecological trait to the recipient insect. *Biol Lett* **7**: 245-248.
- Vala, F., Breeuwer, J.A. & Sabelis, M.W. 2000. *Wolbachia*-induced 'hybrid breakdown' in the two-spotted spider mite *Tetranychus urticae* Koch. *Proc R Soc B Biol Sci* **267**: 1931-1937.
- van Ham, R.C., Kamerbeek, J., Palacios, C., Rausell, C., Abascal, F., Bastolla, U., Fernández, J.M., Jiménez, L., Postigo, M. & Silva, F.J. 2003. Reductive genome evolution in *Buchnera aphidicola*. *Proc Natl Acad Sci USA* **100**: 581-586.

## General introduction

- Vetter, R.S. & Baker, T.C. 1983. Behavioral responses of male *Heliothis virescens* in a sustained-flight tunnel to combinations of seven compounds identified from female sex pheromone glands. *J Chem Ecol* **9**: 747-759.
- Vorholt, J.A. 2012. Microbial life in the phyllosphere. *Nat Rev Microbiol* **10**: 828-840.
- Waldvogel, M. & Gould, F. 1990. Variation in oviposition preference of *Heliothis virescens* in relation to macroevolutionary patterns of Heliothine host range. *Evolution* **44**: 1326-1337.
- Walling, L.L. 2000. The myriad plant responses to herbivores. *J Plant Growth Regul* **19**: 195-216.
- Weeks, A.R., Marec, F. & Breeuwer, J.A. 2001. A mite species that consists entirely of haploid females. *Science* **292**: 2479-2482.
- Werren, J.H. 1997. Biology of *Wolbachia*. *Annu Rev Entomol* **42**: 587-609.
- Werren, J.H., Baldo, L. & Clark, M.E. 2008. *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* **6**: 741-751.
- Willers, J., Schneider, J. & Ramaswamy, S. 1987. Fecundity, longevity and caloric patterns in female *Heliothis virescens*: changes with age due to flight and supplemental carbohydrate. *J Insect Physiol* **33**: 803-808.
- Williams, G.C. 1966. Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am Nat* **100**: 687-690.
- Xie, J., Vilchez, I. & Mateos, M. 2010. *Spiroplasma* bacteria enhance survival of *Drosophila hydei* attacked by the parasitic wasp *Leptopilina heterotoma*. *PLoS ONE* **5**: e12149.
- Xie, J., Butler, S., Sanchez, G. & Mateos, M. 2014. Male killing *Spiroplasma* protects *Drosophila melanogaster* against two parasitoid wasps. *Heredity* **112**: 399-408.
- Xie, R.-R., Chen, X.-L. & Hong, X.-Y. 2011. Variable fitness and reproductive effects of *Wolbachia* infection in populations of the two-spotted spider mite *Tetranychus urticae* Koch in China. *Appl Entomol Zool* **46**: 95-102.
- Zasloff, M. 2002. Antimicrobial peptides of multicellular organisms. *Nature* **415**: 389-395.
- Zhang, Y.K., Ding, X.L., Rong, X. & Hong, X.Y. 2015. How do hosts react to endosymbionts? A new insight into the molecular mechanisms underlying the *Wolbachia*-host association. *Insect Mol Biol* **24**: 1-12.
- Zhang, Z.T. & Zhu, S.Y. 2009. Drosomycin, an essential component of antifungal defence in *Drosophila*. *Insect Mol Biol* **18**: 549-556.
- Zhu, F., Poelman, E.H. & Dicke, M. 2014. Insect herbivore-associated organisms affect plant responses to herbivory. *New Phytol* **204**: 315-321.
- Zilber-Rosenberg, I. & Rosenberg, E. 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* **32**: 723-735.
- Zug, R. & Hammerstein, P. 2015. Bad guys turned nice? A critical assessment of *Wolbachia* mutualisms in arthropod hosts. *Biol Rev* **90**: 89-111.
- Zuk, M. & Stoehr, A.M. 2002. Immune defense and host life history. *Am Nat* **160**: S9-S22.

Variability of bacterial communities in the  
moth *Heliothis virescens* indicates transient  
association with the host

Heike Staudacher, Martin Kaltenpoth, Johannes A.J. Breeuwer,  
Steph B.J. Menken, David G. Heckel & Astrid T. Groot

*Manuscript submitted*

### **Abstract**

Microbes associated with insects can confer a wide range of ecologically relevant benefits to their hosts. Since insect-associated bacteria often increase the nutritive value of their hosts' diets, the study of bacterial communities is especially interesting in species that are important agricultural pests. We investigated the composition of bacterial communities in the noctuid moth *Heliothis virescens* and its variability in relation to developmental stage, diet and population (field and laboratory), using bacterial tag-encoded FLX pyrosequencing of 16S rRNA amplicons. In larvae, bacterial communities differed depending on the food plant on which they had been reared, although the within-group variation between biological replicates was high as well. Moreover, larvae originating from a field or laboratory population did not share any OTUs. Interestingly, *Enterococcus* sp. was found to be the dominant taxon in laboratory-reared larvae, but was completely absent from field larvae, indicating dramatic shifts in microbial community profiles upon cultivation of the moths in the laboratory. Furthermore, microbiota composition varied strongly across developmental stages in individuals of the field population, and we found no evidence for vertical transmission of bacteria from mothers to offspring. Overall, the high variability in bacterial communities suggests that the bacteria that we identified are only loosely and temporarily associated with *H. virescens*.

## **INTRODUCTION**

Symbiotic bacteria of insects provide diverse beneficial services to their hosts, e.g., upgrading of nutrient-poor diets, digestion of refractory food sources and protection of the host against pathogens (e.g., Dillon & Dillon, 2004; Douglas, 2009; Feldhaar, 2011; Engel & Moran, 2013). Therefore, the study of bacterial communities that are associated with an organism of interest has become an integral part of the fundamental and applied research of, in particular, agricultural pests. Investigating bacterial diversity and insect-bacteria associations is greatly facilitated by high-throughput sequencing methods, especially massive parallel amplicon sequencing, that have been developed in the past decade (Margulies *et al.*, 2005; Rothberg & Leamon, 2008; Bosch & McFall-Ngai, 2011; Loman *et al.*, 2012).

Associations of bacterial communities with a host can be stable or dynamic. Stable associations are characterized by the temporal persistence of bacteria in the host across different life stages and across generations (Zilber-Rosenberg & Rosenberg, 2008; Rosenberg *et al.*, 2009; Shade & Handelsman, 2012; Engel & Moran, 2013; Salem *et al.*, 2015). The stability of the community can be indicative of the relevance of bacteria for the ecology and evolution of the host. If bacteria are transmitted vertically from one generation to the next for many generations, host and bacteria can be viewed as a unit of selection, and bacteria are likely beneficial for the host (Zilber-Rosenberg & Rosenberg, 2008; Rosenberg *et al.*, 2009; Oliver *et al.*, 2010).

Another criterion to assess the relevance or the functional role of bacteria for an organism is the variability of bacterial communities in the different environments in which the insect host resides (Robinson *et al.*, 2010; Chandler *et al.*, 2011; Colman *et al.*, 2012; Gayatri Priya *et al.*, 2012; Sullam *et al.*, 2012). Variability of bacterial communities in relation to host diet is of particular interest in agricultural and forest pest insects, because the flexibility of their bacterial communities may enhance the dietary range of phytophagous insects (Dillon & Dillon, 2004; Hosokawa *et al.*, 2007; Douglas, 2009; Tsuchida *et al.*, 2011; Chu *et al.*, 2013; Engel & Moran, 2013). Bacterial communities have been shown to vary depending on diet in several insect species (Broderick *et al.*, 2004; Belda *et al.*, 2011; Chandler *et al.*, 2011; Gayatri Priya *et al.*, 2012) but may also be stable across diets (Sudakaran *et al.*, 2012). Other factors that may cause variation in bacterial communities in insects are the geographic origin of the host population or rearing history of the insect host (Xiang *et al.*, 2006; Adams *et al.*, 2010; Belda *et al.*, 2011).

Studies of bacterial communities in Lepidoptera have accumulated in the past decade, mostly because larvae of many Lepidoptera are major agricultural or forest pests (Broderick *et al.*, 2004; Xiang *et al.*, 2006; Belda *et al.*, 2011; Gayatri Priya *et al.*, 2012; Xia *et al.*, 2013). Bacteria from the genus *Enterococcus* have been repeatedly found in Lepidoptera (Jarosz, 1975; Broderick *et al.*, 2004; Xiang *et al.*, 2006; Brinkmann *et al.*, 2008; Tang *et al.*, 2012; Hammer *et al.*, 2014) and have been shown to be metabolically active in *Manduca sexta* (Brinkmann *et al.*, 2008). Other bacterial candidates that could serve an ecological role in Lepidoptera were isolated from saturniid butterflies and the silkworm *Bombyx mori* (Pinto-Tomas *et al.*, 2007; Prem Anand *et al.*, 2010). These bacteria exhibited, among others, cellulolytic, pectinolytic and xylanolytic activities that might help caterpillars to digest plant material (Pinto-Tomas *et al.*, 2007; Prem Anand *et al.*, 2010). Also, a recent metagenomic study of the bacterial midgut community in the pyralid moth *Ostrinia nubilalis* revealed the presence of bacterial cellulase, amylase,  $\beta$ -galactosidase and  $\beta$ -glucosidase encoding genes, indicating the potential of the gut bacterial community to support digestion in its lepidopteran host (Belda *et al.*, 2011). However, when the midgut is damaged by insecticidal toxins in spores of *Bacillus thuringiensis*, formerly benign, resident bacteria may breach the midgut barrier, enter the hemocoel and participate in the destruction of the host (Broderick *et al.*, 2006, 2009).

The noctuid moth *Heliothis virescens* is a major agricultural pest in North and South America (Fitt, 1989). Its larvae are polyphagous and feed on over 37 plant species from 14 families (Barber, 1937; Stadelbacher, 1981; Fitt, 1989; Waldvogel & Gould, 1990; Sheck & Gould, 1993; Blanco *et al.*, 2007). Among these plants are important economic crops such as cotton (*Gossypium hirsutum*), tobacco (*Nicotiana tabacum*) and chickpea (*Cicer arietinum*) (Barber, 1937; Neunzig, 1969; Fitt, 1989; Blanco *et al.*, 2007). Since bacteria can facilitate host-plant use by herbivorous insects, investigating the bacterial community of *H. virescens* and its variation in

relation to diet could lead to the identification of bacterial candidates that enhance or inhibit its development and thereby its impact as an agricultural pest.

In this study, we identified and compared bacterial communities in different developmental stages (eggs, larvae and female adults) of *H. virescens* from the field and laboratory, and reared on different plant species (cotton, chickpea and tobacco). Additionally, we investigated the transmission of the bacterial community from one generation to the next.

## **MATERIAL AND METHODS**

### ***Ethics statement***

*Heliothis virescens* was collected in North Carolina, USA, where this species is a pest and not protected by law. Eggs of *H. virescens* were collected on private property, for which no permits were required.

### ***Effect of host plant species on bacterial community composition of laboratory-reared larvae***

To investigate whether the bacterial community composition in *H. virescens* larvae changes depending on the food plant species, we used the long-term laboratory-reared ARS strain. This strain was started in 1971 in Washington County, MS, and since then reared in the laboratory at USDA-ARS in Stoneville, MS, and transferred to the Max Planck Institute for Chemical Ecology (MPI-CE), Jena, Germany in 2010, where the larvae were reared on pinto bean diet containing the antibiotic tetracycline hydrochloride (Sigma-Aldrich, The Netherlands) (Burton, 1970). Adults were provided with a 10% (vol/vol) honey water solution. All life stages were kept in climate chambers at a temperature of 25 °C, 60% relative humidity and a light/dark cycle of 16h:8h.

We placed 30 first instars of the ARS strain on whole cotton (*Gossypium hirsutum*), chickpea (*Cicer arietinum*) and tobacco plants (*Nicotiana attenuata*), which were grown in the greenhouse of MPI-CE, Jena, Germany. After nine days, when the larvae were in the second, third or fourth instar, they were taken off the plants and starved for 8 h to empty their gut content and to reduce the amount of chloroplasts, as chloroplast DNA may be amplified in the procedure of 16S rRNA amplicon sequencing. To remove bacteria that resided on the outer cuticle, experimental larvae were washed by submerging them for 10 s in 1% (wt/vol) sodium dodecyl sulfate (SDS) solution and for 10 s in sterile water. Larvae were then put in 1.5-ml Eppendorf tubes, immediately placed on ice and then stored at -80 °C.

### ***Effect of host plant species on bacterial community composition of field larvae***

*Heliothis virescens* eggs were collected in July 2011 in North Carolina, USA, on commercially grown tobacco, *N. tabacum*. Experiments were conducted with larvae, referred to as field larvae, as well as females that developed from these larvae, and

eggs that were laid by these females. The field-collected eggs were divided into three groups in plastic cups (100 ml) which were filled with leaf material of one of the three plant species cotton (*G. hirsutum*), chickpea (*C. arietinum*) and tobacco (*N. tabacum*). After emergence, batches of 20-30 first instar larvae were transferred to whole plants, resulting in 10 batches per host plant species. The plants were kept outdoors until experiments started (i.e., when larvae were placed on them to feed). Each plant was kept in an individual cage made of fine-meshed gauze (diameter: 60 cm, height: 1 m). After larvae had reached the fifth instar, one larva was randomly collected from each plant batch and starved for 8 h. These 30 larvae were washed by submerging them for 10 s in 1% (wt/vol) SDS solution and for 10 s in sterile water. After the wash step, guts were extracted by cutting the larva between the 1st and 2nd thoracic and between the 7th and 8th abdominal segments. Dissected guts were stored at -80 °C until DNA extraction. The remaining larvae were left to pupate in the soil of the potted whole plant and subsequently used in the bacterial transmission experiment (see below).

#### ***Diversity and composition of the bacterial community in adult females and their eggs***

To determine bacterial communities in females and their corresponding eggs, a subset of the field larvae that was not used to determine the larval bacterial community was allowed to pupate in the soil and develop into adults. Upon emergence, males and females that had fed on the same plant species were mated in carton cups (200 ml) in single pairs. The cups contained leaves of the same plant species on which the mating pairs had been as larvae to stimulate, and provide a substrate for, oviposition. After four days, we collected abdomens from three females and 30 of their eggs per plant species group, and thus obtained nine female-egg combinations in total. DNA from eggs was extracted in batches of 30 eggs per female. Eggs were collected from the plant leaves with a brush, transferred to 1.5-ml Eppendorf tubes and placed immediately on ice until storage at -80 °C. Female abdomens were cut off and submerged in 1% (wt/vol) SDS solution for 10 s, followed by a 10 s washing step in sterile water. Abdomens were then transferred to 1.5-ml Eppendorf tubes and placed immediately on ice until storage at -80 °C. All instruments were rinsed with 1% (wt/vol) SDS solution and sterile water between samples. All tissues were collected in 1.5-ml Eppendorf tubes and placed immediately on ice to prevent changes in the bacterial community composition, after which the samples were stored at -80 °C.

#### ***Effect of laboratory-rearing and antibiotics treatment on bacterial communities of field larvae***

To assess whether bacterial communities change after transfer from the field to the laboratory and how the antibiotic tetracycline that is normally part of the laboratory diet affects the bacterial gut community, moths that originated from eggs that were collected in the field to assess microbial communities (see above) were kept in the

laboratory at the University of Amsterdam for three generations (in the following referred to as field-lab larvae). Larvae were kept on artificial pinto-bean diet as described above, but without the addition of antibiotics. Females of the third generation were mated in single pairs in empty 200-ml plastic cups. Eggs of the fourth generation hatched in these cups, and the resulting first instars were transferred to cotton leaves with (AB group) or without tetracycline coating (NoAB group). Every other day, cotton leaves were coated with tetracycline by pipetting 1 ml 0.1% (wt/vol) tetracycline onto the upper leaf surface. Control leaves were treated in the same way with only water. In addition, we wrapped the leaf stems with cotton soaked with 0.1% (wt/vol) tetracycline or water in the treatment or the control group, respectively. When the larvae reached the fifth instar, they were starved for 8 h, then washed by submerging them for 10 s in 1% SDS solution and for 10 s in sterile water, after which their gut was extracted as described above. The guts were stored at -80 °C until DNA extraction.

#### **DNA extraction**

DNA from the field and field-lab population was extracted by grinding samples in 500 µl TES buffer [100 mM tris(hydroxymethyl) aminomethane hydrochloride pH 8, 10 M ethylenediaminetetraacetic acid, 2% sodium dodecyl sulfate] and 4 µl lysozyme from chicken egg white (100 mg/ml) (Sigma-Aldrich), after which the samples were incubated at 37 °C for 30 min. Next we added 2.5 µl proteinase K (20 mg/ml) (Sigma-Aldrich) and incubated the samples for 56 °C overnight. The rest of the extraction procedure was conducted according to the standard CTAB/chloroform protocol as described in Unbehend *et al.* (2013). DNA from the lab-strain larvae was extracted with the Epicenter MasterPure kit (Epicentre Technologies, USA) according to manufacturer's instructions. Additionally, a lysozyme step [4 µl lysozyme (100 mg/ml), incubated at 37 °C for 30 min] was added before the proteinase K step.

#### **Bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP)**

Samples were pooled for sequencing. For the larvae of the laboratory strain, we made three pools, one pool contained DNA of seven larvae that were reared on each of the three plant species (cotton, tobacco or chickpea). For the larvae of the field population, we made six pools of DNA, two pools (biological replicates) for each plant species, to assess the variation between biological replicates; each pool contained the DNA of five larvae. For the field-lab population, we made two pools, one pool of 45 NoAB larvae and one pool of 49 AB larvae. Pooled samples were sent to an external service provider (Molecular Research Lab, MR DNA, Shallowater, TX, USA) for bTEFAP, using the 16S rRNA primers Gray28F (5'-GAGTTTGATC NTGGCTCA-3') and Gray519R (5'-GTNTTACNGCGGCKGCTG-3') (Ishak *et al.*, 2011). A sequencing library was constructed via one-step PCR with 30 cycles, using a mixture of Hot Start and HotStar hi-fidelity polymerases (Qiagen). Sequencing

was performed on a Roche 454 FLX instrument with Titanium reagents and procedures and protocols of Molecular Research LP (<http://www.mrdnlab.com/>).

### ***Sequence analysis***

Analysis of the sequences was performed in QIIME (quantitative insights into microbial ecology), which is a standard pipeline to analyse microbial communities based on 16S rRNA amplicon sequencing (Caporaso *et al.*, 2010a). Raw data were de-noised with `denoise_wrapper` (Reeder & Knight, 2010). Low-quality reads (quality cut-off = 25) and sequences that were shorter than 200 bp or longer than 600 bp were excluded from further analysis. The remaining sequences were clustered into operational taxonomic units (OTUs) with the open-reference OTU picking command, using `uclust` (version 1.2.22) (Edgar, 2010) and applying 97% similarity cut-offs. Sequences were first clustered against a reference data set (<http://greengenes.lbl.gov/>). The sequences that did not cluster with the reference sequences were clustered *de novo* and checked for chimeras with Chimera Slayer (Haas *et al.*, 2011). Identified chimeras were removed from the dataset for downstream analysis.

From each OTU cluster, the most abundant sequence was taken as the representative sequence. Taxonomy was assigned to the representative sequences with the `uclust` consensus taxonomy classifier. The resulting OTU table was manually edited, and global singletons as well as chloroplasts, mitochondria and OTUs that were classified as moth genomic DNA were removed. Representative sequences of the remaining OTUs were aligned with PyNAST using the Greengenes core set (<http://greengenes.lbl.gov/>) as a template (Caporaso *et al.*, 2010b). A phylogenetic tree was constructed with the open source software Fasttree 2.1.3, applying the generalized time-reversible (GTR) model of nucleotide evolution (Price *et al.*, 2010). Local support values for tree splits were calculated with the Shimodaira-Hasegawa test (Shimodaira & Hasegawa, 1999). This phylogenetic tree, which included all identified OTUs, was used to calculate the weighted UniFrac metric (Lozupone & Knight, 2005) which was used as the basis for principal coordinates analysis (PCoA) (see below).

### ***Diversity and phylogenetic analysis***

Diversity analysis of our samples was done with the QIIME pipeline. Since our samples showed unequal sampling depth (see TABLES S2.1-2.3), we investigated alpha- and beta-diversity of the bacterial communities of our samples with rarefied OTU tables. The number of sequences used for rarefaction corresponded to the number of sequences present in the sample with the lowest number of sequences (339 sequences). To assess alpha-diversity, we calculated the Shannon index based on the rarefaction tables (Shannon, 1948). To identify possible clusters based on similarity of bacterial communities among our samples, we produced a PCoA plot on the basis of weighted non-normalized UniFrac distances and with rarefied OTU

tables (Lozupone & Knight, 2005). To assess the robustness of our results, we additionally performed a Jackknife analysis on weighted UniFrac distances, which is included in the QIIME pipeline (see Lozupone & Knight, 2005; Caporaso *et al.*, 2010; Lozupone *et al.*, 2011). Jackknife support is included in the PCoA plots as the size of the ellipsoid areas surrounding the data points.

### ***OTU patterns across samples***

None of the bacterial OTUs was present in all samples, i.e., present in samples of all origins, rearing plants, and life stages. Most common were three OTUs which were present in 65% of the samples. These OTUs were mainly present in females and eggs, which had a high sample size compared to the other groups. To also consider OTUs that were present in the larvae, we filtered for OTUs that were on average represented more than 1% across all samples, resulting in 18 OTUs (i.e., first the percentage of an OTU in a sample was calculated followed by the average of the percentage of the OTU in all samples, see TABLE 2.1). To classify and construct a phylogenetic tree of these 18 OTUs, we first aligned the sequences of our dataset and 16S rRNA reference sequences obtained from Genbank (Benson *et al.*, 2011), using ClustalW (Thompson *et al.*, 1994), after which we constructed a maximum likelihood tree in MEGA 6 (Tamura *et al.*, 2013) with the Tamura-Nei model (Tamura & Nei, 1993), using uniform rate variation and 500 bootstrap replicates. To root the tree, we used three Archaea species as outgroups: *Halobacterium salinarum* (NR\_113428.1), *Sulfolobus acidocaldarius* (NR\_074267.1) and *Pyrolobus fumarii* (NR\_102985.1). Outgroup taxa were excluded from the FIGURE for clarity (FIGURE 2.1).

**TABLE 2.1.** Bacterial OTUs with an average presence of more than 1% across all samples.

#	Family (Genus)	Total Average (29)	Larvae Lab (3)	Larvae Field (6)	Larvae Field- lab (2)	Adults (9)	Eggs (9)	Overall presence > 1%
1	Enterobacteriaceae ( <i>Klebsiella/Enterobacter</i> )	27.63	0.00	0.00	35.62	72.34	35.95	68.97
2	Enterococcaceae ( <i>Enterococcus</i> )	11.40	50.55	0.00	61.70	0.00	0.01	17.24
3	Enterobacteriaceae ( <i>Serratia</i> )	8.57	0.00	0.00	0.01	7.20	22.42	44.83
4	Pseudomonadaceae ( <i>Pseudomonas</i> )	4.82	0.00	0.00	0.77	0.69	19.62	55.17
5	Enterobacteriaceae ( <i>Enterobacter</i> )	3.97	0.00	0.00	0.01	5.06	5.29	20.69
6	Acetobacteraceae ( <i>Asaia</i> )	2.38	49.45	0.00	0.00	0.00	0.00	6.90
7	Enterobacteriaceae (unclassified)	2.15	0.00	7.75	0.00	0.00	0.00	6.90
8	Mycobacteriaceae ( <i>Mycobacterium</i> )	2.15	0.00	0.00	0.00	0.03	4.13	27.59
9	Acidobacteria (unclassified)	2.04	0.00	8.42	0.00	0.00	0.00	3.45
10	Pseudomonadaceae ( <i>Pseudomonas</i> )	1.87	0.00	0.00	0.04	5.32	0.02	3.45

11	Methylobacteriaceae ( <i>Methylobacterium</i> )	1.57	0.00	31.45	0.00	0.00	0.00	13.79
12	Enterobacteriaceae ( <i>Klebsiella/Enterobacter</i> )	1.57	0.00	0.00	0.02	1.92	2.46	13.79
13	Pseudomonadaceae ( <i>Pseudomonas</i> )	1.54	0.00	0.00	0.00	0.17	2.55	20.69
14	Xanthomonadaceae ( <i>Stenotrophomonas</i> )	1.39	0.00	0.00	1.55	4.59	0.20	34.48
15	Paenibacillaceae ( <i>Paenibacillus</i> )	1.12	0.00	18.19	0.00	0.00	0.00	13.79
16	Moraxellaceae ( <i>Acinetobacter</i> )	1.04	0.00	0.00	0.04	0.08	6.53	24.14
17	Sphingomonadaceae ( <i>Sphingomonas</i> )	1.02	0.00	0.00	0.01	0.00	0.16	6.90
18	Comamonadaceae ( <i>Roseateles</i> )	1.00	0.00	8.84	0.07	0.00	0.01	13.79

Taxonomy and percentage of relative abundance in different developmental stages (egg, larva and adult) and populations (laboratory, field and field-lab). Sample sizes are given between brackets. #: OTU identification numbers.

### ***Statistical analysis of similarities between bacterial communities of females and their eggs***

To test whether egg bacterial communities resemble the bacterial community of their mother, we performed a linear mixed model in the package lme4 (Bates *et al.*, 2014) and lmerTest (Kuznetsova *et al.*, 2014) in the software R, version 3.0.2 (R Core Team, 2013). Weighted UniFrac distance values of all 81 female-egg combinations were used as response variable. We used two predictor variables: female-egg combination (eggs belonged to one female or not) and plant species (eggs belonged to the same plant species – cotton, chickpea, or tobacco – as a female or not). To account for repeatedly entering females and eggs into the model, we added egg and female as random factors into the model. Because there was no interaction effect between plant and female-egg combination, we excluded the interaction from the model. Degrees of freedom were approximated using the Satterthwaite method. As a *post hoc* test for the plant effect, we performed planned pairwise comparisons between female-eggs combinations of one plant species group and female-egg combinations in which females did, but eggs did not belong to that respective plant group, by using the multcomp package (Hothorn *et al.*, 2008) in R, applying a Bonferroni correction for multiple comparisons.

## **RESULTS**

### ***Bacterial community composition of *Heliothis virescens****

We received a total of 312,324 sequences, of which 295,289 sequences remained after quality filtering. Other sequences were global singletons or chimeras, derived from chloroplasts and mitochondria or fragments that were identified as part of the moth genome. The bacterial sequences clustered into 566 OTUs, of which 42 were unassigned. An overview of numbers of sequences before and after quality filtering, as well as numbers of OTUs in the different samples is given in TABLES S2.1-2.3.

None of the OTUs in *H. virescens* was present in all samples ( $n = 29$ ), and only three OTUs were found in more than 65% of our samples (65% was the highest percentage of samples in which OTUs were commonly present). When we assessed how many OTUs had an average percentage of more than 1% across all samples, this resulted in 18 OTUs, but only two of these were present with an average percentage of more than 10% across all samples (TABLE 2.1). The phylogenetic analysis revealed that these 18 OTUs belong to the following bacterial families (bacterial orders in parentheses): Acetobacteraceae (Rhodospirillales), Methylobacteriaceae (Rhizobiales), Sphingomonadaceae (Sphingomonadales), Comamonadaceae (Burkholderiales), Enterobacteriaceae (Enterobacteriales), Pseudomonadaceae, Moraxellaceae (both Pseudomonadales), Xanthomonadaceae (Xanthomonadales), Mycobacteriaceae (Actinomycetales), Paenibacillaceae (Bacillales), Enterococcaceae (Lactobacillales); and one OTU belonged to the phylum Acidobacteria (TABLE 2.1 and FIGURE 2.1).

#### ***Variability of bacterial communities between laboratory and field larvae***

We compared bacterial communities between larvae from a long-term laboratory colony and larvae that had been collected in the field as eggs. Laboratory and field larvae had no OTUs in common. The bacterial community of lab larvae was dominated by two OTUs that were classified as *Enterococcus* sp. (Lactobacillales) (#2) and *Asaia* sp. (Rhodospirillales) (#6), while larvae of the field population did not contain any OTUs affiliated with these genera. In fact, field larvae did not contain clearly dominant OTUs, but harboured Enterobacteriales, Burkholderiales or Rhizobiales in relatively high percentages (see below). Furthermore, bacterial communities in the field-lab larvae differed greatly from the bacterial communities of the field larvae: only 1.4% of the OTUs ( $n = 148$ ) of the field larvae were detected in the field-lab larvae that were reared without antibiotics (NoAB). *Enterococcus* sp. (#2) was more abundant in the AB field-lab larvae than in NoAB larvae (FIGURE 2.2).


Larvae of the field population and the laboratory colony also differed in the diversity of their associated bacterial communities. Larvae from the laboratory colony harboured an average of 25 OTUs, while field larvae had an average of 35 OTUs. Additionally, communities of the field larvae showed a greater evenness than those of the laboratory larvae (FIGURE 2.2), which was reflected in the higher Shannon index of the field larvae (FIGURE 2.3).

#### ***Host plant-associated variability of bacterial communities in larvae***


The influence of diet on the bacterial communities of laboratory and field larvae was investigated by comparing bacterial communities in larvae that fed on three different host plant species. In laboratory-reared larvae, we found one OTU (#2) from the family Enterococcaceae dominating the bacterial community in the guts of larvae fed chickpea or tobacco, i.e., 97.8 and 91.5% abundance, respectively (FIGURE 2.2). The



## Bacterial community in *Heliothis virescens*



**FIGURE 2.2.** Relative abundance of bacterial OTUs in *Heliothis virescens* larvae. Bacterial OTUs were combined on the family level. Numbers in brackets indicate how many OTUs one family contains. Rare OTUs that were represented less than 0.5% in at least one sample are not included. Larvae originated from a laboratory colony, from the field or from the field, after which they were reared in the laboratory for four generations (field-lab): AB: larvae received antibiotics treatment, NoAB: larvae received no antibiotics treatment. Field and laboratory larvae were grown on cotton (C), chickpea (Ch) or tobacco (T); square brackets around taxa indicate that the taxon name is not fully established yet.



**FIGURE 2.3.** Rarefaction curves of alpha diversity of the bacterial communities of *Heliothis virescens* larval guts. a) Number of observed bacterial OTUs. b) Shannon index of the bacterial communities. Laboratory (L) and field larvae (F) were fed on cotton (C), chickpea (Ch) or tobacco (T). AB: field larvae that were reared for four generations in the laboratory and received antibiotics during one generation; NoAB: field larvae that were reared for four generations in the laboratory and did not receive antibiotics treatment.

closest hits for OTU #2 were *Enterococcus faecium* (HG937762), *E. gallinarum* (HG937758) and *E. casseliflavus* (KJ528946) with 99% similarity. The larvae fed on cotton also contained OTU #2, but in much lower abundance, i.e., 31.8%, while an OTU of the family Acetobacteraceae (#6) had the highest abundance (67.1%). The closest hit for this OTU was *Asaia lannensis* (KP208318), with 99% similarity.

In the field larvae, we also found differences in bacterial communities depending on the host plant species they fed on. Enterobacteriaceae were the dominant family in the larvae that fed on chickpea and also occurred in larvae that fed on tobacco, while these were rare (< 1%) in the larvae fed on cotton. The larvae fed on tobacco and cotton contained Burkholderiales (Comamonadaceae, e.g., #18, and Oxalobacteraceae), which were rare (< 1%) in larvae fed on chickpea (FIGURE 2.2). Methylobacteriaceae (e.g., #11) were among the most abundant families in the larvae that fed on cotton, while this family was hardly detected in the other two groups. On the genus (and OTU) level, the two biological replicates from each plant group differed greatly from each other (FIGURE S2.1).


### ***Variability of bacterial communities across different life stages of the field population***

We assessed the temporal persistence of bacterial communities in *H. virescens* by comparing bacterial communities of different life stages of the field population (larvae, adults and eggs). Bacterial communities differed greatly across life stages. Females shared 76% of their OTUs (n = 200) with eggs, and eggs shared 52.1% of their OTUs (n = 292) with females. In contrast, larvae only shared 2.7% of their OTUs (n = 148) with females and 6.1% OTUs with eggs. Accordingly, in a PCoA plot, females and eggs partly clustered together, while five of the six larval samples were located in a different area of the plot (FIGURE 2.4). Females of the field population shared 22% of their OTUs (n = 200) with the field-lab larvae that were reared in the laboratory for four generations (the NoAB group), while the eggs had 18.8% of their OTUs (n = 292) in common with the field-lab larvae.

One OTU (#1) from the family Enterobacteriaceae had the highest relative abundance in both females and eggs (FIGURE 2.5). The closest hits for this OTU in NCBI were *Enterobacter ludwigii* (KJ767368), *E. cloacae* (KM817773) and *Klebsiella oxytoca* (KM408615), all with 97% similarity; this was confirmed by its position in the phylogeny (FIGURE 2.1). In the field larvae, OTU #1 was not present, but two other Enterobacteriaceae OTUs were found (#7 and #12), which clustered with OTU #1 in the phylogenetic tree (FIGURES 2.1 and 2.2).

### ***Vertical transmission of bacterial communities***

Based on weighted UniFrac distance values, bacterial communities present in females were not more similar to the bacterial communities of their own eggs than to the bacterial communities of eggs that were not their own ( $F_{1,65.74} = 1.28$ ,  $P = 0.26$ ). However, rearing on different plant species had a significant effect on bacterial com-



**FIGURE 2.4.** Principal coordinates analysis plot of bacterial communities of field larvae, female adults and their eggs. Absence of confidence ellipsoids around the samples indicates good jackknife support for the location of data points in the PCoA plots. Colours indicate food plant; symbols denote developmental stages; lines connect female-egg combinations.


munities in female-egg combinations ( $F_{5,66.53} = 3.17$ ,  $P = 0.013$ ). None of the comparisons for the variable plant species were significant in planned pairwise comparisons with Bonferroni corrections. However, cotton-reared females tended to be more similar to bacterial communities of eggs of their own plant group than to those of a different plant group ( $z = -2.31$ ,  $P = 0.062$ ). This marginal effect was not found for tobacco females and eggs ( $z = 1.46$ ,  $P = 0.43$ ) and was even reversed – bacterial communities of females were more similar to bacterial communities of other eggs than to eggs of their own plant group – for chickpea-reared females ( $z = -2.33$ ,  $P = 0.060$ ). The principal coordinates analysis plot supports these results (see FIGURE 2.4).

## DISCUSSION

In this study we assessed the variability of bacterial communities in the agricultural pest *H. virescens*. We observed that the bacterial communities of different life stages, of different populations (field population or laboratory colony) as well as of larvae that fed on different host plant species, and even biological replicates, were very different.

### *Abundant bacterial strains in Heliothis virescens*

Most of the dominant bacterial families in our study, i.e., Acetobacteraceae, Methylobacteriaceae, Sphingomonadaceae, Comamonadaceae, Enterobacteriaceae, Pseudomonadaceae, Moraxellaceae Xanthomonadaceae, Paenibacillaceae and Enterococcaceae, have been identified in Lepidoptera before (Broderick *et al.*, 2004;



**FIGURE 2.5.** Relative abundance of bacterial OTUs in the abdomen of *Heliothis virescens* females (F) and their eggs (E). Females (F1-F9) and their eggs (E1-E9) are shown next to each other. Females had fed on three different plants as larvae: cotton (C), chickpea (Ch), tobacco (T). Rare OTUs that were represented less than 0.5% in at least one sample are not shown. Brackets around genera signify that there were bacterial candidates with  $\geq 97\%$  identity, but support in the phylogenetic tree was weak [i.e., nodes that divide branches that contain different genera have bootstrap values equal to or below 10 (see FIGURE 2.1)].

Xiang *et al.*, 2006; Robinson *et al.*, 2010; Belda *et al.*, 2011; Pinto-Tomás *et al.*, 2011; Gayatri Priya *et al.*, 2012; Tang *et al.*, 2012; Hammer *et al.*, 2014). This suggests that representatives of these bacterial families may be commonly present in many Lepidoptera and the environments they live in.

### ***Variability of bacterial communities between laboratory and field larvae***

Bacterial communities differed greatly between laboratory and field larvae, in fact no OTUs were shared between the two groups. These large differences in bacterial communities underline the importance of including field populations of an organism in studies of the functional role of its bacterial communities. Bacterial communities of field and laboratory insect populations have for instance been studied in the moths *Helicoverpa armigera* (Xiang *et al.*, 2006) and *Ostrinia nubilalis* (Belda *et al.*,

2011), the mosquito *Anopheles stephensi* (Rani *et al.*, 2009) and several fruitfly species (*Drosophila*) (Chandler *et al.*, 2011). In all these studies, the general finding was that bacterial communities in laboratory colonies are depauperate and dominated by only one (or few) bacterial strain(s), as we also found in *H. virescens*, while communities of field populations tend to be more diverse and contain many bacterial strains with lower and more equal abundances (Xiang *et al.*, 2006; Rani *et al.*, 2009; Belda *et al.*, 2011; Chandler *et al.*, 2011). Differences between laboratory and field populations may be due to differences in environmental conditions. In the field, organisms are probably exposed to a higher diversity of bacterial strains which may vary in space and time, than in the laboratory. For instance, in *H. virescens*, and most moth species, eggs are laid on above-ground parts of plants, where larvae also feed, while the moths pupate in the soil. Emerging adults feed from flowers of many plants and, in the case of the highly polyphagous *H. virescens*, many plant species (Barber, 1937; Stadelbacher, 1981; Fitt, 1989; Waldvogel & Gould, 1990). In contrast, in the laboratory moths are reared in cups of artificial diet, where they also pupate and emerge as adults. Thus, whereas moths in the field may continuously encounter and ingest new strains of bacteria, lab colonies are likely constantly re-infected with the same (small number of) bacterial strains that are dominant under laboratory conditions.

As described in the methods, we used whole (washed) larvae to assess the bacterial communities in the laboratory strain and extracted the guts of larvae to assess bacterial communities in field larvae. Also, laboratory strain larvae were in an earlier instar than field larvae. Even though the differences in treatment and instar might be responsible for part of the differences between field and laboratory larvae in our study, they are unlikely to explain the large qualitative differences with a complete lack of overlap in bacterial communities. Moreover, *Enterococcus* sp. which was the dominant strain in the laboratory larvae was also detected in the field-lab larvae (5th instar), which were kept in the laboratory for four generations and from which only the gut was investigated. Thus the detection of enterococci in laboratory larvae is unlikely to be merely due to methodological differences between laboratory-reared and field larvae.

In our study, *Enterococcus* sp. was the dominant bacterial strain in the laboratory colony and the AB group of the field-lab larvae, and it was the second most abundant strain in the NoAB group of the field-lab larvae. However, this strain was completely absent in the field larvae, suggesting that *Enterococcus* sp. was introduced to the insects in the laboratory. *Enterococcus* sp. has been identified as a dominant microbe in several Lepidopteran gut communities, including *Galleria mellonella*, *Lymantria dispar*, *Helicoverpa armigera*, *Manduca sexta*, *Spodoptera littoralis* and *Heliconius erato* (Jarosz, 1975; Broderick *et al.*, 2004; Xiang *et al.*, 2006; Brinkmann *et al.*, 2008; Tang *et al.*, 2012; Hammer *et al.*, 2014). Interestingly, most of these studies used only laboratory colonies (Jarosz, 1975; Brinkmann *et al.*, 2008; Tang *et al.*, 2012). In *Helicoverpa armigera* and *Ostrinia nubilalis*, where both laboratory and

field populations were investigated (Xiang *et al.*, 2006; Belda *et al.*, 2011), single bacterial strains such as *Enterococcus* sp. and other gram-positive cocci (e.g., *Staphylococcus* sp. in *O. nubilalis*) were found to become dominant in laboratory populations, while they were less abundant in field populations. In *H. erato*, enterococci became more abundant in adults after this species had been in captivity for one generation (from adult to adult) compared to field-collected adults (Hammer *et al.*, 2014). Possibly, enterococci are outcompeted in the field by other bacteria (e.g., Proteobacteria such as *Enterobacter*, *Klebsiella* and *Pseudomonas* species) in the insect gut and/or on the plants while competition changes in favour of enterococci under (more stable) laboratory conditions.

Treating *H. virescens* larvae with tetracycline in the laboratory probably selected for enterococci, as the relative abundance of enterococci was higher in larvae treated with antibiotics than in untreated larvae. This is in line with the finding that many *Enterococcus* strains have developed resistance against tetracycline (Huys *et al.*, 2004; Wilcks *et al.*, 2005; Cauwerts *et al.*, 2007). It is interesting to note, however, that enterococci also became abundant in laboratory-reared larvae in the absence of antibiotics, indicating that antibiotic treatment alone does not account for the shift in microbial communities towards high abundance of *Enterococcus* spp.

#### ***Host plant-associated variability of bacterial communities in larvae***

In both field and laboratory *H. virescens* larvae, bacterial gut communities varied depending on host plant. Diet has been found to influence bacterial gut community composition in larvae of many insect species, e.g., *Drosophila* (Chandler *et al.*, 2011) and several lepidopteran species (Broderick *et al.*, 2004; Gayatri Priya *et al.*, 2012; Tang *et al.*, 2012). Diet might influence bacterial communities in various ways. For example, different bacteria may be present in or on different host plants, the resources provided by the different plants may promote differential bacterial growth and/or secondary plant metabolites may have a selective effect on bacterial communities (Yang *et al.*, 2001; Vorholt, 2012). Additionally, host plants can influence the physiochemical conditions of the larval gut, which possibly results in differential bacterial growth (Schultz & Lechowicz, 1986; Appel & Maines, 1995).

Although we did not sample bacteria from plants on which the larvae had fed, earlier studies have assessed bacterial communities of cotton and tobacco plants, and identified several bacterial families that we also detected in our samples (McInroy & Klopper, 1995; Brinkmann *et al.*, 2008), suggesting that the experimental *H. virescens* larvae had ingested bacteria from their food plants. For instance, Methylobacteriaceae was the most abundant bacterial family in the field larvae fed on cotton in our study, and this bacterial family has been isolated from different tissues of cotton (McInroy & Klopper, 1995; Madhaiyan *et al.*, 2012). Also, bacteria from the order Burkholderiales (Comamonadaceae and Oxalobacteraceae), which were mainly present in the field larvae fed on tobacco in our study, were previously isolated from

tobacco leaves (Brinkmann *et al.*, 2008). Furthermore, *Enterobacter*, *Pseudomonas* and *Serratia* were the most abundant genera in females and eggs in our study, and have been isolated from, among others, cotton tissues and tobacco leaves (McInroy & Klopper, 1995; Junker *et al.*, 2011; Lv *et al.*, 2012; Ma *et al.*, 2013).

Interestingly, in *L. dispar*, bacteria that were isolated from leaves of its host plant aspen, have recently been shown to enhance larval growth of this moth in the presence of phenolic glycosides, a defence metabolite of aspen (Mason *et al.*, 2014). Whether *H. virescens* can profit from bacteria that are transient or reside on its host plants still needs to be investigated.

Notably, there were differences in bacterial community composition between the two biological replicates per plant species in the field larvae, particularly at low taxonomic levels (i.e., genus or OTU). This indicates large individual variation in the bacterial gut community of *H. virescens* larvae in the field, and further suggests that the bacterial community is only loosely associated with this moth.

When investigating microbial communities using bacterial tag-encoded FLX pyrosequencing of 16S rRNA amplicons, it is important to keep in mind that this method can lead to incorrect estimates of OTU numbers (Kunin *et al.*, 2010; Schloss *et al.*, 2011; Pinto & Raskin, 2012). Errors that have been described for this method include PCR bias, sequencing errors (particularly in homopolymeric regions) and the production of chimeric sequences during PCR reactions (Chandler *et al.*, 1997; Margulies *et al.*, 2005; Haas *et al.*, 2011; Schloss *et al.*, 2011). Even though we have implemented in our analyses a number of measures to decrease methodological errors (de-noising raw data, quality cut-off, length limitation of sequences, identification and removal of possible chimeric sequences, removal of global singletons), it is possible that the number of OTUs may still be overestimated and that the diversity patterns that we show on the family level for larvae is actually more realistic than the patterns we encountered on the OTU level (compare FIGURES 2.2 and S2.1).

### ***Variability of bacterial communities across different life stages of the field population***

In *H. virescens*, it seems that bacterial communities are completely restructured or lost during the metamorphosis from crawling larvae to winged adult, because bacterial communities in larvae of the field population differed strongly from those in adult females. One of the few Lepidoptera where bacterial persistence across metamorphosis has been studied is the butterfly *H. erato* (Hammer *et al.*, 2014). In this species, bacterial communities between larvae and adults also differed greatly but more bacterial strains (13%) overlapped between larvae and adults in their study than in our study, in which only 2.7% of OTUs overlapped between larvae and adult females. A similar pattern was found in mosquitoes, in which bacterial composition partly changed during metamorphosis, but many strains were retained (Rani *et al.*, 2009). One reason for the difference in bacterial communities between larvae and adults in Lepidoptera might be that larvae and adults feed on different diets: while

larvae usually feed on the foliage of their host plants, adults often feed only on nectar and in this study on honey water. The gut of lepidopteran larvae is less compartmentalized than the gut of many other insect species, mostly has a very alkaline pH, and is purged before metamorphosis (Nijhout & Williams, 1974; Dow, 1992; Appel & Maines, 1995; Chapman *et al.*, 2013; Engel & Moran, 2013). Together, these characteristics might make it difficult for bacteria to successfully colonize Lepidopteran guts. Moreover, in moths an increase in lysozyme production before and during metamorphosis has been reported, which likely reduces bacterial abundance and diversity especially immediately before and after the pupal stage, i.e., pupating larvae and newly eclosed adults (Russell & Dunn, 1996; Zhang *et al.*, 2009). Upon adult eclosion, bacterial titers may therefore be low and bacterial strains that are different from the larval bacterial community might be able to colonize the adults (see also Hammer *et al.*, 2014).

### ***Vertical transmission of bacterial communities***

We investigated whether bacteria in *H. virescens* are vertically transmitted by comparing bacterial communities of females and their eggs and tested if bacterial communities of eggs were more similar to bacterial communities of their own mother than to any other female. The fact that we did not find this effect suggests that vertical transmission likely does not occur in *H. virescens* at a significant rate. However, since bacterial communities were at least partly similar in females and eggs, we cannot completely exclude the occurrence of vertical transmission. A study using fluorescently-labelled *E. coli* was able to detect a certain amount of mother-offspring transmission in *Galleria mellonella*, although the viability of the bacteria was not assessed (Freitak *et al.*, 2014). Also, a previous study about vertical transmission of *Serratia marcescens* in a laboratory population of *H. virescens* indicated that bacteria can be vertically transmitted in this moth (Sikorowski & Lawrence, 1998).

### ***Conclusions***

The huge variability of bacterial communities that we found between life stages, diets, biological replicates and field and lab populations indicates that the major part of the bacterial communities that we identified in the gut of *H. virescens* is of a transient nature and only loosely associated with its host. In fact, the bacterial communities seem to be entirely restructured during metamorphosis. Our results further suggest that bacterial communities are not transmitted at a significant rate from mothers to eggs in *H. virescens*. Based on these results, it is doubtful that particular bacterial strains that we have identified form a unit of selection with this moth. It is further unlikely that a long-term mutualistic symbiosis between *H. virescens* and bacteria that could facilitate host plant use and adaptation has evolved. Importantly, we found that enterococci are most probably introduced to *H. virescens* larvae in the laboratory. This finding stresses the importance of including field populations when bacterial communities of an organism are to be characterized.

### **Acknowledgements**

We thank Fred Gould and Ed Vargo for the possibility to work in their laboratory in North Carolina and Fred Gould and Clyde Soerenson for their help in collecting *H. virescens* in the field. We thank Pascaline Dumas for her support with phylogenetic analyses and Dennis van Veldhuizen for rearing *H. virescens*. This research was supported by the University of Amsterdam, the Max-Planck-Gesellschaft and was partly funded by the National Science Foundation (awards IOS-1052238 and IOS-1456973).

### **REFERENCES**

- Adams, A.S., Adams, S.M., Currie, C.R., Gillette, N.E. & Raffa, K.F. 2010. Geographic variation in bacterial communities associated with the red turpentine beetle (Coleoptera: Curculionidae). *Environ Entomol* **39**: 406-414.
- Appel, H. & Maines, L. 1995. The influence of host plant on gut conditions of gypsy moth (*Lymantria dispar*) caterpillars. *J Insect Physiol* **41**: 241-246.
- Barber, G.W. 1937. Seasonal availability of food plants of two species of *Heliothis* in eastern Georgia. *J Econ Entomol* **30**: 150-158.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. 2014. lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1-7.
- Belda, E., Pedrola, L., Peretó, J., Martínez-Blanch, J.F., Montagud, A., Navarro, E., Urchueguía, J., Ramón, D., Moya, A. & Porcar, M. 2011. Microbial diversity in the midguts of field and lab-reared populations of the European corn borer *Ostrinia nubilalis*. *PLoS ONE* **6**: e21751.
- Benson, D.A., Karsch-Mizrachi, I., Clark, K., Lipman, D.J., Ostell, J. & Sayers, E.W. 2011. GenBank. *Nucleic Acids Res.*
- Blanco, C.A., Terán-Vargas, A.P., López Jr, J.D., Kauffman, J.V. & Wei, X. 2007. Densities of *Heliothis virescens* and *Helicoverpa zea* (Lepidoptera: Noctuidae) in three plant hosts. *Fla Entomol* **90**: 742-750.
- Bosch, T.C.G. & McFall-Ngai, M.J. 2011. Metaorganisms as the new frontier. *Zoology* **114**: 185-190.
- Brinkmann, N., Martens, R. & Tebbe, C.C. 2008. Origin and diversity of metabolically active gut bacteria from laboratory-bred larvae of *Manduca sexta* (Sphingidae, Lepidoptera, Insecta). *Appl Environ Microbiol* **74**: 7189-7196.
- Broderick, N.A., Raffa, K.F., Goodman, R.M. & Handelsman, J. 2004. Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. *Appl Environ Microbiol* **70**: 293-300.
- Broderick, N.A., Raffa, K.F. & Handelsman, J. 2006. Midgut bacteria required for *Bacillus thuringiensis* insecticidal activity. *Proc Natl Acad Sci USA* **103**: 15196-15199.
- Broderick, N.A., Robinson, C.J., McMahon, M.D., Holt, J., Handelsman, J. & Raffa, K.F. 2009. Contributions of gut bacteria to *Bacillus thuringiensis*-induced mortality vary across a range of Lepidoptera. *BMC Biol* **7**.
- Burton, R.L. 1970. A low-cost artificial diet for the corn earworm. *J Econ Entomol* **63**: 1969-1970.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Pena, A.G., Goodrich, J.K. & Gordon, J.I. 2010a. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* **7**: 335-336.
- Caporaso, J.G., Bittinger, K., Bushman, F.D., DeSantis, T.Z., Andersen, G.L. & Knight, R. 2010b. PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* **26**: 266-267.


- Cauwerts, K., Decostere, A., De Graef, E.M., Haesebrouck, F. & Pasmans, F. 2007. High prevalence of tetracycline resistance in *Enterococcus* isolates from broilers carrying the erm(B) gene. *Avian Pathol* **36**: 395-399.
- Chandler, D., Fredrickson, J. & Brockman, F. 1997. Effect of PCR template concentration on the composition and distribution of total community 16S rDNA clone libraries. *Mol Ecol* **6**: 475-482.
- Chandler, J.A., Lang, J.M., Bhatnagar, S., Eisen, J.A. & Kopp, A. 2011. Bacterial communities of diverse *Drosophila* species: ecological context of a host-microbe model system. *PLoS Genet* **7**: e1002272.
- Chapman, R.F., Simpson, S.J. & Douglas, A.E. 2013. *The insects: structure and function*, 5th edn. Cambridge University Press, Cambridge.
- Chu, C.-C., Spencer, J.L., Curzi, M.J., Zavala, J.A. & Seufferheld, M.J. 2013. Gut bacteria facilitate adaptation to crop rotation in the western corn rootworm. *Proc Natl Acad Sci USA* **110**: 11917-11922.
- Colman, D.R., Toolson, E.C. & Takacs-Vesbach, C.D. 2012. Do diet and taxonomy influence insect gut bacterial communities? *Mol Ecol* **21**: 5124-5137.
- Dillon, R. & Dillon, V. 2004. The gut bacteria of insects: nonpathogenic interactions. *Annu Rev Entomol* **49**: 71-92.
- Douglas, A.E. 2009. The microbial dimension in insect nutritional ecology. *Funct Ecol* **23**: 38-47.
- Dow, J. 1992. pH gradients in Lepidopteran midgut. *J Exp Biol* **172**: 355-375.
- Edgar, R.C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**: 2460-2461.
- Engel, P. & Moran, N.A. 2013. The gut microbiota of insects – diversity in structure and function. *FEMS Microbiol Rev* **37**: 699-735.
- Feldhaar, H. 2011. Bacterial symbionts as mediators of ecologically important traits of insect hosts. *Ecol Entomol* **36**: 533-543.
- Fitt, G.P. 1989. The ecology of *Heliothis* species in relation to agroecosystems. *Annu Rev Entomol* **34**: 17-53.
- Flint, H.J., Duncan, S.H., Scott, K.P. & Louis, P. 2007. Interactions and competition within the microbial community of the human colon: links between diet and health. *Environ Microbiol* **9**: 1101-1111.
- Freitag, D., Schmidtberg, H., Dickel, F., Lochnit, G., Vogel, H. & Vilcinskas, A. 2014. The maternal transfer of bacteria can mediate trans-generational immune priming in insects. *Virulence* **5**: 547-554.
- Gayatri Priya, N., Ojha, A., Kajla, M.K., Raj, A. & Rajagopal, R. 2012. Host plant induced variation in gut bacteria of *Helicoverpa armigera*. *PLoS ONE* **7**: e30768.
- González-Braojos, S., Vela, A.I., Ruiz-de-Castañeda, R., Briones, V. & Moreno, J. 2012. Age-related changes in abundance of enterococci and Enterobacteriaceae in Pied Flycatcher (*Ficedula hypoleuca*) nestlings and their association with growth. *J Ornithol* **153**: 181-188.
- Haas, B.J., Gevers, D., Earl, A.M., Feldgarden, M., Ward, D.V., Giannoukos, G., Ciulla, D., Tabbaa, D., Highlander, S.K., Sodergren, E., Methé, B., DeSantis, T.Z., Consortium, T.H.M., Petrosino, J.F., Knight, R. & Birren, B.W. 2011. Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons. *Genome Res* **21**: 494-504.
- Hall, E.K., Neuhauser, C. & Cotner, J.B. 2008. Toward a mechanistic understanding of how natural bacterial communities respond to changes in temperature in aquatic ecosystems. *ISME J* **2**: 471-481.
- Hammer, T.J., McMillan, W.O. & Fierer, N. 2014. Metamorphosis of a butterfly-associated bacterial community. *PLoS ONE* **9**: e86995.
- Hosokawa, T., Kikuchi, Y., Shimada, M. & Fukatsu, T. 2007. Obligate symbiont involved in pest status of host insect. *Proc R Soc Lond B Biol Sci* **274**: 1979-1984.

- Hothorn, T., Bretz, F. & Westfall, P. 2008. Simultaneous inference in general parametric models. *Biometrical J* **50**: 346-363.
- Huys, G., D'Haene, K., Collard, J.-M. & Swings, J. 2004. Prevalence and molecular characterization of tetracycline resistance in *Enterococcus* isolates from food. *Appl Environ Microbiol* **70**: 1555-1562.
- Ishak, H., Plowes, R., Sen, R., Kellner, K., Meyer, E., Estrada, D., Dowd, S. & Mueller, U. 2011. Bacterial diversity in *Solenopsis invicta* and *Solenopsis geminata* ant colonies characterized by 16S amplicon 454 pyrosequencing. *Microb Ecol* **61**: 821-831.
- Jarosz, J. 1975. Lysozymelike lytic enzyme of *Streptococcus faecalis* and its role in the larval development of wax moth, *Galleria mellonella*. *J Invertebr Pathol* **26**: 275-281.
- Junker, R.R., Loewel, C., Gross, R., Dötterl, S., Keller, A. & Blüthgen, N. 2011. Composition of epiphytic bacterial communities differs on petals and leaves. *Plant Biol* **13**: 918-924.
- Kunin, V., Engelbrekton, A., Ochman, H. & Hugenholtz, P. 2010. Wrinkles in the rare biosphere: pyrosequencing errors can lead to artificial inflation of diversity estimates. *Environ Microbiol* **12**: 118-123.
- Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. 2014. lmerTest: Tests for random and fixed effects for linear mixed effect models (lmer objects of lme4 package). *R package version*: 2.0-11.
- Loman, N.J., Misra, R.V., Dallman, T.J., Constantinidou, C., Gharbia, S.E., Wain, J. & Pallen, M.J. 2012. Performance comparison of benchtop high-throughput sequencing platforms. *Nat Biotech* **30**: 434-439.
- Lozupone, C. & Knight, R. 2005. UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microbiol* **71**: 8228-8235.
- Lozupone, C., Lladser, M. E., Knights, D., Stombaugh, J., & Knight, R. 2011. UniFrac: an effective distance metric for microbial community comparison. *ISME J* **5**: 169.
- Lv, D., Ma, A., Bai, Z., Zhuang, X. & Zhuang, G. 2012. Response of leaf-associated bacterial communities to primary acyl-homoserine lactone in the tobacco phyllosphere. *Res Microbiol* **163**: 119-124.
- Ma, A., Lv, D., Zhuang, X. & Zhuang, G. 2013. Quorum quenching in culturable phyllosphere bacteria from tobacco. *Int J Mol Sci* **14**: 14607-14619.
- Madhaiyan, M., Poonguzhali, S., Senthilkumar, M., Lee, J.-S. & Lee, K.-C. 2012. *Methylobacterium gossypicola* sp. nov., a pink-pigmented, facultatively methylotrophic bacterium isolated from the cotton phyllosphere. *Int J Syst Evol Microbiol* **62**: 162-167.
- Margulies, M., Egholm, M., Altman, W.E., Attiya, S., Bader, J.S., Bemben, L.A., Berka, J., Braverman, M.S., Chen, Y.-J., Chen, Z., Dewell, S.B., Du, L., Fierro, J.M., Gomes, X.V., Godwin, B.C., He, W., Helgesen, S., Ho, C.H., Irzyk, G.P., Jando, S.C., Alenquer, M.L.I., Jarvie, T.P., Jirage, K.B., Kim, J.-B., Knight, J.R., Lanza, J.R., Leamon, J.H., Lefkowitz, S.M., Lei, M., Li, J., Lohman, K.L., Lu, H., Makhijani, V.B., McDade, K.E., McKenna, M.P., Myers, E.W., Nickerson, E., Nobile, J.R., Plant, R., Puc, B.P., Ronan, M.T., Roth, G.T., Sarkis, G.J., Simons, J.F., Simpson, J.W., Srinivasan, M., Tartaro, K.R., Tomasz, A., Vogt, K.A., Volkmer, G.A., Wang, S.H., Wang, Y., Weiner, M.P., Yu, P., Begley, R.F. & Rothberg, J.M. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* **437**: 376-380.
- Mason, C.J., Couture, J.J. & Raffa, K.F. 2014. Plant-associated bacteria degrade defense chemicals and reduce their adverse effects on an insect defoliator. *Oecologia* **175**: 901-910.
- McInroy, J. & Kloepper, J. 1995. Survey of indigenous bacterial endophytes from cotton and sweet corn. *Plant Soil* **173**: 337-342.
- Neunzig, H.H. 1969. Biology of the tobacco budworm and the corn earworm in North Carolina. *North Carolina Exp Station Tech Bull*.
- Nijhout, H.F. & Williams, C.M. 1974. Control of moulting and metamorphosis in the tobacco hornworm, *Manduca sexta* (L.): cessation of juvenile hormone secretion as a trigger for pupation. *J Exp Biol* **61**: 493-501.

- Oliver, K.M., Degnan, P.H., Burke, G.R. & Moran, N.A. 2010. Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. *Annu Rev Entomol* **55**: 247-266.
- Pinto-Tomas, A., Uribe-Lorio, L., Blanco, J., Fontecha, G., Rodríguez, C., Mora, M., Janzen, D., Chavarría, F., Díaz, J. & Sittenfeld, A. 2007. Enzymatic activities of bacteria isolated from the digestive tract of caterpillars and the pupal content of *Automeris zugana* and *Rothschildia lebeau* (Lepidoptera: Saturniidae). *Rev Biol Trop* **55**: 401-415.
- Pinto-Tomás, A.A., Sittenfeld, A., Uribe-Lorio, L., Chavarría, F., Mora, M., Janzen, D.H., Goodman, R.M. & Simon, H.M. 2011. Comparison of midgut bacterial diversity in tropical caterpillars (Lepidoptera: Saturniidae) fed on different diets. *Environ Entomol* **40**: 1111-1122.
- Pinto, A.J. & Raskin, L. 2012. PCR biases distort bacterial and archaeal community structure in pyrosequencing datasets. *PLoS ONE* **7**: e43093.
- Prem Anand, A.A., Vennison, S.J., Sankar, S.G., Gilwax Prabhu, D.I., Vasan, P.T., Raghuraman, T., Jerome Geoffrey, C. & Vendan, S.E. 2010. Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin and starch and their impact on digestion. *J Insect Sci* **10**.
- Price, M.N., Dehal, P.S. & Arkin, A.P. 2010. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS ONE* **5**: e9490.
- Rani, A., Sharma, A., Rajagopal, R., Adak, T. & Bhatnagar, R. 2009. Bacterial diversity analysis of larvae and adult midgut microflora using culture-dependent and culture-independent methods in lab-reared and field-collected *Anopheles stephensi*-an Asian malarial vector. *BMC Microbiol* **9**: 96.
- R Core Team 2013. R: A language and environment for statistical computing., R Foundation for Statistical Computing, Vienna, Austria.
- Reeder, J. & Knight, R. 2010. Rapidly denoising pyrosequencing amplicon reads by exploiting rank-abundance distributions. *Nat Meth* **7**: 668-669.
- Robinson, C., Schloss, P., Ramos, Y., Raffa, K. & Handelsman, J. 2010. Robustness of the bacterial community in the cabbage white butterfly larval midgut. *Microb Ecol* **59**: 199-211.
- Rosenberg, E., Sharon, G. & Zilber-Rosenberg, I. 2009. The hologenome theory of evolution contains Lamarckian aspects within a Darwinian framework. *Environ Microbiol* **11**: 2959-2962.
- Rothberg, J.M. & Leamon, J.H. 2008. The development and impact of 454 sequencing. *Nat Biotech* **26**: 1117-1124.
- Russell, J., Sharp, W. & Baldwin, R. 1979. The effect of pH on maximum bacterial growth rate and its possible role as a determinant of bacterial competition in the rumen. *J Anim Sci* **48**: 251-255.
- Russell, V. & Dunn, P.E. 1996. Antibacterial proteins in the midgut of *Manduca sexta* during metamorphosis. *J Insect Physiol* **42**: 65-71.
- Salem, H., Florez, L., Gerardo, N. & Kaltenpoth, M. 2015. An out-of-body experience: the extracellular dimension for the transmission of mutualistic bacteria in insects. *Proc R Soc B Biol Sci* **282**: 20142957.
- Schloss, P.D., Gevers, D. & Westcott, S.L. 2011. Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PLoS ONE* **6**: e27310.
- Schultz, J.C. & Lechowicz, M.J. 1986. Hostplant, larval age, and feeding behavior influence midgut pH in the gypsy moth (*Lymantria dispar*). *Oecologia* **71**: 133-137.
- Shade, A. & Handelsman, J. 2012. Beyond the Venn diagram: the hunt for a core microbiome. *Environ Microbiol* **14**: 4-12.
- Shannon, C.E. 1948. The mathematical theory of communication. *Bell Syst Tech J* **27**: 379-423.
- Sheck, A.L. & Gould, F. 1993. The genetic basis of host range in *Heliothis virescens*: larval survival and growth. *Entomol Exp Appl* **69**: 157-172.

- Shimodaira, H. & Hasegawa, M. 1999. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol Biol Evol* **16**: 1114-1116.
- Sikorowski, P. & Lawrence, A. 1998. Transmission of *Serratia marcescens* (Enterobacteriaceae) in adult *Heliothis virescens* (Lepidoptera: Noctuidae) laboratory colonies. *Biol Control* **12**: 50-55.
- Stadelbacher, E. 1981. Role of early-season wild and naturalized host plants in the buildup of the F1 generation of *Heliothis zea* and *H. virescens* in the Delta of Mississippi. *Environ Entomol* **10**: 766-770.
- Sudakaran, S., Salem, H., Kost, C. & Kaltenpoth, M. 2012. Geographical and ecological stability of the symbiotic mid-gut microbiota in European firebugs, *Pyrrhocoris apterus* (Hemiptera, Pyrrhocoridae). *Mol Ecol* **21**: 6134-6151.
- Sullam, K.E., Essinger, S.D., Lozupone, C.A., O'Connor, M.P., Rosen, G.L., Knight, R.O.B., Kilham, S.S. & Russell, J.A. 2012. Environmental and ecological factors that shape the gut bacterial communities of fish: a meta-analysis. *Mol Ecol* **21**: 3363-3378.
- Tamura, K. & Nei, M. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* **10**: 512-526.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Mol Biol Evol* **30**: 2725-2729.
- Tang, X., Freitak, D., Vogel, H., Ping, L., Shao, Y., Cordero, E.A., Andersen, G., Westermann, M., Heckel, D.G. & Boland, W. 2012. Complexity and variability of gut commensal microbiota in polyphagous Lepidopteran larvae. *PLoS ONE* **7**: e36978.
- Thompson, J.D., Higgins, D.G. & Gibson, T.J. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**: 4673-4680.
- Tsuchida, T., Koga, R., Matsumoto, S. & Fukatsu, T. 2011. Interspecific symbiont transfection confers a novel ecological trait to the recipient insect. *Biol Lett* **7**: 245-248.
- Unbehend, M., Hänniger, S., Meagher, R.L., Heckel, D.G. & Groot, A.T. 2013. Pheromonal divergence between two strains of *Spodoptera frugiperda*. *J Chem Ecol* **39**: 364-376.
- Vorholt, J.A. 2012. Microbial life in the phyllosphere. *Nat Rev Microbiol* **10**: 828-840.
- Waldvogel, M. & Gould, F. 1990. Variation in oviposition preference of *Heliothis virescens* in relation to macroevolutionary patterns of heliothine host range. *Evolution*: 1326-1337.
- Wilcks, A., Andersen, S.R. & Licht, T.R. 2005. Characterization of transferable tetracycline resistance genes in *Enterococcus faecalis* isolated from raw food. *FEMS Microbiol Lett* **243**: 15-19.
- Xia, X., Zheng, D., Zhong, H., Qin, B., Gurr, G.M., Vasseur, L., Lin, H., Bai, J., He, W. & You, M. 2013. DNA sequencing reveals the midgut microbiota of diamondback moth, *Plutella xylostella* (L.) and a possible relationship with insecticide resistance. *PLoS ONE* **8**: e68852.
- Xiang, H., Wei, G.-F., Jia, S., Huang, J., Miao, X.-X., Zhou, Z., Zhao, L.-P. & Huang, Y.-P. 2006. Microbial communities in the larval midgut of laboratory and field populations of cotton bollworm (*Helicoverpa armigera*). *Can J Microbiol* **52**: 1085-1092.
- Yang, C.-H., Crowley, D.E., Borneman, J. & Keen, N.T. 2001. Microbial phyllosphere populations are more complex than previously realized. *Proc Natl Acad Sci USA* **98**: 3889-3894.
- Zhang, Y., Huang, J., Zhou, B., Zhang, C., Liu, W., Miao, X. & Huang, Y. 2009. Up-regulation of lysozyme gene expression during metamorphosis and immune challenge of the cotton bollworm, *Helicoverpa armigera*. *Arch Insect Biochem Physiol* **70**: 18-29.
- Zilber-Rosenberg, I. & Rosenberg, E. 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* **32**: 723-735.

## SUPPORTING INFORMATION



**FIGURE S2.1.** Relative abundance of bacterial OTUs in *Heliothis virescens* larvae. OTUs are identified at the genus level. Rare OTUs that were represented less than 0.5% in at least one sample are not included. Larvae originated from a laboratory colony, from the field or were collected from the field as eggs, after which they were reared in the laboratory for four generations (field-lab): AB: larvae received antibiotics treatment, NoAB: larvae received no antibiotics treatment. Field and laboratory larvae were grown on cotton (C), chickpea (Ch) or tobacco (T). Brackets around genera signify that there were bacterial candidates with  $\geq 97\%$  identity, but support in the phylogenetic tree was weak [i.e., nodes that divide branches that contain different genera have bootstrap values equal to or below 10 (see FIGURE 2.1)].

## Bacterial community in *Heliothis virescens*

**TABLE S2.1.** Statistics of bacterial tag-encoded FLX amplicon sequencing and number of OTUs in *Heliothis virescens* larvae. The bacterial communities of field larvae (F), laboratory larvae (L) and field-lab (AB/NoAB) larvae were characterized. The latter was reared in the laboratory for four generations after collection in the field. AB: treated with antibiotics (tetracycline); NoAB: not treated with antibiotics. Larvae of the field and laboratory strain were fed on three different plant species: cotton (C), chickpea (Ch), and tobacco (T). Field larvae: two sequencing pools per plant species; laboratory and field-lab larvae: one pool per plant species; # = number, Qual. seqs = quality filtered sequences.

Sample	Laboratory larvae			Field larvae						Field-lab larvae	
	CL	ChL	TL	CF1	CF2	ChF1	ChF2	TF1	TF2	AB	NoAB
# Reads	30475	6917	5614	33880	2226	2590	2835	5356	3189	11464	10344
# Qual. seqs.	30400	6750	5112	20963	1520	2318	2791	5333	3189	11443	10341
# OTUs	27	21	27	55	23	55	53	23	16	64	90

**TABLE S2.2.** Statistics of bacterial tag-encoded FLX amplicon sequencing and number of OTUs in individual *Heliothis virescens* females. Females fed on three plant species as larvae: cotton (Cot), chickpea (Chi), tobacco (Tob); # = number, Qual. seqs = quality filtered sequences.

Sample	Cot-fem1	Cot-fem2	Cot-fem3	Chi-fem4	Chi-fem5	Chi-fem6	Tob-fem7	Tob-fem8	Tob-fem9
# Reads	7586	16398	7374	15853	13047	9509	13254	2895	16990
# Qual. seqs.	7586	16396	7373	15852	13047	9507	13156	2895	16990
# OTUs	33	44	34	65	72	65	54	38	57

**TABLE S2.3.** Statistics of bacterial tag-encoded FLX amplicon sequencing and number of OTUs in *Heliothis virescens* eggs. Mothers of these eggs had fed on three different plant species as larvae: cotton (Cot), chickpea (Chi), tobacco (Tob); # = number, Qual. seqs = quality filtered sequences.

Sample	Cot-egg1	Cot-egg2	Cot-egg3	Chi-egg4	Chi-egg5	Chi-egg6	Tob-egg7	Tob-egg8	Tob-egg9
# Reads	1346	18062	8461	1724	8033	466	27786	8890	19760
# Qual. seqs.	1241	17888	8384	503	7587	339	27784	8855	19746
# OTUs	37	108	81	48	134	30	82	70	124

## Sex-specific trade-offs between immune response and reproduction in a moth

Andrea Barthel\*, Heike Staudacher\*, Antje Schmaltz,  
David G. Heckel & Astrid T. Groot

*\*Contributed equally to this study*

*BMC Evolutionary Biology 15: 282 (2015)*

### **Abstract**

Immune response induction benefits insects in combatting infection by pathogens. However, organisms have a limited amount of resources available and face the dilemma of partitioning resources between immunity and other life-history traits. Since males and females differ in their life histories, sex-specific resource investment strategies to achieve an optimal immune response following an infection can be expected. We investigated immune response induction of females and males of *Heliothis virescens* in response to the entomopathogenic bacterium *Serratia entomophila*, and its effects on mating success and the female sexual signal. We found that females had higher expression levels of immune-related genes after bacterial challenge than males. However, males maintained a higher baseline expression of immune-related genes than females. The increased investment in immunity of female moths was negatively correlated with mating success and the female sexual signal. Male mating success was unaffected by bacterial challenge. Our results show that the sexes differed in their investment strategies: females invested in immune defense after a bacterial challenge, indicating facultative immune deployment, whereas males had higher baseline immunity than females, indicating immune maintenance. Interestingly, these differences in investment were reflected in the mate choice assays. As female moths are the sexual signallers, females need to invest resources in their attractiveness. However, female moths appeared to invest in immunity at the cost of reproductive effort.

### **INTRODUCTION**

In ecological immunology, mounting a robust immune response against pathogens is considered a necessary but costly trait in terms of energy and resources. The effectiveness of the immune response thus depends on access to limited resources and on investment required by other costly life history traits, such as reproduction (Zuk, 1990; Sheldon & Verhulst, 1996; Zuk & Stoehr, 2002; Bonneaud *et al.*, 2003; Jacobs & Zuk, 2011). Many studies on the relationship between immunity and reproduction have attempted to link variation in immune responses with variation in reproductive success (Siva-Jothy, 2000; Rantala *et al.*, 2002; Ahtiainen *et al.*, 2005; Jacobs & Zuk, 2002). There is evidence that trade-offs exist between the two (McKean & Nunnery, 2001; Rolff & Siva-Jothy, 2002). For example, in *Drosophila melanogaster*, an increased sexual activity reduced the ability of the immune system to eliminate bacteria (McKean & Nunnery, 2001). However, other studies reported a positive correlation between immune response and reproduction (Birkhead *et al.*, 1998; Apanius & Nisbet, 2006; Lee, 2006). For example, mated females of *Gryllus texensis* are less susceptible to *S. marcescens* infection than virgins, suggesting that immune function and reproduction are positively correlated in these crickets (Shoemaker & Adamo, 2007). Furthermore, a trade-off may not be present in general when two fitness-related traits do not share important resources, or if the resources are not limiting (Moret & Schmid-Hempel, 2000; Lee, 2006).

Since males and females differ in their life histories, sex-specific resource investment strategies to achieve an optimal immune response following an infection can be expected (Folstad & Karter, 1992; Rolff, 2002; Zuk & Stoehr, 2002). In invertebrate species, males have been found to invest fewer resources in immune response than females (Rheins & Karp, 1985; Gray, 1998; Rolff, 2001; Rolff & Siva-Jothy, 2002). These sex-specific differences in immunity have been related to Bateman's principle, i.e., females gain fitness by maximizing their lifespan through immunity investment, assuming that higher immunity increases longevity, which in turn increases the time for egg production and oviposition (Rolff, 2002). Males gain fitness by increasing their mating frequency, and should thus invest in increasing their mating rates instead of immunity (Bateman, 1948; Trivers & Willard, 1973; Zuk, 1990; Rolff, 2002; Roth *et al.*, 2011). Bateman's principle is based on the generalization that females make a large investment into offspring because they have large immobile gametes, whereas males make a small investment into offspring because they produce small and mobile gametes (Bateman, 1948). However, Bateman's principle as well as its consequences for investment in immune defense activation was shown to be reversed in species in which sex-roles are reversed and males invest more into offspring than females, e.g., by investing into parental care (Aisenberg & Peretti, 2011; Roth *et al.*, 2011).

As in other Lepidoptera, both *Heliothis virescens* females and males invest substantially into their offspring (Fye & McAda, 1972; Proshold *et al.*, 1982; Blanco *et al.*, 2009). Female moths spend energetic and nutritional resources into the production of up to 1500 eggs, whereas males invest resources to produce a spermatophore, comprising up to 5% of their body mass (Fye & McAda, 1972; Proshold *et al.*, 1982; Blanco *et al.*, 2009). Males as well as females in *H. virescens* can mate multiple times in their lifetime, which is about 30 days in the laboratory at 25 °C (Proshold *et al.*, 1982; Willers *et al.*, 1987; Fitt, 1989; Blanco *et al.*, 2009). However, unlike many animal species in which males can mate multiple times per day every day, *H. virescens* males and females only mate at most once per night (Raina & Stadelbacher, 1990; Heath *et al.*, 1991; Blanco *et al.*, 2009). Consequently, the number of matings is not only limited in females, but also in males. Thus, the male's capacity to increase its number of matings is tightly linked to male longevity, because every night means one more mating opportunity. Therefore, both *H. virescens* males and females can be expected to invest into immunity in response to an immune challenge.

Investment in immunity can be intimately linked to sexual attractiveness. In many species, the attractiveness of a potential mate is determined by the quality of a sexual signal (Domb & Pagel, 2001; Scheuber *et al.*, 2003; Groot *et al.*, 2014). Parasite-mediated sexual selection indicates that sexual signals honestly reflect the quality of the signaler to the receiver (Hamilton & Zuk, 1982). In this scenario, honest signals are costly and condition dependent, and animals in good condition are able to afford to produce a higher quality signal, whereas animals in poor condition cannot invest much

in sexual signalling (Zahavi, 1975; Sheldon & Verhulst, 1996; Rantala *et al.*, 2003). Most studies that have investigated the relationship between reproductive success and immune response used species where males are the sexual signalers, competing for the attention of choosy females, e.g., birds, crickets or wolf-spiders (Faivre *et al.*, 2003; Jacot *et al.*, 2004; Peters *et al.*, 2004; Ahtiainen *et al.*, 2005; Spencer *et al.*, 2005; Shaw *et al.*, 2011). In moths, however, females produce a sex pheromone to attract males from a distance, i.e., hundreds of meters up to kilometers (Roelofs *et al.*, 1974; Tumlinson *et al.*, 1975; Klun *et al.*, 1980; Teal *et al.*, 1981; Vetter & Baker, 1983). In *H. virescens*, males have been shown to differentiate between sex pheromone blends that differ in their qualitative and/or quantitative composition (Vetter & Baker, 1983; Groot *et al.*, 2009, 2014). Thus, female fitness in this species is likely also determined by quality of the female signal, because it is crucially important to attract males. Whether the cost of immunity is associated with the quality of female signals in moths has not been determined.

In moths, the female sex pheromone has been shown to be an important sexual selection signal: females with a signal that deviates from the population mean attract significantly fewer males, so that these signals seem to be under stabilizing selection (Lofstedt, 1993; Cossé *et al.*, 1995; Linn *et al.*, 1997; Zhu *et al.*, 1997). However, only a few studies have considered the female sex pheromone as an honest signal that is costly to produce and plastic depending on the condition of the females (Groot *et al.*, 2010; Harari *et al.*, 2011). A decrease in sex pheromone production or quality following infection might provide a further indication that the female pheromone of *H. virescens* is indeed an honest signal that can indicate the quality of the female to the male.

In this study, we investigated Bateman's principle and immunity as well as the correlation between immunity and mating success in both sexes of *H. virescens* and between immunity and sexual signalling in females. We adapted the predictions for Bateman's principle and immunity to the life history of *H. virescens* as follows. We hypothesize that females and males invest similarly in immunity, because longevity is equally important to the fitness of both sexes, i.e., for females to increase the time to produce and lay eggs and for males to increase their number of matings. We therefore predict that (1) immune gene expression will be similarly induced in males and females in response to an immune challenge. We also hypothesize that sexual attractiveness and immunity compete for the same resource pool and thus are negatively linked. Therefore, we further predict that an induced immune response (2) elicits reduced mating success in both sexes and (3) negatively affects the female sexual signal and sexual signalling behaviour.

## **MATERIAL AND METHODS**

### ***Ethics statement***

Eggs of the moth *H. virescens* were collected on private property in North Carolina, USA and no sampling permissions were required. *Heliothis virescens* is not an

endangered species in the USA and is not protected by law. No ethical approval was required to work with this species in our study.

### ***Insects and bacterial culture***

*Heliothis virescens* (JEN2; collected in 1988 in Clayton, North Carolina) was reared in environmental chambers at 26 °C, 60% humidity, with a reverse 16:8 hours light-dark cycle (scotophase starting at 8 am and ending at 16 pm). Larvae of *H. virescens* were fed on artificial pinto bean diet (Burton, 1970). Pupae were collected, separated by sex, and placed in cups individually. Adults were provided with a 10% honey-water solution. One to four day old virgin adults were used in all experiments of this study.

To induce an immune response in the moths we used freeze-dried cells of the entomopathogenic bacterium *Serratia entomophila*. This strain was shown to be deadly for *H. virescens* larvae and to induce hemocyte apoptosis in *H. virescens* larvae in an earlier study (Barthel *et al.*, 2014). *Serratia entomophila* was obtained from the Department of Bioorganic Chemistry (MPICE, Jena, Germany). Bacteria were grown at 30 °C and 250 rpm in Caso medium. Overnight cultures of *S. entomophila* were centrifuged and the resulting supernatant was discarded. The extracted bacterial pellet was frozen at -20 °C. Samples were then frozen and dried in a lyophilisator at -80 °C for 5 days to kill the bacteria. Lyophilized cells of *S. entomophila* were stored at -20 °C. Dead cells of the entomopathogen were used in this study to measure the effects of immune defense, because this eliminates the confounding effects of metabolism and dynamics of a living pathogen in the moth (Moret & Schmid-Hempel, 2000).

### ***Activation of the immune system response by bacterial challenge in adult moths***

For all our experiments, we challenged the moths by injecting adult males and females with bacteria. Injection of bacteria into the abdominal cavity mimics that bacteria enter the body via wounds in the cuticle, and has been commonly used to measure molecules that are involved in the immune response (Shelby & Popham, 2008). By using injections, we ensured a defined and equal immune response induction in all adult moths throughout all experiments. To induce an immune response in adult moths in all experiments of this study, adult moths were injected with 4 µg / 4 µl lyophilized cells of *S. entomophila* diluted in 1x phosphate buffered saline (PBS). As control treatments, adults were either injected with 4 µl pure PBS (referred to as PBS-injected or wounded) or were not injected at all (referred to as non-injected or control). All injections were conducted using a 10-µl Hamilton syringe and were performed at the onset of photophase, approximately 16-20 h before the start of the experiments.

#### ***(1) Effect of immune challenge on immune gene expression in females and males***

To determine the level and extent of immune response induction in both sexes, we assessed the expression of immune-related genes in female and male adult moths. An induced immune response in insects includes the expression of genes encoding a variety of antimicrobial peptides such as lysozyme, gloverin and hemolin to combat

infections (Hoffmann, 1995). Besides antimicrobial peptides, phenoloxidase (PO) activating enzymes and heat shock proteins (Hsp) are important compounds of an efficient immune response in insects (Hoffmann, 1995; Robert, 2003; Cerenius *et al.*, 2008). For our study we measured the expression levels of five immune related genes, i.e., heat shock protein 70, phenoloxidase activating factor, lysozyme, hemolin and gloverin. These genes have been found to be differentially regulated between control and *S. entomophila*-injected *H. virescens* larvae in a preliminary microarray study (unpublished data). We evaluated the expression differences of these 5 genes between *S. entomophila*-injected, PBS-injected and non-injected moths. Injections were carried out as described above. RNA extraction started 3 to 4 h after the onset of scotophase (20 h after injections). Moths were flash-frozen in liquid nitrogen and stored at -80 °C until RNA extraction. For each treatment, three replicates of each five bodies were used. Total RNA extraction was performed using TRIzol® (Invitrogen) according to the manufacturer's protocol. To ensure that the isolated RNA was free of genomic DNA we performed a DNase treatment by adding 10 µl of Turbo DNase buffer and 1 µl of Turbo DNase enzyme (Ambion). Samples were then incubated for 30 min at 37 °C. RNA was additionally cleaned with the RNeasy MinElute Cleanup-Kit (Qiagen). To check the quality and concentration of the total RNA, isolated RNA was measured by ultraviolet (UV) detection using NanoDrop ND-1000 (Thermo Scientific). First-strand cDNA was synthesized using Verso™ SYBR Green 2-Step QRT-PCRKit Plus ROX Vial (Thermo Scientific, ABgene, UK) according to the manufacturer's instructions, starting with 900 ng of total RNA. Quantitative real-time PCR analysis was performed on a Stratagene Mx3000P QPCR System. Reagents were purchased from ABgene (Thermo Scientific) and used according to the manufacturer's specifications. The PCR reaction for comparative quantification was run at 95 °C for 15 min and 40 cycles at 95 °C for 15 s, 58 °C for 30 s and 72 °C for 30 s. All PCR reactions were performed in technical duplicates, using three biological replicates for each treatment. The efficiency of each primer pair was calculated using the software program LinRegPCR (Ramakers *et al.*, 2003). To evaluate gene expression, ribosomal protein S18 (RpS18), based on a sequence from an In-house database, was used as a reference. Gene expression is given as copy number per 1000 molecules RpS18. All primers used in this study are shown in TABLE S3.1.

### ***(2) Effect of immune-challenge on the mating success of females and males***

To investigate the consequence of an induced immune response on mating success in *H. virescens* females and males, we conducted mate choice experiments. Climate and light conditions in the experimental room were the same as for the rearing, i.e., 26 °C, 60% relative humidity, and a reverse 16:8 hours light-dark cycle (lights off at 8am). Experiments were conducted in square gauze cages (33 x 33 x 33 cm). Each cage contained three adult moths, one chooser and two potential mates of the opposite sex (referred to as potential mate 1 and 2). Mate choice experiments were

conducted in six assays, three male and three female choice assays (TABLE S3.2). In assay 1 and 4, one of the two potential mates in one cage was injected with *S. entomophila* to trigger an immune response, whereas the other potential mate was not injected. To test whether wounding alone could influence mating behaviour, we conducted assays 2, 3, 5 and 6 as controls. The choice in these assays was between *S. entomophila*-injected and wounded potential partners (assay 2 and 5) or between wounded and control potential partners (assay 3 and 6). Choosers were not injected in any of the assays (1-6). The different mate choice experiments were randomly spread over several days, and at least two different assays were conducted on a single day, to avoid block or day effects. Fifty cages were observed in one night. To distinguish between the two potential mates in one cage, one was marked with a waterproof black marker, which was alternated between the two differentially treated potential mates to exclude bias due to marking. The moths were placed in the cages at the end of the photophase and 16 h after injection. Experiments started 150 min after the start of the scotophase, because females and males start to be reproductively active after ~2.5 h into scotophase (Proshold, 1983; Mistrot Pope *et al.*, 1984; Heath *et al.*, 1991). All cages were checked every 30 min for copulation events until 450 min into scotophase, after which time point no further activity occurred. Mating behaviour was observed with the use of a red LED light (Sigma LED safety light). Only first matings in a cage were recorded.

To check whether mating behaviour was associated with the survival of adult moths for three days, we also recorded adult mortality daily for three days following the mate choice experiments. For these three days, all moths were kept individually in small plastic beakers (25 ml) and were provided with a 10% honey-water solution.

### ***(3a) Effect of immune-challenge on the female sexual signal***

To assess the relationship between an induced immune response and the female sex pheromone profile, we injected virgin *H. virescens* with bacteria, with PBS, or we did not inject them, as described above, and analyzed their sex pheromone profile. Females for this experiment were not used in the mate choice experiments. We extracted sex pheromone glands 20 h after the injections (see above) when the virgin females were 2 days old, and between 3 and 4 h after the onset of scotophase. Glands were dissected with microscissors (FST instruments) and incubated for 30 min in conical vials containing 50  $\mu$ l of hexane and 125 ng of the internal standard pentadecane to dissolve the sex pheromone. All pheromone samples were analyzed using a HP7890 gas chromatograph (GC) with a 7683 automatic injector. For the GC analysis, the hexane solution was evaporated with N<sub>2</sub> to 2  $\mu$ l. A volume of 4  $\mu$ l (2  $\mu$ l sample and 2  $\mu$ l octane) was injected into a HP7890 gas chromatograph (GC) with a splitless inlet. The GC was equipped with a DB-WAXetr (extended temperature range) column of 30 m x 0.25 mm x 0.5  $\mu$ m and was coupled with a flame ionization detector (FID). For further information on the GC analysis see Groot *et al.* (2010). Sex pheromone peaks were identified by comparing the retention times of our

samples with the retention times of synthetic compounds (Pherobank, The Netherlands) of the sex pheromone blend of *H. virescens*. Pheromone peaks were integrated manually.

### ***(3b) Effect of immune challenge on female signalling behaviour***

To test whether the frequency and temporal pattern of female signalling, referred to as ‘calling’, was affected by induction of the immune response, we recorded the temporal patterns of calling behaviour of female moths. As the female sex pheromone is produced in a specialized gland that is located around the ovipositor (Jefferson *et al.*, 1968), females were recorded as calling when the ovipositor with gland was clearly extruded from the female abdomen (see also FIGURE 3.4). Injection experiments were done as described above. Two-day old virgin females were placed separately in transparent plastic cups (500 ml) at the end of the photophase (16 h after injection procedure). Calling behaviour experiments were conducted on three consecutive days. Each experimental day, females of all three treatments were tested to avoid block effects. In total, 178 females were observed, 34 at day one, 72 at day two and three, and females were used only once. As female sex pheromone is emitted at night, experiments started at the onset of the scotophase (at 8 am) and the experimental room was kept dark during the experimental period. Calling behaviour (yes or no) was recorded every 5 min until 450 min into scotophase, using a red light torch.

### ***Statistical analysis***

Expression levels of immune-related genes were calculated relative to a housekeeping gene, the ribosomal protein S18 (RpS18), to adjust for possible differences among the samples which are not due to the treatments. Adjusted expression levels were then analyzed with one-way ANOVAs and consecutive Least-Squares means (LS-means) pairwise comparisons with Tukey adjustment. Data were log-transformed to obtain normality of residuals.

Female and male mate choice were tested with two-sided binomial tests. To statistically test the influence of immune system activation on the female sex pheromone composition, we calculated the relative percentage of each sex pheromone compound in the blend. Females that had a total sex pheromone amount of less than 25 ng were excluded from the analysis, because this was the threshold of accuracy of integration of the pheromone peaks. Moreover, due to a dirt peak in the GC run we had to exclude the minor compound Z7-16:Ald from our data set. Relative amounts of the compounds in the pheromone blend are not independent from each other. We therefore divided them by the minor compounds Z9-16:Ald to be able to perform multivariate analysis (Groot *et al.*, 2010, 2014). We chose Z9-16:Ald as denominator, because this compound is not known to be relevant to attract males and did not differ between treatments in preliminary analyses. We then log<sub>10</sub>-transformed the data and performed a MANOVA analysis with the ratios of the five remaining compounds as

response variable and treatment (*S. entomophila*-, PBS- or non-injected) as predictor variable. Since we detected an overall treatment effect in the MANOVA analysis (Pillai's trace = 0.31, df = 2,  $P = 0.00035$ ), we further analyzed the single compounds using ANOVA, followed by LS-means pairwise comparisons with Tukey adjustment. We additionally analyzed the ratio between the relative amounts of 16:Ald and Z11-16:Ald, because Z11-16:Ald is the major sex pheromone component and essential to attract males (Vetter & Baker, 1983; Groot *et al.*, 2014) and the ratio 16:Ald / Z11-16:Ald was shown to have biological relevance to attract males as well: females with low ratios of 16:Ald to Z11-16:Ald were more attractive for males than females with high ratio (Groot *et al.*, 2014). To meet the assumption of normally distributed residuals, we log-transformed the ratio. We then conducted LS-means pairwise comparisons between our three treatments with Tukey adjustment for multiple comparisons with the ratio as response and treatment as predictor variable.

The influence of treatments on the calling behaviour of females was tested with a generalized linear mixed model using Poisson distribution in the glmmADMB package version 0.8.0 (Skaug *et al.*, 2011; Fournier *et al.*, 2012) in the software R. To account for temporal autocorrelation, we summed up the calling events of individual females in blocks of 30 min, such that females calling in a period of 30 min were counted once for this period (even if they called more than one time in these 30 min), which resulted in a smooth parabolic curve (FIGURE 3.4). We used calling as the response variable and treatment, time and time squared as fixed predictor variables. To account for repeated measurements over time, we added individual females as a random effect to the model, time was added as a random slope (time|individual female). Data were not overdispersed (sum of squared Pearson residuals / residuals degrees of freedom = 0.85). An interaction effect of time and treatment was tested but left out of the final model, because it was not significant. The overall significance of treatment was tested with a Chi-square test, using the car package. All statistical analyses were conducted in the program R, version 3.0.2 (R Core Team, 2013).


## RESULTS

### **(1) Effect of immune challenge on immune gene expression in females and males**

Males had higher baseline expression levels of immune-related genes than females (FIGURE 3.1a). Hsp 70, lysozyme and gloverin exhibited significantly higher expression in control males than in control females, whereas PO activating factor and hemolin were similarly expressed in both sexes. When *S. entomophila* was injected into female and male moths, transcription levels of immune-related genes were induced in both sexes compared to the control (FIGURE 3.1b,c). In females, a significant increase in transcript levels of all tested immune-related genes was observed after injection of *S. entomophila* compared to wounded and control females (FIGURE 3.1b). In males, only lysozyme was significantly upregulated after injection with *S. entomophila* compared to wounded and control males (FIGURE 3.1c). Levels of gloverin and hemolin, which were significantly upregulated in males upon *S. entomophila* injection, were also upregulated in wounded

## Immune response versus reproduction in a moth


males compared to control males. The transcript level of PO activating factor was significantly higher in wounded males compared to control males but not compared to *S. entomophila*-injected individuals. Overall, immune-related genes in females were induced only in response to bacterial injections, whereas in males immune-related genes were similarly induced after wounding and bacterial injections (see TABLE S3.3 for significance values).



**FIGURE 3.1.** Expression level of immune-related genes in non-injected, *Serratia entomophila*- or PBS-injected *Heliothis virescens* females and males. Expression levels were compared between control females and males (a) and among all treatments in females (b) and males (c). Values are given as logarithmic copy number per 1000 molecules Rps18. Bars represent the mean of 3 biological replicates with corresponding standard errors. Different letters above the bars represent significant differences at a level of  $P < 0.05$ , based on ANOVA and LS-means pairwise comparisons with Tukey adjustment.




## Immune response versus reproduction in a moth



**FIGURE 3.3.** Influence of immune defense activation on sex pheromone composition of *Heliothis virescens* females. a) Relative amounts of five compounds scaled to Z9-16:Ald. b) Ratio between 16:Ald and Z11-16:Ald. Non-injected n = 38; PBS-injected n = 25; *Serratia entomophila*-injected n = 38. Different letters above the bars indicate significant differences between treatments at a level of  $P < 0.05$  based on LS-means pairwise comparisons with Tukey adjustment for multiple comparisons.

### (3b) Effect of immune-challenge on female signalling behaviour

The number of females calling was not affected by the different treatments ( $\chi^2 = 4.85$ ,  $df = 2$ ,  $P = 0.088$ ). We did not find an interaction effect of treatment and time, suggesting that calling timing was not affected by the immune challenge (FIGURE 3.4). The calling behaviour of all females followed similar temporal and frequency patterns with an ‘on and off’ calling behaviour throughout scotophase. The majority of all females started calling 120 min into scotophase and stopped calling after 420 min, with a peak of calling activity between 240 and 360 min.



**FIGURE 3.4.** Calling activity of virgin *Heliothis virescens* females during scotophase. Individual calling behaviour was registered every 5 min, calling events of 30 min periods were grouped together. Calling behaviour is depicted as the percentage of females that called per time interval (non-injected n = 58; PBS-injected n = 60; *Serratia entomophila*-injected n = 60). The overall treatment effect was tested with a generalized linear mixed model (glmmADMB in R). Photo: calling *H. virescens* female, arrow: extruded ovipositor with sex pheromone gland.

## DISCUSSION

Our study revealed a negative correlation between immune system activation and reproduction in the moth *H. virescens* on a physiological as well as a behavioural level. However, in contrast to our hypothesis, we found that females and males invest differently in immunity: males had higher baseline immunity than females, indicating immune maintenance, while immune system activation was higher in females than in males in response to a bacterial challenge, indicating immune deployment. Our hypothesis that sexual attractiveness and immunity compete for the same resource pool and thus negatively affect each other was partly confirmed: mating success was reduced in *S. entomophila*-injected females, but not in *S. entomophila*-injected males. In addition, injection with *S. entomophila* caused females to modify their sex pheromone profile.

### ***(1) Effect of immune challenge on immune gene expression in females and males***

When assessing the immune status of organisms, it is important to distinguish between immune maintenance, which implies physiological costs to keep the immune system at a certain level of readiness, and immune deployment, which implies physiological costs of immune system activation to combat pathogens (Schmid-Hempel, 2011). We tested this by measuring the expression level of immune-related genes in adults upon challenge with *S. entomophila*, and found that immune deployment was higher in *H. virescens* females than in males. This is in line with reports from other insects. For example, in the scorpionfly *Panorpa vulgaris*, females had a higher lysozyme-like activity and a higher number of (phagocytically active) hemocytes than males (Kurtz *et al.*, 2000). In the grasshopper *Melanoplus sanguinipes*, adult females showed higher PO activities than males in response to a fungus infection (Gillespie & Khachatourians, 1992).

Interestingly, our immune gene study showed that immune maintenance was higher in males than in females. If such an investment increases their longevity, male fitness is likely to be increased: *Heliothis virescens* males only mate at most once per night, so that every additional night of being alive means another mating opportunity (Proshold *et al.*, 1982; Raina & Stadelbacher, 1990). Additionally, *H. virescens* males invest in their offspring by transferring resources for a spermatophore to the female, which can make up to 5% of their body weight and contains not only sperm, but can also contain one third of the male's zinc supplies as well as sugars and proteins (Engebretson & Mason, 1980; Blanco *et al.*, 2009). Male investment into offspring has been shown to affect their investment in immunity in various species. For instance, in the pipefish *Syngnathus typhle*, a sex role reversed species where males invest more in offspring than females, males showed higher innate and adaptive immunity than females (Roth *et al.*, 2011). In another species with sex role reversal, the spider *Allocosa brasiliensis*, males showed a higher encapsulation response than females in response to a nylon filament implant (Aisenberg & Peretti, 2011). Interestingly, in *H. virescens*, which neither shows conventional sex roles nor sex role reversal but an investment of both sexes into

offspring, we found that both sexes invested into immunity in different ways: males had higher immune maintenance, while females showed higher immune deployment.

The immune system of *H. virescens* females also seemed more specific than that of males, because immune gene expression levels differed more between wounding and bacterial challenge in females than in males (FIGURE 3.1). To save resources, it is likely important for animals to distinguish between wounding and bacterial challenge, because the activation of an immune response is costly and can negatively affect other life-history traits (Stearns, 1992; Zuk & Stoehr, 2002; Schmid-Hempel, 2005). This negative correlation may affect female fitness more than male fitness in *H. virescens*, because females are the sexual signalers, whereas males have been shown to be choosy (Vetter & Baker, 1983; Groot *et al.*, 2014). Therefore, access to males is important for female fitness, and females likely need to invest resources in their attractiveness (Groot *et al.*, 2010; Harari *et al.*, 2011). Consequently, females may have evolved a more specific immune response than males, which is only activated in response to cues of pathogens, whereas males may afford constitutively high levels of immune defense and react stronger to minor cues of disease, like wounding.

### **(2) Effect of immune challenge on mating success of females and males**

Since behaviour is intimately linked to the physiology of insects, we hypothesized that sexual attractiveness and immunity compete for the same resource pool and thus influence each other negatively. In the mate choice experiments, we found male choice but not female choice. Specifically, we found in the male choice experiment that *H. virescens* males mated less with *S. entomophila*-injected females and chose more control females in a two-choice test. Males thus seem to choose the healthier female, as would be expected by parasite-mediated sexual selection theory, which states that individuals should prefer parasite-free mates because the genes for resistance would be transmitted to their offspring (Hamilton & Zuk, 1982; Jacobs & Zuk, 2011). Additionally, the transmission avoidance hypothesis postulates that individuals should avoid parasitized mates to avoid getting infected themselves (Clayton, 1991; Møller & Gregersen, 1994; Jacobs & Zuk, 2011). Furthermore, the resource provisioning model assumes that parasite-free individuals should be better at providing resources to mates and their offspring (Clayton, 1991; Jacobs & Zuk, 2011). Therefore, males of *H. virescens* may choose healthier females to either avoid getting infected themselves or ensure that their offspring are provided with optimal resources. However, we also found that males did not distinguish between wounded females and *S. entomophila*-injected females or wounded females and control females. This may be explained by comparing expression levels in our immune gene expression assays, where we found that wounding also induced an immune response in females of *H. virescens*, e.g., in the expression of lysozyme. Additionally, the differences in immune response induction between control and wounded, or wounded and *S. entomophila*-injected females were not as big as between control and *S. entomophila*-injected females (FIGURE 3.1b). These smaller differences might

not have been enough to be recognized by the males. Furthermore, it is known that stress factors, like wounding, induce increases in octopamine which leads to changes in the level of juvenile hormone and thus can inhibit reproduction (Gruntenko & Rauschenbach, 2008; Adamo & Baker, 2011). This phenomenon may explain why males do not distinguish between wounded females and *S. entomophila*-injected females, since males may detect a stress response in both types of females and thus these females appear to be similar.

We can exclude the possibility that male choice was due to an acute risk of *S. entomophila*-injected females to die (in the course of three days), because in all three treatments females survived for more than three days.

In the female mate choice experiments, all types of males were chosen similarly often, indicating that females did not distinguish between differentially treated males. As males reacted less, physiologically, to the bacterial challenge than females, in terms of up-regulation of immune-related genes, it is likely that all males appeared similar to the females in the female mate-choice experiments.

In addition, oxidative stress generated by immune system activation may cause a negative relationship between immunity and attractiveness (van Schantz *et al.*, 1999; Alonso-Alvarez *et al.*, 2007; Jacobs & Zuk, 2011). Since the immune deployment was lower in *H. virescens* males than females, males may suffer less oxidative damage and thus appear healthy and still attractive for females (van Schantz *et al.*, 1999). Furthermore, heat shock proteins (HSPs) were found to be upregulated under stress conditions in insects (Kaufmann, 1990; Feder & Hofmann, 1999). Our results that males do not upregulate HSP 70 in response to *S. entomophila*, suggest that these males may not be stressed and might thus explain why females are not able to differentiate between healthy and sick males in our mate choice experiments. On the other hand, females were found to upregulate the expression of HSP 70 in response to *S. entomophila*. Males may recognize the stress response in infected females and thus avoid sick females, as was found in our mate choice experiment.

In *Tenebrio molitor* juvenile hormone (JH) was found to induce attractiveness, while suppressing the expression of phenoloxidase (Rolff & Siva-Jothy, 2002). Our finding that the expression level of a phenoloxidase activating factor remains constant in males, but significantly increases in females in response to injection with *S. entomophila*, suggests that immunity is negatively correlated with reproduction in females, but not in males. Another explanation for our results might be that the pathogen applied in our study does not cause enough harm in males for choosy females to benefit by selecting males with genes for parasite resistance or the ability to provide optimal resources for their offspring (Jacobs & Zuk, 2011).

### **(3) Effect of immune challenge on sexual signal and calling behaviour in females**

We found the immune system activation in *H. virescens* females to be associated with an altered sex pheromone profile. The ratio of 16:Ald to Z11-16:Ald was significantly higher in *S. entomophila*-injected females than in wounded or control

females. Z11-16:Ald is the major sex pheromone component in the *H. virescens* sex pheromone blend and is essential to attract males (Vetter & Baker, 1983; Groot *et al.*, 2014). As Groot *et al.* (2014) showed that females with higher ratios of 16:Ald / Z11-16:Ald (and thus less of the major component) were less attractive to males in the field than females with lower ratios (Groot *et al.*, 2014), these results indicate that infected females likely attract fewer males than healthy females under field conditions. Our results are consistent with studies by Worden *et al.* (2000), who found that an immune challenge reduces the attractiveness and reproductive success in grain beetles (Worden *et al.*, 2000). Therefore, our findings corroborate the hypothesis that production of pheromones is a condition dependent sexual trait (Penn & Potts, 1998; Rantala *et al.*, 2003).

Interestingly, the changes in the sexual signal towards a less attractive profile did not coincide with a change in calling behaviour. These results indicate that only female immunity and the quality of the female sexual signal are linked, and suggest that the female sex pheromone in moths is an honest signal that indicates the quality of a female to the male, which also fits the framework of parasite mediated sexual selection theory (Hamilton & Zuk, 1982).

Since the pheromone profile differed between wounded and *S. entomophila*-injected females, while males did not mate more with wounded than with *S. entomophila*-injected females in the mate choice experiments, the altered pheromone profile cannot fully explain our mate choice results. Most likely, at close range in mating cages, other traits than the female long-range sex pheromone are important for male choice and female mating success. In moths, males produce and emit a courtship pheromone that is likely important for female choice (Birch *et al.*, 1990; Hillier & Vickers, 2004). However, a male courtship pheromone cannot explain male choice for specific females. Perhaps cuticular hydrocarbon profile (CHCs) plays a role in mate choice at close range in *H. virescens*, as has been found in many insect species, e.g., *Drosophila*, crickets and beetles (Howard *et al.*, 2003; Peterson *et al.*, 2007; Thomas & Simmons, 2009). CHCs have hardly been investigated in moths, with a few exceptions (Jurenka & Subchev, 2000; Piskorski *et al.*, 2010) and their role in moth sexual communication is still unknown.

In this study we found evidence for a negative correlation between the expression of immune-related genes and mating success as well as the female sexual signalling. This negative correlation between the two traits strongly suggests that there is a trade-off between immunity and reproduction in *H. virescens* females. However, testing theories relating to trade-offs between reproduction and immunity require large-scale experiments that assay immune parameters and reproductive processes in the same individuals, while we tested different groups of moths in our experiment.

### **Conclusions**

In the noctuid moth *H. virescens*, we found that investment in immunity is linked to the life history trajectories of males and females, and can negatively impact sexual

attractiveness and reproduction. Immune response is important for both sexes in *H. virescens*, because both females and males mate only once per night and invest substantially into their offspring. Accordingly, in our immune gene expression assays, we found that both males and females invested in immunity. However, the sexes differed in their investment strategies: females invested in immune defense after a bacterial challenge, whereas males had higher baseline immunity than females. Interestingly, these differences in investment were reflected in the mate choice assays. Males chose more for non-injected females than for *S. entomophila*-injected females, whereas females did not show a preference for differentially treated males. Furthermore, *S. entomophila*-injected females had an altered sexual signal compared to non-injected females. Thus, there is a negative correlation between immune system activation and reproduction in females. We did not find this pattern in males, probably because *H. virescens* males invest in immunity maintenance. It will be interesting to determine whether these sex-specific differences in the type of immunity, i.e., immune deployment in females and immune maintenance in males are a general phenomenon.

### **Acknowledgements**

We would like to thank Anna Schwind, Anne Hahn, Stephanie Ley and Susanne Donnerhacke for their help with the insect rearing and all behavioural experiments. We also wish to thank Yukie Sato and Bram Knegt for their advice on statistical analyses and Steph Menken for helpful comments on this manuscript. This research was supported by the National Science Foundation (awards IOS-1052238 and IOS-1456973), the University of Amsterdam and the Max-Planck-Gesellschaft.

### **REFERENCES**

- Adamo, S.A. & Baker, J.L. 2011. Conserved features of chronic stress across phyla: the effects of long-term stress on behavior and the concentration of the neurohormone octopamine in the cricket, *Gryllus texensis*. *Horm Behav* **60**: 478-483.
- Ahtiainen, J.J., Alatalo, R.V., Kortet, R. & Rantala, M.J. 2005. A trade-off between sexual signalling and immune function in a natural population of the drumming wolf spider *Hygrolycosa rubrofasciata*. *J Evol Biol* **18**: 985-991.
- Aisenberg, A. & Peretti, A.V. 2011. Sexual dimorphism in immune response, fat reserves and muscle mass in a sex role reversed spider. *Zoology* **114**: 272-275.
- Alonso-Alvarez, C., Bertrand, S., Faivre, B., Chastel, O. & Sorci, G. 2007. Testosterone and oxidative stress: the oxidation handicap hypothesis. *Proc R Soc B Biol Sci* **274**: 819-825.
- Apanius, V. & Nisbet, I.C. 2006. Serum immunoglobulin G levels are positively related to reproductive performance in a long-lived seabird, the common tern (*Sterna hirundo*). *Oecologia* **147**: 12-23.
- Barthel, A., Kopka, I., Vogel, H., Zipfel, P., Heckel, D.G. & Groot, A.T. 2014. Immune defence strategies of generalist and specialist insect herbivores. *Proc R Soc B Biol Sci* **281**.
- Bateman, A.J. 1948. Intra-sexual selection in *Drosophila*. *Heredity* **2**: 349-368.
- Birch, M., Poppy, G. & Baker, T. 1990. Scents and eversible scent structures of male moths. *Annu Rev Entomol* **35**: 25-54.
- Birkhead, T., Fletcher, F. & Pellatt, E. 1998. Sexual selection in the zebra finch *Taeniopygia guttata*: condition, sex traits and immune capacity. *Behav Ecol Sociobiol* **44**: 179-191.

## Immune response versus reproduction in a moth

- Blanco, C.A., Guadalupe Rojas, M., Groot, A.T., Morales-Ramos, J. & Abel, C.A. 2009. Size and chemical composition of *Heliothis virescens* (Lepidoptera: Noctuidae) spermatophores. *Ann Entomol Soc Am* **102**: 629-637.
- Bonneaud, C., Mazuc, J., Gonzalez, G., Haussy, C., Chastel, O., Faivre, B. & Sorci, G. 2003. Assessing the cost of mounting an immune response. *Am Nat* **161**: 367-379.
- Burton, R.L. 1970. A low cost artificial diet for the corn earworm. *J Econ Entomol* **63**: 1969-1970.
- Cerenius, L., Lee, B.L. & Soderhall, K. 2008. The proPO-system: pros and cons for its role in invertebrate immunity. *Trends Immunol* **29**: 263-271.
- Clayton, D. 1991. The influence of parasites on host sexual selection. *Parasitol Today* **7**: 329-334.
- Cory, J.S. & Hoover, K. 2006. Plant-mediated effects in insect-pathogen interactions. *Trends Ecol Evol* **21**: 278-286.
- Cossé, A., Campbell, M., Glover, T., Linn Jr, C., Todd, J., Baker, T. & Roelofs, W. 1995. Pheromone behavioral responses in unusual male European corn borer hybrid progeny not correlated to electrophysiological phenotypes of their pheromone-specific antennal neurons. *Experientia* **51**: 809-816.
- Domb, L.G. & Pagel, M. 2001. Sexual swellings advertise female quality in wild baboons. *Nature* **410**: 204-206.
- Engebretson, J.A. & Mason, W.H. 1980. Transfer of Zn at mating in *Heliothis virescens*. *Environ Entomol* **9**: 119-121.
- Faivre, B., Gregoire, A., Preault, M., Cezilly, F. & Sorci, G. 2003. Immune activation rapidly mirrored in a secondary sexual trait. *Science* **300**: 103-103.
- Feder, M.E. & Hofmann, G.E. 1999. Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology. *Annu Rev Physiol* **61**: 243-282.
- Fitt, G.P. 1989. The ecology of *Heliothis* species in relation to agroecosystems. *Annu Rev Entomol* **34**: 17-52.
- Folstad, I. & Karter, A.J. 1992. Parasites, bright males, and the immunocompetence handicap. *Am Nat* **139**: 603-622.
- Fournier, D.A., Skaug, H.J., Ancheta, J., Ianelli, J., Magnusson, A., Maunder, M.N., Nielsen, A. & Sibert, J. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim Methods Softw* **27**: 233-249.
- Fye, R.E. & McAda, W. 1972. *Laboratory studies on the development, longevity, and fecundity of six lepidopterous pests of cotton in Arizona*. Agricultural Research Service, US Dept. of Agriculture.
- Gillespie, J.P. & Khachatourians, G.G. 1992. Characterization of the *Melanoplus sanguinipes* hemolymph after infection with *Beauveria bassiana* or wounding. *Comp Biochem Physiol B Biochem Mol Biol* **103**: 455-463.
- Gray, D.A. 1998. Sex differences in susceptibility of house crickets, *Acheta domesticus*, to experimental infection with *Serratia liquefaciens*. *J Invertebr Pathol* **71**: 288-289.
- Groot, A.T., Inglis, O., Bowdridge, S., Santangelo, R.G., Blanco, C., Lopez, J.D., Jr., Teran Vargas, A., Gould, F. & Schal, C. 2009. Geographic and temporal variation in moth chemical communication. *Evolution* **63**: 1987-2003.
- Groot, A.T., Claßen, A., Staudacher, H., Schal, C. & Heckel, D.G. 2010. Phenotypic plasticity in sexual communication signal of a noctuid moth. *J Evol Biol* **23**: 2731-2738.
- Groot, A.T., Schöfl, G., Inglis, O., Donnerhacke, S., Classen, A., Schmalz, A., Santangelo, R.G., Emerson, J., Gould, F., Schal, C. & Heckel, D.G. 2014. Within-population variability in a moth sex pheromone blend: genetic basis and behavioural consequences. *Proc R Soc B Biol Sci* **281**: 20133054
- Gruntenko, N. & Rauschenbach, I.Y. 2008. Interplay of JH, 20E and biogenic amines under normal and stress conditions and its effect on reproduction. *J Insect Physiol* **54**: 902-908.


- Hamilton, W.D. & Zuk, M. 1982. Heritable true fitness and bright birds: a role for parasites? *Science* **218**: 384-387.
- Harari, A.R., Zahavi, T. & Thiéry, D. 2011. Fitness cost of pheromone production in signaling female moths. *Evolution* **65**: 1572-1582.
- Heath, R.R., McLaughlin, J.R., Proshold, F. & Teal, P.E.A. 1991. Periodicity of female sex pheromone titer and release in *Heliothis subflexa* and *H. virescens* (Lepidoptera: Noctuidae). *Ann Entomol Soc Am* **84**: 182-189.
- Hillier, N. & Vickers, N. 2004. The role of heliothine hairpencil compounds in female *Heliothis virescens* (Lepidoptera: Noctuidae) behavior and mate acceptance. *Chem Senses* **29**: 499-511.
- Hoffmann, J.A. 1995. Innate immunity of insects. *Curr Opin Immunol* **7**: 4-10.
- Howard, R.W., Jackson, L.L., Banse, H. & Blows, M.W. 2003. Cuticular hydrocarbons of *Drosophila birchii* and *D. serrata*: identification and role in mate choice in *D. serrata*. *J Chem Ecol* **29**: 961-976.
- Jacobs, A.C. & Zuk, M. 2011. Sexual selection and parasites. In: *Ecoimmunology* (G.E. Demas & J. Nelson, eds), p. 468.
- Jacot, A., Scheuber, H. & Brinkhof, M.W.G. 2004. Costs of an induced immune response on sexual display and longevity in field crickets. *Evolution* **58**: 2280-2286.
- Jefferson, R., Shorey, H. & Rubin, R. 1968. Sex pheromones of noctuid moths. XVI. The morphology of the female sex pheromone glands of eight species. *Ann Entomol Soc Am* **61**: 861-865.
- Jurenka, R.A. & Subchev, M. 2000. Identification of cuticular hydrocarbons and the alkene precursor to the pheromone in hemolymph of the female gypsy moth, *Lymantria dispar*. *Arch Insect Biochem Physiol* **43**: 108-115.
- Kaufmann, S.H. 1990. Heat shock proteins and the immune response. *Immunol Today* **11**: 129-136.
- Klun, J.A., Bierlleonhardt, B.A., Plimmer, J.R., Sparks, A.N., Primiani, M., Chapman, O.L., Lepone, G. & Lee, G.H. 1980. Sex pheromone chemistry of the female tobacco budworm moth, *Heliothis virescens*. *J Chem Ecol* **6**: 177-183.
- Kurtz, J., Wiesner, A., Götz, P. & Sauer, K.P. 2000. Gender differences and individual variation in the immune system of the scorpionfly *Panorpa vulgaris* (Insecta: Mecoptera). *Dev Comp Immunol* **24**: 1-12.
- Lee, K.A. 2006. Linking immune defenses and life history at the levels of the individual and the species. *Integr Comp Biol* **46**: 1000-1015.
- Linn, C., Young, M.S., Gendle, M., Glover, T.J. & Roelofs, W.L. 1997. Sex pheromone blend discrimination in two races and hybrids of the European corn borer moth, *Ostrinia nubilalis*. *Physiol Entomol* **22**: 212-223.
- Lofstedt, C. 1993. Moth pheromone genetics and evolution. *Philos Trans R Soc B Biol Sci* **340**: 167-177.
- McKean, K.A. & Nunney, L. 2001. Increased sexual activity reduces male immune function in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **98**: 7904-7909.
- Mistrot Pope, M., Gaston, L. & Baker, T. 1984. Composition, quantification, and periodicity of sex pheromone volatiles from individual *Heliothis zea* females. *J Insect Physiol* **30**: 943-945.
- Møller, A.P. & Gregersen, J. 1994. *Sexual selection and the barn swallow*. Oxford University Press, Oxford.
- Moret, Y. & Schmid-Hempel, P. 2000. Survival for immunity: The price of immune system activation for bumblebee workers. *Science* **290**: 1166-1168.
- Penn, D. & Potts, W.K. 1998. Chemical signals and parasite-mediated sexual selection. *Trends Ecol Evol* **13**: 391-396.
- Peters, A., Delhey, K., Denk, A.G. & Kempenaers, B. 2004. Trade-offs between immune investment and sexual signaling in male mallards. *Am Nat* **164**: 51-59.

## Immune response versus reproduction in a moth

- Peterson, M., Dobler, S., Larson, E., Juárez, D., Schlarbaum, T., Monsen, K. & Francke, W. 2007. Profiles of cuticular hydrocarbons mediate male mate choice and sexual isolation between hybridising *Chrysochus* (Coleoptera: Chrysomelidae). *Chemoecology* **17**: 87-96.
- Piskorski, R., Trematerra, P. & Dorn, S. 2010. Cuticular hydrocarbon profiles of codling moth larvae, *Cydia pomonella* (Lepidoptera: Tortricidae), reflect those of their host plant species. *Biol J Linn Soc* **101**: 376-384.
- Proshold, F.I., Karpenko, C.P. & Graham, C.K. 1982. Egg production and oviposition in the tobacco budworm: effect of age at mating. *Ann Entomol Soc Am* **75**: 51-55.
- Proshold, F.I. 1983. Release of backcross insects on St. Croix, U.S. Virgin Islands, to suppress the tobacco budworm (Lepidoptera: Noctuidae): infusion of sterility into a native population. *J Econ Entomol* **76**: 1353-1359.
- Raina, A.K. & Stadelbacher, E.A. 1990. Pheromone titer and calling in *Heliothis virescens* (Lepidoptera, Noctuidae): Effect of mating with normal and sterile backcross males. *Ann Entomol Soc Am* **83**: 987-990.
- Ramakers, C., Ruijter, J.M., Deprez, R.H.L. & Moorman, A.F.M. 2003. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci Lett* **339**: 62-66.
- Rantala, M.J., Jokinen, I., Kortet, R., Vainikka, A. & Suhonen, J. 2002. Do pheromones reveal male immunocompetence? *Proc R Soc B Biol Sci* **269**: 1681-1685.
- Rantala, M.J., Kortet, R., Kotiaho, J.S., Vainikka, A. & Suhonen, J. 2003. Condition dependence of pheromones and immune function in the grain beetle *Tenebrio molitor*. *Funct Ecol* **17**: 534-540.
- R Core Team 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rheins, L.A. & Karp, R.D. 1985. Effect of gender on the inducible humoral immune response to honeybee venom in the american cockroach (*Periplaneta americana*). *Dev Comp Immunol* **9**: 41-49.
- Robert, J. 2003. Evolution of heat shock protein and immunity. *Dev Comp Immunol* **27**: 449-464.
- Roelofs, W.L., Hill, A.S., Carde, R.T. & Baker, T.C. 1974. 2 Sex pheromone components of tobacco budworm moth, *Heliothis virescens*. *Life Sci* **14**: 1555-1562.
- Rolff, J. 2001. Effects of age and gender on immune function of dragonflies (Odonata, Lestidae) from a wild population. *Can J Zool* **79**: 2176-2180.
- Rolff, J. 2002. Bateman's principle and immunity. *Proc R Soc B Biol Sci* **269**: 867-872.
- Rolff, J. & Siva-Jothy, M.T. 2002. Copulation corrupts immunity: a mechanism for a cost of mating in insects. *Proc Natl Acad Sci USA* **99**: 9916-9918.
- Roth, O., Scharsack, J.P., Keller, I. & Reusch, T.B.H. 2011. Bateman's principle and immunity in a sex-role reversed pipefish. *J Evol Biol* **24**: 1410-1420.
- Scheuber, H., Jacot, A. & Brinkhof, M.W.G. 2003. Condition dependence of a multicomponent sexual signal in the field cricket *Gryllus campestris*. *Anim Behav* **65**: 721-727.
- Schmid-Hempel, P. 2005. Evolutionary ecology of insect immune defenses. *Annu Rev Entomol* **50**: 529-551.
- Schmid-Hempel, P. 2011. *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford University Press, New York.
- Shaw, K.L., Ellison, C.K., Oh, K.P. & Wiley, C. 2011. Pleiotropy, 'sexy' traits, and speciation. *Behav Ecol* **22**: 1154-1155.
- Shelby, K.S. & Popham, H.J.R. 2008. Cloning and characterization of the secreted hemocytic prophenoloxidasases of *Heliothis virescens*. *Arch Insect Biochem Physiol* **69**: 127-142.
- Sheldon, B.C. & Verhulst, S. 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol Evol* **11**: 317-321.
- Shoemaker, K.L. & Adamo, S.A. 2007. Adult female crickets, *Gryllus texensis*, maintain reproductive output after repeated immune challenges. *Physiol Entomol* **32**: 113-120.

- Siva-Jothy, M.T. 2000. A mechanistic link between parasite resistance and expression of a sexually selected trait in a damselfly. *Proc R Soc B Biol Sci* **267**: 2523-2527.
- Skaug, H., Fournier, D., Nielsen, A., Magnusson, A. & Bolker, B. 2011. glmmADMB. Generalized Linear Mixed Models using AD Model Builder. R package version 0.8.0.
- Spencer, K.A., Buchanan, K.L., Leitner, S., Goldsmith, A.R. & Catchpole, C.K. 2005. Parasites affect song complexity and neural development in a songbird. *Proc R Soc B Biol Sci* **272**: 2037-2043.
- Stearns, S.C. 1992. *The evolution of life histories*. Oxford University Press, Oxford.
- Teal, P.E.A., McLaughlin, J.R. & Tumlinson, J.H. 1981. Analysis of the reproductive behavior of *Heliothis virescens* (f) (Lepidoptera, Noctuidae) under laboratory conditions. *Ann Entomol Soc Am* **74**: 324-330.
- Thomas, M.L. & Simmons, L.W. 2009. Male dominance influences pheromone expression, ejaculate quality, and fertilization success in the Australian field cricket, *Teleogryllus oceanicus*. *Behav Ecol* **20**: 1118-1124.
- Trivers, R.L. & Willard, D.E. 1973. Natural selection of parental ability to vary sex ratio of offspring. *Science* **179**: 90-92.
- Tumlinson, J.H., Hendricks, D.E., Mitchell, E.R., Doolittle, R.E. & Brennan, M.M. 1975. Isolation, identification and synthesis of the sex pheromone of the tobacco budworm. *J Chem Ecol* **1**: 203-214.
- Vetter, R.S. & Baker, T.C. 1983. Behavioral responses of male *Heliothis virescens* in a sustained flight tunnel to combinations of 7 compounds identified from female sex pheromone glands. *J Chem Ecol* **9**: 747-759.
- von Schantz, T., Bensch, S., Grahn, M., Hasselquist, D. & Wittzell, H. 1999. Good genes, oxidative stress and condition-dependent sexual signals. *Proc R Soc B Biol Sci* **266**: 1-12.
- Willers, J., Schneider, J. & Ramaswamy, S. 1987. Fecundity, longevity and caloric patterns in female *Heliothis virescens*: changes with age due to flight and supplemental carbohydrate. *J Insect Physiol* **33**: 803-808.
- Worden, B.D., Parker, P.G. & Pappas, P.W. 2000. Parasites reduce attractiveness and reproductive success in male grain beetles. *Anim Behav* **59**: 543-550.
- Zahavi, A. 1975. Mate selection—a selection for a handicap. *J Theor Biol* **53**: 205-214.
- Zhu, J., Chastain, B.B., Spohn, B.G. & Haynes, K.F. 1997. Assortative mating in two pheromone strains of the cabbage looper moth, *Trichoplusia ni*. *J Insect Behav* **10**: 805-817.
- Zuk, M. 1990. Reproductive strategies and disease susceptibility: An evolutionary viewpoint. *Parasitol Today* **6**: 231-233.
- Zuk, M. & Stoehr, A.M. 2002. Immune defense and host life history. *Am Nat* **160**: S9-S22.

**SUPPORTING INFORMATION**



**FIGURE S3.1.** Pheromone composition of *Heliothis virescens* females that were non-injected (n = 38), injected with *Serratia entomophila* (n = 38) or with PBS (n = 25). Z7-16:Ald was excluded from the analysis, see main text for explanation.

**TABLE S3.1.** Primer sequences used in qRT-PCR analysis.

Gene name	Accession number	Forward primer (5'-3')	Reverse primer (5'-3')
Heat shock protein 70	ACV32640	GTGCTCAGGATCATCAACGA	AGGTCGAAGATGAGCAGGTT
PO activating factor	ACI32835	TGATCAGCCACAGCGTAAAG	CTCCGAATTTGGTCTCTCCA
Lysozyme	AAD00078	CGCTAGAAAGACGGACAAGG	CATTTCAGCGCAAGTGACAT
Gloverin	ACR78446	AGCAGCTTCTTGGGAGGAC	TCCTTATGGACATCAAGAGCAT
Hemolin	ACC91897	TTCCTGAGCCGAAGAATCAC	AATGTTACGCCAACACCACA
Ribosomal protein S18 (RpS18)	In-house database	GCGTGCTGGAGAATGTACTG	GCCTGTGAGGAACCACTCT

**TABLE S3.2.** Experimental setup for mate choice experiments.

Assay	Chooser	Potential mate 1	Potential mate 2	Sample size [n]
Male choice	1 ♂ Non-injected <sup>1</sup>	♀ <i>S. entomophila</i>	♀ Non-injected <sup>1</sup>	43
	2 ♂ Non-injected <sup>1</sup>	♀ <i>S. entomophila</i>	♀ PBS <sup>2</sup>	38
	3 ♂ Non-injected <sup>1</sup>	♀ Non-injected <sup>1</sup>	♀ PBS <sup>2</sup>	41
Female choice	4 ♀ Non-injected <sup>1</sup>	♂ <i>S. entomophila</i>	♂ Non-injected <sup>1</sup>	58
	5 ♀ Non-injected <sup>1</sup>	♂ <i>S. entomophila</i>	♂ PBS <sup>2</sup>	48
	6 ♀ Non-injected <sup>1</sup>	♂ Non-injected <sup>1</sup>	♂ PBS <sup>2</sup>	39

<sup>1</sup>Referred to as control individuals; <sup>2</sup>Referred to as wounded individuals

**TABLE S3.3.** Pairwise comparisons (LS-means with Tukey adjustment) of the expression level of immune-related genes between all treatments. Male and female moths were injected with *Serratia entomophila* (SER), PBS (PBS) or non-injected (NON); n = 3 for all groups. The overall treatment effect was tested by one-way-ANOVA with  $P < 0.001$  for all genes.

Gene	Sex	Comparison	df	t-ratio	P	
Heat shock protein 70	♀	PBS – NON	12	-2.125	0.33	
		SER – PBS	12	-11.837	<0.001***	
		SER – NON	12	-13.963	<0.001***	
	♂	PBS – NON	12	-1.310	0.77	
		SER – PBS	12	0.477	1.00	
		SER – NON	12	-0.833	0.96	
		♀ vs ♂	NON-NON	12	-27.368	<0.001***
	Gloverin	♀	PBS – NON	12	-1.693	0.56
			SER – PBS	12	-11.711	<0.001***
			SER – NON	12	-13.404	<0.001***
♂		PBS – NON	12	-3.601	0.03*	
		SER – PBS	12	-2.652	0.16	
		SER – NON	12	-6.253	<0.001***	
		♀ vs ♂	NON-NON	12	-4.060	0.02*
Hemolin	♀	PBS – NON	12	-0.297	1.00	
		SER – PBS	12	-6.106	<0.001***	
		SER – NON	12	-6.403	<0.001***	
	♂	PBS – NON	12	-3.988	0.02*	
		SER – PBS	12	-1.371	0.74	
		SER – NON	12	-5.359	0.002**	
		♀ vs ♂	NON-NON	12	0.245	1.00
Lysozyme	♀	PBS – NON	12	-7.291	<0.001***	
		SER – PBS	12	-7.858	<0.001***	
		SER – NON	12	-15.149	<0.001***	
	♂	PBS – NON	12	-2.523	0.19	
		SER – PBS	12	-4.752	0.005**	
		SER – NON	12	-7.275	<0.001***	
		♀ vs ♂	NON-NON	12	-9.007	<0.001***
Phenoloxidase activating factor	♀	PBS – NON	12	-0.686	0.98	
		SER – PBS	12	-4.609	0.006**	
		SER – NON	12	-5.294	0.002**	
	♂	PBS – NON	12	-3.505	0.04*	
		SER – PBS	12	3.107	0.08	
		SER – NON	12	-0.398	1.00	
		♀ vs ♂	NON-NON	12	-2.547	0.19

**TABLE S3.4a.** Overall treatment effect for differences in *Heliothis virescens* female sex pheromone between *Serratia entomophila*-injected (n = 38), non-injected (n = 38) and PBS-injected (n = 25) females.

MANOVA	df	Pillai's trace	Approximate F	Pr(>F)
Treatment	2	0.31	3.45	0.00035
Residuals	98			

Immune response versus reproduction in a moth

**TABLE S3.4b.** Effect of treatment on individual compounds of *Heliothis virescens* female sex pheromone tested with ANOVAs and LS-means pairwise comparisons (with Tukey adjustment) for compounds with treatment effect of  $P < 0.05$ . SER: *Serratia entomophila*-injected (n = 38), NON: non-injected (n = 38), PBS: PBS-injected (n = 25)

ANOVAs, individual compounds				LS means pairwise comparisons (Tukey adjusted)			
Compound	df	F	Pr(>F)	Comparison	df	t-ratio	P
14:ALD	2	2.08	0.13	SER-NON	98	-1.40	0.35
				SER-PBS		-1.95	0.13
				PBS-NON		0.71	0.76
Z9:14:ALD	2	0.61	0.55	SER-NON	98	0.97	0.60
				SER-PBS		0.91	0.64
				PBS-NON		-0.05	0.64
16:ALD	2	1.05	0.36	SER-NON	98	-0.14	0.99
				SER-PBS		-1.36	0.37
				PBS-NON		1.23	0.44
Z11-16:ALD	2	1.47	0.24	SER-NON	98	1.67	0.22
				SER-PBS		1.08	0.53
				PBS-NON		0.40	0.91
Z11-16:OH	2	4.0	0.021*	SER-NON	98	2.82	0.016*
				SER-PBS		1.47	0.31
				PBS-NON		1.04	0.55

**TABLE S3.4c.** 16:ALD /Z11-16:ALD ratio in *Heliothis virescens* female sex pheromone. ANOVA for overall treatment effect and LS-means pairwise comparisons with Tukey adjustment between females of different treatment groups, SER: *Serratia entomophila*-injected (n = 38), NON: non-injected (n = 38), PBS: PBS-injected (n = 25).

Tested Ratio	df	F	Pr(>F)	LS means pairwise comparisons (Tukey adjusted)			
				Comparison	df	t-ratio	P
16:ALD / Z11-16:ALD	2	6.74	0.0018**	SER-NON	98	2.45	0.042*
				SER-PBS		3.54	0.018**
				PBS-NON		1.36	0.37

# Effects of immune challenge on the oviposition strategy of a noctuid moth

Heike Staudacher, Steph B.J. Menken & Astrid T. Groot

*Journal of Evolutionary Biology* 28: 1568-1577 (2015)

**Abstract**

Infections can have detrimental effects on the fitness of an animal. Reproducing females may therefore be sensitive to cues of infection and be able to adaptively change their oviposition strategy in the face of infection. As one possibility, females could make a terminal investment and shift reproductive effort from future to current reproduction as life expectancy decreases. We hypothesized that females of the noctuid moth *Heliothis virescens* make a terminal investment and adapt their oviposition timing as well as their oviposition site selectivity in response to an immune challenge. We indeed found that females that were challenged with the bacterial entomopathogen *Serratia entomophila* laid more eggs than control females one night after the challenge. Additionally, bacteria-challenged females were less discriminating between oviposition sites than control females. Whereas control females preferred undamaged over damaged plants, immune-challenged females did not differentiate between the two. These results indicate that terminal investment is part of the life history of *H. virescens* females. Moreover, our results suggest that a strategy of terminal investment in *H. virescens* oviposition represents a fitness trade-off for females: in the face of infection, an increase in oviposition rate enhances female fitness, whereas low oviposition site selectivity reduces female fitness.

**INTRODUCTION**

Pathogens are virtually always present in the environment of organisms and can have detrimental effects on their fitness when infection occurs (Grenfell & Dobson, 1995; Poulin, 2007; Schmid-Hempel, 2011). Under pathogen pressure, natural selection should favour adaptive changes in response to an immune challenge to minimize the costs of infection and maximize lifetime reproductive success (e.g., Adamo, 1999; Agnew *et al.*, 2000; Bonneaud *et al.*, 2004; Javoš & Tamaru, 2004).

One possibility for organisms to adapt their life history strategy is terminal investment, that is the increase of current reproductive effort as life expectancy decreases (Clutton-Brock, 1984). Current reproductive output is expected to trade off with future reproductive output and is therefore in most cases not maximized (Williams, 1966). However, if life expectancy decreases, no resources need to be saved for future reproduction and investment in current reproduction is predicted to increase (Part *et al.*, 1992; Polak & Starmer, 1998; Velando *et al.*, 2006). Terminal investment may be apparent in increased courtship activity, a temporary increase in numbers of offspring and/or investment in offspring survival (Part *et al.*, 1992; Polak & Starmer, 1998; Adamo, 1999; Bonneaud *et al.*, 2004; Creighton *et al.*, 2009). For example, a temporary increase in numbers of eggs in response to an immune challenge has been found in crickets and freshwater snails (Minchella & Loverde, 1981; Adamo, 1999). Studies of several bird species showed an increase in parental care in response to an immune challenge and/or because of ageing (Part *et al.*, 1992; Hanssen, 2006; Velando *et al.*, 2006).

Nonsocial herbivorous insects do not generally provide parental care to their offspring (Janz, 2002). However, in many species, females show oviposition site choice that can be viewed as an investment in offspring survival, as well as a form of parental care (Wiklund & Persson, 1983; Janz, 2002; Lefèvre *et al.*, 2010). Oviposition site choice includes finding and selecting host plants on which offspring can obtain realized fitness and avoiding nonhost or herbivore-infested plants on which offspring cannot (Rothschild & Schoonhoven, 1977; Renwick, 1989; Nishida *et al.*, 1990; Tingle & Mitchell, 1991; De Moraes *et al.*, 2001; Kessler & Baldwin, 2001). Avoiding herbivore-infested plants may be especially important for offspring survival, and thus female fitness, because of enhanced intra- and interspecific competition on such plants (Denno *et al.*, 1995; Kaplan & Denno, 2007), indirect plant defences, such as the emission of volatiles that attract natural enemies of herbivores (Turlings *et al.*, 1990; McCall *et al.*, 1993; De Moraes *et al.*, 1998; Thaler, 1999) or the induced synthesis and enhanced accumulation of secondary plant metabolites that may render the plant unpalatable (e.g., Schoonhoven *et al.*, 2005).

It has long been suggested that when available oviposition time is limited, ovipositing females become less selective and more likely to accept lower-ranked hosts or even non-hosts (Jaenike, 1978; Courtney & Courtney, 1982). As infection can be a cue for shortened life expectancy, oviposition site selectivity and thus investment in offspring survival can be expected to decrease with infection. Moreover, if oviposition rate actually increases in response to infection, ovipositing females become more time-limited. There is evidence that terminal investment behaviour in herbivorous insects in response to cues for shortened life expectancy indeed results in increased plant acceptance (Javošič & Tammaru, 2004). However, to our knowledge, cues of shortened life expectancy have not yet been shown to lower oviposition site selectivity.

In this study, we investigated oviposition timing and oviposition site preference in response to an immune challenge in the generalist herbivore *Heliothis virescens* (Lepidoptera, Noctuidae). Females of this moth species can lay up to about 1500 eggs in their lifetime of about 30 days in the laboratory at 25 °C (Proshold *et al.*, 1982; Willers *et al.*, 1987; Fitt, 1989). Females oviposit throughout the night and lay their eggs singly on plants (Fitt, 1989; Ramaswamy, 1990). *Heliothis virescens* larvae are cannibalistic from their 3rd instar onwards (Gould *et al.*, 1980), so that avoiding to oviposit on plants that are infested with conspecific larvae is likely adaptive for *H. virescens* females. Accordingly, *H. virescens* females have been shown to avoid *Nicotiana tabacum* plants on which conspecific larvae had fed (De Moraes *et al.*, 2001).

Even though *H. virescens* can be regarded an r-strategist, oviposition site choice can be considered an investment into offspring survival, because females need to invest time and resources into finding and selecting an optimal site (Pianka, 1970; Wiklund & Persson, 1983; Willers *et al.*, 1987; Janz, 2002). As infection may limit the time that is available for oviposition, we hypothesized that *H. virescens* females would increase their oviposition rate after a bacterial challenge (H1). Secondly, we

predicted that terminal investment would be apparent in oviposition site selectivity, such that bacteria-challenged females are less discriminating in their choice of oviposition site than control females (H2).

## **MATERIAL AND METHODS**

### ***Insects and bacterial culture***

*Heliothis virescens* was collected in July 2011 in North Carolina, USA, and reared in climate chambers at 25 °C, 60% relative humidity and a light-dark cycle of 10L:14D with lights on at 11 am. Larvae were grown on artificial pinto bean diet (Burton, 1970). Adults were provided with a 10% (wt/vol) sugar-water solution.

To induce an immune response in *H. virescens* females, we used the entomopathogenic bacterium *Serratia entomophila*, which was obtained from the Department of Bioorganic Chemistry (MPICE, Jena, Germany). *Serratia entomophila* was grown overnight in CASO medium at 30 °C on a shaker set at 250 r.p.m. After one night, cultures were centrifuged and the supernatant was discarded. To investigate effects of immune challenge without the confounding effects of the dynamics of a living pathogen, it is common to use immune elicitors like lipopolysaccharides (LPS) (Moret & Schmid-Hempel, 2000; Korner & Schmid-Hempel, 2004) or dead bacterial cells (Haine *et al.*, 2008; Cotter *et al.*, 2010). We therefore killed the bacteria by freezing and drying the samples in a lyophilisator at -80 °C for 5 days. To confirm that bacteria were dead, we streaked them out on Luria-Bertani (LB) agar plates. Lyophilized cells of *S. entomophila* were stored at -20 °C until used in the experiments.

### ***Immune activation via bacterial challenge in mated females***

In all experiments, two groups of females were tested; one group was injected with lyophilized cells of *S. entomophila* to induce an immune response [4 µl of a 1µg/µl solution of bacteria diluted in 1x phosphate buffered saline (PBS)]. This entomopathogenic bacterium was shown to be lethal for *H. virescens* larvae and to induce hemocyte apoptosis in *H. virescens* larvae (Barthel *et al.*, 2014). The other group of females was the control, which was injected with 4 µl of 1x PBS solution. All females were mated in single pair matings one night before they were used in the oviposition assay (night zero). Matings were observed to ensure that females were mated. At the onset of the photophase and between 15 and 17 h before the start of the experiments, the mated females were injected into their abdominal cavity with a 10-µl Hamilton syringe. Previously, we found that injecting lyophilized *S. entomophila* cells in this way elicits an immune response in *H. virescens* females (A. Barthel, H. Staudacher, A. Schmaltz, D.G. Heckel, A.T. Groot, unpublished data). Injecting bacteria has been commonly used to investigate the effects of an immune challenge and as such mimics the process of bacterial cells breaking through the cuticle (Shelby & Popham, 2008). Furthermore, injection ensures that equal amounts of bacterial and PBS solution are used for each female and in all experiments.

### ***H1: Females increase their oviposition rate after a bacterial challenge***

To test the hypothesis that females would make a terminal investment and increase their oviposition rate (that we define here as mean number of eggs/ female/ night) after a bacterial challenge, we conducted oviposition assays with two groups of 2- to 9-day-old *H. virescens* females. One group was injected with *S. entomophila* (n = 27), the other group of females served as the control and was PBS-injected (n = 25), as described above. Experimental females of different ages were distributed evenly between the two treatment groups. The mated and injected females were placed in paper cups (200 ml) at the beginning of the first night after mating (night one), and provided with one dental stick that was soaked in 10% sugar water, which was renewed every night. Cups were closed with transparent gauze. During this experiment, the females were kept without plants. For each female, eggs (all eggs in a cup) were counted at the end of each consecutive night until she died. In the course of the experiment females died, which was recorded daily. Females that did not lay eggs during the experiment were excluded. Also with these exclusions, the age distribution of females remained similar between the two treatment groups (W = 284.5,  $P = 0.33$ , Wilcoxon rank sum test with continuity correction) (see FIGURE S4.1 for age distribution of females in the two treatment groups).

*Statistical analysis.* Differences in number of oviposited eggs between *S. entomophila*-injected and control females for night one were tested with a linear model. Number of eggs at night one was used as response variable and treatment and female age at the start of the experiment as explanatory variables. Additionally, to test whether bacterial challenge had an influence on oviposition rate over time, we analysed the data from the first 13 nights. We chose the period of 13 nights, because after that period there were fewer than 10 surviving females in the *S. entomophila*-injected group left. We used a generalized linear mixed model with negative binomial distribution in the R package glmmADMB (version 0.8.0) (Fournier *et al.*, 2012; Skaug *et al.*, 2014). Number of eggs per female per night was used as response variable. Night and treatment as well as the interaction effect between night and treatment were used as explanatory variables. Female age at the start of the experiment was included as fixed effect in the model. To account for re-peated measurements, we added individual females as random effect to the model, because the eggs of each female were counted every night. To account for temporal autocorrelation, we added night as a random effect slope to the model (night | individual female). To improve the model fit, we included experimental night square and zero inflation in the model which lowered the AIC (Akaike information criterion).

We used a linear model to test for differences in total number of eggs (number of eggs that females laid after injection until they died) between *S. entomophila*-injected and PBS-injected females. Total number of eggs was used as response variable. Treatment and female age at the start of the experiment were used as explanatory variables. To analyse the survival of control and *S. entomophila*-injected females, we used weighted Cox regression in the package coxphw in R (Heinze *et al.*, 2014), with

censoring applied to females that did not die by day 13, and using average hazard ratio (AHR) as template for the case of nonproportional hazards (Schemper, 1992; Schemper *et al.*, 2009; Heinze *et al.*, 2014). Treatment and age were used as explanatory variables. One female of the *S. entomophila* group escaped at night three and was excluded from the models of survival and number of total eggs. To test whether female age and treatment influenced the death of females two nights after injection (when many females were found dead), we constructed a generalized linear model with binomial distribution, using likelihood-ratio as test statistic. Survival [n(alive) = 41] or nonsurvival [n(dead) = 10] for longer than night two was used as response variable, female age and treatment served as explanatory variables.

To determine whether an actually shortened life span affected the number of eggs oviposited in night one after the bacterial challenge, we applied another linear model to the data of the *S. entomophila*-injected group. Number of eggs was used as response variable, and female age as well as survival [n(alive) = 16] or nonsurvival [n(dead) = 10] for longer than night two after injection were used as explanatory variables.

The response variables in the linear models were square-root-transformed when it improved the residual structure of the models. Interaction effects of treatment and female age were tested for all linear models, but were excluded when they were not significant.

## ***H2: Bacteria-challenged females are less discriminating in their choice of oviposition site than control females***

To test the hypothesis that immune system activation influences female oviposition preference, we conducted dual-choice oviposition assays with a different group of mated females, that were *S. entomophila*-injected or PBS-injected (control) as described above. The light-dark cycle of the moth rearing for this experiment was L16:D8 with lights on at 6 am. Mated females were given a choice between a damaged and an undamaged plant. We used *Nicotiana attenuata* plants, which is one of the natural host plants of *H. virescens*. To generate damaged plants, five *H. virescens* 3rd instar larvae were placed on the plants for 48 h and removed from the plants right before the start of the experiment. Undamaged plants were left untreated. All plants were eight to nine weeks old and in their flowering stage, as females mainly lay their eggs on tobacco buds and flowers. Experiments were conducted in cages (2.0 x 0.83 x 1.0 m), which contained one damaged plant, one undamaged plant and one mated female which was 2-8 days old at the night the oviposition experiment started. Age did not differ significantly between the treatments ( $W = 269$ ,  $P = 0.70$ , Wilcoxon rank sum test with continuity correction; for age structure see FIGURE S4.2). Damaged and undamaged plants were placed at a distance of 1.5 m from each other. Each plant was only used once for testing one female. Positions of damaged and undamaged plants in the cages were switched every night to avoid directional effects. The experiment was conducted under natural light conditions during eight nights in May 2014 (sunrise ~5:35 and sunset ~21:40 hours) and two nights in June 2014 (sunrise 5:18 and sunset 22:06 hours) at 25 °C.

Each night, 2-8 females were tested and each female was tested only once, that is for one night. Each night we tested the same number of control and *S. entomophila*-injected females. Females were released in the middle of the cages 1 h before dusk and were removed from the cages on the next day. Eggs were counted on the undamaged and the damaged plant as well as on the sides, bottom and top of the cages, on the day after the experimental night. We will refer to eggs that were not found on plants but anywhere else in the cage as off-plant eggs.

*Statistical analysis.* To test differences in preference between control and *S. entomophila*-injected females for oviposition site (damaged plant, undamaged plant or off-plant), we performed mixed-design ANOVAs with one between variable (female age) and one within variable (oviposition site) to account for the paired character of oviposition site. The analysis was performed separately for control and *S. entomophila*-injected females. The interaction effect between female age and treatment was not significant for *S. entomophila*-injected females and thus excluded from the model for this group. To compare number of eggs between the three oviposition sites, we performed LS-means pairwise comparisons with Tukey correction based on the above-described ANOVA models. Differences in total number of eggs between control and *S. entomophila*-injected females were tested with a linear model, using number of eggs as response variable and treatment and female age as explanatory variables. All analyses were conducted with R version 3.0.2 (R Core Team, 2013).

## RESULTS

### ***H1: Females increase their oviposition rate after a bacterial challenge***


*Serratia entomophila*-injected females laid on average significantly more eggs than control females one night after the immune challenge ( $F_{1,49} = 7.66$ ,  $P = 0.0079$ , FIGURE 4.1a,b). Female age at the start of the experiment did not significantly affect the number of eggs one night after injection ( $F_{1,49} = 2.48$ ,  $P = 0.12$ , FIGURE S4.3).

When we analysed the effect of treatment over time, we found an overall significant interaction effect of night and treatment on the number of eggs per female per night ( $\chi^2 = 4.56$ ,  $df = 1$ ,  $P = 0.033$ , FIGURE 4.1b). *Serratia entomophila*-injected females laid on average more eggs per night than control females from night one to five with an exception of night three. Control females laid on average more eggs per night from night six to night 13 (FIGURE 4.1b).


Total number of eggs laid by control females did not differ significantly from the total number of eggs laid by *S. entomophila*-injected females ( $F_{1,48} = 2.10$ ,  $P = 0.15$ ). Female age at the start of the experiment had a marginally significant effect on the total number of eggs of the females ( $F_{1,48} = 3.95$ ,  $P = 0.053$ , FIGURE 4.2).

*Serratia entomophila*-injected females died significantly earlier than control females after injection ( $z = -2.25$ ,  $P = 0.024$ , FIGURE 4.3). Of the *S. entomophila*-injected females, 38.5% died two nights after the injection. Female age at the start of the experiment did not have a significant effect on the mortality of females (i.e., num-

## Immune challenge and oviposition strategy in a moth




**FIGURE 4.1.** Mean ( $\pm$  SE) number of eggs oviposited per female in response to an immune challenge. a) Number per female 24h after immune challenge; different letters above the bars indicate significant differences at a level of  $\alpha \leq 0.01$  (linear model). b) Daily number per female in the course of 13 nights with polynomial trendline (trendline is only for visualization and was not used for calculations); females were PBS-injected ( $n = 25$  at night zero) or *Serratia entomophila*-injected ( $n = 27$  at night zero).




**FIGURE 4.2.** Total number of eggs laid by *Serratia entomophila*-injected ( $n = 26$ ) and PBS-injected ( $n = 25$ ) females after injection and mating. Boxes span the 25-75 percentiles, lines in the boxes represent the medians, whiskers span the 10-90 percentiles and circles represent data points outside this range; linear model, n.s. = not significant.

ber of days from injection to death) ( $z = 1.07$ ,  $P = 0.29$ ). Female age at the start of the experiment also did not affect the number of females that died two nights after injection ( $\chi^2 = 0.69$ ,  $df = 1$ ,  $P = 0.41$ ). Treatment did have a significant effect on the number of females that were found dead at night two ( $\chi^2 = 16.5$ ,  $df = 1$ ,  $P < 0.001$ , FIGURE 4.3). The number of eggs at night one that were laid by *S. entomophila*-injected females which survived longer than two nights after injection did not differ significantly from the number of eggs that were laid by nonsurvivors of *S. entomophila* injection ( $F_{1,24} = 1.62$ ,  $P = 0.22$ , FIGURE 4.4).



**FIGURE 4.3.** Survival probability of PBS-injected ( $n = 25$ ) and *Serratia entomophila*-injected females ( $n = 26$ ) on 27 consecutive nights after injection; weighted Cox regression.




**FIGURE 4.4.** Number of eggs per *Serratia entomophila*-injected female one night after injection. Boxes span the 25-75 percentiles, lines in the boxes represent the medians, whiskers span the 10-90 percentiles and circles represent data points outside this range. Dead: females were dead at night two after injection ( $n = 10$ ), alive: females survived longer than night two after injection ( $n = 16$ ); n.s. = not significant.

## ***H2: Bacteria-challenged females are less discriminating in their choice of oviposition site than control females***

In control females, the number of eggs differed significantly between oviposition sites (undamaged, damaged plant or off-plant) ( $F_{2,46} = 4.24$ ,  $P = 0.020$ , FIGURE 4.5a). We also detected a significant interaction effect between oviposition site and female age on the number of oviposited eggs ( $F_{2,46} = 13.06$ ,  $P < 0.001$ , FIGURE S4.4). In the pairwise comparison of oviposition sites, control females oviposited more eggs on undamaged than on damaged plants ( $t_{46} = 2.61$ ,  $P = 0.032$ ) or off-plants ( $t_{46} = 2.42$ ,  $P = 0.050$ ), and

a similar number of eggs on damaged plants and off-plant ( $t_{46} = 0.191$ ,  $P = 0.98$ , FIGURE 4.5a). In the control females, age did not have a significant effect on the number of eggs oviposited ( $F_{1,23} = 1.38$ ,  $P = 0.25$ ). In *S. entomophila*-injected females, the number of oviposited eggs did not differ between oviposition sites ( $F_{2,44} = 2.06$ ,  $P = 0.14$ , FIGURE 4.5b), and there was no interaction effect between female age and oviposition site ( $F_{2,42} = 1.51$ ,  $P = 0.23$ , FIGURE S4.5). The total number of eggs was not affected by female age when the two treatment groups were combined ( $F_{1,45} = 0.014$ ,  $P = 0.92$ ).



**FIGURE 4.5.** Oviposition site preference (undamaged plants, damaged plants and off-plant) in *Serratia entomophila*-injected and control *Heliothis virescens* females. a) Number of eggs per female on the three oviposition sites in control females ( $n = 25$ ). Undamaged vs damaged plants:  $P = 0.032$ , undamaged plants vs off-plant:  $P = 0.050$ . b) Number of eggs per female on the three oviposition sites in *S. entomophila*-injected females ( $n = 23$ ), overall effect: not significant. Boxes span the 25-75 percentiles, lines in the boxes represent the medians, whiskers span the 10-90 percentiles and circles represent data points outside this range.

## DISCUSSION

In this study, we found evidence for terminal investment in the oviposition behaviour of *H. virescens* females in response to immune challenge with the bacterial entomopathogen *S. entomophila*. We confirmed our hypotheses that (H1) *H. virescens* females increase their oviposition rate after bacterial challenge and (H2) bacteria-challenged *H. virescens* females are less discriminating in their choice of oviposition site than control females.

### **H1: Females increase their oviposition rate after a bacterial challenge**

The findings that *S. entomophila*-injected females oviposited more eggs earlier and especially laid more eggs on the first night after a bacterial challenge provide further evidence for the terminal investment hypothesis. Indication for terminal investment in nonsocial herbivorous insects has been found earlier in the cricket *Acheta domesticus*, which increased its egg output in response to immune challenge

(Adamo, 1999). Similarly, in the moth *Scotopteryx chenopodiata*, oviposition rate of injured females was also higher than that of females without injuries (Javoš & Tammaru, 2004). In our study, the change in oviposition strategy in response to bacterial challenge was underlined by the fact that the total number of eggs laid by *S. entomophila*-injected and control females did not differ significantly, even though *S. entomophila*-injected females lived fewer days. This suggests that females compensate for a shortened lifetime with a shift of reproductive output from future to current reproduction in response to an immune challenge, which is in accordance with the terminal investment strategy (Williams, 1966; Clutton-Brock, 1984).

The fact that *S. entomophila* injection was deadly for many females in our study suggests a cost of immune response, which has been shown to reduce life span (Sheldon & Verhulst, 1996; Moret & Schmid-Hempel, 2000). Moreover, cytotoxic substances that are produced in the course of immune defence are possibly harmful for host tissue as well (Nappi & Vass, 1993; Zuk & Stoehr, 2002; Schmid-Hempel, 2005). Even though longevity was found to be affected by immune response in many studies (Moret & Schmid-Hempel, 2000; Armitage *et al.*, 2003; Javoš & Tammaru, 2004), mortality is usually not as high as in our study. However, Krams *et al.* (2014) also found very high mortality in larvae of the moth *Galleria mellonella* in response to an immune challenge by nylon bead implantation: larvae which were grown on high energy food showed shorter developmental time, weaker encapsulation response and higher mortality in response to the challenge than larvae grown on low-energy food. As the shorter developmental time was associated with weak encapsulation response, the authors argue that low encapsulation response was likely responsible for high mortality in the high-energy food group (Krams *et al.*, 2014). Thus, possibly more complex relationships between life history traits and immune defence underlie the high mortality of *H. virescens* females in response to immune challenge.

## ***H2: Bacteria-challenged females are less discriminating in their choice of oviposition site than control females***

As control females preferred undamaged over damaged plants and off-plant oviposition sites whereas *S. entomophila*-injected females did not differentiate between oviposition sites, we conclude that oviposition site selectivity can be lowered by cues of shortened life expectancy in herbivorous insects if females are given a choice between sites of different quality. Our results are in line with previous studies that investigated host plant acceptance and oviposition delay without providing a choice. For example, in the moth *S. chenopodiata*, survival of females was associated with oviposition latency on plants of different host quality (Javoš & Tammaru, 2004): survivors of experimentally applied injuries showed a small oviposition latency on superior hosts, but a large oviposition latency on inferior hosts, whereas the reverse was found for moths that laid eggs but did not survive until the end of the experiment (Javoš & Tammaru, 2004). We also confirmed the model prediction that oviposition site selectivity decreases when there is a cue that oviposition time is reduced (Jaenike, 1978; Courtney & Courtney, 1982).

### ***Trade-off between early oviposition and oviposition site preference***

The fact that a cue for shortened life expectancy provoked a temporary increase in oviposition rate in *H. virescens* females after a bacterial challenge indicates a change in oviposition strategy. This change likely optimizes female fitness when infected with pathogens (Minchella & Loverde, 1981; Adamo, 1999; Bonneaud *et al.*, 2004). However, a cue for shortened life expectancy also decreased female oviposition site selectivity, as we found differences in oviposition site choice between *S. entomophila*-injected and control females. A decrease in oviposition site selectivity is expected to reduce female fitness, because oviposition site selectivity has been shown to affect offspring survival in herbivorous insects (Singer, 1972; Rausher, 1982; Gripenberg *et al.*, 2010). The nonpreference for undamaged or damaged plants in *S. entomophila*-injected *H. virescens* females is possibly due to an increased pressure to oviposit early (before death) and indicates that increased egg output after an immune challenge may be linked to lowered oviposition site preference. Terminal investment in this moth may thus be characterized by a trade-off between early oviposition and oviposition site selectivity which likely translates into a fitness trade-off for the females.

The trade-off that we found in our experiments is not necessarily typical of all types of infections. In our experiments, we only used one strain of a pathogenic bacterium to induce an immune defence and one population of *H. virescens* as host. As genotype-specific host-parasite interactions are widespread in nature, it should be stressed that our results cannot be generalized for interactions of *H. virescens* with different parasites or even different strains of the same bacterium used in this study (Schmid-Hempel & Ebert, 2003; de Roode & Altizer, 2010). We chose to conduct the experiments with *S. entomophila*, because *S. entomophila* injection has been shown to induce the immune system of *H. virescens* in a previous study (Barthel *et al.*, 2014).

### ***The influence of female age on oviposition behaviour***

The finding that female age did not affect the number of eggs laid one night after mating could be explained by the fact that in *H. virescens* mating stimulates egg maturation and oviposition (Proshold *et al.*, 1982; Ramaswamy *et al.*, 1997; Zeng *et al.*, 1997). Virgin females lay far fewer eggs than mated females and mating has a particularly stimulating effect on oviposition one day after mating (Proshold *et al.*, 1982). In our experiments, the effect of female age on total number of eggs was marginally significant, which is comparable to the results of Proshold *et al.* (1982) who found that the total number of eggs laid depended on the age at which females were mated. As in our experiments the number of females that were less than three and more than six days old was very small, we could not firmly test homogeneity in the number of eggs laid across age groups.

Interestingly, when we investigated oviposition site preference, we did not find an interaction effect between female age and oviposition site in *S. entomophila*-injected moths, but we did detect such an interaction in the control females. This result indicates that older females discriminate less between oviposition sites than

young females. Since lifetime expectancy generally decreases with age, this age effect fits the prediction that oviposition site selectivity decreases with less time to oviposit (Jaenike, 1978). Hence, bacterial challenge and age seem to similarly induce terminal investment behaviour: possibly, age did not affect oviposition site choice in *S. entomophila*-injected females, because terminal investment was induced by bacterial challenge in females of all ages in this group. Since our experiments were not designed to measure the effect of age, future experiments are needed to investigate the effect of age on oviposition choice in *H. virescens*.

### ***Presence versus absence of plants in oviposition experiments***

When we tested oviposition preference between damaged and undamaged plants, both *S. entomophila*-injected and control females laid about three times as many eggs compared to *S. entomophila*-injected females in the first experiment, where plants were not involved. The presence of tobacco (*N. tabacum*) or tobacco leaf extracts is known to stimulate oviposition in mated *H. virescens* females (Jackson *et al.*, 1984; Mitchell *et al.*, 1990; Ramaswamy *et al.*, 1997): females were found to lay about three times as many eggs on cloth treated with tobacco leaf extract than on untreated cloth (Mitchell *et al.*, 1990). Conversely, Proshold *et al.* (1982) found that *H. virescens* females in mating cups without plant stimuli laid about 200 eggs per female one day after mating, which is more than three times the number that we encountered during oviposition without plant stimuli. These differences may be due to adaptation to host plant-free laboratory rearing conditions, which may inadvertently have selected for females that oviposit more readily on artificial substrate in the absence of plant stimuli: Proshold *et al.* (1982) used a *H. virescens* strain that had been reared in the laboratory for > 60 generations, whereas our oviposition timing experiment was performed with moths that were collected from the field as eggs in 2011 and reared in the laboratory for only 14 generations.

Interestingly, when we tested for an increase in the number of eggs after a bacterial challenge without plants, *S. entomophila*-injected females laid more eggs than control females one night after injection, whereas when we tested oviposition site preference, control and *S. entomophila*-injected females laid a similar total number of eggs. The absence of a suitable host plant can cause a delay in oviposition in moths (Leather & Burnand, 1987; Tammaru & Javoiš, 2000). Oviposition delay was also shown for *S. chenopodiata* on an inferior host plant for survivors of injury, but not for moths that laid eggs but did not survive until the end of the experiment (Javoiš & Tammaru, 2004). Hence, control females in our experiments possibly delayed oviposition only in the absence of a suitable host plant, whereas *S. entomophila*-injected *H. virescens* females laid high numbers of eggs one night after injection even in the absence of a suitable host. These findings are also in accordance with the terminal investment strategy.

### **Conclusions**

We conclude that *H. virescens* females are able to adapt their oviposition strategy by shifting their egg output from future to current reproduction, when survival prospects are compromised by infection. Moreover, by this shift, immune-challenged females seem to be able to compensate for their shorter lifetime by ovipositing more eggs earlier in life, as total number of eggs was not different between immune-challenged and control females. As immune-challenged females were less selective for oviposition site than control females, we show that oviposition site selectivity is another trait that can be affected by an immune challenge in herbivorous insects. As oviposition timing and plant selectivity are likely linked in herbivorous insects, we suggest that there is a fitness trade-off between making a terminal investment by laying more eggs early at the expense of oviposition site selectivity.

### **Acknowledgements**

We thank Andrea Barthel for providing the lyophilized *Serratia entomophila* cells. This study was supported by the University of Amsterdam and partly funded by the National Science Foundation (award IOS-1052238).

### **REFERENCES**

- Adamo, S.A. 1999. Evidence for adaptive changes in egg laying in crickets exposed to bacteria and parasites. *Anim Behav* **57**: 117-124.
- Agnew, P., C. Koella, J. & Michalakis, Y. 2000. Host life history responses to parasitism. *Microbes Infect* **2**: 891-896.
- Armitage, S.A.O., Thompson, J.J.W., Rolff, J. & Siva-Jothy, M.T. 2003. Examining costs of induced and constitutive immune investment in *Tenebrio molitor*. *J Evol Biol* **16**: 1038-1044.
- Barthel, A., Kopka, I., Vogel, H., Zipfel, P., Heckel, D.G. & Groot, A.T. 2014. Immune defence strategies of generalist and specialist insect herbivores. *Proc Biol Sci* **281**: 20140897.
- Bonneaud, C., Mazuc, J., Chastel, O., Westerdahl, H. & Sorci, G. 2004. Terminal investment induced by immune challenge and fitness traits associated with major histocompatibility complex in the house sparrow. *Evolution* **58**: 2823-2830.
- Burton, R.L. 1970. A low-cost artificial diet for the corn earworm. *J Econ Entomol* **63**: 1969-1970.
- Clutton-Brock, T.H. 1984. Reproductive effort and terminal investment in iteroparous animals. *Am Nat* **123**: 212-229.
- Cotter, S.C., Topham, E., Price, A.J.P. & Kilner, R.M. 2010. Fitness costs associated with mounting a social immune response. *Ecol Lett* **13**: 1114-1123.
- Courtney, S.P. & Courtney, S. 1982. The 'edgeeffect' in butterfly oviposition: causality in *Anthocharis cardamines* and related species. *Ecol Entomol* **7**: 131-137.
- Creighton, J.C., Heflin, N.D. & Belk, M.C. 2009. Cost of reproduction, resource quality, and terminal investment in a burying beetle. *Am Nat* **174**: 673-684.
- De Moraes, C.M., Lewis, W.J., Pare, P.W., Alborn, H.T. & Tumlinson, J.H. 1998. Herbivore-infested plants selectively attract parasitoids. *Nature* **393**: 570-573.
- De Moraes, C.M., Mescher, M.C. & Tumlinson, J.H. 2001. Caterpillar-induced nocturnal plant volatiles repel conspecific females. *Nature* **410**: 577-580.


- de Roode, J.C. & Altizer, S. 2010. Host–parasite genetic interactions and virulence–transmission relationships in natural populations of monarch butterflies. *Evolution* **64**: 502–514.
- Denno, R.F., McClure, M.S. & Ott, J.R. 1995. Interspecific interactions in phytophagous insects: competition reexamined and resurrected. *Ann Rev Entomol* **40**: 297–331.
- Fitt, G.P. 1989. The ecology of *Heliothis* species in relation to agroecosystems. *Ann Rev Entomol* **34**: 17–53.
- Fournier, D.A., Skaug, H.J., Ancheta, J., Ianelli, J., Magnusson, A., Maunder, M.N., Nielsen, A. & Sibert, J. 2012. AD Model Builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim Method Softw* **27**: 233–249.
- Gould, F., Holtzman, G., Rabb, R.L. & Smith, M. 1980. Genetic variation in predatory and cannibalistic tendencies of *Heliothis virescens* strains. *Ann Entomol Soc Am* **73**: 243–250.
- Grenfell, B.T. & Dobson, A.P. 1995. *Ecology of infectious diseases in natural populations*. Cambridge University Press, Cambridge, UK.
- Gripenberg, S., Mayhew, P.J., Parnell, M. & Roslin, T. 2010. A meta-analysis of preference–performance relationships in phytophagous insects. *Ecol Lett* **13**: 383–393.
- Haine, E.R., Pollitt, L.C., Moret, Y., Siva-Jothy, M.T. & Rolff, J. 2008. Temporal patterns in immune responses to a range of microbial insults (*Tenebrio molitor*). *J Insect Physiol* **54**: 1090–1097.
- Hanssen, S.A. 2006. Costs of an immune challenge and terminal investment in a long-lived bird. *Ecology* **87**: 2440–2446.
- Heinze, G., Ploner, M. & Dunkler, D. 2014. coxphw: weighted estimation in Cox regression. *R package version 3.0.0*.
- Jackson, D.M., Severson, R.F., Johnson, A.W., Chaplin, J.F. & Stephenson, M.G. 1984. Ovipositional response of tobacco budworm moths (Lepidoptera: Noctuidae) to cuticular chemical isolates from green tobacco leaves. *Environ Entomol* **13**: 1023–1030.
- Jaenike, J. 1978. On optimal oviposition behavior in phytophagous insects. *Theor Popul Biol* **14**: 350–356.
- Janz, N. 2002. Evolutionary ecology of oviposition strategies. In: *Chemoecology of insect eggs and egg deposition* (M. Hilker & T. Meiners, eds), pp. 349–376. Blackwell, Berlin.
- Javoš, J. & Tammaru, T. 2004. Reproductive decisions are sensitive to cues of life expectancy: the case of a moth. *Anim Behav* **68**: 249–255.
- Kaplan, I. & Denno, R.F. 2007. Interspecific interactions in phytophagous insects revisited: a quantitative assessment of competition theory. *Ecol Lett* **10**: 977–994.
- Kessler, A. & Baldwin, I.T. 2001. Defensive function of herbivore-induced plant volatile emissions in nature. *Science* **291**: 2141–2144.
- Korner, P. & Schmid-Hempel, P. 2004. In vivo dynamics of an immune response in the bumble bee *Bombus terrestris*. *J Invertebr Pathol* **87**: 59–66.
- Krams, I., Kecko, S., Kangassalo, K., Moore, F.R., Jankevics, E., Inashkina, I., Krama, T., Lietuvietis, V., Meija, L. & Rantala, M.J. 2014. Effects of food quality on trade-offs among growth, immunity and survival in the greater wax moth *Galleria mellonella*. *Insect Sci* **22**: 431–439.
- Leather, S.R. & Burnand, A.C. 1987. Factors affecting life-history parameters of the pine beauty moth, *Panolis flammea* (D&S): the hidden costs of reproduction. *Funct Ecol* **1**: 331–338.
- Lefèvre, T., Oliver, L., Hunter, M.D. & de Roode, J.C. 2010. Evidence for trans-generational medication in nature. *Ecol Lett* **13**: 1485–1493.
- McCall, P., Turlings, T.J., Lewis, W.J. & Tumlinson, J. 1993. Role of plant volatiles in host location by the specialist parasitoid *Microplitis croceipes* cresson (Braconidae: Hymenoptera). *J Insect Behav* **6**: 625–639.
- Minchella, D.J. & Loverde, P.T. 1981. A cost of increased early reproductive effort in the snail *Biomphalaria glabrata*. *Am Nat* **118**: 876–881.

## Immune challenge and oviposition strategy in a moth


- Mitchell, E.R., Tingle, P.C. & Heath, R.R. 1990. Ovipositional response of three *Heliothis* species (Lepidoptera: Noctuidae) to allelochemicals from cultivated and wild host plants. *J Chem Ecol* **16**: 1817-1827.
- Moret, Y. & Schmid-Hempel, P. 2000. Survival for immunity: the price of immune system activation for bumblebee workers. *Science* **290**: 1166-1168.
- Nappi, A.J. & Vass, E. 1993. Melanogenesis and the generation of cytotoxic molecules during insect cellular immune reactions. *Pigm Cell Res* **6**: 117-126.
- Nishida, R., Ohsugi, T., Fukami, H. & Nakajima, S. 1990. Oviposition deterrent of a Rutaceae-feeding swallowtail butterfly, *Papilio xuthus*, from a non-host rutaceous plant, *Orixa japonica*. *Agr Biol Chem* **54**: 1265-1270.
- Part, T., Gustafsson, L. & Moreno, J. 1992. 'Terminal Investment' and a sexual conflict in the collared flycatcher (*Ficedula albicollis*). *Am Nat* **140**: 868-882.
- Pianka, E.R. 1970. On r-and K-selection. *Am Nat* **104**: 592-597.
- Polak, M. & Starmer, W.T. 1998. Parasite-induced risk of mortality elevates reproductive effort in male *Drosophila*. *Proc Biol Sci* **265**: 2197-2201.
- Poulin, R. 2007. *Evolutionary ecology of parasites*. Princeton University Press, New Jersey.
- Proshold, F.I., Karpenko, C.P. & Graham, C.K. 1982. Egg production and oviposition in the tobacco budworm: effect of age at mating. *Ann Entomol Soc Am* **75**: 51-55.
- Ramaswamy, S. 1990. Periodicity of oviposition, feeding, and calling by mated female *Heliothis virescens* in a field cage. *J Insect Behav* **3**: 417-427.
- Ramaswamy, S.B., Shu, S., Park, Y.I. & Zeng, F. 1997. Dynamics of juvenile hormone-mediated gonadotropism in the Lepidoptera. *Arch Insect Biochem* **35**: 539-558.
- Rausher, M.D. 1982. Population differentiation in *Euphydryas editha* butterflies: larval adaptation to different hosts. *Evolution* **36**: 581-590.
- R Core Team 2013. R: *A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Renwick, J.A.A. 1989. Chemical ecology of oviposition in phytophagous insects. *Experientia* **45**: 223-228.
- Rothschild, M. & Schoonhoven, L.M. 1977. Assessment of egg load by *Pieris brassicae* (Lepidoptera: Pieridae). *Nature* **266**: 352-355.
- Schemper, M. 1992. Cox analysis of survival data with non-proportional hazard functions. *Statistician* **41**: 455-465.
- Schemper, M., Wakounig, S. & Heinze, G. 2009. The estimation of average hazard ratios by weighted Cox regression. *Stat Med* **28**: 2473-2489.
- Schmid-Hempel, P. & Ebert, D. 2003. On the evolutionary ecology of specific immune defence. *Trends Ecol Evol* **18**: 27-32.
- Schmid-Hempel, P. 2005. Evolutionary ecology of insect immune defenses. *Ann Rev Entomol* **50**: 529-551.
- Schmid-Hempel, P. 2011. *Evolutionary Parasitology*. Oxford University Press, New York.
- Schoonhoven, L.M., van Loon, J.J.A. & Dicke, M. 2005. *Insect-Plant Biology*, 2nd edn. Oxford University Press, Oxford.
- Shelby, K.S. & Popham, H.J.R. 2008. Cloning and characterization of the secreted hemocytic prophenoloxidasases of *Heliothis virescens*. *Arch Insect Biochem* **69**: 127-142.
- Sheldon, B.C. & Verhulst, S. 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol Evol* **11**: 317-321.
- Singer, M.C. 1972. Complex components of habitat suitability within a butterfly colony. *Science* **176**: 75-77.
- Skaug, H., Fournier, D., Bolker, B., Magnusson, A. & Nielsen, A. 2014. *Generalized linear mixed models using AD Model Builder*. R package version 0.8.0.
- Tammaru, T. & Javoš, J. 2000. Responses of ovipositing moths (Lepidoptera: Geometridae) to host plant deprivation: life-history aspects and implications for population dynamics. *Environ Entomol* **29**: 1002-1010.

- Thaler, J.S. 1999. Jasmonate-inducible plant defences cause increased parasitism of herbivores. *Nature* **399**: 686-688.
- Tingle, F.C. & Mitchell, E.R. 1991. Effect of oviposition deterrents from elderberry on behavioral responses by *Heliothis virescens* to host-plant volatiles in flight tunnel. *J Chem Ecol* **17**: 1621-1631.
- Turlings, T.C.J., Tumlinson, J.H. & Lewis, W.J. 1990. Exploitation of herbivore-induced plant odors by host-seeking parasitic wasps. *Science* **250**: 1251-1253.
- Velando, A., Drummond, H. & Torres, R. 2006. Senescent birds redouble reproductive effort when ill: confirmation of the terminal investment hypothesis. *Proc Biol Sci* **273**: 1443-1448.
- Wiklund, C. & Persson, A. 1983. Fecundity, and the relation of egg weight variation to offspring fitness in the speckled wood butterfly *Pararge aegeria*, or why don't butterfly females lay more eggs? *Oikos* **40**: 53-63.
- Willers, J.L., Schneider, J.C. & Ramaswamy, S.B. 1987. Fecundity, longevity and caloric patterns in female *Heliothis virescens*: changes with age due to flight and supplemental carbohydrate. *J Insect Physiol* **33**: 803-808.
- Williams, G.C. 1966. Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am Nat* **100**: 687-690.
- Zeng, F., Shu, S., Park, Y. & Ramaswamy, S. 1997. Vitellogenin and egg production in the moth, *Heliothis virescens*. *Arch Insect Biochem* **34**: 287-300.
- Zuk, M. & Stoehr, A.M. 2002. Immune defense and host life history. *Am Nat* **160**: S9-S22.


**SUPPORTING INFORMATION**




**FIGURE S4.1.** Age structure of control (n = 25) and *Serratia entomophila*-injected females (n = 27) at night one after injection and mating in the oviposition timing experiment.




**FIGURE S4.2.** Age structure of control (n = 25) and *Serratia entomophila*-injected females (n = 23) in the experiment for oviposition site selectivity.



**FIGURE S4.3.** Effect of female age on the number of eggs at night one after injection and mating in the oviposition timing experiment in control (n = 25) and *Serratia entomophila*-injected (n = 27) females.



**FIGURE S4.4.** Oviposition site choice (damaged plants, undamaged plants, off-plant) in different age groups in control females (n = 25).



**FIGURE S4.5.** Oviposition site choice (damaged plants, undamaged plants, off-plant) in different age groups in *Serratia entomophila*-injected females (n = 23).



The bacterial symbionts *Wolbachia*,  
*Cardinium* and *Spiroplasma* affect gene  
expression and survival of their spider mite  
host and impact distinct induced responses in  
plants

Heike Staudacher\*, Bernardus C.J. Schimmel\*, Mart M. Lamers,  
Nicky Wybouw, Astrid T. Groot & Merijn R. Kant

*\*Contributed equally to this study*

**Abstract**

Herbivore-associated microbial symbionts may facilitate plant utilization of their hosts directly by providing nutrients or by supporting the breakdown of food. Symbionts may also indirectly facilitate plant utilization by manipulating plant physiology to their host's benefit. Here we investigated the role of the symbiotic prokaryotes *Wolbachia* (W), '*Candidatus Cardinium*' (C) and *Spiroplasma* (S) in plant utilization by two strains of the spider mite *Tetranychus urticae*: Santpoort-2, an 'inducer' strain of plant defence in tomato (*Solanum lycopersicum*), and DeLier-1, a 'suppressor' strain of plant defence. We sequenced the bacterial communities of both strains and found that the inducer strain carries *Cardinium* and *Spiroplasma* (C+S+), while the suppressor strain carries *Wolbachia* and *Spiroplasma* (W+S+). After mites were treated with antibiotics to remove these symbionts, we obtained mite lines for 5 groups: W+S+, W-S+, W-S- for the suppressor and C+S+, C-S- for the inducer strain. In the suppressor strain, we found that the absence of *Wolbachia* correlated with lower mite survival on tomato, while the absence of *Spiroplasma* did not affect mite survival. In the inducer strain, food intake, survival and fecundity were increased when *Cardinium* and *Spiroplasma* were removed from the host. Using microarrays, we determined to what extent the absence of symbionts affected the mite's transcriptome. Interestingly, many of the mite genes that were differentially regulated in mites with or without bacteria encode proteins that play a role in digestion and detoxification or are predicted to be secreted, some of them from the salivary gland. Specifically, in the suppressor strain we found a reduced expression of these genes when *Wolbachia* was absent, while in the inducer strain the absence of *Cardinium* and *Spiroplasma* had the opposite effect. Finally, we investigated to what extent the antibiotics-treated mites induced or suppressed plant responses compared to the non-treated control strains. Excluding *Wolbachia* from the suppressor mites while retaining *Spiroplasma* (W-S+) resulted in a dramatically reduced accumulation of several jasmonic acid (JA)-intermediates, such as 12-oxo-phytodienoic acid (OPDA), but did not affect the end product JA-Ile nor the expression of defence marker-genes downstream of JA. We did however detect an increased salicylic acid (SA) accumulation, but without a uniform effect on the downstream marker genes. Removal of both (endo)symbionts from the suppressor strain yielded an intermediate plant phenotype. Feeding by C-S- inducer mites resulted in higher levels of abscisic acid (ABA) and lower expression of SA marker genes compared to feeding by C+S+ mites, while SA levels were unaffected. Our data suggest that *Wolbachia* is beneficial for the suppressor mites, by possibly enhancing the ability of its host to deal with toxic secondary metabolites in its food. In contrast, *Cardinium* seems to negatively affect the performance of the inducer mites, possibly by inducing expression of (SA-responsive) pathogenesis-related genes in tomato leaflets, while simultaneously inhibiting detoxification processes in the mite. *Spiroplasma* might counteract these effects to some extent through direct interactions with *Wolbachia* and *Cardinium*. In conclusion, spider mite (endo)symbionts seem to interact directly and indirectly with their host. While *Wolbachia* has the characteristics of a mutualist, *Cardinium* has the characteristics of a parasite.

## INTRODUCTION

Herbivores face the challenge to digest, absorb and utilize plant material in order to grow, develop and reproduce. In many cases, plant material is poor or unbalanced in nutrients or contains structural molecules that are hard to digest like cellulose or lignin (Schoonhoven *et al.*, 2005; Karban & Baldwin, 2007). Moreover, herbivores have to cope with plant defences ranging from mechanical barriers, such as thorns and trichomes, to the production of poisonous substances (Schoonhoven *et al.*, 2005; Karban & Baldwin, 2007; Mithöfer & Boland, 2012). Not only herbivores, but also microbes have evolved to consume plant material either as (biotrophic or necrotrophic) pathogens or as detritivores. In some cases, herbivores and microbes have established symbioses (i.e., the living together of dissimilar species; De Bary, 1879) that benefit both partners (mutualistic symbiosis). For instance, some herbivores provide nutrients and shelter to microbes, while in return they make use of the huge metabolic capabilities of these microbes to feed on otherwise unpalatable plants or plant parts, thereby expanding their niche space (Douglas, 2009; Feldhaar, 2011; Engel & Moran, 2013; Hansen & Moran, 2014).

Plant usage by herbivores may be facilitated either by beneficial bacterial symbionts directly, or indirectly via the host plant (Barbosa *et al.*, 1991; Frago *et al.*, 2012; Casteel & Hansen, 2014). For example, direct facilitation may occur via bacteria that upgrade low quality food, by producing essential amino acids or vitamins that the host diet lacks, or by the production of enzymes which enhance the digestion of refractory food sources (Douglas, 2009; Feldhaar, 2011; Engel & Moran, 2013). In contrast, indirect facilitation may occur via interaction of symbiotic bacteria with the plant to benefit their host's fitness (Frago *et al.*, 2012; Hansen & Moran, 2014). For example, bacteria associated with oral secretions, such as that of the Colorado potato beetle (*Leptinotarsa decemlineata*), were shown to alter plant resistance and turn the host plant into better food (Chung *et al.*, 2013).

Not all bacterial symbionts are (always) beneficial for their hosts. In fact, with respect to the effect on host fitness, a symbiosis can span the entire range from beneficial to the microbe but harmful to the host (parasitism), via a neutral relationship (commensal symbiont) to true mutualism. Moreover, symbiosis can have mixed effects and may change over (evolutionary) time, depending on environmental conditions (Werren *et al.*, 2008; Zug & Hammerstein, 2015). Among arthropods, the most prevalent microbial symbionts are so-called reproductive parasites such as *Wolbachia* (Rickettsiales), 'Candidatus Cardinium' (Cytophagales) and *Spiroplasma* (Entomoplasmatales) (Duron *et al.*, 2008). Reproductive parasites commonly secure their prevalence in a host population by increasing the proportion of infected females through various mechanisms including cytoplasmic incompatibility, feminization, parthenogenesis or male killing (Werren, 1997; Duron *et al.*, 2008; Werren *et al.*, 2008; Engelstädter & Hurst, 2009). In most cases, these manipulations by the symbiont are not beneficial to the host, yet very effective for the persistence of the symbiont. This was demonstrated by a recent study in which

approximately 40% of all terrestrial arthropod species was estimated to be infected with *Wolbachia* (Zug & Hammerstein, 2012). However, direct beneficial effects of reproductive manipulators on host fitness are thought to mediate their spread within populations as well, especially when manipulation of host reproduction is weak (Hoffmann *et al.*, 1998; Fry *et al.*, 2004). Accordingly, evidence has accumulated that reproductive manipulators, which have long been considered parasites, can benefit their hosts or vectors in various ways (Casteel & Hansen, 2014; Sugio *et al.*, 2015; Zug & Hammerstein, 2015). For instance, reproductive manipulators have been shown to protect their host against parasitoids, predators and bacterial or viral pathogens (Hedges *et al.*, 2008; Teixeira *et al.*, 2008; Moreira *et al.*, 2009; Jaenike *et al.*, 2010a; Xie *et al.*, 2010, 2014; Walker *et al.*, 2011; Hamilton *et al.*, 2014). Beside protection, *Wolbachia* is known to function as nutritional mutualist in filarial nematodes (Foster *et al.*, 2005), while some examples also exist for arthropod hosts (Brownlie *et al.*, 2009; Hosokawa *et al.*, 2010; Unckless & Jaenike, 2012). Interestingly, infection of arthropods with *Wolbachia* has even been associated with the manipulation of plant physiology (Barr *et al.*, 2010; Kaiser *et al.*, 2010; Body *et al.*, 2013; Robert *et al.*, 2013).

The two-spotted spider mite *Tetranychus urticae* can harbour several (endo)symbiotic bacteria that are known as reproductive manipulators in mites (Breeuwer & Jacobs, 1996; Gotoh *et al.*, 2003, 2007; Enigl & Schausberger, 2007). However, the infection status was shown to vary widely among and within spider mite populations (Breeuwer & Jacobs, 1996; Gotoh *et al.*, 2007). *Tetranychus urticae* is a highly polyphagous pest species found on over 1100 plant species worldwide, including economically important crops like tomato, cucumber, strawberry, bean and cotton (Bolland *et al.*, 1998; Grbic *et al.*, 2011; Migeon *et al.*, 2011). Plants have evolved a wide array of defences, which are organized by the action of several phytohormones in which jasmonic acid (JA) and salicylic acid (SA) are the two central players (Erb *et al.*, 2012; Pieterse *et al.*, 2012). Defence against biotrophic pathogens is orchestrated by SA, while jasmonates, in particular jasmonic acid-isoleucine (JA-Ile), are crucial for defence against herbivores and pathogens with a necrotrophic lifestyle (Kessler *et al.*, 2004; Staswick & Tiryaki, 2004; Glazebrook, 2005). However, like other piercing-sucking arthropods (Walling, 2000), spider mites induce these hormones simultaneously, albeit JA-mediated defences seem to be most important for increased resistance against mites (Li *et al.*, 2002; Ament *et al.*, 2004; Kant *et al.*, 2004, 2008; Zhurov *et al.*, 2014; Alba *et al.*, 2015). Recently, we isolated mites from natural *T. urticae* populations and demonstrated that some of them suppressed JA-mediated defences of tomato to uphold a relatively high reproductive performance on this hostile plant (Alba *et al.*, 2015). One of these suppressor strains, designated as the DeLier-1 strain, was characterized in more detail and shown to significantly reduce JA- and SA-mediated defences that were elicited by the 'normal' inducer strain Santpoort-2 (Alba *et al.*, 2015).

Against the background that (endo)symbiotic bacteria can influence host fitness in various ways, we investigated the bacterial communities that are associated with the *T. urticae* plant defence suppressor strain DeLier-1 and the defence inducer strain Santpoort-2, hereafter referred to as the suppressor and the inducer strain, respectively. The suppressor strain was found to contain *Wolbachia* and *Spiroplasma*, while the inducer strain harboured ‘*Candidatus Cardinium*’ and *Spiroplasma*. We subsequently treated both mite strains with antibiotics to kill the bacteria and tested how their presence was correlated with the performance and transcriptome of the mites. Moreover, we investigated induced plant responses in tomato (*Solanum lycopersicum*) by analyzing phytohormone profiles and expression levels of defence-related genes after infestation of the plants with antibiotics-treated and untreated control mites of both strains that differed in their bacterial community. Together, this enabled us to assess to which extent mite endosymbionts impact mites directly or indirectly via the plant.

## MATERIAL AND METHODS

### *Plants*

Tomato (*S. lycopersicum* cv. Castlemart) and bean (*Phaseolus vulgaris* cv. Speedy) were germinated and grown in a greenhouse (25/18 °C day/night temperature, 16L:8D photoperiod, 50-60% relative humidity [RH]). Experiments involving plants were carried out in a climate room (default settings: 25 °C, 16L:8D photoperiod, 60% RH, 300  $\mu\text{E m}^{-2} \text{s}^{-1}$ ), to which plants were transferred seven days in advance.

### *Spider mites*

We used spider mites from the *T. urticae* strains Santpoort-2 (‘inducer’) and DeLier-1 (‘suppressor’). The Santpoort-2 mites have been described before as inducers of tomato JA- and SA defences, to which they are also susceptible (Kant, 2006; Alba *et al.*, 2015), while DeLier-1 mites suppress these defences (Alba *et al.*, 2015). Spider mites from both strains were reared separately on detached bean leaflets in a climate room. For all plant infestation experiments and mite performance assays, we used age-equilibrated adult females.

### ***I. Bacterial communities of a Tetranychus urticae suppressor and inducer strain, in antibiotics-treated and non-treated mite lines***

Results from a preliminary assessment of the presence of bacteria in the two mite strains indicated that they harboured different endosymbiotic bacteria. Suppressor mites contained *Wolbachia* sp., while *Candidatus Cardinium* (referred to as *Cardinium*) was identified in inducer mites. In addition, *Spiroplasma* sp. was found in both mite strains (data not shown).

#### *Ia. Antibiotics treatments and nomenclature of mite lines*

We treated mites from both strains with antibiotics to remove *Wolbachia*, *Cardinium*

and *Spiroplasma* bacteria. In short, offspring from randomly selected mated adult females ('founder mites') was divided over two treatments: (i) antibiotics-treated and (ii) untreated controls. For the antibiotics treatment, adult female spider mites were first kept on bean leaf discs placed on cotton wool soaked with tetracycline hydrochloride (Sigma-Aldrich, St Louis, MO, USA) for 2-3 days, after which they were transferred to new leaf discs on water-saturated cotton wool to produce eggs. Two days later, adult females were individually sampled in Eppendorf tubes and stored at -80 °C until DNA was extracted for diagnostic PCRs to establish the bacterial infection status of the mites (see below). The eggs on the leaf discs were allowed to hatch and mature in a climate room, after which the antibiotics treatment was repeated. In parallel, untreated control mites were kept on leaf discs placed on water-saturated cotton wool and after egg production mites were sampled for diagnostic PCRs as described below. Three subsequent generations of mites were treated in this way, but with increasing concentrations of tetracycline (i.e., 0.15, 0.20 and 0.30% vol/vol) to obtain mites free of *Wolbachia*, *Cardinium* and *Spiroplasma* as assessed via diagnostic PCR. From generation 4 onwards, all mites (antibiotics-treated or not) were reared on untreated detached bean leaflets to accommodate larger populations (from here on referred to as 'lines').

During the antibiotics treatment, we kept track of the individual mites and their offspring and only kept those lines (antibiotics-treated versus control) that both originated from the same 'founder mite', i.e., these were 'sister lines'. This was done to minimize genetic variation between antibiotics-treated and untreated control lines. Following these criteria, we obtained four lines for the suppressor strain, designated as line 1, 2, 3 and 4. Each of the four lines had three sublines: W+S+ contained both *Wolbachia* and *Spiroplasma*; W-S+ was free of *Wolbachia*, but contained *Spiroplasma*, W-S- was free of *Wolbachia* and *Spiroplasma*. We did not manage to obtain W+S- sublines. For the inducer strain, we obtained four lines as well, which were designated as lines 5, 6, 7 and 8. Each of the four lines had two sublines: C+S+ contained *Cardinium* and *Spiroplasma*, and C-S- was free of *Cardinium* and *Spiroplasma*. Sublines with the same respective bacteria will be referred to as 'groups'. In the suppressor strain the groups were W+S+, W-S+ and W-S-, in the inducer strain the groups were C+S+ and C-S (see FIGURE 5.1 and TABLE 5.1 for an overview of the mite lines and their bacterial communities). Mites from each strain and subline were regularly checked for their bacterial infection status by diagnostic PCR and kept on untreated detached bean leaflets for approximately 15 generations before they were used for the plant infestation assay and mite fecundity tests.

### *Ib. Illumina sequencing*

To assess the presence of *Wolbachia*, *Cardinium*, *Spiroplasma* and other potentially present bacteria in mites from the five groups (W-S-, W-S+, W+S+, C-S- and C+S+) that were used for the plant infestation assay, we sampled five tomato-habituated mites per subline (as described above) for Illumina sequencing. DNA was extracted

from single mites using a fast Chelex method modified from Breeuwer & Jacobs (1996). To isolate the DNA, a single mite was ground and homogenized in 100  $\mu$ l sterile 5% (wt/vol) Chelex (Sigma-Aldrich) with a sterile pestle, after which 2.5  $\mu$ l proteinase K (20 mg/ml, Sigma-Aldrich) was added. Samples were then incubated at 56 °C for 1 h, followed by incubation at 95 °C for 8 min to complete the DNA extraction. DNA from the five mites from the same subline was pooled to form one sample. DNA concentration was adjusted to 25-35 ng/ $\mu$ l per sample. In total, 20 (pooled) samples were sent for sequencing, one for each subline. Amplification and sequencing of the 16S rRNA gene fragment was done by LGC Genomics (Berlin, Germany) using an Illumina MiSeq sequencer (2 x 250 bp paired-end reads; Illumina, San Diego, CA, USA) and the universal primers 341F and 785R (modified from Klindworth *et al.*, 2012, see TABLE S5.1). Since the Chelex method does not yield highly pure DNA, the initial PCR amplification of the 16S rRNA genetic region was done on 20 times diluted DNA. Furthermore, the PCR was run with 35 instead of the usual 30 cycles.

Sequences were provided as adapter clipped FASTQ files and analysed in QIIME (quantitative insights into microbial ecology), which is a standard pipeline for microbial community analysis (Caporaso *et al.*, 2010a). First, forward and reverse reads were joined with the `join_paired_ends.py` algorithm. Joined sequences were quality filtered, applying a Phred threshold of 20. Subsequently, sequences were clustered into operational taxonomic units (OTUs) with the open reference OTU picking command, applying the `uclust` algorithm (Edgar, 2010) and 97% similarity cut-offs. First, sequences were clustered against the reference Greengenes 16S rRNA gene database (<http://greengenes.lbl.gov/>; DeSantis *et al.*, 2006). Sequences that did not cluster with the reference sequences were clustered *de novo*. The most abundant sequence from each OTU cluster was taken as representative sequence. Representative sequences were aligned with PyNAST, using the Greengenes core set as a template (Caporaso *et al.*, 2010b). PyNAST-aligned sequences were checked for chimeras with Chimera Slayer (Haas *et al.*, 2011). Identified chimeras were removed from *de novo* clustered sequences for downstream analysis. Taxonomy was assigned to the representative sequences using the `uclust` consensus taxonomy classifier (Edgar, 2010). The resulting OTU table was manually edited; global singletons and sequences identified as chloroplast and mitochondrial DNA were removed from the dataset. For the graph (FIGURE 5.1), we show OTUs that were present at > 0.5% in at least one of the sublines.

### *Ic. Diagnostic PCRs on mites*

To verify the presence or absence of the most common endosymbionts in mites, viz. *Wolbachia*, *Cardinium* and *Spiroplasma* (Breeuwer & Jacobs, 1996; Gotoh *et al.*, 200, 20073; Enigl & Schausberger, 2007), we performed diagnostic PCRs on DNA extracted from spider mites using genus-specific bacterial primers (TABLE S5.1). Adult female mites, by default not surface-sterilized, were sampled in Eppendorf

tubes and their DNA was extracted using the fast Chelex method as described above. Samples were stored at 4 °C until they were used for PCR.

One  $\mu\text{l}$  of the mite-derived DNA solution (approximately 30–40 ng of DNA) was used as template in a 10.5  $\mu\text{l}$  PCR reaction, further containing 2  $\mu\text{l}$  5x Phire Hot Start Buffer (Thermo Fisher Scientific, Waltham, MA, USA), 2  $\mu\text{l}$  dNTPs (1 mM), 0.5  $\mu\text{l}$  of each primer (10  $\mu\text{M}$  each) and 0.1  $\mu\text{l}$  Phire Hot Start DNA Polymerase (Thermo Fisher Scientific). PCR amplification of *T. urticae actin* was used as a positive control for DNA quality. The PCRs were run on a Bio-Rad T100 thermal cycler (Bio-Rad Laboratories, Hercules, CA, USA) with the following cycling conditions; an initial denaturation for 30 s at 98 °C, followed by 32 cycles of 5 s denaturation at 98 °C, 5 s primer annealing at 52 °C (for *Wolbachia* and *Spiroplasma*), 57 °C (for *Cardinium*), or 58 °C (for *T. urticae actin*) and 10 s elongation at 72 °C; finalized with a 1 min elongation step at 72 °C. PCR-generated amplicons were analysed by agarose-gel electrophoresis. Primer specificity was confirmed by Sanger sequencing of the amplicons.

## **II. Effects of *Wolbachia*, *Cardinium* and *Spiroplasma* on spider mites**

### *Ila. Spider mite performance assay*

To establish whether (endo)symbionts had an effect on mite performance, we assessed spider mite fecundity and survival on wild type tomato plants. For the experiment, an ‘egg-wave’ (see Alba *et al.*, 2015) was generated by allowing random adult females from each strain to produce eggs on the adaxial surface of detached bean leaflets, which had been put flat on wet cotton wool. After 48 h of egg production, all mites were removed from the leaflets and collected in Eppendorf tubes (25–35 mites from the same subline were pooled), flash-frozen in liquid nitrogen and stored at -80 °C, until their DNA was extracted for diagnostic PCRs. The eggs were allowed to hatch and mature in a climate room for another 9 days. The bean leaflets with mites were then transferred to leaves of 21-day-old tomato plants to habituate the mites to tomato. Three days later, the then  $1 \pm 1$ -day-old adult female mites were collected from the tomato leaves and transferred to ‘new’ 21-day-old tomato plants for the mite performance assay. Plants were infested with 5 mites per leaflet; 3 leaflets per plant; 3–6 plants per treatment. A lanolin (Sigma-Aldrich) barrier was made around the petiole to prevent the mites from escaping. After 4 days, the number of eggs produced by the mites, as well as the number of alive, dead and missing (i.e., migrated) mites, was recorded using a stereo microscope (Leica MZ6, Leica Microsystems, Wetzlar, Germany). This experiment was repeated 2–3 times for all four lines of the suppressor strain. However, for the inducer strain, populations from the C-S- subline of line 6, as well as both sublimes of line 7 went (for yet unknown reasons) extinct before we could complete the performance assays. Data from the C+S+ subline of line 6 was included for analysis.

### *Statistical analysis of spider mite performance assay*

To test the effect of *Wolbachia*, *Cardinium* and/or *Spiroplasma* on mite oviposition, we constructed one linear mixed effect models (LMM) for each mite strain

(suppressor and inducer) in the lme4 package (Bates *et al.*, 2013), using either ‘average number of eggs per number of females that were originally put on the leaves’ (5 mites), or ‘the average number of eggs per number of females that survived until the end of the experiment’ as response variable. To test effects of bacteria on survival of the mites, we used generalized linear mixed models in the lme4 package using a binomial distribution to analyse the proportion of mites that were dead or alive at the end of the oviposition experiment (i.e., after 4 days). In the models of oviposition and survival, we used ‘bacterial group’ as explanatory variable and ‘line’ was added as random effect. Additionally, since experiments were spread over different experimental days, and in total 6-9 plants were used per line (with three leaflets per plant), we added a nested random effect with ‘leaflet’ nested in ‘plant’, nested in ‘day’ (1 | day/plant/leaflet) to the model. Pairwise comparisons for suppressor strain were done using Tukey contrasts in the multcomp package (Hothorn *et al.*, 2008) and applying Holm adjustments to account for multiple comparisons. All analyses were performed using the statistical software R 3.0.2 (R Core Team, 2013).

#### *Iib. Spider mite RNA isolation and microarray analysis*

To determine if the presence of *Wolbachia*, *Cardinium* and/or *Spiroplasma* in the mites is associated with gene expression of their host, we isolated RNA from the same mites that were used for the plant infestation assay (see below) and analysed it by means of a microarray. After 7 days of tomato infestations, mites were sampled, as described below in the ‘plant infestation assay’ section. All mites from the same plant (maximum  $3 \times 15 = 45$ ; dead and alive) were pooled to form one biological replicate. Total RNA was isolated from the pooled spider mites using the Direct-zol RNA MiniPrep kit (Zymo Research, Irvine, CA, USA), according to the instructions of the manufacturer. RNA integrity was checked by agarose-gel electrophoresis and a NanoDrop spectrophotometer (ND-1000, Thermo Fisher Scientific) was used to assess RNA purity and quantity.

For the microarray hybridizations, equal amounts of RNA (400 ng), derived from the five biological replicates of each subline, were pooled to form one sample. In total there were twelve samples for the suppressor strain (4 lines x 3 sublimes) and eight samples for the inducer strain (4 lines x 2 sublimes), which were hybridized separately on two arrays, i.e., one with suppressor strain samples, the other with inducer strain samples.

The RNA integrity of each sample was verified with the 2200 TapeStation system (Agilent Technologies, Santa Clara, CA, USA) using the RNA ScreenTape (Agilent Technologies). The amount of RNA per  $\mu\text{l}$  was measured on a NanoDrop spectrophotometer (ND-2000). Per sample, 100 ng of total RNA, combined with Spike A, was amplified and labeled according to the Agilent Two-Colour Microarray-Based Gene Expression Analysis guide version 6.6 (G4140-90050, Agilent Technologies) using the Low Input Quick Amp Labeling Kit (Agilent Technolo-

gies). For the common reference, an equimolar pool of all samples was made and 100 ng samples were amplified similarly as the test samples with the exception that Spike B was used. Synthesized antisense RNA (aRNA) was purified with the EZNA MicroElute RNA Clean Up Kit (Omega Bio-Tek, Norcross, GA, USA). The NanoDrop ND-2000 was used to assess aRNA quantity and CyDye incorporation.

Each hybridization mixture was made up from 1.1 µg Test (Cy3) and 1.1 µg Reference (Cy5) sample. Samples were dried and 1.98 µl water was added. The hybridization cocktail was made according to the manufacturer's instructions (NimbleGen Arrays User's Guide – Gene Expression Arrays version 5.0, Roche NimbleGen, Basel, Switzerland) and 7.2 µl of this mix was added to each sample. The samples were incubated for 5 min at 65 °C and 5 min at 42 °C prior to loading onto a 12 x 135k microarray (Roche NimbleGen), custom designed as described in the next paragraph (Kant *et al.*, 2004, 2008; Alba *et al.*, 2015). Microarrays were hybridized for 20 h at 42 °C with the NimbleGen Hybridization System (Roche NimbleGen). Afterwards, the slides were washed according to the NimbleGen Arrays User's Guide (Gene Expression Arrays version 6.0, Roche NimbleGen) and scanned with an Agilent DNA microarray scanner G2565CA (Agilent Technologies). Feature extraction was performed with NimbleScan v2.6 (Roche NimbleGen).

Microarrays were designed using normalized transcriptomes of *T. urticae* strains Houten-1 ('KOP'; Kant *et al.*, 2004), Santpoort-1 ('KMT'; Kant *et al.*, 2008) and DeLier-1 (Alba *et al.*, 2015). Transcriptomes had been prepared from a normalized shotgun cDNA library via random priming (Houten-1) and three normalized 3-prime libraries cDNA (Houten-1, Santpoort-1 and DeLier-1) via 5' sequencing of poly A-selected fragments of 600-800 nts long via 454 Titanium sequencing (Eurofins, Germany) delivering 1,416,647 reads in total with an average length of 330 nts. For the microarray probe design, we assembled only the 744,649 3' fragment reads, because these will contain more cDNA-specific sequence information than the shotgun reads. These fragment reads were assembled into 16,372 isotigs (with an average length of 571 nts; N50 = 597 nts), using Newbler (using a 99% identity setting and a minimum of 40 nts overlap), from which we could extract 14,335 unique 60-mer probes. After assembly, 167,643 3' reads remained as singleton (i.e., reads that could not be assembled), from which another 120,000 unique 60-mer probes were extracted. Together, these were printed in 12-fold on a 12 x 135k array in Nimblegen format.

Probe sequences were (re)mapped to the latest annotation of the *T. urticae* genome (December 2014) using Bowtie2-2.10 with the 'very sensitive' preset option (Langmead *et al.*, 2009). Of the 134,672 probes on each array, 30,759 aligned with the *T. urticae* genome (Grbic *et al.*, 2011), representing 9,424 (51%) of the 18,414 *in silico* predicted protein-coding sequences. Prior to gene expression analysis, signal intensity data were log<sub>2</sub>-transformed and normalized (Loess and Aquantile). The biological replication within each mite strain and its cured sublimes were assessed by a PCA-analysis, using the `prcomp` function in R. The first two principal components, which explained a cumulative proportion of the total variance of 65.6 and 32.6% for

the inducer and suppressor strain, respectively, were used to explore the data and to identify outliers. Next, a linear model of a non-connected design was fitted to the processed data, using *limma* (Smyth, 2004; Smyth & Altman, 2013). An empirical Bayes approach (Smyth, 2004) was applied to assess relative transcript levels and the associated *P*-values between different treatments. When multiple probes aligned to the same mite gene, the expression values of the individual probes were averaged. To control for false discovery rate, obtained *P*-values were adjusted for multiple testing according to Benjamini and Hochberg's step-up procedure (FDR; Benjamini & Hochberg, 1995). A transcription heat map was constructed, using the relative gene-expression levels obtained in *limma* (bioproject website). All analyses, except for the genome alignment with Bowtie2-2.10 (Linux), were performed using the statistical software R 3.0.2 (R Core Team, 2013). Biological functions were ascribed to the DEG lists as described in Wybouw *et al.* (2015).

### ***III. Effects of spider-mite associated Wolbachia, Cardinium and Spiroplasma on mite-plant interactions***

#### ***IIIa. Plant infestation assay***

To measure phytohormone levels and plant defence gene expression upon mite-inflicted feeding damage, tomato plants were infested with spider mites, as described before (Alba *et al.*, 2015), with an additional habituation step (of 2 days) on tomato to minimize possible effects of the previous diet (i.e., bean) on mite behaviour, mite performance, and/or induced/suppressed tomato defences. Dietary effects are known to persist for at least 48 h, after which they diminish rapidly (Storms, 1971). For the experiment, we used age-equilibrated adult female spider mites obtained from an egg-wave. The eggs were allowed to hatch and mature in a climate room for another 12 days. The bean leaflets with mites were then taken from the cotton wool and placed upside-down on leaves of 28-day-old tomato plants to infest these (i.e., habituation step). Two days later, the  $3 \pm 1$ -day-old adult female mites were collected and transferred to 21-day-old tomato plants for the plant infestation assay, according to our standard infestation protocol (Alba *et al.*, 2015); 15 mites per leaflet, 3 leaflets per plant. To prevent the mites from escaping, a lanolin (Sigma-Aldrich) barrier was made around the petiolule, which was also applied to uninfested control leaflets. A total of 5 plants was infested per mite subline. To verify the bacterial infection status of each strain, 5 tomato-habituated mites per strain were individually collected in Eppendorf tubes, flash-frozen in liquid nitrogen and stored at  $-80\text{ }^{\circ}\text{C}$  until DNA was extracted for PCR amplification and subsequent Illumina sequencing of the 16S rRNA genetic region (see 'Illumina sequencing' section).

At 7 days post-infestation (dpi), mites and tomato leaflets were harvested separately. First, spider mites were removed from the leaflets and collected in Eppendorf tubes (all mites obtained from the same plant were pooled), flash-frozen in liquid nitrogen and stored at  $-80\text{ }^{\circ}\text{C}$  until their RNA was extracted for microarray analysis (see 'Spider mite RNA isolation and microarray analysis' section). A

vacuum pump, sterile 1 ml pipet tip, and mite-proof gauze were used to quickly sample the mites without touching and hence mechanically damaging the leaflets. Subsequently, the mite-cleared leaflets were excised without the petiole. The three detached leaflets obtained from the same plant, along with a scale marker, were then aligned on black paper, gently covered with a thin glass plate to flatten them out, and photographed with a Canon EOS 300D DSLR camera (Canon, Tokyo, Japan) equipped with a Canon EF-S 18-55 mm lens to enable *in silico* calculation of spider mite-inflicted feeding damage, using Adobe Photoshop CS6 Extended (Adobe Systems, San Jose, CA, USA) as described by Kant *et al.* (2004). Finally, the leaflets were flash-frozen in liquid nitrogen and stored at -80 °C until we extracted their phytohormones and RNA. The 3 leaflets obtained from the same plant were pooled to form 1 biological replicate. In total, it took less than 5 min per plant to complete these 3 steps and harvest the leaflets. Care was taken to not damage them. Except for removal of the mites, uninfested control leaflets were processed in the same way.

### *IIIb. Isolation of phytohormones and analysis by means of LC-MS/MS*

Per sample, 200-300 mg of frozen leaf material was homogenized (Precellys 24, Bertin Technologies, Aix-en-Provence, France) in 1 ml of ethyl acetate which had been spiked with D<sub>6</sub>-SA and D<sub>5</sub>-JA (C/D/N Isotopes, Pointe-Claire, Quebec, Canada) as internal standards with a final concentration of 100 ng/ml. Tubes were centrifuged at 13,000 rpm (15,493 x g; Sigma 3-30KS; SIGMA Laborzentrifugen, Osterode am Harz, Germany) for 10 min at 4 °C and the supernatant (the ethyl acetate phase) was transferred to new tubes. The pellet was re-extracted with 0.5 ml of ethyl acetate (without internal standards) and centrifuged again at 13,000 rpm for 10 min at 4 °C. Both supernatants were combined and then evaporated to dryness on a vacuum concentrator (CentriVap Centrifugal Concentrator, Labconco, Kansas City, MO, USA) at 30 °C. The residue was re-suspended in 0.1 ml of 70% methanol (vol/vol), centrifuged at 14,800 rpm (20,081 x g) for 15 min at 4 °C, and the supernatants were transferred to glass vials and then analysed using a liquid chromatography tandem mass spectrometry (LC-MS/MS) system (Varian 320-MS LC/MS, Agilent Technologies). A serial dilution of pure standards of abscisic acid (ABA), traumatic acids, 12-oxo-phytodienoic acid (OPDA), jasmonic acid (JA), jasmonic acid-isoleucine (JA-Ile) and salicylic acid (SA) was run separately. We injected 10 µl of each sample onto a Kinetix 5u C18 100A column (C18 phase, 5 µm particle size, 100Å pore size, 50 × 2.1 mm; Phenomenex, Torrance, CA, USA) equipped with a Phenex-RC guard cartridge (Phenomenex). The mobile phase comprised of solvent A (0.05% formic acid in LC-MS-grade water; Sigma-Aldrich) and solvent B (0.05% formic acid in LC-MS-grade methanol; Sigma-Aldrich). The program, with a constant flow rate of 0.2 ml min<sup>-1</sup>, was set as follows: (i) 95% solvent A/5% solvent B for 1 min 30 s; (ii) followed by 6 min in which solvent B gradually increased till 98%; (iii) continuing with 98% solvent B for 5 min; (iv) then a rapid (in 1 min) but gradual decrease returning to 95% solvent A/5% solvent B until the end of the run. A

negative electrospray ionization mode was used for detection. LC-MS/MS parameters, e.g., analysed compounds, their parent ions, daughter ions, and collision energies used in these analyses, are listed in TABLE S5.1. FIGURE S5.1 shows a comprehensive overview of the most important compounds and their ‘position’ within the plant JA and SA defence response pathways induced by *T. urticae*. For all oxylipins and ABA, we used D<sub>5</sub>-JA to estimate the recovery rate and their *in planta* concentrations were subsequently quantified using the external standard series. For SA we used D<sub>6</sub>-SA to estimate the recovery rate and it was quantified using the external standard. Phytohormone amounts were expressed as ng per gram fresh mass leaf material (ng/g fresh weight). Compounds for which a pure standard was not available (i.e., 13(*S*)-hydroperoxyoctadecatrienoic acid (13-HPOT), 3-oxo-2-(2-pentenyl)-cyclopentane-1-octanoic acid (OPC-8), 3-oxo-2-(2-pentenyl)-cyclopentane-1-hexanoic acid (OPC-6) and C<sub>12</sub>-derivatives of the hydroperoxide lyase (HPL) pathway) were analysed similarly, but their amounts were expressed as ‘Ion Count’ per gram fresh mass leaf material (IC/g fresh weight).

### *IIIc. Gene expression analysis by quantitative reverse-transcriptase PCR (qRT-PCR)*

To determine the effect of mite-associated bacteria on defence gene expression, we performed qRT-PCRs on plant defence marker genes. Therefore, total RNA was isolated from the tomato leaf tissue that was used for phytohormone isolation, using the hot phenol method (Verwoerd *et al.*, 1989). RNA integrity was checked by agarose-gel electrophoresis and a NanoDrop spectrophotometer (ND-1000, Thermo Fisher Scientific) was used to assess RNA purity and quantity. Per sample, 3 µg DNase (Ambion, Austin, TX, USA)-treated RNA was used as template for reverse transcription and first strand cDNA synthesis using RevertAid H Minus Reverse Transcriptase (Thermo Fisher Scientific). For gene expression analysis, 1 µl of 10x diluted cDNA (i.e., the equivalent of 7.5 ng total RNA) served as template in a 20 µl qRT-PCR using the 5x HOT FIREPol EvaGreen qPCR Mix Plus (ROX) kit (Solis Biodyne, Tartu, Estonia) and the ABI 7500 Real-Time PCR system (Applied Biosystems, Foster City, CA, USA), according to the instructions of the manufacturers. We monitored expression of genes involved in JA biosynthesis: *Allene Oxide Synthase 1 (AOS1)* (Howe *et al.*, 2000; Sivasankar *et al.*, 2000) and *OPDA reductase 3 (OPR3)* (Strassner *et al.*, 2002), as well as JA-defence marker genes; *Jasmonate-inducible Protein-21 (JIP-21)* (Lisón *et al.*, 2006), *Threonine Deaminase-2 (TD-2)* (Gonzales-Vigil *et al.*, 2011), *Proteinase Inhibitor IIc (PI-IIc)* (Gadea *et al.*, 1996), SA-defence marker genes; *Pathogenesis-related protein 1a (PR-1a)* (Tornero *et al.*, 1997), *PR-P6* (van Kan *et al.*, 1992), and finally putative OPDA-responsive genes; *Tomato Wound-induced 1 (TWI-1)* (Truesdale *et al.*, 1996), *Alcohol Dehydrogenase (ADH)* (Tieman *et al.*, 2007), an uncharacterized *Gluthathione S-transferase* (‘*GST6*’) (Solyc06g009020.2) and an uncharacterized *Glutaredoxin* (‘*GRX*’) (Solyc07g053550.1). See FIGURE S5.1 for a schematic overview of the plant defence

response against mites, which includes the marker genes whose expression levels were analysed in this study. The amino acid sequences of established *Arabidopsis thaliana* OPDA-responsive genes (ORGs; (Taki *et al.*, 2005; Mueller *et al.*, 2008; Park *et al.*, 2013)) were used to identify their putative tomato homologs: At2g15480 (*AtUGT73B5*) for *SLTWI-1*, At1g09500 (*AtCAD*) for *SLADH*, At2g47730 (*AtGST6*) for *SIGST6*, and At1g28480 (*AtGRX480*) for *SIGRX*. *Actin* was used as a reference gene to normalize expression data and hence correct for variance in quantity of cDNA input. Standard dilution series of selected samples were included with each qRT-PCR run to calculate primer efficiency. PCR-generated amplicons were sequenced to verify primer specificity. Gene identifiers, primer sequences and references are listed in TABLE S5.2. The normalized expression (NE) data were calculated by the  $\Delta Ct$  method as described before (Alba *et al.*, 2015):  $NE = (PE_{\text{target}}^{Ct_{\text{target}}}) / (PE_{\text{reference}}^{Ct_{\text{reference}}})$ , in which PE is the primer efficiency and Ct the number of cycles to reach the cycle threshold value.

*Statistical analysis of the plant infestation assay data (phytohormones, qRT-PCR, feeding damage)*

To test the effect of bacteria on phytohormone levels, tomato gene expression and amount of spider mite-inflicted feeding damage, we constructed LMM using the package lme4 (Bates *et al.*, 2013). The respective amounts of phytohormones (ng/g FW or IC/g FW), normalized gene expression (NE) or total amount of feeding damage (mm<sup>2</sup>) for three leaflets of one plant combined were used as response variable, while ‘presence of bacteria’ (i.e., W/S/C) was used as explanatory variable. To test the level of induction or suppression as compared to uninfested control plants, in the case of phytohormones and gene expression this explanatory variable also included control plants that were not infested. Since we had four lines which were present as sublimes in all mite-bacteria groups, we added ‘line’ as a random factor to the model to account for variation between the lines. The response variables were transformed using log, sqrt or 1/sqrt if applicable for meeting the assumptions of homogeneity of variance and normality of residuals required for LMM. For pairwise comparisons, we used Tukey contrasts with Holm adjustment for multiple comparisons in the multcomp package (Hothorn *et al.*, 2008). To assess how the phytohormones SA and OPA correlated with expression levels of putative ORGs (*TWI-1*, *GRX*, *GST-6*, *ADH* and *OPR3*), we calculated linear correlations between phytohormone amounts and the NE level of these genes using the R package Hmisc 3.15 (Harrel *et al.*, 2015). Further, we calculated linear correlations between SA amounts and NE levels of the SA marker genes *PR-P6* and *PR-1a*. *P*-values of correlations were adjusted for multiple testing, using the Holm method. All analyses were performed using the statistical software R 3.0.2 (R Core Team, 2013).

## RESULTS

### *I. Bacterial communities of a Tetranychus urticae suppressor and inducer strain, in antibiotics-treated and non-treated mite lines*

Adult female mites, obtained from laboratory populations of either the *T. urticae* suppressor or *T. urticae* inducer strain (Alba *et al.*, 2015), harboured different bacterial communities, as was determined by Illumina sequencing of the 16S rRNA derived PCR products (FIGURE 5.1, TABLE 5.1). Most evidently, in the suppressor strain a high percentage of the reads corresponded with the endosymbiotic bacterium *Wolbachia* sp. (W) (Rickettsiaceae), with an average of 30.41% ( $\pm$  13.58 SD). *Wolbachia* was identified in the inducer strain as well, albeit at low relative levels with an average of 0.25% ( $\pm$  0.32 SD). In contrast, in the inducer strain a high percentage of reads corresponded to the endosymbiotic bacterium ‘*Candidatus Cardinium*’ (C) (Bacteroidaceae), with an average of 29.05% ( $\pm$  8.72 SD). *Cardinium* was also found in the suppressor strain, but at low relative levels with an average of 0.0032% ( $\pm$  0.0037 SD). In addition, the same *Spiroplasma* sp. (S) (Spiroplasmataceae) OTU was present in similar relative amounts in both mite strains: an average of 4.03% ( $\pm$  1.06 SD) in the suppressor mites and 4.42% ( $\pm$  2.67 SD) in inducer mites. Therefore, the suppressor strain was classified as W+S+ and the inducer strain as C+S+.

To assess the effect(s) of the bacteria on mite performance, transcriptome and induced plant responses, mites from both strains were treated with antibiotics (tetracycline) as described above. The antibiotics treatments successfully cleared *Wolbachia* from the suppressor mites, as only very few (0-6) reads were detected per line in the W-S+ and W-S- groups (FIGURE 5.1, TABLE 5.1). *Spiroplasma* was completely removed in the latter group, but was relatively more abundant in the W-S+ group than in the W+S+ one, i.e., on average at 7.98% ( $\pm$  1.71 SD). The antibiotics treatment was also successful in the inducer mites (FIGURE 5.1). Only a small fraction of the reads (< 0.04%) recovered from tetracycline-treated C-S- mites corresponded to *Cardinium* or *Spiroplasma* (TABLE 5.1). Moreover, presence of *Wolbachia*, *Spiroplasma* or *Cardinium* was no longer detectable in the antibiotics-treated sublines by means of PCR, using bacteria-specific primers on DNA of individual mites, which was also used for Illumina sequencing (data not shown).


Besides *Wolbachia*, *Cardinium* and *Spiroplasma*, bacteria from the families Enterobacteriaceae (Enterobacteriales) and Pseudomonadaceae (Pseudomonadales) were also present in varying amounts in all groups and sublines of both mite strains (FIGURE 5.1). Other bacterial families reached high relative abundances in some of the sublines, e.g., two different Oxalobacteraceae (Burkholderiales), one in line 5 subline C-S-, the other line 6 subline C+S+ and line 2 subline W+S+; Sphingobacteriaceae (Sphingobacteriales) in line 2 subline W+S+, or Nocardiaceae in line 4 subline W-S+. Taken together, this demonstrates that we cleared the mites from *Wolbachia*, *Spiroplasma* or *Cardinium*, but not from all bacteria residing in or on the mites, possibly because we had not surface-sterilized mites before assaying them.

Therefore, we only focused on the consistent presence/absence of *Wolbachia*, *Spiroplasma* or *Cardinium* in all our analyses, to which we refer when we use the terms ‘bacteria’ or ‘mite-associated bacteria’.

## II. Effects of *Wolbachia*, *Cardinium* and *Spiroplasma* on spider mites

### IIa. Effects of *Wolbachia*, *Cardinium* and *Spiroplasma* on spider mite performance

To assess if the mite-associated bacteria were correlated with the fitness of their host, we determined the number of eggs and survival of adult female mites by performing oviposition assays on tomato. For the suppressor mites, we found an overall significant effect of the factor ‘bacterial group’ (i.e., presence/absence of *Wolbachia* and/or *Spiroplasma*) on mite survival (FIGURE 5.2a). More W+S+ mites



**FIGURE 5.1.** Bacterial community composition derived from Illumina 16S rRNA amplicon-sequencing in two strains of the spider mite *Tetranychus urticae* which had been treated with tetracycline and hence did (+) or did not (-) contain *Wolbachia* (W), *Spiroplasma* (S) and/or *Candidatus Cardinium* (C); Plant defence suppressor strain DeLier-1 with three mite groups: W+S+, W-S+ and W-S-; Plant defence inducer strain Santpoort-2 with two mite groups: C+S+ and C-S-; 4 lines per group. Bacterial operational taxonomic units (OTUs) were combined at the family level. Numbers behind family names indicate how many OTUs of one family were combined. Rare OTUs that were overall represented less than 0.5% are not shown; square brackets around taxa indicate that the taxon name is not fully established yet.

**TABLE 5.1.** Antibiotics treatments of two strains of the spider mite *Tetranychus urticae* resulted in the (near) complete removal of their associated bacteria *Wolbachia*, *Spiroplasma* and/or *Candidatus Cardinium*, as demonstrated by Illumina 16S rRNA amplicon-sequencing. The *T. urticae* plant defence suppressor strain DeLier-1 contained *Wolbachia* and *Spiroplasma* (W+S+) and tetracycline treatments either removed only *Wolbachia* or both bacteria, yielding the groups W-S+ and W-S-, respectively. The plant defence inducer strain Santpoort-2 harboured *Cardinium* and *Spiroplasma* (C+S+), which were both removed by the tetracycline treatments, yielding the group of C-S- mites. Each group is represented by four independent mite lines. Note that mites from lines with the same number are ‘sister lines’ which originate from the same untreated ‘founder mite’, hence each DeLier-1 line consists of three sublines (W+S+, W-S+, W-S-) and each Santpoort-2 line consists of two sublines (C+S+ and C-S-). Shown are the overall total number of Illumina reads obtained per subline, as well as the total number of reads corresponding to all 11 *Wolbachia* operational taxonomic units (OTUs), all 3 *Spiroplasma* OTUs and the only *Cardinium* OTU identified. The 16S rRNA amplification was done with universal 341F and 785R primers, modified from Klindworth *et al.* (2012), see TABLE S5.2.

Mite strain				Suppressor strain				
Group	W+S+			W-S+				
Line	1	2	3	4	1	2	3	4
Total	3,164	14,303	34,118	19,747	4,371	16,637	27,939	8,812
<i>Wolbachia</i>	1,272	2,131	7,940	8,546	1	-	6	1
<i>Spiroplasma</i>	141	350	1,605	890	440	1,120	2,422	567
<i>Cardinium</i>	-	1	2	-	-	1	2	-
Group	W-S-							
Line	1	2	3	4				
Total	4,975	20,054	10,241	8,906				
<i>Wolbachia</i>	3	4	2	1				
<i>Spiroplasma</i>	-	-	-	-				
<i>Cardinium</i>	-	1	-	-				
Mite strain				Inducer strain				
Group	C+S+			C-S-				
Line	5	6	7	8	5	6	7	8
Total	26,384	18,367	48,325	28,966	39,276	22,648	30,608	18,702
<i>Wolbachia</i>	7	5	108	204	100	110	66	63
<i>Spiroplasma</i>	1,159	165	3,555	1,467	1	1	-	6
<i>Cardinium</i>	7,443	3,643	19,744	7,904	2	1	3	-


survived until the end of the experiment (4 days) compared to both W-S+ and W-S- mites, while survival did not differ between the latter two groups, with the exception of line 2 (FIGURE S5.2a). There was no statistical difference in number of eggs between the mites from each group when we calculated the total number of eggs produced per number of females that were initially put on each leaflet (5 mites), (FIGURE 5.3a). This suggests that the fewer W-S+ and W-S- mites that did survive produced more eggs. Hence, we determined the total number of eggs produced per female that survived, and here the factor ‘bacterial group’ had an overall significant effect on number of eggs ( $F_{2,207} = 3.14$ ,  $P = 0.045$ ) (data not shown in graph). However, *post hoc* tests with Holm correction for multiple testing indicated no

significant difference between surviving W+S+ mites laying fewer eggs than W-S+ ( $z = -2.00$ ,  $P = 0.09$ ) and W-S- ( $z = -2.30$ ,  $P = 0.06$ ).


Between the two groups of the inducer strain, significantly more mites of the C-S- group generally survived until the end of the experiment (FIGURE 5.2b), but this was not consistent among the lines (FIGURE S5.2b). The number of eggs between C+S+ and C-S- mites was also significantly different: overall, C-S- mites produced more eggs than C+S+ mites (FIGURE 5.3b). This difference was significant for line 5 ( $F_{1,29} = 14.71$ ,  $P < 0.001$ ), but not for line 8 ( $F_{1,32} = 1.21$ ,  $P = 0.27$ ), although both lines followed the same trend (FIGURE S5.3). The oviposition calculated per surviving mite did not differ between the C+S+ and C-S- groups ( $F_{1,76} = 3.34$ ,  $P = 0.07$ ) (data not shown in graph), possibly due to the lower survival of C+S+ mites. However, the data from the oviposition assays with the inducer mites were inconclusive, because the C-S- subline of line 6, as well as both sublines of line 7, went extinct before the performance assays were completed.

### *Iib. Effects of Wolbachia, Cardinium and Spiroplasma on the mite's transcriptome*

In the microarray analysis, we found significant differences in expression levels of mite genes across the five groups (W+S+, W-S+, W-S-, C+S- and C-S-). Even though the microarray represented only 9,424 (51%) of the in total 18,414 *in silico* predicted protein-coding sequences (Grbic *et al.*, 2011), we identified 201 (1.1% of the total) differentially expressed genes (DEGs; absolute fold change (FC) > 1, Ben-



**FIGURE 5.2.** Survival, migration and mortality in two strains of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain *Wolbachia* (W), *Spiroplasma* (S) and/or *Candidatus Cardinium* (C); (a) Plant defence suppressor strain DeLier-1 with three mite groups: W+S+, W-S+ and W-S-. (b) Plant defence inducer strain Santpoort-2 with two mite groups: C+S+ and C-S-. Different letters above the bars indicate significant differences at a level of  $P \leq 0.05$ , after applying a generalized linear mixed model followed by Tukey multiple comparisons with Holm adjustment. n.s. = not significant



**FIGURE 5.3.** Reproductive performance (number of eggs produced per female in four days) in two strains of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain *Wolbachia* (W), *Spiroplasma* (S) and/or *Candidatus Cardinium* (C); (a) plant defence suppressor strain DeLier-1 with three mite groups: W+S+, W-S+ and W-S-. (b) plant defence inducer strain Santpoort-2 with two mite groups: C+S+ and C-S-. Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside of this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$ , after applying a linear mixed model followed by Tukey multiple comparisons with Holm adjustment. n.s. = not significant

jamini-Hochberg false discovery rate (FDR)-adjusted  $P \leq 0.05$ ) across all groups. Overall, differences were relatively small, i.e., absolute FCs ranged from +1.1 to +5.2 for upregulated genes and from -1.1 to -3.4 for downregulated genes respectively, with only 23 genes displaying an absolute FC  $> 2$  ( $P \leq 0.05$ , after  $P$ -value adjustment). The most pronounced transcriptional differences were found between W+S+ and W-S+ mites of the suppressor strain with 152 DEGs (FIGURE S5.4), 30 of which were up-regulated (see TABLE S5.3a for the top-20) and 122 were down-regulated (see TABLE S5.3b for the top-20) in W-S+ mites. Comparing the transcriptome of W+S+ and W-S- mites resulted in 50 DEGs: 14 of them were up-regulated (TABLE S5.4a), while 36 were down-regulated (see TABLE S5.4b for the top-20) in W-S- mites. Out of the 50 DEGs, 29 also differed between W+S+ and W-S+ mites (FIGURE S5.4), hence the other 21 correlated with the presence of *Spiroplasma*. The transcript levels of only three genes (2 up-, 1 down-regulated in W-S-) differed significantly between W-S- and W-S+ mites (TABLE S5.5, FIGURE S5.4). Based on the principal component analysis, one of the inducer C-S- samples was identified as an outlier and therefore excluded from further analysis. Using the remaining samples, we identified 33 DEGs between the two groups of the inducer strain C+S+ and C-S-; 23 were higher expressed in C-S- mites (TABLE S5.6a), while 10 were lower expressed (TABLE S5.6b). Five of the 33 DEGs were also significantly affected by *Wolbachia* and/or *Spiroplasma* in the suppressor mites (TABLE S5.6a). Many of the 201 DEGs, i.e., 36 genes (17.9%), had no homology with any of the genes characterized so far, hence their function in the

mite physiology remains unknown. Among the DEGs in both mite strains, those encoding proteins involved in digestion, metabolic detoxification (phase I), binding/conjugation of xenobiotics (phase II detoxification), transport of xenobiotics (phase III detoxification) or encoding putative (salivary gland-derived) secreted proteins were highly abundant (TABLE 5.2). Genes related to immune-responses (Grbic *et al.*, 2011), of which the expression is associated with the presence of *Wolbachia* in dipterans (Kambris *et al.*, 2009; Moreira *et al.*, 2009), were not differentially expressed among the mite groups (data not shown).

### ***III. Effects of spider mite-associated Wolbachia, Cardinium and Spiroplasma on mite-plant interactions***

#### ***IIIa. Effects of mite-associated Wolbachia, Cardinium and Spiroplasma on tomato induced responses***

For the suppressor mite infested plants, the most consistent differences in phytohormone profiles were found between leaflets infested with W-S+ mites and those infested with W-S- and/or W+S+ mites (FIGURES 5.4 and S5.5). The (combined) presence of *Wolbachia* and/or *Spiroplasma* correlated with phytohormone levels in tomato as follows. The accumulation of the JA-precursor OPDA was correlated with mite-associated bacteria (FIGURE 5.4b). Feeding by W+S+ mites induced the accumulation of OPDA, as did feeding by W-S- mites. However, leaflets infested with W-S- mites accumulated significantly lower amounts of OPDA than W+S+ mites. Leaflets infested with the third group of mites, W-S+, accumulated significantly less OPDA than W-S- and W+S+ infested leaflets and did not accumulate more OPDA than uninfested control leaflets. This clear pattern was detected in leaflets infested with all four suppressor mite lines (FIGURE S5.5a). A similar pattern was found for several other oxylipins as well, i.e., lowest hormone levels in W-S+ compared to W+S+ and W-S- infested leaflets. First, for the JA-precursor OPC8 (FIGURE 5.4c), which is the direct downstream product of OPDA, and to a lesser extent for OPC6 (FIGURE S5.5b), but not for JA itself (FIGURE S5.5c), nor for its biologically active conjugate JA-Ile (FIGURES 5.4d and S5.5d). Second, we found the pattern for traumatic acid (FIGURE 5.4e) and 9-OH-traumatins (FIGURES 5.4f and S5.5e), which are C<sub>12</sub>-derivatives of the HPL pathway, derived from the same 13-HPOT pool as JA (FIGURE S5.1). The accumulation of 13-HPOT, one of the first intermediates in the biosynthesis process of oxylipin signalling molecules (FIGURE S5.1) (Wasternack, 2007), was not correlated with mite-associated bacteria (FIGURE 5.4a).

The levels of SA in infested leaflets followed a pattern that appeared opposite of that of OPDA: amounts of SA in leaflets that were infested with W-S+ mites were significantly higher than in leaflets infested with either W-S- or W+S+ mites, but were not higher than in uninfested control leaflets (FIGURES 5.4g and S5.5f). The accumulation of ABA was not affected by infestations with mites from any of the suppressor strain groups (FIGURE 5.4h).

**TABLE 5.2.** Overview of differentially expressed genes (DEGs; absolute fold change > 1;  $P \leq 0.05$ , after  $P$ -value adjustment) in adult female *Tetranychus urticae* suppressor (DeLier-1) and inducer (Santpoort-2) mites after selective removal of their associated bacteria *Wolbachia* sp. (W), *Spiroplasma* sp. (S) and/or *Candidatus Cardinium* (C) and feeding from tomato (*Solanum lycopersicum*) for seven days. Shown are the DEGs encoding proteins that are involved in digestion or detoxification, or are (putatively) secreted.

▲, up-regulated in mites of the first-mentioned bacterial group; ▼, down-regulated in mites of the first-mentioned bacterial group; +, present; -, absent


Biological process	Gene family	Suppressor strain			Inducer strain	Unique	Total*
		W-S+ vs W+S+	W-S- vs W+S+	W-S- vs W-S+	C-S- vs C+S+		
		▲/▼	▲/▼	▲/▼	▲/▼		
<b>Digestion</b>							
	Cathepsin cysteine peptidases	0 / 7	0 / 2	-	-	7	49
	Legumain cysteine peptidases	0 / 5	0 / 2	-	-	5	12
	Cystatins	-	2 / 0	1 / 0	2 / 0	2	11
<b>Detoxification</b>							
Phase I	Cytochrome P450 monooxygenases	1 / 4	0 / 3	-	2 / 0	9	59
	Carboxyl/choline esterases (CCEs)	0 / 7	0 / 1	-	1 / 0	8	51
	Intradial ring-cleavage dioxygenases	0 / 2	-	-	-	2	15
Phase II	Lipocalins	0 / 6	-	-	-	6	41
	Glutathione S-transferases (GSTs)	0 / 2	0 / 1	-	-	2	26
	UDP-glycosyltransferases (UGTs)	-	-	-	1 / 0	1	51
Phase III	ABC-transporters	0 / 1	0 / 1	-	1 / 0	3	88
	MFS-transporters	0 / 1	-	-	1 / 1	3	97
	total	1 / 35	2 / 10	1 / 0	8 / 1	48	
<b>Unknown (secreted)</b>							
	Unknown (various)	2 / 9	2 / 6	1 / 1	3 / 1	16	
	total DEGs	152	50	3	33	201	

All *T. urticae* multi-gene families implicated in digestion and detoxification have been identified and/or manually annotated (Grbic *et al.*, 2011; Santamaria *et al.*, 2012; Dermauw *et al.*, 2013; Ahn *et al.*, 2014; Wybouw *et al.*, 2015). \*indicates the total number of genes (of the indicated gene family) that is represented on the microarray

The phytohormone profiles of inducer mite (C+S+ and C-S-) infested leaflets were as follows (see FIGURES 5.5 and S5.6). The levels of OPDA, JA-Ile and SA were induced by C+S+ and C-S- mites compared to uninfested controls, but did not differ between the two groups. However, two of the C<sub>12</sub>-derivatives of the HPL pathway, i.e., traumatic acid (FIGURES 5.5c and S5.6c) and 9,12-OH-(10E)-dodecenoic acid (FIGURE 5.5d), accumulated to higher amounts in C-S- infested than in C+S+ infested leaflets. ABA levels followed the same pattern (FIGURES 5.5f and S5.6d).


Previously, suppression of plant defences by spider mites was shown to act downstream of phytohormones (Alba *et al.*, 2015). We therefore augmented the phytohormone data with the expression data of downstream marker genes (FIGURE S5.1) with qRT-PCRs. Upon infestation with suppressor mites, the amounts of OPDA

Mite-associated bacteria




in W-S+ infested leaflets differed from those in W+S+ and W-S- infested leaflets, with the latter having intermediate levels (FIGURE 5.4b). Transcript levels of *AOS1*, which encodes an enzyme involved in the biosynthesis of OPDA, were higher in suppressor mite-infested leaflets compared to uninfested controls, but there was no difference among the mite groups (FIGURE 5.6a). In contrast, expression of *OPR3*, which acts directly downstream of OPDA in the JA biosynthesis pathway, was significantly higher in W-S+ than in W+S+ or W-S- infested leaflets (FIGURES 5.6b and S5.7a). The same pattern was found for multiple putative OPDA-responsive genes (ORGs); *TWI-1*, *ADH*, *GST6* and *GRX* (FIGURES 5.6c-f and S5.7b-e, respectively). The expression levels of these ORGs were thus negatively correlated with OPDA amounts and positively with SA amounts, while the coefficient of determination ( $R^2$ ) values were similar (FIGURE S5.8), except for expression levels of *ADH*, which correlated with OPDA ( $R^2 = 0.42$ ,  $P < 0.0001$ ) but not with SA ( $R^2 = 0.09$ ,  $P = 0.22$ ). Analogous to the overall induction of JA-Ile by W+S+, W-S+ and W-S- mites (FIGURE 5.4d), expression of the JA-defence marker genes *TD-2* (FIGURE 5.6g) and *PI-IIc* (FIGURE 5.6h) were induced in all three groups compared to the control, but did not differ among them. However, transcripts of *JIP-21* accumulated to significantly higher levels in W-S- infested leaflets than in those of W+S+ and W-S+ (FIGURES 5.6i and S5.7f). As for the SA-defence marker genes, *PR-P6* transcript accumulation was higher in W-S+ leaflets than in W+S+ and W-S- leaflets (like SA itself) (FIGURE 5.6j). Here it has to be noted that this *PR-P6* expression pattern was visually clear in two lines, but not in the other two (FIGURE S5.7g). We did not find any significant differences in the expression of *PR-1a* in leaflets infested with the different groups of suppressor mites, nor between infested and uninfested leaflets (FIGURE 5.6k). Accordingly, SA amounts showed a weak but significant correlation with expression levels of *PR-P6* (FIGURE S5.9a) and did not correlate with *PR-1a* transcript levels (FIGURE S5.9b).

**FIGURE 5.4.** Phytohormone amounts in tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with a plant defence suppressor strain DeLier-1 of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain the bacteria *Wolbachia* (W) and *Spiroplasma* (S). Control plants were not infested. Phytohormones for which we assayed include (a) 13(*S*)-hydroperoxyoctadecatrienoic acid (13-HPOT), (b) 12-oxo-phytodienoic acid (OPDA), (c) 3-oxo-2-(2-pentenyl)-cyclopentane-1-octanoic acid (OPC8), (d) jasmonic acid-isoleucine (JA-Ile), (e) traumatic acid, (f) 9-OH-traumatol, (g) free salicylic acid (SA), (h) abscisic acid (ABA); Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$ , after applying a linear mixed model followed by Tukey multiple comparisons with Holm adjustment. Phytohormone amounts are presented as nanogram (ng) per gram fresh leaf weight (g FW). Compounds for which we did not have a pure standard are presented as ion counts (IC / g FW). n.s. = not significant



**FIGURE 5.5.** Phytohormone amounts in tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with a plant defence inducer strain Santpoort-2 of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain the bacteria *Candidatus* Cardinium (C) and *Spiroplasma* (S). Control plants were not infested. Phytohormones for which we assayed include (a) 12-oxo-phytodienoic acid (OPDA), (b) jasmonic acid-isoleucine (JA-Ile), (c) traumatic acid, (d) 9,12-OH-(10E)- dodecenoic acid, (e) free salicylic acid (SA) and (f) abscisic acid (ABA) (f); Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside of this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$ , after applying a linear mixed model followed by Tukey multiple comparisons with Holm adjustment. Phytohormone amounts are presented as nanogram (ng) per gram fresh leaf weight (g FW). Compounds for which we did not have a pure standard are presented as ion counts (IC / g FW). n.s. = not significant




**FIGURE 5.6.** Normalized expression (NE) of plant defence related genes obtained via qRT-PCR in tomato (*Solanum lycopersicum*) leaflets after infestation (7 days) with a plant defence suppressor strain DeLier-1 of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain the bacteria *Wolbachia* (W) and *Spiroplasma* (S). (a) *Allene oxide synthase 1* (*AOS1*), (b) *12-oxophytodienoate Reductase 3* (*OPR3*), (c) *Tomato wound-induced 1* (*TWI-1*), (d) *Alcohol dehydrogenase* (*ADH*), (e) *Glutathione S-transferase 6* (*GST6*), (f) *Glutaredoxin* (*GRX*), (g) *Threonine Deaminase-2* (*TD-2*), (h) *Proteinase Inhibitor IIc* (*PI-IIc*), (i) *Jasmonate-inducible protein-21* (*JIP-21*), (j) *Pathogenesis-related protein 6* (*PR-P6*), (k) *Pathogenesis-related protein 1a* (*PR-1a*); Gene expression levels were normalized to the levels of tomato actin (TABLE S5.2). Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside of this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$ , after applying a linear mixed model followed by Tukey multiple comparisons with Holm adjustment. n.s. = not significant.

In the leaflets infested with the inducer mites, none of the JA-defence marker genes (*TD-2*, *PI-I1c* and *JIP-21*) were differentially expressed ( $P < 0.05$ ) between C+S+ and C-S- infested leaflets (FIGURE 5.7a-c). Interestingly, transcripts of both SA-defence marker genes, *PR-P6* and *PR-1a*, were more abundant in C+S+ infested leaflets than in C-S- infested ones (FIGURES 5.7d-e and S5.10a,b).


### IIIb. Effects of *Wolbachia*, *Cardinium* and *Spiroplasma* on the amount of feeding damage inflicted by the spider mites

A possible explanation for the observed differences in phytohormone and gene expression levels, which were correlated with the presence of mite-associated bacteria might be of behavioural nature, i.e., mites that feed more cause more damage to the plant and therefore might elicit stronger induced responses. To test if the magnitude of induced responses was correlated with the amount of feeding of the mites, we quantified the amount of spider mite-inflicted feeding damage (recognizable as chlorotic spots) on the same leaflets that were used for phytohormone extractions and



**FIGURE 5.7.** Normalized expression (NE) of plant defence related genes obtained via qRT-PCR in tomato (*Solanum lycopersicum*) leaflets after infestation (7 days) with a plant defence inducer strain Santpoort-2 of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain the bacteria *Candidatus* *Cardinium* (C) and *Spiroplasma* (S). (a) *Threonine Deaminase-2* (*TD-2*), (b) *Proteinase Inhibitor I1c* (*PI-I1c*), (c) *Jasmonate-inducible protein-21* (*JIP-21*), (d) *Pathogenesis-related protein 6* (*PR-P6*), (e) *Pathogenesis-related protein 1a* (*PR-1a*); Gene expression levels were normalized to the levels of tomato actin (TABLE S5.2). Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside of this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$ , after applying a linear mixed models followed by Tukey multiple comparisons with Holm adjustment. n.s. = not significant

tomato RNA isolation for qRT-PCRs. Overall, there was no significant difference in feeding damage between the three groups (W+S+, W-S+, W-S-) of suppressor mites (FIGURE 5.8a). In contrast, there was a clear difference in the amount of feeding damage caused by the two groups of inducer mites and in the type of damage these inflicted. Not only did C-S- mites feed significantly more than the C+S+ mites (FIGURE 5.8b), feeding by C+S+ mites resulted in rusty red/brown ‘scars’ on the leaflets (FIGURE 5.8c), while those infested with C-S- mites had clear white scars (FIGURE 5.8d). This feeding damage phenotype was evident for three out of four lines, it was less clear for line 7 (FIGURE S5.11).




**FIGURE 5.8.** Feeding damage (mm<sup>2</sup>) on tomato (*Solanum lycopersicum*) leaflets inflicted by two strains of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain the bacteria *Wolbachia* (W), *Spiroplasma* (S) and/or *Candidatus Cardinium* (C). (a) Plant defence suppressor strain DeLier-1 with three mite groups: W+S+, W-S+ and W-S-. (b) Plant defence inducer strain Santpoort-2 with two mite groups: C+S+ and C-S-. Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside of this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$ , n.s. = not significant after applying a linear mixed model. (c) Typical rusty red/brown scars inflicted by feeding of C+S+ mites of the inducer strain and (d) typical white scars inflicted by feeding of C-S- mites of the inducer strain.

## DISCUSSION

In this study we showed that *T. urticae* females of the plant suppressor strain DeLier-1, contained *Wolbachia* (W) and *Spiroplasma* (S) bacteria, while those from the plant defence inducer strain Santpoort-2, harboured *Candidatus Cardinium* (C) and also *Spiroplasma* (i.e., the same OTU as in suppressor mites). We determined the effects of these well-known reproductive parasites on (a) the spider mites and (b) on induced plant responses, after removing them via antibiotic treatment. In the spider mites, we showed I) that the presence of the (endo)symbionts correlated positively with the survival, but not oviposition of the suppressor mites, while it correlated negatively with both survival and oviposition of the inducer mites. II) We showed that a subset of spider mite genes that are predicted to be predominantly involved in digestion and detoxification processes, as well as genes encoding putative secreted (salivary gland) proteins, were differentially expressed in mites, with opposite trends occurring in the suppressor versus the inducer mites. III) Plant responses to mite infestations differed between the mite groups that did or did not harbour *Wolbachia*, *Cardinium* and/or *Spiroplasma*. With the suppressor strain the most prominent result was that the amount of OPDA was lowest in W-S+ infested compared to W-S- and W+S+ leaflets, while SA followed an opposite pattern (highest amount in W-S+ infested leaflets). With the inducer strain the most interesting result was that SA-marker genes were expressed at lower levels in C-S- inducer mites, than in C+S+ infested leaflets. Moreover, the antibiotics-treated inducer mites (C-S-) consumed significantly more plant material and caused visually different feeding scars compared to the non-treated (C+S+) mites. Remarkably, well-known upstream-downstream relationships between phytohormones and expression of their marker genes were in some cases not found in our study, as discussed below. Finally, we showed IV) the combined presence of *Wolbachia* and *Spiroplasma* has consequences for mite gene expression and induced plant responses. Together, our results indicate that different (endo)symbiotic bacteria may have distinct consequences for their host, as is summarized in FIGURE 5.9 and discussed below.

Importantly, it has to be noted that although the antibiotic treatments resulted in the (near-) complete removal of *Wolbachia*, *Cardinium* and/or *Spiroplasma* from the mites and we found their absence/presence to correlate well with certain parameters (e.g., mite performance, mite transcriptomic responses and induced plant responses), various other bacteria were (randomly) present in/on antibiotics-treated and non-treated mites as well. These bacterial strains were not restricted to a certain group of

**FIGURE 5.9.** Schematic and simplified overview of the most important findings of this study. Adult female spider mites (*Tetranychus urticae*) from the plant defence suppressor strain DeLier-1 and the defence inducer strain Santpoort-2 were treated with antibiotics to remove their associated bacteria *Wolbachia* sp. (W), *Spiroplasma* sp. (S) and/or *Candidatus Cardinium* (C), after which the indicated mite and plant (tomato; *Solanum lycopersicum*) parameters were assayed. For a more detailed description we refer to the Results and Discussion sections. med, intermediate; +, bacteria present; -, bacteria absent.



mites and are thus not likely to be responsible for the effects that we found between the mite groups. In addition, tetracycline treatments may have had effects on the mites other than the removal of bacteria. Direct toxic effects of tetracycline, such as inhibition of mitochondrial functioning, are unlikely to play a role in our study, because we started experiments at least 15 generations after the antibiotics treatments. Selective effects, in which tetracycline treatment has selected e.g., for more toxin resistant mites, cannot be ruled out. However, subjectively we did not notice higher mortality in the tetracycline-treated lines compared to non-treated lines.

### ***I. Effects of Wolbachia, Spiroplasma and Cardinium on spider mite performance***

When we investigated the performance of suppressor mites in a four days trial, survival was highest in mites containing *Wolbachia* and *Spiroplasma*. The presence or absence of *Spiroplasma* did not affect mite survival. The *Wolbachia*-mediated effect on suppressor mite survival could have resulted from direct effects on host gene transcript levels, from indirect effects via induced or suppressed plant defences, or a combination hereof, as discussed below. Surprisingly, we did not find that mite fecundity was higher in the presence of *Wolbachia* than in its absence, indicating that the *Wolbachia*-containing survivors may have laid eggs at a lower frequency than mites without *Wolbachia*. Future experiments should focus on the lifetime production of eggs by mites with and without symbionts, since differences in survival are expected to have a significant effect on reproductive performance in the long run.

The performance of inducer mites was negatively correlated with the presence of *Cardinium* and *Spiroplasma*. Both mite survival and fecundity were higher for C-S- mites than for the C+S+ mites. Unfortunately, the performance assay for the inducer strain could not be completed, because the C-S- subline of line 6, as well as both sublines of line 7, went extinct before the assays started. However, the results of the plant infestation assay and mite transcriptome analysis, which were performed with all lines, were all in line with our hypothesis that *Cardinium* and/or *Spiroplasma* have a negative influence on the performance of the inducer mites: expression of SA-marker genes in C-S- infested leaflets was reduced compared to the C+S+ infested leaflets, even though C-S- mites inflicted almost twice as much damage. Furthermore, on the microarray the expression of digestion and detoxification genes was higher in C-S- mites than in C+S+ ones.

Previous studies indicate that fitness effects of reproductive parasites may strongly depend on the genotype of symbiont and host and whether symbionts optimize their prevalence in a population by reproductive manipulation and/or by positively affecting host fitness (Bordenstein & Werren, 2000; Fry *et al.*, 2004; Zug & Hammerstein, 2015). Accordingly, the reported effects of *Cardinium*, *Wolbachia* and *Spiroplasma* on host fitness are diverse. Whereas *Cardinium* and/or *Spiroplasma* negatively affected the fitness of mites in our experiments, an earlier study with a different *T. urticae* strain found that *Cardinium* did not alter survival or

egg hatchability of its host, nor did it manipulate its reproduction (Gotoh *et al.*, 2007). Studies of other arthropods indicate that infection with *Cardinium* can have all possible effects on fitness parameters of its host (Weeks & Stouthamer, 2004; Ros & Breeuwer, 2009; White *et al.*, 2011; Stefanini & Duron, 2012). The same is true for *Wolbachia* infection, which seemed to positively affect *T. urticae* fitness in our study. In *T. urticae*, *Wolbachia* was previously shown to have positive, negative or no effect on fitness parameters of this mite species (Vala *et al.*, 2000; Perrot-Minnot *et al.*, 2002; Xie *et al.*, 2011). Such varying effects of *Wolbachia* are also known in *Drosophila melanogaster* (Hoffmann *et al.*, 1994; Fry *et al.*, 2002). Effects of *Spiroplasma* on animal host fitness parameters were previously found to be negative or neutral (Ebbert, 1991; Fukatsu *et al.*, 2001; Montenegro *et al.*, 2006; Anbutsu & Fukatsu, 2011).

## **II. Effects of *Wolbachia*, *Spiroplasma* and *Cardinium* on the mite's transcriptome**

Previous studies have established that (endo)symbiotic bacteria can affect gene expression of their host. Some data has been published on *Cardinium* (Nakamura *et al.*, 2011) and *Spiroplasma* (Hutchence *et al.*, 2011), but most research has focused on *Wolbachia* (Kambris *et al.*, 2009; Moreira *et al.*, 2009; Hussain *et al.*, 2011; Zhang *et al.*, 2013; Mayoral *et al.*, 2014).

In *T. urticae*, *Wolbachia* has been shown to impact gene expression of its host: sex-specific differences in transcript levels of up to 251 protein-coding genes (Zhang *et al.*, 2015), as well as up to 91 miRNAs (Rong *et al.*, 2014) were found when comparing *Wolbachia*-infected and uninfected female mites. In our microarray analysis with antibiotics-treated and non-treated adult female suppressor and inducer mites, we identified 201 differentially regulated spider mite genes among the mites from all groups (i.e., W+S+, W-S+, W-S- and C-S-, C+S+), i.e., roughly 1% of the 18,414 *in silico* predicted protein-coding sequences (Grbic *et al.*, 2011) of which about 50% were present on our array. Consistent with the data of Zhang *et al.* (2015), none of these mite genes putatively involved in immunity (Grbic *et al.*, 2011) were differentially regulated. This is in agreement with previous findings that the immune system is not activated in response to *Wolbachia* in arthropod host species that are naturally infected by this bacterium, but can be activated in species that are naturally *Wolbachia*-free and are experimentally infected, as was shown in *Aedes aegypti* (Rances *et al.*, 2012; Zug & Hammerstein, 2015).

Among the DEGs, there was a high abundance of genes that implicated the mite-associated bacteria in direct or indirect regulation of digestive and detoxification processes of their host. This is again similar to the findings of Zhang *et al.* (2015). However, when comparing the gene IDs of the DEGs of our study and that of Zhang *et al.* (2015) we found only five genes with an identical ID. Moreover, according to the data of Zhang *et al.* (2015), two of these five genes were differentially regulated in males but not females, while the remaining three genes were inversely regulated on our arrays. Notably, Zhang *et al.* (2015) worked with a

Chinese mite strain that was sampled from bean, whereas we worked with Dutch mite lines that were sampled from tomato, while both data sets were mapped onto the *T. urticae* London genome (Grbic *et al.*, 2011) for identification of DEGs. Possibly this is not accurate enough for identifying the homologous loci, but only for identifying homologous gene families. Additionally, *Wolbachia* strains may have differed between the two studies or *Wolbachia* may interact differently with mites on bean than with mites on tomato.

Absence of only *Wolbachia* from suppressor mites (W-S+) correlated with reduced transcript accumulation of 35 out of 36 DEGs, which belong to 9 gene families involved in digestion and in detoxification of xenobiotics. This result indicates that *Wolbachia* may have a direct effect on mites and may impact their capability to digest food and their ability to catabolize secondary metabolites. It has been shown before for various strains of *Wolbachia* that they can metabolically assist their host, in particular in filarial nematodes in which *Wolbachia* represents an obligate nutritional mutualist, for instance by providing its host with the energy source ATP (Darby *et al.*, 2012) or essential coenzymes, the cofactor heme and nucleotides, in return for amino acids (Foster *et al.*, 2005). With respect to arthropod hosts, *Wolbachia* is indispensable for the bedbug *Cimex lectularius*, because it provides essential B vitamins (Hosokawa *et al.*, 2010). Furthermore, in the parasitic wasp *Asobara tabida* and the fruitfly *D. melanogaster*, *Wolbachia* influences iron (and hence redox) homeostasis, which benefits its host in perturbed iron environments (Brownlie *et al.*, 2009; Kremer *et al.*, 2009). Hence, in suppressor mites, *Wolbachia* possibly performs early catabolic steps of (plant) nutrients or secondary metabolites and delivers intermediates which the mite can further process using the genes listed in TABLE 5.2. If so, this may explain why Zhang *et al.* (2015) observed similar functional processes but different loci in mites in correlation with the absence or presence of *Wolbachia*. In this scenario, *Wolbachia* may act as a nutritional mutualist when mites feed from tomato, which might explain the higher survival of suppressor mites that contained *Wolbachia*.

Strikingly, our data reveals fewer DEGs (roughly 65% less) when comparing W+S+ mites with W-S- (instead of W-S+) mites, while transcriptomic differences between W-S+ and W-S- mites were virtually absent. This suggests a direct antagonistic interaction between *Wolbachia* and *Spiroplasma*. Unfortunately we did not obtain a W+S- line to substantiate this. Possibly, part of the *Wolbachia*-associated transcriptional response of the mite is a direct consequence of the presence of *Spiroplasma*. It has been suggested before that *Spiroplasma* negatively affects the abundance of *Wolbachia* within the same host (Goto *et al.*, 2006). If so, *Wolbachia* may chemically combat and/or constrain *Spiroplasma* and *vice versa*, leaving the mite to clean up ‘the waste’ using the genes listed in TABLE 5.2.

The transcriptome of *Cardinium*- and *Spiroplasma*-free inducer mites also differed in the expression of genes involved in digestion, detoxification and transport of xenobiotics as compared to the mites that still harboured both bacteria. However,

we found fewer DEGs and the directionality of transcriptional change was largely opposite from what we found for the suppressor strain. Expression of cysteine peptidase and lipocalin encoding genes for instance, which responded strongest to the absence of *Wolbachia*, was unaffected in C-S- inducer mites. In fact, the only overlap with suppressor mites was the increased transcript abundance of two cystatin encoding genes in C-S- versus C+S+ mites. When compared to W+S+ mites, these genes were also induced in W-S- mites, but not in W-S+ ones. Together, this suggests that these cystatin genes respond to the presence of *Spiroplasma*. Other DEGs (encoding two CYPs, a CCE, an UGT, an ABC- and two MFS-transporters) seem to respond to the presence of *Cardinium* in inducer mites, mainly because they followed a trend opposite to when *Wolbachia* was present in the suppressor strain: all but one MFS-transporter were up-regulated in C-S- mites compared to C+S+ ones. We did not manage to obtain inducer strain lines with only *Spiroplasma* or only *Cardinium*, therefore it remains difficult to pinpoint the effects of each bacterial strain more precisely. All in all, the transcriptome data suggest *Cardinium* to have a negative effect on the mite, possibly by reducing the mite's capability to detoxify and transport harmful secondary metabolites. This is consistent with our results for mite performance, in which C+S+ mites preformed worse than C-S- mites.

Among the 201 DEGs, the ones encoding (putative) secreted proteins were also highly abundant. Again this is in agreement with the data from Zhang *et al.* (2015). Four of our identified DEGs were predicted to encode proteins that are secreted from the mite salivary gland and are thus thought to indeed directly interact with the plant. Moreover, *Wolbachia* was found to be localized in the gnathosoma (mouth and feeding parts) of both male and female *T. urticae* (Zhao *et al.*, 2013), from where the proteins could also be secreted to the plant. Mite salivary gland-secreted proteins might be recognized by the plant and induce anti-herbivore defences (i.e., act as elicitors) or they may act as effectors and interfere with plant defences (Villarreal *in prep*). Pathogens (Dou & Zhou, 2012; Lo Presti *et al.*, 2015) and nematodes (Goverse & Smant, 2014) are well-known for their use of effector molecules to subvert plant immunity and while it is likely that many arthropods have evolved them as well, empirical evidence is still scarce (Kant *et al.*, 2015). In some cases defence suppression and the secretion of effectors by arthropods was reported to depend on associated microbes. For instance, the leafhopper *Macrostelus quadrilineatus* vectors phytoplasma bacteria, that after being transmitted to the host plant, produce and secrete molecules that suppress JA-defences, which is beneficial for the leafhopper (Sugio *et al.*, 2011). Recently, the saliva of the whitefly *Bemisia tabaci* was shown to contain a small (< 3 kDa) non-proteinaceous compound(s) responsible for the effective suppression of JA-defences via manipulation of JA-SA crosstalk (Su *et al.*, 2015). Defence suppression was only detected with saliva from whiteflies that harboured the endosymbiont *Hamiltonella defensa*, substantiating a relationship between the presence of the bacterium and the production of the effector(s) (Su *et al.*, 2015). Our microarray data indicates that mite-associated

bacteria can affect the host's transcript accumulation of genes encoding secreted (salivary gland) proteins and may in this way affect plant defence responses.

Together, the transcriptional differences that we observed between mites with *Wolbachia*, *Cardinium* and/or *Spiroplasma* bacteria or without them, may have several, not mutually exclusive, causes. First of all, they may reflect processes in the host associated with suitability for bacterial colonization, as was shown in *A. aegypti* (Hussain *et al.*, 2011) and this could be a direct consequence of bacterial manipulation of host transcripts e.g., via RNA-interference (RNAi). Secondly, it may reflect responses of the host to bacterial metabolites, either plant-derived or not, eligible for further metabolization or catabolization. Such metabolites may be nutritional, may be plant toxins or be generated by multiple bacterial strains competing for space or nutrients. Thirdly, they may reflect a response of the mite to differences in plant tissue quality, e.g., due to differentially induced plant responses (discussed in the next section), directly or indirectly caused by differences in the host's bacterial composition and possibly mediated by alterations in the host's secretome.

### ***III. Effects of Wolbachia, Cardinium and Spiroplasma on tomato induced responses***

When we investigated the effect of *Wolbachia* and *Spiroplasma* infection of the suppressor strain on tomato induced responses, the most striking finding was that the JA-precursor OPDA did not significantly accumulate in W-S+ infested leaflets compared to uninfested leaflets and W+S+ infested leaflets, while OPDA levels were intermediate in leaflets infested with W-S- mites. Thus, the presence of *Wolbachia* in suppressor mites is correlated with enhanced OPDA accumulation, while the presence of *Spiroplasma* in these mites is correlated with suppression of OPDA accumulation. Note that the OPDA (or any other plant response) phenotype cannot be explained by the amount of damage inflicted due to mite feeding, as this was equal among all four groups. Expression of *OPR3* (Strassner *et al.*, 2002) was highest in W-S+ infested leaflets, thus an increased conversion rate of OPDA might explain the reduced OPDA accumulation in the respective leaflets. However, the immediate downstream product of OPDA, OPC8 showed the same pattern as OPDA. Surprisingly, the end product of the oxylipin pathway, JA-Ile, which is considered the main biologically active molecule (Fonseca *et al.*, 2009; Wasternack & Hause, 2013), did not show any pattern that was correlated with bacterial presence, neither did JA-responsive defence marker genes. In addition, the accumulation profiles of two C<sub>12</sub>-derivatives of the hydroperoxide lyase (HPL)-pathway, traumatic acid and 9-OH-traumatol, closely resembled that of OPDA across the treatments. This is notable, because both the JA and the HPL pathway represent branches of the LOX pathway that use the same substrate, i.e., 13-HPOT (FIGURE S5.1; Wasternack, 2007). Moreover, levels of the substrate 13-HPOT as well as transcript levels of *AOS1*, which encodes the first of two enzymes

responsible for turnover of 13-HPOT into OPDA (Howe *et al.*, 2000; Sivasankar *et al.*, 2000), were not correlated with the presence of bacteria. Although gene transcript levels do not necessarily reflect protein levels and/or enzyme activity (Sullivan & Green, 1993), it remains unknown how accumulation of some wound hormones is affected by spider mite-associated bacteria, while other oxylipin pathway intermediates/products are not. One explanation for the altered OPDA amounts might be that *Wolbachia* and *Spiroplasma* differentially affect conjugation of OPDA with other cellular compounds. In Arabidopsis (*Arabidopsis thaliana*) and tobacco (*Nicotiana tabacum*) for example, OPDA is known to form conjugates with glutathione (GSH) (Davoine *et al.*, 2006; Mueller *et al.*, 2008). Although we did not detect OPDA-GSH in tomato leaflets, it is possible that other conjugates (that we did not search for) were formed.

Unexpectedly, expression levels of putative OPDA-responsive genes (ORGs) in suppressor mite infested leaflets negatively correlated with OPDA amounts. While OPDA induces expression of the ORGs *UGT73B5*, *CAD*, *GST6* and *GRX480* in Arabidopsis (Taki *et al.*, 2005; Mueller *et al.*, 2008; Park *et al.*, 2013), the expression of their respective putative tomato homologs *TWI-1*, *ADH*, *GST6* and *GRX* all followed exactly the opposite pattern in our experiments. The fact that all four tomato genes had identical expression profiles (we found this for more putative ORGs, data not shown) and that their sequences and gene ontologies were conserved between Arabidopsis and tomato (52-62% identical, 69-77% similar at the amino acid level), strongly suggests these genes are indeed functional homologs. An explanation for the inverse correlation of ORG transcript levels and OPDA amounts might be that these genes are regulated by one or more signals other than OPDA. This might for instance be attributed to SA, since we found significantly higher levels of SA in W-S+ than in W+S+ and W-S- infested leaflets and thus SA amounts and ORG transcript levels were positively correlated. Interestingly, some Arabidopsis ORGs have been shown to be SA responsive (Uquillas *et al.*, 2004; Langlois-Meurinne *et al.*, 2005; Ndamukong *et al.*, 2007). In tomato, the ORG *TWI-1* is also responsive to SA (O'Donnell *et al.*, 1998). This suggests that SA might (co-)regulate the expression of the putative tomato ORGs.

SA amounts correlated only weakly with expression levels of *PR-P6* and did not correlate with *PR-1a* expression levels at all. Interestingly, *PR-1a* was not induced compared to the control in leaflets infested with suppressor mites, while it was induced in leaflets infested with inducer mites. This weak or lack of correlation with PR gene expression might thus be indicative of defence suppression by suppressor mites, which was demonstrated to occur downstream of phytohormone accumulation, e.g., at the *PR-1a* transcript level (Alba *et al.*, 2015).

Remarkably, plant response patterns (in which W-S+ differed from W+S+ and W-S-) were not reflected in mite performance (mite survival was lower for W-S+ and W-S- than for W+S+ mites), possibly because *Wolbachia* and *Spiroplasma* do not seem to have a significant effect on the JA defence responses to which the mites are susceptible

(Alba *et al.*, 2015). However, differences in OPDA amounts might have a significant impact on the fitness of mites independently from JA, because accumulation of OPDA itself is sufficient to confer resistance to several herbivores and pathogens as has been demonstrated in various studies (Stintzi *et al.*, 2001; Stotz *et al.*, 2011; Park *et al.*, 2013; Bosch *et al.*, 2014; Scalschi *et al.*, 2015; Guo *et al.*, 2014). We can therefore not exclude, that plant responses, that were altered in correlation to bacteria, affect fitness of the suppressor strain in other traits than we have measured.

For the inducer strain, the results form a clearer picture. Leaflets infested with C-S- mites contained higher amounts of ABA, traumatic acid and 9,12-OH-(10E)-dodecenoic acid, while transcript levels of SA-responsive *PR-P6* and *PR-1a* were reduced. Consistent with these results, earlier studies found that feeding by arthropods with bacterial symbionts was associated with increased amounts of SA and higher expression levels of *PR* genes in host plants (Chung *et al.*, 2013; Su *et al.*, 2015). ABA appears to negatively regulate SA defences in tomato, in particular by inhibiting expression of *PR-1a* (Audenaert *et al.*, 2002). The reduced expression of *PR* genes in C-S- samples compared to the C+S+ samples might thus be explained by the negative action of ABA. Since these results were paralleled by different feeding intensities of mites with or without these bacteria, *Cardinium* and/or *Spiroplasma* may negatively affect mite fitness via induction of plant defences.

This hypothesis is complemented by different feeding scar phenotypes of C-S- and C+S+ mites. Whereas, feeding by C+S+ mites resulted in rusty red/brown scars, infestation with C-S- mites yielded white scars. Similar scar phenotypes have been reported before for the Kanzawa spider mite *T. kanzawai*. In that case, red scars were associated with increased SA amounts of bean leaves as well as increased expression of a SA defence marker gene (Matsushima *et al.*, 2006), which is in agreement with our findings. Notably, the scar colour resulting from Kanzawa mite feeding was found to be determined by the mite genotype, with dominance of the red phenotype over the white one and no maternal effect (Yano *et al.*, 2003). Thus, maternally inherited symbionts cannot be the cause of differentially coloured scars produced by *T. kanzawai* feeding. Nevertheless, since we did not select for scar phenotype and the genetic background of the mite lines used in our experiments was equal among the C+S+ and C-S- groups, we suggest that the red scars in our case did probably not have a genetic basis but were caused by the presence of a bacterium and concomitant induction of SA-responses. Most likely the presence of *Cardinium*, resulted in rusty red/brown scars, because we did not observe this scar phenotype with W-S+ suppressor mites that contained the same *Spiroplasma* OTU.

#### ***IV. The combined presence of Wolbachia and Spiroplasma bacteria has consequences for mite gene expression and induced plant responses***

Within a host, the various symbionts that (can) co-occur possibly interact, which may affect host and bacterial fitness in various ways. Our results suggest that a single or double infection status of suppressor mites differentially affects host gene expression

as well as induced plant responses, with *Wolbachia* demonstrating characteristics of a mutualist, while the effect of *Spiroplasma* on the mite was less clear, but its presence (partially) antagonized *Wolbachia*-associated responses. This could have resulted from competition between the two symbionts, for instance for space and/or resources. Microbial competition for space (i.e., the ovaries) within the mosquito hosts *Anopheles stephensi* and *Aedes aegypti* was reported to occur between *Wolbachia* and an unrelated *Asaia* bacterium (Hughes *et al.*, 2014; Rossi *et al.*, 2015). It is thus possible that *Wolbachia* spatially displaces *Spiroplasma* or *vice versa*, i.e., away from organs where they may affect induced plant responses or mite digestive processes, like the salivary glands or gut epithelial cells. It would therefore be interesting to determine the spatial localization of both bacteria in single and double infected mites using *in situ* hybridizations. Beside the proper localization, symbiont densities are another important parameter for fidelity of their vertical transmission as well as their effect on host fitness (Rousset *et al.*, 1999; Goto *et al.*, 2006; Oliver *et al.*, 2006; Unckless *et al.*, 2009). Our Illumina MiSeq analysis showed that the relative abundance (as well as the total number of reads) of *Spiroplasma* in the W-S+ lines was higher than in the W+S+ lines, suggesting that *Wolbachia* may negatively affect the abundance of *Spiroplasma* in suppressor mites. However, since 16S amplicon sequencing is only a semi-quantitative method, bacterial abundance should be assessed by means of qRT-PCRs to test this hypothesis.

How co-infecting symbionts affect each other's densities depends on both the species and genotype of host and symbiont (Bordenstein & Werren, 2000; Kondo *et al.*, 2005). For instance, in double or triple infections of wasps, fruitflies or moths with different *Wolbachia* genotypes, none of the tested *Wolbachia* strains influenced densities of the other *Wolbachia* strain(s) (Rousset *et al.*, 1999; Ikeda *et al.*, 2003; Mouton *et al.*, 2003). In contrast, in the beetle *Callosobruchus chinensis* one *Wolbachia* strain suppressed a second one (Kondo *et al.*, 2005). In *D. melanogaster* infected with *Wolbachia* and *Spiroplasma*, the latter negatively affected *Wolbachia* densities, but *Wolbachia* did not affect *Spiroplasma* densities (Goto *et al.*, 2006).

In addition, fitness effects of mixed infections with different bacterial species on hosts are diverse. For instance, female *Bryobia sarothamni* mites that were doubly infected with *Wolbachia* and *Cardinium* had a higher fecundity than singly or uninfected females (Ros & Breeuwer, 2009). In contrast, pea aphids that were doubly infected with *H. defensa* and *Serratia symbiotica* had a lower fecundity than singly or uninfected aphids (Oliver *et al.*, 2006). Infection of *D. melanogaster* with *Wolbachia*, *Spiroplasma* or both did not affect fitness parameters of the flies (Montenegro *et al.*, 2006). A mutualistic interaction has been proposed for *Wolbachia* and *Spiroplasma* that co-infect the fruitfly *D. neotestaceae* (Jaenike *et al.*, 2010b). The frequency of co-infected fruitflies in sampled populations and estimated evolutionary age of the two bacteria that had co-infected the flies, were seen as a possible basis for the evolution of a mutualism between the bacteria (Jaenike *et al.*, 2010b). The fact that *Wolbachia* and *Spiroplasma* bacteria in the

suppressor strain seemed to differentially influence mite survival, transcriptomic responses and induced plant responses, emphasizes the importance of considering interaction effects of multiple bacteria in/on one host.

Unfortunately, we did not obtain all bacterial combinations in *T. urticae* to disentangle the exact effects of single bacterial strains and their combined effects on mite fitness, induced plant responses and mite gene expression. For instance, we did not test mites that were infected with only *Wolbachia*, hence we can only indirectly infer the role of *Wolbachia* in our experiments from comparisons between W+S+, W-S+ and W-S- mites. Moreover, for the inducer strain the roles of *Spiroplasma* and *Cardinium* remain difficult to interpret because we did not have C-S+ or C+S- mites. The same *Spiroplasma* OTU occurred in both the inducer and suppressor mites. If we assume that *Spiroplasma* in inducer mites had the same effect on host fitness parameters as in suppressor mites, we could attribute the negative effects of bacteria in inducer mites to *Cardinium*. However, *Spiroplasma* may interact with *Cardinium* in the mites. Furthermore, its effects on host fitness may depend on mite genotype (Schmid-Hempel, 2011).

### Conclusions

We assessed the influence of the (endo)symbiotic bacteria *Wolbachia*, *Cardinium* and *Spiroplasma* on the performance and transcriptome of their respective hosts, the *T. urticae* suppressor (W+S+) and inducer (C+S+) strains. Furthermore, we tested how these spider mite-associated bacteria affect the induced responses of tomato plants triggered by feeding mites. Our data indicate that the different (endo) symbiotic bacteria may have distinct consequences for their host (see FIGURE 5.9):

1. The presence of *Wolbachia* correlates positively with survival of suppressor mites, possibly via manipulation of OPDA and SA-related responses. We did not observe changes in JA- and SA marker gene expression levels. Further, the presence of *Wolbachia* correlates positively with transcript levels of suppressor mite genes involved in digestion and detoxification of xenobiotics. This might reflect responses of the mite to symbiont-generated or -modified metabolites, either plant-derived (transcriptome profiles of the suppressor strain resemble the OPDA phenotype) or not, that are suitable for further processing by the mite.
2. The presence of *Cardinium* and *Spiroplasma* correlates negatively with the feeding activity, survival and fecundity of inducer mites, while it correlates positively with the induction of SA-marker-gene expression as well as with the frequently observed rusty scars of infested leaves. This indicates a distinct and clearly visible (in)direct impact of the presence of *Cardinium* and *Spiroplasma* on the mite's host plant while simultaneously their presence correlates with inhibition of digestion and detoxification processes in the mite.
3. There is no clear correlation between the absence or presence of *Spiroplasma* and survival of suppressor mites, while the induced plant responses of leaflets fed on by mites with or without *Spiroplasma* are ambiguous. The data suggests

a complex interaction between *Wolbachia* and *Spiroplasma* in which the latter may suppress OPDA-related responses that are induced in the presence of *Wolbachia*. In addition, some mite digestion and detoxification genes that respond positively to the presence of *Wolbachia* seem to be slightly suppressed when only *Spiroplasma* is present. Unfortunately, we did not obtain W+S- lines to disentangle this interaction.

4. The same *Spiroplasma* OTU was present in suppressor and inducer mites. Hence, although we did not obtain singly-infected inducer lines, a comparison of the results between the mite strains suggests that the presence of *Spiroplasma* in C+S+ inducer mites is not related with the observed negative effects on its host (e.g., induction of SA defences, impaired expression of detoxification genes and reduced feeding). However, we cannot exclude that *Spiroplasma* affects the suppressor mites differently than the inducer mites.

In conclusion, according to our data, *Wolbachia* has the characteristics of a mutualist, while *Cardinium* has the characteristics of a parasite. The role of *Spiroplasma* remains unclear, but it does seem to interact – directly or indirectly – with *Wolbachia* at the level of mite gene expression and induced plant responses (note that these two responses might be causally linked).

## REFERENCES

- Ahn, S.-J., Dermauw, W., Wybouw, N., Heckel, D.G. & Van Leeuwen, T. 2014. Bacterial origin of a diverse family of UDP-glycosyltransferase genes in the *Tetranychus urticae* genome. *Insect Biochem Mol Biol* **50**: 43-57.
- Alba, J.M., Schimmel, B.C.J., Glas, J.J., Ataide, L.M.S., Pappas, M.L., Villarroel, C.A., Schuurink, R.C., Sabelis, M.W. & Kant, M.R. 2015. Spider mites suppress tomato defenses downstream of jasmonate and salicylate independently of hormonal crosstalk. *New Phytol* **205**: 828-840.
- Ament, K., Kant, M.R., Sabelis, M.W., Haring, M.A. & Schuurink, R.C. 2004. Jasmonic acid is a key regulator of spider mite-induced volatile terpenoid and methyl salicylate emission in tomato. *Plant Physiol* **135**: 2025-2037.
- Anbutsu, H. & Fukatsu, T. 2011. *Spiroplasma* as a model insect endosymbiont. *Environ Microbiol Rep* **3**: 144-153.
- Audenaert, K., De Meyer, G.B. & Höfte, M.M. 2002. Abscisic acid determines basal susceptibility of tomato to *Botrytis cinerea* and suppresses salicylic acid-dependent signaling mechanisms. *Plant Physiol* **128**: 491-501.
- Barbosa, P., Krischik, V.A. & Jones, C.G. 1991. *Microbial mediation of plant-herbivore interactions*. John Wiley & Sons, New York, USA.
- Barr, K.L., Hearne, L.B., Briesacher, S., Clark, T.L. & Davis, G.E. 2010. Microbial symbionts in insects influence down-regulation of defense genes in maize. *PLoS ONE* **5**: e11339.
- Bates, D., Maechler, M., Bolker, B. & Walker, S. 2013. lme4: Linear mixed-effects models using Eigen and S4. *R package version 1*.
- Benjamini, Y. & Hochberg, Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Methodol* **57**: 289-300.
- Body, M., Kaiser, W., Dubreuil, G., Casas, J. & Giron, D. 2013. Leaf-miners co-opt microorganisms to enhance their nutritional environment. *J Chem Ecol* **39**: 969-977.
- Bolland, H.R., Gutierrez, J. & Flechtmann, C.H. 1998. *World catalogue of the spider mite family (Acari: Tetranychidae)*. Brill, Leiden, the Netherlands.

- Bordenstein, S.R. & Werren, J.H. 2000. Do *Wolbachia* influence fecundity in *Nasonia vitripennis*? *Heredity* **84**: 54-62.
- Bosch, M., Wright, L.P., Gershenson, J., Wasternack, C., Hause, B., Schaller, A. & Stintzi, A. 2014. Jasmonic acid and its precursor 12-oxophytodienoic acid control different aspects of constitutive and induced herbivore defenses in tomato. *Plant Physiol* **166**: 396-410.
- Breeuwer, J.A.J. & Jacobs, G. 1996. *Wolbachia*: intracellular manipulators of mite reproduction. *Exp Appl Acarol* **20**: 421-434.
- Brownlie, J.C., Cass, B.N., Riegler, M., Witsenburg, J.J., Iturbe-Ormaetxe, I., McGraw, E.A. & O'Neill, S.L. 2009. Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS Pathol* **5**: e1000368.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Pena, A.G., Goodrich, J.K. & Gordon, J.I. 2010a. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* **7**: 335-336.
- Caporaso, J.G., Bittinger, K., Bushman, F.D., DeSantis, T.Z., Andersen, G.L. & Knight, R. 2010b. PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* **26**: 266-267.
- Casteel, C.L. & Hansen, A.K. 2014. Evaluating insect-microbiomes at the plant-insect interface. *J Chem Ecol* **40**: 836-847.
- Chung, S.H., Rosa, C., Scully, E.D., Peiffer, M., Tooker, J.F., Hoover, K., Luthe, D.S. & Felton, G.W. 2013. Herbivore exploits orally secreted bacteria to suppress plant defenses. *Proc Natl Acad Sci USA* **110**: 15728-15733.
- Darby, A.C., Armstrong, S.D., Bah, G.S., Kaur, G., Hughes, M.A., Kay, S.M., Koldkjær, P., Rainbow, L., Radford, A.D. & Blaxter, M.L. 2012. Analysis of gene expression from the *Wolbachia* genome of a filarial nematode supports both metabolic and defensive roles within the symbiosis. *Genome Res* **22**: 2467-2477.
- Davoine, C., Falletti, O., Douki, T., Iacazio, G., Ennar, N., Montillet, J.-L. & Triantaphylidès, C. 2006. Adducts of oxylipin electrophiles to glutathione reflect a 13 specificity of the downstream lipoxygenase pathway in the tobacco hypersensitive response. *Plant Physiol* **140**: 1484-1493.
- De Bary, A. 1879. *The phenomenon of symbiosis*. Karl J. Trubner, Strasbourg, Germany.
- Dermauw, W., Osborne, E.J., Clark, R.M., Grbić, M., Tirry, L. & Van Leeuwen, T. 2013. A burst of ABC genes in the genome of the polyphagous spider mite *Tetranychus urticae*. *BMC Genomics* **14**: 317.
- DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi, D., Hu, P. & Andersen, G.L. 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* **72**: 5069-5072.
- Dou, D. & Zhou, J.-M. 2012. Phytopathogen effectors subverting host immunity: different foes, similar battleground. *Cell Host Microbe* **12**: 484-495.
- Douglas, A. 2009. The microbial dimension in insect nutritional ecology. *Funct Ecol* **23**: 38-47.
- Duron, O., Bouchon, D., Boutin, S., Bellamy, L., Zhou, L., Engelstädter, J. & Hurst, G.D. 2008. The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol* **6**: 27.
- Ebbert, M.A. 1991. The interaction phenotype in the *Drosophila willistoni*-*Spiroplasma* symbiosis. *Evolution* **45**: 971-988.
- Edgar, R.C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**: 2460-2461.
- Engel, P. & Moran, N.A. 2013. The gut microbiota of insects—diversity in structure and function. *FEMS Microbiol Rev* **37**: 699-735.
- Engelstädter, J. & Hurst, G.D. 2009. The ecology and evolution of microbes that manipulate host reproduction. *Annu Rev Ecol Evol Syst* **40**: 127-149.

- Erb, M., Meldau, S. & Howe, G.A. 2012. Role of phytohormones in insect-specific plant reactions. *Trends Plant Sci* **17**: 250-259.
- Feldhaar, H. 2011. Bacterial symbionts as mediators of ecologically important traits of insect hosts. *Ecol Entomol* **36**: 533-543.
- Fonseca, S., Chini, A., Hamberg, M., Adie, B., Porzel, A., Kramell, R., Miersch, O., Wasternack, C. & Solano, R. 2009. (+)-7-iso-Jasmonoyl-L-isoleucine is the endogenous bioactive jasmonate. *Nat Chem Biol* **5**: 344-350.
- Foster, J., Ganatra, M., Kamal, I., Ware, J., Makarova, K., Ivanova, N., Bhattacharyya, A., Kapatral, V., Kumar, S. & Posfai, J. 2005. The *Wolbachia* genome of *Brugia malayi*: endosymbiont evolution within a human pathogenic nematode. *PLoS Biol* **3**: e121.
- Frago, E., Dicke, M. & Godfray, H.C.J. 2012. Insect symbionts as hidden players in insect-plant interactions. *Trends Ecol Evol* **27**: 705-711.
- Fry, A.J., Rand, D.M. & Poulin, R. 2002. *Wolbachia* interactions that determine *Drosophila melanogaster* survival. *Evolution* **56**: 1976-1981.
- Fry, A., Palmer, M. & Rand, D. 2004. Variable fitness effects of *Wolbachia* infection in *Drosophila melanogaster*. *Heredity* **93**: 379-389.
- Fukatsu, T., Tsuchida, T., Nikoh, N. & Koga, R. 2001. *Spiroplasma* symbiont of the pea aphid, *Acyrtosiphon pisum* (Insecta: Homoptera). *Appl Environ Microbiol* **67**: 1284-1291.
- Gadea, J., Mayda, M.E., Conejero, V. & Vera, P. 1996. Characterization of defense-related genes ectopically expressed in viroid-infected tomato plants. *Mol Plant Microbe Interact* **9**: 409-415.
- Glazebrook, J. 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annu Rev Phytopathol* **43**: 205-227.
- Gonzales-Vigil, E., Bianchetti, C.M., Phillips, G.N. & Howe, G.A. 2011. Adaptive evolution of threonine deaminase in plant defense against insect herbivores. *Proc Natl Acad Sci USA* **108**: 5897-5902.
- Goto, S., Anbutsu, H. & Fukatsu, T. 2006. Asymmetrical interactions between *Wolbachia* and *Spiroplasma* endosymbionts coexisting in the same insect host. *Appl Environ Microbiol* **72**: 4805-4810.
- Gotoh, T., Noda, H. & Ito, S. 2007. *Cardinium* symbionts cause cytoplasmic incompatibility in spider mites. *Heredity* **98**: 13-20.
- Goverse, A. & Smant, G. 2014. The activation and suppression of plant innate immunity by parasitic nematodes. *Annu Rev Phytopathol* **52**: 243-265.
- Grbic, M., Van Leeuwen, T., Clark, R.M., Rombauts, S., Rouze, P., Grbic, V., Osborne, E.J., Dermauw, W., Thi Ngoc, P.C., Ortego, F., Hernandez-Crespo, P., Diaz, I., Martinez, M., Navajas, M., Sucena, E., Magalhaes, S., Nagy, L., Pace, R.M., Djuranovic, S., Smagghe, G., Iga, M., Christiaens, O., Veenstra, J.A., Ewer, J., Villalobos, R.M., Hutter, J.L., Hudson, S.D., Velez, M., Yi, S.V., Zeng, J., Pires-daSilva, A., Roch, F., Cazaux, M., Navarro, M., Zhurov, V., Acevedo, G., Bjelica, A., Fawcett, J.A., Bonnet, E., Martens, C., Baele, G., Wissler, L., Sanchez-Rodriguez, A., Tirry, L., Blais, C., Demeestere, K., Henz, S.R., Gregory, T.R., Mathieu, J., Verdon, L., Farinelli, L., Schmutz, J., Lindquist, E., Feyereisen, R. & Van de Peer, Y. 2011. The genome of *Tetranychus urticae* reveals herbivorous pest adaptations. *Nature* **479**: 487-492.
- Guo, H.-M., Li, H.-C., Zhou, S.-R., Xue, H.-W. & Miao, X.-X. 2014. Cis-12-oxo-phytyldienoic acid stimulates rice defense response to a piercing-sucking insect. *Mol Plant* **7**: 1683-1692.
- Haas, B.J., Gevers, D., Earl, A.M., Feldgarden, M., Ward, D.V., Giannoukos, G., Ciulla, D., Tabbaa, D., Highlander, S.K. & Sodergren, E. 2011. Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons. *Genome Res* **21**: 494-504.
- Hamilton, P.T., Leong, J.S., Koop, B.F. & Perlman, S.J. 2014. Transcriptional responses in a *Drosophila* defensive symbiosis. *Mol Ecol* **23**: 1558-1570.

- Hansen, A.K. & Moran, N.A. 2014. The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol Ecol* **23**: 1473-1496.
- Harrel, F.E., Dupont, C. & al, e. 2015. Hmisc: Harrell Miscellaneous. R package version 3.15-0. <http://CRAN.R-project.org/package=Hmisc>.
- Hedges, L.M., Brownlie, J.C., O'Neill, S.L. & Johnson, K.N. 2008. *Wolbachia* and virus protection in insects. *Science* **322**: 702-702.
- Hoffmann, A.A., Clancy, D.J. & Merton, E. 1994. Cytoplasmic incompatibility in Australian populations of *Drosophila melanogaster*. *Genetics* **136**: 993-999.
- Hoffmann, A.A., Hercus, M. & Dagher, H. 1998. Population dynamics of the *Wolbachia* infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* **148**: 221-231.
- Hosokawa, T., Koga, R., Kikuchi, Y., Meng, X.-Y. & Fukatsu, T. 2010. *Wolbachia* as a bacteriocyte-associated nutritional mutualist. *Proc Natl Acad Sci USA* **107**: 769-774.
- Hothorn, T., Bretz, F. & Westfall, P. 2008. Simultaneous inference in general parametric models. *Biometrical J* **50**: 346-363.
- Howe, G.A., Lee, G.I., Itoh, A., Li, L. & DeRocher, A.E. 2000. Cytochrome P450-dependent metabolism of oxylipins in tomato. Cloning and expression of allene oxide synthase and fatty acid hydroperoxide lyase. *Plant Physiol* **123**: 711-724.
- Hughes, G.L., Dodson, B.L., Johnson, R.M., Murdock, C.C., Tsujimoto, H., Suzuki, Y., Patt, A.A., Cui, L., Nossa, C.W. & Barry, R.M. 2014. Native microbiome impedes vertical transmission of *Wolbachia* in *Anopheles* mosquitoes. *Proc Natl Acad Sci USA* **111**: 12498-12503.
- Hussain, M., Frentiu, F.D., Moreira, L.A., O'Neill, S.L. & Asgari, S. 2011. *Wolbachia* uses host microRNAs to manipulate host gene expression and facilitate colonization of the dengue vector *Aedes aegypti*. *Proc Natl Acad Sci USA* **108**: 9250-9255.
- Hutchence, K.J., Fischer, B., Paterson, S. & Hurst, G.D. 2011. How do insects react to novel inherited symbionts? A microarray analysis of *Drosophila melanogaster* response to the presence of natural and introduced *Spiroplasma*. *Mol Ecol* **20**: 950-958.
- Ikeda, T., Ishikawa, H. & Sasaki, T. 2003. Regulation of *Wolbachia* density in the mediterranean flour moth, *Ephestia kuehniella*, and the almond moth, *Cadra cautella*. *Zoolog Sci* **20**: 153-157.
- Jaenike, J., Unckless, R., Cockburn, S.N., Boelio, L.M. & Perlman, S.J. 2010a. Adaptation via symbiosis: recent spread of a *Drosophila* defensive symbiont. *Science* **329**: 212-215.
- Jaenike, J., Stahlhut, J.K., Boelio, L.M. & Unckless, R.L. 2010b. Association between *Wolbachia* and *Spiroplasma* within *Drosophila neotestacea*: an emerging symbiotic mutualism? *Mol Ecol* **19**: 414-425.
- Kaiser, W., Huguet, E., Casas, J., Commin, C. & Giron, D. 2010. Plant green-island phenotype induced by leaf-miners is mediated by bacterial symbionts. *Proc R Soc B Biol Sci* **277**: 2311-2319.
- Kambis, Z., Cook, P.E., Phuc, H.K. & Sinkins, S.P. 2009. Immune activation by life-shortening *Wolbachia* and reduced filarial competence in mosquitoes. *Science* **326**: 134-136.
- Kant, M.R., Ament, K., Sabelis, M.W., Haring, M.A. & Schuurink, R.C. 2004. Differential timing of spider mite-induced direct and indirect defenses in tomato plants. *Plant Physiol* **135**: 483-495.
- Kant, M.R. 2006. *The consequences of herbivore variability for direct and indirect defenses of plants*, PhD Thesis, Amsterdam, The Netherlands.
- Kant, M.R., Sabelis, M.W., Haring, M.A. & Schuurink, R.C. 2008. Intraspecific variation in a generalist herbivore accounts for differential induction and impact of host plant defences. *Proc R Soc B Biol Sci* **275**: 443-452.
- Kant, M.R., Jonckheere, W., Knecht, B., Lemos, F., Liu, J., Schimmel, B.C.J., Villarroel, C.A., Ataide, L.M.S., Dermauw, W., Glas, J.J., Egas, M., Janssen, A., Van Leeuwen, T., Schuurink, R.C., Sabelis, M.W. & Alba, J.M. 2015. Mechanisms and ecological

- consequences of plant defence induction and suppression in herbivore communities. *Ann Bot* **115**: 1015-1051.
- Karban, R. & Baldwin, I.T. 2007. *Induced responses to herbivory*. University of Chicago Press.
- Kessler, A., Halitschke, R. & Baldwin, I.T. 2004. Silencing the jasmonate cascade: induced plant defenses and insect populations. *Science* **305**: 665-668.
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. & Glöckner, F.O. 2012. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res*: gks808.
- Kondo, N., Shimada, M. & Fukatsu, T. 2005. Infection density of *Wolbachia* endosymbiont affected by co-infection and host genotype. *Biol Lett* **1**: 488-491.
- Kremer, N., Voronin, D., Charif, D., Mavingui, P., Mollereau, B. & Vavre, F. 2009. *Wolbachia* interferes with ferritin expression and iron metabolism in insects. *PLoS Pathol* **5**: e1000630.
- Langlois-Meurinne, M., Gachon, C.M. & Saindrenan, P. 2005. Pathogen-responsive expression of glycosyltransferase genes UGT73B3 and UGT73B5 is necessary for resistance to *Pseudomonas syringae* pv tomato in *Arabidopsis*. *Plant Physiol* **139**: 1890-1901.
- Langmead, B., Trapnell, C., Pop, M. & Salzberg, S.L. 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10**: R25.
- Li, C., Williams, M.M., Loh, Y.-T., Lee, G.I. & Howe, G.A. 2002. Resistance of cultivated tomato to cell content-feeding herbivores is regulated by the octadecanoid-signaling pathway. *Plant Physiol* **130**: 494-503.
- Lisón, P., Rodrigo, I. & Conejero, V. 2006. A novel function for the cathepsin D inhibitor in tomato. *Plant Physiol* **142**: 1329-1339.
- Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A., Reissmann, S. & Kahmann, R. 2015. Fungal effectors and plant susceptibility. *Annu Rev Plant Biol* **66**: 513-545.
- Matsushima, R., Ozawa, R., Uefune, M., Gotoh, T. & Takabayashi, J. 2006. Intraspecific variation in the Kanzawa spider mite differentially affects induced defensive response in lima bean plants. *J Chem Ecol* **32**: 2501-2512.
- Mayoral, J.G., Etebari, K., Hussain, M., Khromykh, A.A. & Asgari, S. 2014. *Wolbachia* infection modifies the profile, shuttling and structure of microRNAs in a mosquito cell line. *PLoS ONE* **9**(4): e96107.
- Migeon, A., Nouguié, E. & Dorkeld, F. 2011. Spider Mites Web: a comprehensive database for the Tetranychidae. In: *Trends Acarol*, pp. 557-560. Springer.
- Mithöfer, A. & Boland, W. 2012. Plant defense against herbivores: chemical aspects. *Annu Rev Plant Biol* **63**: 431-450.
- Montenegro, H., Petherick, A., Hurst, G. & Klaczko, L. 2006. Fitness effects of *Wolbachia* and *Spiroplasma* in *Drosophila melanogaster*. *Genetica* **127**: 207-215.
- Moreira, L.A., Iturbe-Ormaetxe, I., Jeffery, J.A., Lu, G., Pyke, A.T., Hedges, L.M., Rocha, B.C., Hall-Mendelin, S., Day, A. & Riegler, M. 2009. A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and *Plasmodium*. *Cell* **139**: 1268-1278.
- Mouton, L., Henri, H., Bouletreau, M. & Vavre, F. 2003. Strain-specific regulation of intracellular *Wolbachia* density in multiply infected insects. *Mol Ecol* **12**: 3459-3465.
- Mueller, S., Hilbert, B., Dueckershoff, K., Roitsch, T., Krischke, M., Mueller, M.J. & Berger, S. 2008. General detoxification and stress responses are mediated by oxidized lipids through TGA transcription factors in *Arabidopsis*. *Plant Cell* **20**: 768-785.
- Nakamura, Y., Gotoh, T., Imanishi, S., Mita, K., Kurti, T. & Noda, H. 2011. Differentially expressed genes in silkworm cell cultures in response to infection by *Wolbachia* and *Cardinium* endosymbionts. *Insect Mol Biol* **20**: 279-289.


- Ndamukong, I., Abdallat, A.A., Thurow, C., Fode, B., Zander, M., Weigel, R. & Gatz, C. 2007. SA-inducible *Arabidopsis* glutaredoxin interacts with TGA factors and suppresses JA-responsive PDF1.2 transcription. *Plant J* **50**: 128-139.
- O'Donnell, P.J., Truesdale, M.R., Calvert, C.M., Dorans, A., Roberts, M.R. & Bowles, D.J. 1998. A novel tomato gene that rapidly responds to wound-and pathogen-related signals. *Plant J* **14**: 137-142.
- Oliver, K.M., Moran, N.A. & Hunter, M.S. 2006. Costs and benefits of a superinfection of facultative symbionts in aphids. *Proc R Soc B Biol Sci* **273**: 1273-1280.
- Park, S.-W., Li, W., Viehhauser, A., He, B., Kim, S., Nilsson, A.K., Andersson, M.X., Kittle, J.D., Ambavaram, M.M. & Luan, S. 2013. Cyclophilin 20-3 relays a 12-oxo-phytodienoic acid signal during stress responsive regulation of cellular redox homeostasis. *Proc Natl Acad Sci USA* **110**: 9559-9564.
- Perrot-Minnot, M.J., Cheval, B., Migeon, A. & Navajas, M. 2002. Contrasting effects of *Wolbachia* on cytoplasmic incompatibility and fecundity in the haplodiploid mite *Tetranychus urticae*. *J Evol Biol* **15**: 808-817.
- Pieterse, C.M., Van der Does, D., Zamioudis, C., Leon-Reyes, A. & Van Wees, S.C. 2012. Hormonal modulation of plant immunity. *Annu Rev Cell Dev Biol* **28**: 489-521.
- Rances, E., Ye, Y.H., Woolfit, M., McGraw, E.A. & O'Neill, S.L. 2012. The relative importance of innate immune priming in *Wolbachia*-mediated dengue interference. *PLoS Pathol* **8**: e1002548.
- R Core Team 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Version 3.0.2.
- Robert, C.A., Frank, D.L., Leach, K.A., Turlings, T.C., Hibbard, B.E. & Erb, M. 2013. Direct and indirect plant defenses are not suppressed by endosymbionts of a specialist root herbivore. *J Chem Ecol* **39**: 507-515.
- Rong, X., Zhang, Y.-K., Zhang, K.-J. & Hong, X.-Y. 2014. Identification of *Wolbachia*-responsive microRNAs in the two-spotted spider mite, *Tetranychus urticae*. *BMC Genomics* **15**: 1122.
- Ros, V. & Breeuwer, J. 2009. The effects of, and interactions between, *Cardinium* and *Wolbachia* in the doubly infected spider mite *Bryobia sarothamni*. *Heredity* **102**: 413-422.
- Rossi, P., Ricci, I., Cappelli, A., Damiani, C., Ulissi, U., Mancini, M.V., Valzano, M., Capone, A., Epis, S. & Crotti, E. 2015. Mutual exclusion of *Asaia* and *Wolbachia* in the reproductive organs of mosquito vectors. *Parasit Vectors* **8**: 278.
- Rousset, F., Braig, H.R. & O'Neill, S.L. 1999. A stable triple *Wolbachia* infection in *Drosophila* with nearly additive incompatibility effects. *Heredity* **82**: 620-627.
- Santamaría, M.E., Hernández-Crespo, P., Ortego, F., Grbic, V., Grbic, M., Diaz, I. & Martinez, M. 2012. Cysteine peptidases and their inhibitors in *Tetranychus urticae*: a comparative genomic approach. *BMC Genomics* **13**: 307.
- Scalschi, L., Sanmartín, M., Camañes, G., Troncho, P., Sánchez-Serrano, J.J., García-Agustín, P. & Vicedo, B. 2015. Silencing of OPR3 in tomato reveals the role of OPDA in callose deposition during the activation of defense responses against *Botrytis cinerea*. *Plant J* **81**: 304-315.
- Schmid-Hempel, P. 2011. *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford University Press, New York.
- Schoonhoven, L.M., van Loon, J.J.A. & Dicke, M. 2005. *Insect-plant biology, 2nd edn*. Oxford University Press, Oxford.
- Sivasankar, S., Sheldrick, B. & Rothstein, S.J. 2000. Expression of allene oxide synthase determines defense gene activation in tomato. *Plant Physiol* **122**: 1335-1342.
- Smyth, G.K. 2004. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* **3**: 1-25.
- Smyth, G.K. & Altman, N.S. 2013. Separate-channel analysis of two-channel microarrays: Recovering inter-spot information. *BMC Bioinformatics* **14**: 165.

- Staswick, P.E. & Tiryaki, I. 2004. The oxylipin signal jasmonic acid is activated by an enzyme that conjugates it to isoleucine in *Arabidopsis*. *Plant Cell* **16**: 2117-2127.
- Stefanini, A. & Duron, O. 2012. Exploring the effect of the *Cardinium* endosymbiont on spiders. *J Evol Biol* **25**: 1521-1530.
- Stintzi, A., Weber, H., Reymond, P., Browse, J. & Farmer, E.E. 2001. Plant defense in the absence of jasmonic acid: The role of cyclopentenones. *Proc Natl Acad Sci USA* **98**: 12837-12842.
- Storms, J.J.H. 1971. Some physiological effects of spider mite infestation on bean plants. *Neth J Plant Pathol* **77**: 154-167.
- Stotz, H.U., Jikumaru, Y., Shimada, Y., Sasaki, E., Stingl, N., Mueller, M.J. & Kamiya, Y. 2011. Jasmonate-dependent and COI1-independent defense responses against *Sclerotinia sclerotiorum* in *Arabidopsis thaliana*: Auxin is part of COI1-independent defense signaling. *Plant Cell Physiol* **52**: 1941-1956.
- Strassner, J., Schaller, F., Frick, U.B., Howe, G.A., Weiler, E.W., Amrhein, N., Macheroux, P. & Schaller, A. 2002. Characterization and cDNA-microarray expression analysis of 12-oxophytodienoate reductases reveals differential roles for octadecanoid biosynthesis in the local versus the systemic wound response. *Plant J* **32**: 585-601.
- Su, Q., Oliver, K.M., Xie, W., Wu, Q., Wang, S. & Zhang, Y. 2015. The whitefly-associated facultative symbiont *Hamiltonella defensa* suppresses induced plant defences in tomato. *Funct Ecol* **29**: 1007-1018.
- Sugio, A., Kingdom, H.N., MacLean, A.M., Grieve, V.M. & Hogenhout, S.A. 2011. Phytoplasma protein effector SAP11 enhances insect vector reproduction by manipulating plant development and defense hormone biosynthesis. *Proc Natl Acad Sci USA* **108**: 1254-1263.
- Sugio, A., Dubreuil, G., Giron, D. & Simon, J.-C. 2015. Plant-insect interactions under bacterial influence: ecological implications and underlying mechanisms. *J Exp Bot* **66**: 467-478.
- Sullivan, M.L. & Green, P.J. 1993. Post-transcriptional regulation of nuclear-encoded genes in higher plants: the roles of mRNA stability and translation. *Plant Mol Biol* **23**: 1091-1104.
- Taki, N., Sasaki-Sekimoto, Y., Obayashi, T., Kikuta, A., Kobayashi, K., Ainai, T., Yagi, K., Sakurai, N., Suzuki, H. & Masuda, T. 2005. 12-oxo-phytodienoic acid triggers expression of a distinct set of genes and plays a role in wound-induced gene expression in *Arabidopsis*. *Plant Physiol* **139**: 1268-1283.
- Teixeira, L., Ferreira, Á. & Ashburner, M. 2008. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol* **6**: e1000002.
- Tieman, D.M., Loucas, H.M., Kim, J.Y., Clark, D.G. & Klee, H.J. 2007. Tomato phenylacetaldehyde reductases catalyze the last step in the synthesis of the aroma volatile 2-phenylethanol. *Phytochemistry* **68**: 2660-2669.
- Tornero, P., Gadea, J., Conejero, V. & Vera, P. 1997. Two PR-1 genes from tomato are differentially regulated and reveal a novel mode of expression for a pathogenesis-related gene during the hypersensitive response and development. *Mol Plant Microbe Interact* **10**: 624-634.
- Truesdale, M.R., Doherty, H.M., Loake, G.J., McPherson, M.J., Roberts, M. & Bowles, D.J. 1996. Molecular cloning of a novel wound-induced gene from tomato: *twi1*. *Plant Physiol* **112**: 446.
- Unckless, R.L., Boelio, L.M., Herren, J.K. & Jaenike, J. 2009. *Wolbachia* as populations within individual insects: causes and consequences of density variation in natural populations. *Proc R Soc B Biol Sci* **276**: 2805-2811.
- Unckless, R.L. & Jaenike, J. 2012. Maintenance of a male-killing *Wolbachia* in *Drosophila innubila* by male-killing dependent and male-killing independent mechanisms. *Evolution* **66**: 678-689.


- Uquillas, C., Letelier, I., Blanco, F., Jordana, X. & Holuigue, L. 2004. NPR1-independent activation of immediate early salicylic acid-responsive genes in *Arabidopsis*. *Mol Plant Microbe Interact* **17**: 34-42.
- Vala, F., Breeuwer, J.A. & Sabelis, M.W. 2000. *Wolbachia*-induced 'hybrid breakdown' in the two-spotted spider mite *Tetranychus urticae* Koch. *Proc R Soc B Biol Sci* **267**: 1931-1937.
- van Kan, J.A., Joosten, M.H., Wagemakers, C.A., van den Berg-Velthuis, G.C. & de Wit, P.J. 1992. Differential accumulation of mRNAs encoding extracellular and intracellular PR proteins in tomato induced by virulent and avirulent races of *Cladosporium fulvum*. *Plant Mol Biol* **20**: 513-527.
- Verwoerd, T.C., Dekker, B. & Hoekema, A. 1989. A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Res* **17**: 2362.
- Walker, T., Johnson, P., Moreira, L., Iturbe-Ormaetxe, I., Frentiu, F., McMeniman, C., Leong, Y.S., Dong, Y., Axford, J. & Kriesner, P. 2011. The wMel *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti* populations. *Nature* **476**: 450-453.
- Walling, L.L. 2000. The myriad plant responses to herbivores. *J Plant Growth Regul* **19**: 195-216.
- Wasternack, C. 2007. Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Ann Bot* **100**: 681-697.
- Wasternack, C. & Hause, B. 2013. Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. An update to the 2007 review in *Annals of Botany*. *Ann Bot* **111**: 1021-1058.
- Weeks, A.R. & Stouthamer, R. 2004. Increased fecundity associated with infection by a Cytophaga-like intracellular bacterium in the predatory mite, *Metaseiulus occidentalis*. *Proc R Soc B Biol Sci* **271**: 193-195.
- Werren, J.H. 1997. Biology of *Wolbachia*. *Annu Rev Entomol* **42**: 587-609.
- Werren, J.H., Baldo, L. & Clark, M.E. 2008. *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* **6**: 741-751.
- White, J., Kelly, S., Cockburn, S., Perlman, S. & Hunter, M. 2011. Endosymbiont costs and benefits in a parasitoid infected with both *Wolbachia* and *Cardinium*. *Heredity* **106**: 585-591.
- Wybouw, N., Zhurov, V., Martel, C., Bruinsma, K.A., Hendrickx, F., Grbić, V. & Van Leeuwen, T. 2015. Adaptation of a polyphagous herbivore to a novel host plant extensively shapes the transcriptome of herbivore and host. *Mol Ecol*. **24**: 4647-4663
- Xie, J., Vilchez, I. & Mateos, M. 2010. *Spiroplasma* bacteria enhance survival of *Drosophila hydei* attacked by the parasitic wasp *Leptopilina heterotoma*. *PLoS ONE* **5**.
- Xie, J., Butler, S., Sanchez, G. & Mateos, M. 2014. Male killing *Spiroplasma* protects *Drosophila melanogaster* against two parasitoid wasps. *Heredity* **112**: 399-408.
- Xie, R.-R., Chen, X.-L. & Hong, X.-Y. 2011. Variable fitness and reproductive effects of *Wolbachia* infection in populations of the two-spotted spider mite *Tetranychus urticae* Koch in China. *Appl Entomol Zool* **46**: 95-102.
- Yano, S., Kanaya, M. & Takafuji, A. 2003. Genetic basis of color variation in leaf scars induced by the Kanzawa spider mite. *Entomol Exp Appl* **106**: 37-44.
- Zhang, G., Hussain, M., O'Neill, S.L. & Asgari, S. 2013. *Wolbachia* uses a host microRNA to regulate transcripts of a methyltransferase, contributing to dengue virus inhibition in *Aedes aegypti*. *Proc Natl Acad Sci USA* **110**: 10276-10281.
- Zhang, Y.K., Ding, X.L., Rong, X. & Hong, X.Y. 2015. How do hosts react to endosymbionts? A new insight into the molecular mechanisms underlying the *Wolbachia*-host association. *Insect Mol Biol* **24**: 1-12.
- Zhao, D.-X., Zhang, X.-F., Chen, D.-S., Zhang, Y.-K. & Hong, X.-Y. 2013. *Wolbachia*-host interactions: host mating patterns affect *Wolbachia* density dynamics. *PLoS ONE* **8**: e66373.


- Zhurov, V., Navarro, M., Bruinsma, K.A., Arbona, V., Santamaria, M.E., Cazaux, M., Wybouw, N., Osborne, E.J., Ens, C. & Rioja, C. 2014. Reciprocal responses in the interaction between *Arabidopsis* and the cell-content-feeding chelicerate herbivore spider mite. *Plant Physiol* **164**: 384-399.
- Zug, R. & Hammerstein, P. 2012. Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected. *PLoS ONE* **7**: e38544.
- Zug, R. & Hammerstein, P. 2015. Bad guys turned nice? A critical assessment of *Wolbachia* mutualisms in arthropod hosts. *Biol Rev* **90**: 89-111.

SUPPORTING INFORMATION




**FIGURE S5.1.** Schematic and simplified overview of tomato (*Solanum lycopersicum*) defence responses typically induced by spider mite (*Tetranychus urticae*) feeding and which were monitored in this study. Infestation with inducer *T. urticae* Santpoort-2 mites (inset) results in the simultaneous, enhanced accumulation of the phytohormones jasmonic acid (JA) and salicylic acid (SA) as well as transcript levels of their downstream marker genes. JA biosynthesis originates from linolenic acid, which is converted to 13-HPOT by LOX. This compound then serves either as substrate for the enzyme HPL to generate various C<sub>12</sub> traumatic acids, or for AOS to ultimately yield JA-Ile. Arrows indicate biosynthetic steps carried out by a single enzyme, some of them are shown in red letters. Note that SA can be synthesized via different pathways and how this is done upon mite infestation is still unclear. Dashed arrows symbolize transcriptional activation of marker genes (which are in italic face). Amounts of linolenic acid and HPL transcripts were not assayed in this study.






**FIGURE S5.2.** Survival, migration and mortality in two strains of the spider mite *Tetranychus urticae* which did (+) or did not (-) contain *Wolbachia*, *Spiroplasma* and/or *Candidatus Cardinium* after feeding on tomato (*Solanum lycopersicum*) for 4 days. (a) Four lines of the plant defence suppressor strain DeLier-1 were tested (1, 2, 3 and 4). Each line was subdivided into three mite groups: W+S+, W-S+ and W-S-; (b) Two lines (5 and 8) were tested for the plant defence inducer strain Santpoort-2 with two mite groups: C+S+ and C-S-. Additionally we tested C+S+ of line 6. C-S- of line 6 as well as both groups of line 7 went extinct before the experiment.





**FIGURE S5.3.** Reproductive performance (number of eggs produced per female in four days) of two lines (5, and 8) of a plant defence inducer strain Santpoort-2 of the spider mite *Tetranychus urticae*. Each of the lines was subdivided into two groups which did (+) or did not (-) contain the bacteria *Candidatus Cardinium* and *Spiroplasma*: C+S+ (grey) and C-S- (white). Additionally we tested C+S+ of line 6; the C-S- groups of line 6 as well as both groups of line 7 went extinct before the experiment. Boxes span the 25-75 percentiles, horizontal lines in the boxes represent medians, whiskers span 1.5\*IQR (interquartile range), dots represent data points outside of this range. Different letters above the boxes indicate significant differences at a level of  $P \leq 0.05$  (tested per line) after applying a linear mixed model.

## Mite-associated bacteria




**FIGURE S5.4.** Transcription heatmap of differentially expressed genes in the *T. urticae* suppressor strain DeLier-1 after removal of *Wolbachia* (W-S+) and *Wolbachia* and *Spiroplasma* (W-S-) as compared to transcription levels of the original strain (W+S+) that contained both bacteria. A total of 173 differentially expressed genes were detected after *Wolbachia* (W-S+) and *Wolbachia-Spiroplasma* removal (W-S-) and between the two treatments, mutually (FDR-corrected  $P \leq 0.05$ ). Using Euclidean distance metrics, genes were clustered based on their relative transcript levels in the W-S+ vs W+S+ and W-S- vs W+S+ comparisons. In the left sidebar, orange and green rectangles indicate genes that code for enzymes involved in the xenobiotic and digestive pathways of *T. urticae*, respectively.






**FIGURE S5.5.** Phytohormone amounts in tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with four lines (1, 2, 3 and 4) of a plant defence suppressor strain DeLier-1 of the spider mite *Tetranychus urticae*. Each of the lines was subdivided into three groups which did (+) or did not (-) contain the bacteria *Wolbachia* and *Spiroplasma*: W+S+ (dark grey), W-S+ (light grey) and W-S- (white). Control plants were not infested (darkest grey). Tested phytohormones included (a) 12-oxo-phytodienoic acid (OPDA), (b) 3-oxo-2-(2-pentenyl)-cyclopentane-1-hexanoic acid (OPC6), (c) jasmonic acid (JA) (d) jasmonic acid-isoleucine (JA-Ile), (e) 9-OH-traumatoin, (f) free salicylic acid (SA); Circles represent individual data points, horizontal lines indicate the medians. Phytohormone amounts are presented as nanogram (ng) per gram fresh leaf weight (g FW). Compounds for which we did not have a pure standard are presented as ion counts (IC /g FW).




**FIGURE S5.6.** Phytohormone amounts in tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with four lines (5, 6, 7 and 8) of a plant defence inducer strain Santpoort-2 of the spider mite *Tetranychus urticae*. Each of the lines was subdivided into two groups which did (+) or did not (-) contain the bacteria *Candidatus Cardinium* and *Spiroplasma*: C+S+ (grey), C-S- (white). Control plants were not infested (dark grey). Tested phytohormones included (a) jasmonic acid-isoleucine (JA-Ile), (b) free salicylic acid (SA) (c) traumatic acid, (d) abscisic acid (ABA) (f); Circles represent individual data points, horizontal lines indicate the medians. Phytohormone amounts are presented as nanogram (ng) per gram fresh leaf weight (g FW). Compounds for which we did not have a pure standard are presented as ion counts (IC /g FW).

Mite-associated bacteria




**FIGURE S5.7.** Normalized expression (NE) of plant defence related genes (qRT-PCR) in tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with four lines (1, 2, 3 and 4) of a plant defence suppressor strain DeLier-1 of the spider mite *Tetranychus urticae*. Each of the lines was subdivided into three groups which did (+) or did not (-) contain the bacteria *Wolbachia* and *Spiroplasma*: W+S+ (dark grey), W-S+ (light grey) and W-S- (white). Control plants were not infested (darkest grey). (a) *12-oxophytodienoate Reductase 3 (OPR3)*, (b) *Tomato wound-induced 1 (TWI-1)*, (c) *Alcohol dehydrogenase (ADH)*, (d) *Glutathione S-transferase 6 (GST6)*, (e) *Glutaredoxin (GRX)*, (f) *Jasmonate-inducible protein-21 (JIP-21)*, (g) *Pathogenesis-related protein 6 (PR-P6)*; Circles represent individual data points, horizontal lines indicate the medians.




**FIGURE S5.8.** Correlations between phytohormone amounts of salicylic acid (SA) and 12-oxo-phytodienoic acid (OPDA) and the normalized expression levels of putative OPDA responsive genes (*TWI-1*, *GRX*, *GST-6*, *ADH* and *OPR3*) measured in tomato leaflets after 7 days of infestations with the plant defence suppressor strain DeLier-1 of the spider mite *Tetranychus urticae*. *TWI-1*: Tomato wound-induced 1, *GRX*: Glutaredoxin, *ADH*: Alcohol dehydrogenase, *GST6*: Glutathione S-transferase 6;  $P$ -values were adjusted for multiple comparisons with the holm method.


## Mite-associated bacteria



**FIGURE S5.9.** Correlations between phytohormone amounts of salicylic acid (SA) and the normalized expression levels of the two SA marker genes *PR-P6* and *PR-1a*, measured in tomato leaflets after 7 days of infestations with the plant defence suppressor strain DeLier-1 of the spider mite *Tetranychus urticae*. *PR-P6*: *Pathogenesis-related protein 6*, *PR-1a*: *Pathogenesis-related protein 1a*; *P*-values were adjusted for multiple comparisons with the holm method.



**FIGURE S5.10.** Normalized expression (NE) of plant defence related genes (qRT-PCR) in tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with four lines (5, 6, 7 and 8) of a plant defence inducer strain Santpoort-2 of the spider mite *Tetranychus urticae*. Each of the lines was subdivided into two groups which did (+) or did not (-) contain the bacteria *Candidatus Cardinium* and *Spiroplasma*: C+S+ (grey) and C-S- (white). Control plants were not infested (dark grey). (a) *Pathogenesis-related protein 6* (*PR-P6*), (b) *Pathogenesis-related protein 1a* (*PR-1a*); Circles represent individual data points, horizontal lines indicate the medians.



**FIGURE S5.11.** Feeding damage (mm<sup>2</sup>) on tomato (*Solanum lycopersicum*) leaflets after 7 days of infestation with four lines (5, 6, 7 and 8) of a plant defence inducer strain Santpoort-2 of the spider mite *Tetranychus urticae*. Each of the lines was subdivided into two groups which did (+) or did not (-) contain the bacteria *Candidatus Cardinium* and *Spiroplasma*: C+S+ (grey) and C-S- (white). Circles represent individual data points, horizontal lines indicate the medians.

**TABLE S5.1.** Parameters used for detection of phytohormones and related compounds by LC-MS/MS

Compound	Q <sub>1</sub> <sup>*1</sup>	Q <sub>3</sub> <sup>*2</sup>	CID <sup>*3</sup>	CE <sup>*4</sup>	Detected	Reference
SA	137	93	-35	15.0	yes	Wu <i>et al.</i> , 2007
D <sub>6</sub> -SA (internal standard)	141	97	-35	15.0	yes	Alba <i>et al.</i> , 2015
ABA	263	153	-35	9.0	yes	Bonaventure, 2011
13-HPOT	309	291	-35	5.0	yes	Kallenbach <i>et al.</i> , 2010 <sup>*5</sup>
dinor-OPDA	263	165	-35	18.0	no	Bao <i>et al.</i> , 2014
OPDA	291	165	-35	18.0	yes	Koo <i>et al.</i> , 2009
OPDA-GSH	598	306	-35	18.0	no	Dąbrowska <i>et al.</i> , 2009
OPC-8	293	275	-35	12.0	yes	Kallenbach <sup>*5</sup> ; Bao <i>et al.</i> , 2014
OPC-6	265	247	-35	12.0	yes	Kallenbach <sup>*5</sup>
OPC-4	237	219	-35	12.0	no	Kallenbach <sup>*5</sup>
JA	209	59	-35	12.0	yes	Wu <i>et al.</i> , 2007
D <sub>3</sub> -JA (internal standard)	213	61	-35	12.0	yes	Alba <i>et al.</i> , 2015
12-OH-JA	225	59	-35	19.0	no	Stitz <i>et al.</i> , 2011
12-OH-JA-Ile	338	130	-35	19.0	yes	Stitz <i>et al.</i> , 2011
12-COOH-JA-Ile	352	130	-35	19.0	no	Stitz <i>et al.</i> , 2011
12-oxo-(9Z)-dodecenoic acid	211	183	-35	13.5	no	Kallenbach <i>et al.</i> , 2011
(2E-) and (3Z)-dodecenedioic acid (traumatic acids)	227	183	-35	13.0	yes	Kallenbach <i>et al.</i> , 2011
12-OH-(9Z)-dodecenoic acid	213	183	-35	15.5	no	Kallenbach <i>et al.</i> , 2011
9-OH-12-oxo-(10E)-dodecenoic acid (9-OH-traumatin)	227	209	-35	9.5	yes	Kallenbach <i>et al.</i> , 2011
4-OH- (2E)-dodecenedioic acid (4-OH-traumatic acid)	243	225	-35	9.5	yes	Kallenbach <i>et al.</i> , 2011
9,12-OH-(10E)-dodecenoic acid	229	211	-35	9.5	yes	Kallenbach <i>et al.</i> , 2011
9,12-OH-(10E)-dodecanoic acid	231	213	-35	9.5	no	Kallenbach <i>et al.</i> , 2011
9-OH-traumatin-GSH	534	306	-35	16.5	no	Kallenbach <i>et al.</i> , 2011
JA-Ala	280	88	-35	12.0	no	Koo <i>et al.</i> , 2009
JA-Gly	266	74	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Val	308	116	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Ile / JA-Leu	322	130	-35	19.0	yes	Wu <i>et al.</i> , 2007
JA-Pro	306	114	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Ser	296	104	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Thr	310	118	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Met	340	148	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-His	346	154	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Phe	356	164	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Tyr	372	180	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Trp	395	203	-35	12.0	no	Wang <i>et al.</i> , 2007
JA-Gln	337	145	-35	19.0	no	Stitz <i>et al.</i> , 2011
JA-ACC	292	100	-35	12.0	yes	Wang <i>et al.</i> , 2007
JA-glucose	417	209	-35	18.0	no	VanDoorn <i>et al.</i> , 2011

\*<sup>1</sup> Q<sub>1</sub> mass = Molecular Weight - 1 = [M - H]<sup>-</sup> (m/z); \*<sup>2</sup> CID Q<sub>3</sub> mass (of daughter with 100% intensity) (m/z); \*<sup>3</sup> Capillary Collision-Induced Dissociation (V); \*<sup>4</sup> Collision Energy (V); \*<sup>5</sup> personal communication.

TABLE S5.2. Nucleotide sequence of primers used for PCR and (g)RT-PCR analysis.

Target organism(s)	Target gene	Name	Gene identifier (BOGAS/ITAG2.3)	Forward primer 5' → 3'	Reverse primer 5' → 3'	References
<i>Wolbachia pipiensis</i>	<i>16S rRNA</i>	<i>16S ribosomal RNA</i>	-	TTGTAGCCTCTATGGTAACT	GAATAGGATAGATTTTCATGT	O'Neill <i>et al.</i> , 1992
<i>Cardinium</i> sp. (CLO); cytophaga like organism)	<i>16S rRNA</i>	<i>16S ribosomal RNA</i>	-	GCGGTGTAAAAATGAGCGTG	ACCTMTTCTTAACCTCAAGCCT	Weeks <i>et al.</i> , 2003
<i>Sproplasma</i> sp.	<i>dnaA</i>	<i>DnaA</i>	-	ATTCTTCAGTAAAAATGCTTTGGA	ACACATTTACTTCATGCTATTGA	Fukatsu <i>et al.</i> , 2001
Bacteria (general)	<i>16S rRNA</i>	<i>16S ribosomal RNA</i>	-	TCCTACGGGNGGCWGCAG	TGACTACHVGGGTATCTAAKCC	Klindworth <i>et al.</i> , 2012
<i>Tetranychus urticae</i>	<i>Actin</i>	<i>Actin</i>	Tetur03g09480	CAGCCATGTATGTGGCATC	AAATCACGACAGCCAAATC	Feng <i>et al.</i> , 2010
<i>Solanum lycopersicum</i>	<i>AOSI</i>	<i>Allene Oxide Synthase 1</i>	Solye04g079730.1	AACAGTGTCCCGGAAAAGAC	AATGGAGATGCACCGACTTC	Howe <i>et al.</i> , 2000; Sivasankar <i>et al.</i> , 2000
	<i>OPR3</i>	<i>OPDA reductase 3</i>	Solye07g007870.2	GATCCAGTTGTGGGATACACAG	GCCCAACAAAATCAGGTTTC	Strassner <i>et al.</i> , 2002
	<i>TW1-1</i>	<i>Tomato Wound-induced 1</i>	Solye01g107820.2	CATCTTACAATGGATGGGCTAC	CGAGATGATTTGATCTTGGATTTC	Truesdale <i>et al.</i> , 1996
	<i>ADH</i>	<i>Alcohol Dehydrogenase</i>	Solye01g087640.2	GTCTTGAGTTGAGCGGTGAAGG	CAGGCCTAGTGTATTCGGTTTC	Tieman <i>et al.</i> , 2007
	<i>GST6</i>	<i>Gluthathione S-transferase</i>	Solye06g009020.2	GTGAAGAGCTTGTTCGATGC	CCTTCTTTCAACGGCGATAC	This study
	<i>GRX</i>	<i>Glutaredoxin</i>	Solye07g053550.1	ATGATGCAACAAAGCACTTCC	GATGATGTCGATCAACTCTTGG	This study
	<i>JIP-2/1</i>	<i>Jasmonate-inducible protein 2/1</i>	Solye03g098790.1	ACTCGTCTGTGCTTTGTCC	CCCAAGAGGATTTTTCGTGA	Lisón <i>et al.</i> , 2006
	<i>TD2</i>	<i>Threonine Deaminase-2</i>	Solye09g008670.2	TGCCGTTAAAAATGTCACCA	ACTGGCGATGCCAAAATATC	Chen <i>et al.</i> , 2005
	<i>PR-1a</i>	<i>Pathogenesis-related protein 1a</i>	Solye09g007010.1	TGGTGGTTCATTTCTTGCACACTAC	ATCAATCCGATCCACTATCATTTTA	van Kan <i>et al.</i> , 1992
	<i>PR-P6</i>	<i>Pathogenesis-related protein P6</i>	Solye00g174340.1	GTACTGCACTCTTCTTGTTTCCA	TAGATAAGTCTTGTGATGTGCC	van Kan <i>et al.</i> , 1992
	<i>P1-Iic</i>	<i>Proteinase Inhibitor Iic</i>	Solye03g020050.2	CAGGATGTACGACGGTGTTC	GAGTTGCAACCCCTCTCCITG	Gaddea <i>et al.</i> , 1996
	<i>Actin</i>	<i>Actin</i>	Solye03g078400.2	TCAGCACATTCACAGCAGATGT	AACAGACAGGACACTCGCACT	Tomato Genome Consortium, 2012

**TABLE S5.3a.** List of top-20 genes significantly up-regulated in adult female *Tetranychus urticae* suppressor (DeLier-1) W-S+ mites as compared to W+S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after  $P$ -value adjustment).

#	Locus identifier	Annotation (BOGAS)	Fold change	$P$
1	tetur06g06585	Hypothetical protein (snosnR61, small nucleolar RNA)	1.81	0.0086
2	tetur18g02170	Hypothetical protein (no hits found)	1.54	0.0031
3	tetur24g01690	Hypothetical protein (similar to sorting nexin 13)	1.49	0.0261
4	tetur05g04420	Hypothetical protein (choline/ethanolaminephosphotransferase 1)	1.45	0.0344
5	tetur31g00280 <sup>d</sup>	CSPCA1: CUB domain-containing Secreted Protein Conserved in Arthropods	1.44	0.0185
6	tetur39g00780	Hypothetical protein (vitellogenin1)	1.35	0.0449
7	tetur04g06140 <sup>d</sup>	Hypothetical protein (IscW_ISCW012949)	1.34	0.0140
8	tetur05g02950	CYP389C2: Cytochrome P450	1.31	0.0127
9	tetur23g01800	Hypothetical protein (Rho GTPase-activating protein 29)	1.30	0.0295
10	tetur09g04980	Hypothetical protein (no hits found)	1.29	0.0069
11	tetur10g04770	Hypothetical protein (leukocyte elastase inhibitor)	1.28	0.0216
12	tetur60g00070	Hypothetical protein (TcasGA2_TC006277)	1.25	0.0184
13	tetur07g01880	SSPE2: Small Secreted Protein, Family E	1.25	0.0143
14	tetur06g03890	Hypothetical protein (trichohyalin, putative)	1.24	0.0483
15	tetur05g08770 <sup>d</sup>	Hypothetical protein (intracellular protein transport)	1.23	0.0014
16	tetur05g03440	Hypothetical protein (transmembrane and coiled-coil domains protein 2-like)	1.22	0.0414
17	tetur37g01020	Hypothetical protein (M-phase inducer phosphatase)	1.21	0.0086
18	tetur23g01870	Hypothetical protein (tyrosine kinase, putative)	1.21	0.0146
19	tetur02g10760	Hypothetical protein (no hits found)	1.20	0.0190
20	tetur06g01790	Hypothetical protein (mRNA splicing protein SMN)	1.20	0.0344

<sup>d</sup> these genes were significantly up-regulated in *T. urticae* suppressor W-S- mites as compared to W+S+ mites.

**TABLE S5.3b.** List of top-20 genes significantly down-regulated in adult female *Tetranychus urticae* suppressor (DeLier-1) W-S+ mites as compared to W+S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after  $P$ -value adjustment).

#	Locus identifier	Annotation (BOGAS)	Fold change	$P$
1	tetur16g03730 <sup>d</sup>	Hypothetical protein (similar to WT Secreted Protein 12)	-3.44	0.0020
2	tetur16g03680 <sup>d</sup>	TuPap-31: Cathepsin L	-3.28	0.0014
3	tetur16g03740 <sup>d</sup>	WTSP12: WT Secreted Protein 12	-3.25	0.0014
4	tetur16g03620 <sup>d</sup>	Hypothetical protein (similar to WT Secreted Protein 12)	-2.74	0.0032
5	tetur196g00010 <sup>d</sup>	Hypothetical protein (Legumain)	-2.64	0.0025
6	tetur44g00311	Hypothetical protein (transposon)	-2.58	0.0121
7	tetur32g02297	SSPF13: Small Secreted Protein, Family F	-2.56	0.0223
8	tetur16g03670	TuLeg-13: Legumain	-2.52	0.0031
9	tetur16g03610	WTSP4: WT Secreted Protein, pseudogene	-2.44	0.0020
10	tetur31g00810	Hypothetical protein (similar to methyl-accepting chemotaxis sensory transducer, extracellular)	-2.43	0.0321
11	tetur16g03770	TuPap-44: Cathepsin L	-2.35	0.0086
12	tetur128g00030	Hypothetical protein (Serp13 inhibitory serine protease inhibitor, putative secreted salivary gland protein)	-2.32	0.0365
13	tetur31g00830	Hypothetical protein (methyl-accepting chemotaxis sensory transducer, extracellular)	-2.17	0.0310

## Mite-associated bacteria

14	tetur01g06610 <sup>d</sup>	TuLeg-17: Legumain	-2.11	0.0062
15	tetur39g00220	Hypothetical protein (extracellular)	-2.08	0.0143
16	tetur11g05410	Hypothetical protein (MFS-transporter)	-2.00	0.0030
17	tetur06g03540	TuLeg-7: Legumain	-1.99	0.0044
18	tetur01g00490	ID-RCD: intradiol ring-cleavage dioxygenase	-1.96	0.0101
19	tetur33g01640	Hypothetical protein (extracellular)	-1.96	0.0179
20	tetur03g02710 <sup>d</sup>	Hypothetical protein (phospholipid scramblase 2-like)	-1.89	0.0014

<sup>d</sup> these genes were significantly down-regulated in *T. urticae* suppressor W-S- mites as compared to W+S+ mites.

**TABLE S5.4a.** List of all genes significantly up-regulated in adult female *Tetranychus urticae* suppressor (DeLier-1) W-S- mites as compared to W+S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after *P*-value adjustment).

W, *Wolbachia* sp.; S, *Spiroplasma* sp.; +, present; -, absent

#	Locus identifier	Annotation (BOGAS)	Fold change	<i>P</i>
1	tetur06g06650 <sup>y</sup>	TuPap-19: Cystatin (oncocystatin)	3.60	0.0197
2	tetur06g01060 <sup>b y</sup>	TuCPI-22: Cystatin (cystatin precursor)	2.03	0.0008
3	tetur02g15167	SP68: Serine Protease (Peptidase S1)	1.63	0.0443
4	tetur15g00310	CSPA1: Cell Surface Protein with GTPLASR(K,R,S) repeats	1.63	0.0247
5	tetur11g00700	Hypothetical protein (tribbles homolog 2, serine/threonine-protein kinase)	1.55	0.0151
6	tetur17g02060 <sup>b</sup>	Hypothetical protein (secreted salivary gland peptide, putative)	1.54	0.0002
7	tetur31g00280 <sup>d</sup>	CSPCA1: CUB domain-containing Secreted Protein Conserved in Arthropods	1.53	0.0176
8	tetur15g00320	CSPA2: Cell Surface Protein with GTPLASR(K,R,S) repeats	1.49	0.0443
9	tetur04g06140 <sup>d</sup>	Hypothetical protein (IscW_ISCW012949)	1.36	0.0197
10	tetur11g00270	Hypothetical protein (signal recognition particle 14 kDa protein)	1.33	0.0493
11	tetur11g04330	Hypothetical protein (ADP-ribosylation factor GTPase-activating protein 3)	1.24	0.0324
12	tetur05g08770 <sup>d</sup>	Hypothetical protein (no hits found)	1.19	0.0151
13	tetur04g05590	TuGR62: Chemosensory Receptor, Gustatory receptor family	1.17	0.0392
14	tetur14g02440	Hypothetical protein (An09g06400, <i>Aspergillus niger</i> chitinase)	1.15	0.0369

<sup>b</sup> these genes were significantly up-regulated in *T. urticae* suppressor W-S- mites as compared to W-S+ mites. <sup>d</sup> these genes were significantly up-regulated in suppressor W-S+ mites as compared to W+S+ mites. <sup>y</sup> these genes were significantly up-regulated in *T. urticae* inducer (Santpoort-2) C-S- mites as compared to C+S+ mites. C, *Candidatus* Cardinium

**TABLE S5.4b.** List of top-20 genes significantly down-regulated in adult female *Tetranychus urticae* suppressor (DeLier-1) W-S- mites as compared to W+S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after *P*-value adjustment).

W, *Wolbachia* sp.; S, *Spiroplasma* sp.; +, present; -, absent

#	Locus identifier	Annotation (BOGAS)	Fold change	<i>P</i>
1	tetur03g08800 <sup>b d</sup>	SSPB2: Small secreted protein, family B	-3.36	0.0000
2	tetur03g05070	CYP392D8: Cytochrome P450	-2.90	0.0008
3	tetur16g03730 <sup>d</sup>	Hypothetical protein (no hits found)	-2.53	0.0315
4	tetur03g09961	CYP392D7: Cytochrome P450	-2.34	0.0197

5	tetur16g03620 <sup>d</sup>	Hypothetical protein (similar to WT Secreted Protein 12)	-2.21	0.0392
6	tetur16g03740 <sup>d</sup>	WTSP12: WT Secreted Protein 12	-2.13	0.0498
7	tetur16g03680 <sup>d</sup>	TuPap-31: Cathepsin L	-2.12	0.0443
8	tetur03g08810 <sup>d</sup>	SSPB1: Small Secreted Protein, Family B	-2.04	0.0151
9	tetur19g00010 <sup>d</sup>	Hypothetical protein (Legumain)	-2.01	0.0455
10	tetur01g06610 <sup>d</sup>	TuLeg-17: Legumain	-2.00	0.0199
11	tetur03g02710 <sup>d</sup>	Hypothetical protein (phospholipid scramblase 2-like)	-1.59	0.0197
12	tetur03g07920 <sup>d</sup>	TuGSTd06: Glutathione S-transferase, delta class	-1.58	0.0262
13	tetur28g01730 <sup>d</sup>	Hypothetical protein (3-oxoacyl-acyl-carrier-protein reductase)	-1.58	0.0315
14	tetur06g02930 <sup>d</sup>	TuPap-45: fibronase precursor, cathepsin L	-1.55	0.0186
15	tetur01g03150 <sup>d</sup>	CCEinc-01: Carboxyl/choline esterase, incomplete	-1.50	0.0455
16	tetur22g02500 <sup>d</sup>	Hypothetical protein (No hits found)	-1.50	0.0151
17	tetur16g03750 <sup>d</sup>	Hypothetical protein (3-Pan modules Cell Surface Protein, antigen-2-like)	-1.50	0.0460
18	tetur03g05030	CYP392D6: Cytochrome P450	-1.44	0.0239
19	tetur02g10560 <sup>d</sup>	AGO-1G: ortholog of Argonaute-1 ( <i>Drosophila melanogaster</i> )	-1.44	0.0307
20	tetur01g03410 <sup>d</sup>	Hypothetical protein (apical endosomal glycoprotein)	-1.40	0.0418

<sup>b</sup> this gene was significantly down-regulated in *T. urticae* suppressor W-S- mites as compared to W-S+ mites. <sup>d</sup> these genes were significantly down-regulated in suppressor W-S+ mites as compared to W+S+ mites.

**TABLE S5.5.** List of all genes significantly up- and down-regulated respectively, in adult female *Tetranychus urticae* suppressor (DeLier-1) W-S- mites as compared to W-S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after *P*-value adjustment).

W, *Wolbachia* sp.; S, *Spiroplasma* sp.; +, present; -, absent

#	Locus identifier	Annotation (BOGAS)	Fold change	<i>P</i>
1	tetur06g01060 <sup>v</sup>	TuCPI-22: Cystatin (cystatin precursor)	1.84	0.0187
2	tetur17g02060 <sup>v</sup>	Hypothetical protein (secreted salivary gland peptide, putative)	1.39	0.0187
1	tetur03g08800 <sup>x</sup>	SSPB2: Small secreted protein, family B	-1.96	0.0235

<sup>v</sup> these genes were significantly up-regulated in *T. urticae* suppressor W-S- mites as compared to W+S+ mites and in inducer (Santpoort-2) C-S- mites as compared to C+S+ mites. <sup>x</sup> this gene was significantly down-regulated in both suppressor W-S- and W-S+ mites as compared to W+S+ mites. C, *Candidatus Cardinium*

**TABLE S5.6a.** List of all genes significantly up-regulated in adult female *Tetranychus urticae* inducer (Santpoort-2) C-S- mites as compared to C+S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after *P*-value adjustment).

C, *Candidatus Cardinium*; S, *Spiroplasma* sp.; +, present; -, absent

#	Locus identifier	Annotation (BOGAS)	Fold change	<i>P</i>
1	tetur11g05780 <sup>o</sup>	Hypothetical protein (no hits found)	5.17	0.0046
2	tetur06g06650 <sup>h</sup>	TuPap-19: Cystatin (onchocystatin)	5.03	0.0001
3	tetur06g01060 <sup>p</sup>	TuCPI-22: Cystatin (cystatin precursor)	3.02	0.0001
4	tetur11g05760	TuCCE-33: Carboxyl/choline esterase (putative esterase)	2.84	0.0288
5	tetur37g00580	Hypothetical protein (4-nitrophenylphosphatase)	2.32	0.0008
6	tetur08g08060	Hypothetical protein (secreted salivary gland peptide, putative)	2.11	0.0008

## Mite-associated bacteria

7	tetur17g02060 <sup>P</sup>	Hypothetical protein (secreted salivary gland peptide, putative)	1.96	0.0004
8	tetur01g02310	5TM1: 5-pass Trans-membrane protein of unknown function	1.80	0.0337
9	tetur11g00550	Hypothetical protein (MFS-transporter)	1.72	0.0046
10	tetur08g07240	Hypothetical protein (secreted salivary gland peptide, putative)	1.61	0.0066
11	tetur11g05000 <sup>o</sup>	CYP385C2: Cytochrome P450	1.61	0.0313
12	tetur23g00470	Hypothetical protein (no hits found)	1.58	0.0108
13	tetur03g05040	Hypothetical protein (Cytochrome P450 – fragment)	1.55	0.0398
14	tetur33g01340	Hypothetical protein (ribosomal protein)	1.47	0.0017
15	tetur03g04990	CYP392D2: Cytochrome P450	1.46	0.0143
16	tetur02g09850	TuUT11: UDP-glycosyltransferase (UGT)	1.45	0.0185
17	tetur01g10390	TuABCC-02: ABC-transporter, class C	1.43	0.0064
18	tetur02g02190	Hypothetical protein (tRNA)	1.41	0.0100
19	tetur10g01570	TuCAS: $\beta$ -cyanoalanine synthase	1.38	0.0100
20	tetur13g03000	Hypothetical protein (no hits found)	1.36	0.0023
21	tetur89g00030	Hypothetical protein (no hits found)	1.33	0.0307
22	tetur11g05670	Hypothetical protein (ornithine aminotransferase)	1.32	0.0145
23	tetur11g05680	Hypothetical protein (vacuolar protein sorting 13 homolog D)	1.29	0.0337

<sup>h</sup> this gene was significantly up-regulated in *T. urticae* suppressor (DeLier-1) W-S- mites as compared to W+S+ mites. <sup>o</sup> these genes were significantly down-regulated in suppressor W-S+ mites as compared to W+S+ mites. <sup>p</sup> these genes were significantly up-regulated in suppressor W-S- mites as compared to both W+S+ and W-S+ mites. W, *Wolbachia* sp.

**TABLE S5.6b.** List of all genes significantly down-regulated in adult female *Tetranychus urticae* inducer (Santpoort-2) C-S- mites as compared to C+S+ mites, after feeding from tomato (*Solanum lycopersicum*) for seven days ( $P \leq 0.05$ , after  $P$ -value adjustment).

C, *Candidatus* Cardinium; S, *Spiroplasma* sp.; +, present; -, absent

#	Locus identifier	Annotation (BOGAS)	Fold change	$P$
1	tetur11g05550	Hypothetical protein (MFS-transporter)	-1.59	0.0219
2	tetur18g02050	Hypothetical protein (no hits found)	-1.55	0.0369
3	tetur11g05720	PLAT10: Lipase/lipoxygenase	-1.47	0.0066
4	tetur02g13460	Hypothetical protein (very acidic salivary protein)	-1.46	0.0337
5	tetur01g13560	ZN207: Zinc Finger protein 207	-1.37	0.0436
6	tetur28g01570	Hypothetical protein (deoxyhypusine synthase)	-1.36	0.0185
7	tetur11g06310	Hypothetical protein (galactose-binding domain-like)	-1.36	0.0079
8	tetur05g03510	Hypothetical protein (ribosome production factor 2 homolog)	-1.35	0.0143
9	tetur15g00160	Hypothetical protein (no hits found)	-1.30	0.0329
10	tetur27g02556	SLCP1: secreted low complexity protein, putative	-1.29	0.0313

## SUPPLEMENTARY REFERENCES

- Alba, J.M., Schimmel, B.C.J., Glas, J.J., Ataide, L.M.S., Pappas, M.L., Villarreal, C.A., Schuurink, R.C., Sabelis, M.W. & Kant, M.R. 2015. Spider mites suppress tomato defenses downstream of jasmonate and salicylate independently of hormonal crosstalk. *New Phytol* **205**: 828-840.
- Bao, J., Gao, X. & Jones, A.D. 2014. Unusual negative charge-directed fragmentation: collision-induced dissociation of cyclopentenone oxylipins in negative ion mode. *Rapid Commun Mass Sp* **28**: 457-464.

- Bonaventure, G. 2011. The *Nicotiana attenuata* LECTIN RECEPTOR KINASE 1 is involved in the perception of insect feeding. *Plant Signal Behav* **6**: 2060-2063.
- Chen, H., Wilkerson, C.G., Kuchar, J.A., Phinney, B.S. & Howe, G.A. 2005. Jasmonate-inducible plant enzymes degrade essential amino acids in the herbivore midgut. *Proc Natl Acad Sci USA* **102**: 19237-19242.
- Consortium, T.G. 2012. The tomato genome sequence provides insights into fleshy fruit evolution. *Nature* **485**: 635-641.
- Dąbrowska, P., Freitak, D., Vogel, H., Heckel, D.G. & Boland, W. 2009. The phytohormone precursor OPDA is isomerized in the insect gut by a single, specific glutathione transferase. *Proc Natl Acad Sci USA* **106**: 16304-16309.
- Feng, H., Wang, L., Liu, Y., He, L., Li, M., Lu, W. & Xue, C. 2010. Molecular characterization and expression of a heat shock protein gene (HSP90) from the carmine spider mite, *Tetranychus cinnabarinus* (Boisduval). *J Insect Sci* **10**: 112
- Fukatsu, T., Tsuchida, T., Nikoh, N. & Koga, R. 2001. *Spiroplasma* symbiont of the pea aphid, *Acyrtosiphon pisum* (Insecta: Homoptera). *Appl Environ Microbiol* **67**: 1284-1291.
- Gadea, J., Mayda, M.E., Conejero, V. & Vera, P. 1996. Characterization of defense-related genes ectopically expressed in viroid-infected tomato plants. *Mol Plant Microbe Interact* **9**: 409-415.
- Howe, G.A., Lee, G.I., Itoh, A., Li, L. & DeRocher, A.E. 2000. Cytochrome P450-dependent metabolism of oxylipins in tomato. Cloning and expression of allene oxide synthase and fatty acid hydroperoxide lyase. *Plant Physiol* **123**: 711-724.
- Kallenbach, M., Alagna, F., Baldwin, I.T. & Bonaventure, G. 2010. *Nicotiana attenuata* SIPK, WIPK, NPR1, and fatty acid-amino acid conjugates participate in the induction of jasmonic acid biosynthesis by affecting early enzymatic steps in the pathway. *Plant Physiol* **152**: 96-106.
- Kallenbach, M., Gilardoni, P.A., Allmann, S., Baldwin, I.T. & Bonaventure, G. 2011. C<sub>12</sub> derivatives of the hydroperoxide lyase pathway are produced by product recycling through lipoxygenase-2 in *Nicotiana attenuata* leaves. *New Phytol* **191**: 1054-1068.
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. & Glöckner, F.O. 2012. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res*: gks808.
- Koo, A.J., Gao, X., Daniel Jones, A. & Howe, G.A. 2009. A rapid wound signal activates the systemic synthesis of bioactive jasmonates in *Arabidopsis*. *Plant J* **59**: 974-986.
- Lisón, P., Rodrigo, I. & Conejero, V. 2006. A novel function for the cathepsin D inhibitor in tomato. *Plant Physiol* **142**: 1329-1339.
- O'Neill, S.L., Giordano, R., Colbert, A., Karr, T.L. & Robertson, H.M. 1992. 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proc Natl Acad Sci USA* **89**: 2699-2702.
- Sivasankar, S., Sheldrick, B. & Rothstein, S.J. 2000. Expression of allene oxide synthase determines defense gene activation in tomato. *Plant Physiol* **122**: 1335-1342.
- Stitz, M., Gase, K., Baldwin, I.T. & Gaquerel, E. 2011. Ectopic expression of AtJMT in *Nicotiana attenuata*: creating a metabolic sink has tissue-specific consequences for the jasmonate metabolic network and silences downstream gene expression. *Plant Physiol* **157**: 341-354.
- Strassner, J., Schaller, F., Frick, U.B., Howe, G.A., Weiler, E.W., Amrhein, N., Macheroux, P. & Schaller, A. 2002. Characterization and cDNA-microarray expression analysis of 12-oxophytodienoate reductases reveals differential roles for octadecanoid biosynthesis in the local versus the systemic wound response. *Plant J* **32**: 585-601.
- Tieman, D.M., Loucas, H.M., Kim, J.Y., Clark, D.G. & Klee, H.J. 2007. Tomato phenylacetaldehyde reductases catalyze the last step in the synthesis of the aroma volatile 2-phenylethanol. *Phytochemistry* **68**: 2660-2669.

## Mite-associated bacteria

- Truesdale, M.R., Doherty, H.M., Loake, G.J., McPherson, M.J., Roberts, M. & Bowles, D.J. 1996. Molecular cloning of a novel wound-induced gene from tomato: *twi1*. *Plant Physiol* **112**: 446.
- van Kan, J.A., Joosten, M.H., Wagemakers, C.A., van den Berg-Velthuis, G.C. & de Wit, P.J. 1992. Differential accumulation of mRNAs encoding extracellular and intracellular PR proteins in tomato induced by virulent and avirulent races of *Cladosporium fulvum*. *Plant Mol Biol* **20**: 513-527.
- VanDoorn, A., Bonaventure, G., Schmidt, D.D. & Baldwin, I.T. 2011. Regulation of jasmonate metabolism and activation of systemic signaling in *Solanum nigrum*: COI1 and JAR4 play overlapping yet distinct roles. *New Phytol* **190**: 640-652.
- Wang, L., Halitschke, R., Kang, J.-H., Berg, A., Harnisch, F. & Baldwin, I.T. 2007. Independently silencing two JAR family members impairs levels of trypsin proteinase inhibitors but not nicotine. *Planta* **226**: 159-167.
- Weeks, A.R., Velten, R. & Stouthamer, R. 2003. Incidence of a new sex-ratio-distorting endosymbiotic bacterium among arthropods. *Proc R Soc B: Biol Sci* **270**: 1857-1865.
- Wu, J., Hettenhausen, C., Meldau, S. & Baldwin, I.T. 2007. Herbivory rapidly activates MAPK signaling in attacked and unattacked leaf regions but not between leaves of *Nicotiana attenuata*. *Plant Cell* **19**: 1096-1122.

6

## General discussion

Virtually all macro-organisms are associated with microbes. If microbe-host associations persist across generations, a host and its associated microbes (holobiont) can be viewed as a unit of selection (Margulis & Chapman, 1998; Rohwer *et al.*, 2002; Zilber-Rosenberg & Rosenberg, 2008; Guerrero *et al.*, 2013). Natural and sexual selection acts on the holobiont as if it were an integrated unit and the formation of microbe-host associations is often considered a driver of evolution (Zilber-Rosenberg & Rosenberg, 2008; Oliver *et al.*, 2010; Guerrero *et al.*, 2013). Hosts containing beneficial microbes may gain a selective advantage compared to hosts which do not contain these microbes (Zilber-Rosenberg & Rosenberg, 2008). One of the most prominent examples of beneficial microbes in arthropods is the obligate intracellular bacterium *Buchnera aphidicola*, which provides essential amino acids to its aphid host (Buchner, 1965; Shigenobu *et al.*, 2000; Baumann, 2005). Microbes are further known to support the breakdown of recalcitrant food in termites (Cleveland, 1923; Inoue *et al.*, 1997; Warnecke *et al.*, 2007) and have been shown to protect their host against pathogens (Teixeira *et al.*, 2008; Moreira *et al.*, 2009; Koch & Schmid-Hempel, 2011). Microbes with detrimental effects on their hosts promote the evolution of host defences, such as an immune system or behavioural adaptations that reduce the frequency or impact of infection (Schmid-Hempel, 2011). Since immune defence is costly, having and using an immune system can negatively affect other life history traits (Sheldon & Verhulst, 1996).

In my thesis, I investigated bacterial communities of two arthropod herbivores and the impact of bacteria or immune challenge on the life history of their hosts. In the following I will discuss the findings of my thesis in the light of present literature on (1) non-pathogenic host-microbe interactions and (2) pathogenic effects of microbes on arthropods.

### ***1. Non-pathogenic associations of bacteria with herbivores and their role in plant utilization***

Herbivore-associated bacteria may support the host plant use of their herbivore hosts via different mechanisms, such as the breakdown of nutrients (1.1) and the alteration of plant physiology to the host's benefit (1.2) (Dillon & Dillon, 2004; Douglas, 2009; Engel & Moran, 2013; Hansen & Moran, 2014). In this section, I will discuss these aspects in relation to my findings in *Heliothis virescens* and *Tetranychus urticae* (CHAPTERS 2 and 5). As it is unlikely that microbes in these hosts synthesize and/or provide nutrients, I will not discuss the possibility that bacteria can synthesize and provide nutrients to the host that are lacking its diet.

#### ***1.1. Breakdown of plant chemicals and nutrients by herbivore-associated bacteria***

In *H. virescens* there was high variability in bacterial communities depending on diet, developmental stage, rearing background (laboratory or field population) and even between biological replicates, indicating that the bacterial community of this moth is only transiently associated with its host (CHAPTER 2). In Lepidoptera,

evidence that bacteria may enhance host fitness in a long-term, stable symbiosis is scarce, even though bacterial communities of many lepidopteran species have been investigated. High alkalinity and the tubular gut structure without pouches, which are characteristic of lepidopteran guts, may impede a stable colonization by microbes (Broderick *et al.*, 2004; Engel & Moran, 2013). Moreover, holometabolous metamorphosis of Lepidoptera, during which the gut is purged, may further complicate the establishment of stable bacterial communities (Moll *et al.*, 2001; Rani *et al.*, 2009; Hammer *et al.*, 2014). Nevertheless, Lepidoptera may benefit from bacteria that are present in their guts. For instance, bacteria isolated from lepidopteran guts possess enzymes with cellulolytic, pectinolytic or xylanolytic activity which can break down plant material (Pinto-Tomas *et al.*, 2007; Anand *et al.*, 2010; Belda *et al.*, 2011). Similar metabolic capabilities can also (at least partly) be expected for the bacterial community of *H. virescens*, because many of the bacterial genera that were found associated with this moth have been identified before in Lepidoptera (CHAPTER 2).

Irrespective of the stability of an association, insects that take up bacteria from the environment may benefit from traits that bacteria have evolved independently of their host associations (Oliver *et al.*, 2010). For example, some phyllosphere bacteria can use methanol as carbon source, fix nitrogen or break down secondary plant metabolites (Sy *et al.*, 2005; Frnkranz *et al.*, 2008; Vorholt, 2012; Mason *et al.*, 2014). As an example of the latter, the growth of larvae of the gypsy moth, *Lymantria dispar*, was enhanced in the presence of bacteria from its host plant aspen, when phenolic glycosides, the defensive metabolites of aspen, were present in the larval diet (Mason *et al.*, 2014). The bacteria that were detected in *H. virescens* could have been taken up from the plants on which the caterpillars were reared (CHAPTER 2). In line with benefitting from plant-associated bacteria, the variability of bacterial communities in *H. virescens* could be viewed as flexibility, which may support this generalist herbivore in feeding on different host plants.

*Wolbachia* are intracellular bacteria that are commonly found in arthropods and nematodes. They are well known as reproductive parasites, but they are also able to deliver metabolic benefit to their hosts (Duron *et al.*, 2008; Werren *et al.*, 2008; Zug & Hammerstein, 2015). For instance, in *Drosophila melanogaster*, *Wolbachia* appears to be involved in iron homeostasis (Brownlie *et al.*, 2009). Furthermore, *Wolbachia* in filarial nematodes was suggested to provide nutrients or ATP to the host (Foster *et al.*, 2005; Darby *et al.*, 2012), and in the bedbug *Cimex lectularius*, *Wolbachia* likely provides its host with vitamin B (Hosokawa *et al.*, 2010). In the mite *T. urticae*, digestion- and detoxification-related genes of the mite were upregulated in the presence of *Wolbachia* (CHAPTER 5). One possible explanation for this finding is that *Wolbachia* performs early catabolic steps, which could alter digestion processes in the mites. Notably, in contrast to some of the above-mentioned examples, *Wolbachia* is not an obligate symbiont of the *T. urticae* strains that were tested, because mites were viable and could reproduce without this

bacterium (Staudacher and Schimmel, personal observation). Nevertheless, *Wolbachia* may increase mite fitness by performing metabolic steps that reduce the costs of digestion for the mite.

### *1.2. Effects of herbivore-associated bacteria on host plant physiology*

Herbivore-associated microbes can affect plant physiology. For instance, the presence of *Wolbachia* in the lepidopteran leafminer *Phyllonorycter blancardella* is associated with the preservation of photosynthetically active and nutrient rich regions (so-called ‘green-islands’) in otherwise senescing leaves (Kaiser *et al.*, 2010). Furthermore, herbivore-associated microbes may alter induced plant resistance in favour of their herbivore hosts (Chung *et al.*, 2013; Su *et al.*, 2015). Bacteria can reduce the jasmonic acid (JA) defence responses of plants, that are commonly induced by herbivore feeding (Schoonhoven *et al.*, 2005; Karban & Baldwin, 2007). Herbivores are susceptible to JA-responses, which can cause reduced larval growth, amount of feeding, survival and oviposition of the herbivores and may increase the attraction of their natural enemies (Howe *et al.*, 1996; Li *et al.*, 2002; Thaler *et al.*, 2002). JA-responses can be suppressed by salicylic acid (SA)-responses, which may be induced by microbes (JA-SA crosstalk) (Pieterse *et al.*, 2012; Thaler *et al.*, 2012). For instance, the Colorado potato beetle, *Leptinotarsa decemlineata*, secretes in its oral secretions *Pseudomonas*, *Stenotrophomonas* and *Enterobacter* to the plants, which suppresses JA-related plant responses via JA-SA crosstalk (Chung *et al.*, 2013). The moth *H. virescens* contained bacteria that belonged to the same genera as those in the Colorado potato beetle (CHAPTER 2). Importantly, the beetles also contained Enterobacteriaceae and one *Pseudomonas* strain that did not suppress defences, so that the capacity to suppress plant defences seems to be strain specific (Chung *et al.*, 2013). Plant defence suppression may thus not necessarily occur in other herbivores that contain bacteria of the same genus as bacteria in the Colorado potato beetle, such as *H. virescens*.

In the mite *T. urticae*, we tested whether the presence of *Wolbachia*, *Spiroplasma* and *Cardinium* affected induced plant responses after feeding of two strains of mites: a plant defence inducer strain (Santpoort-2), which induces JA- and SA-responses upon feeding, and a plant defence suppressor strain, which suppresses such defences (Alba *et al.*, 2015; CHAPTER 5). The suppressor strain contained *Wolbachia* and *Spiroplasma*, whereas the inducer strain contained *Cardinium* and *Spiroplasma*. Both mite strains were cleared from these bacteria with antibiotics. Additionally, there was one group of suppressor mites that only contained *Spiroplasma*. When testing the suppressor strain of plant defences, SA accumulated significantly more in leaflets infested with mites containing only *Spiroplasma* compared to leaflets infested with mites containing *Wolbachia* and *Spiroplasma* or none of the two bacteria. This suggests that *Spiroplasma* induced SA-responses, while *Wolbachia* antagonizes this effect. SA induction did not result in suppression of JA-responses via JA-SA crosstalk. However, for the precursor of JA, 12-*oxo*-

phytyldienoic acid (OPDA), the pattern that we found for SA was nearly reversed, suggesting that *Spiroplasma* suppressed OPDA-responses, while *Wolbachia* seemed to induce it. Even though OPDA may also mediate plant resistance, independently from JA (Stintzi *et al.*, 2001; Bosch *et al.*, 2014; Scalschi *et al.*, 2015), the differences in plant responses depending on bacteria in the mites were not correlated to mite fitness in the suppressor strain. Possibly, changes in plant physiology are primarily beneficial for the bacteria, but not for the herbivorous host: bacteria that suppress plant defences may be exposed to fewer secondary plant metabolites that are ingested by their host herbivores, which often have antimicrobial activity (Wallace, 2004; Karamanoli *et al.*, 2005; Karban & Baldwin, 2007).

In the plant defence inducer strain of *T. urticae*, the presence of *Cardinium* and *Spiroplasma* in the mites was associated with higher transcript accumulation of SA marker genes. This observation was in line with the finding that *Cardinium* and/or *Spiroplasma* negatively affected mite fitness (CHAPTER 5). Bacteria such as *Wolbachia*, *Cardinium* and *Spiroplasma* secure their persistence in a host population by manipulating host reproduction to increase the number of infected females and/or by increasing the fitness of the host (Werren *et al.*, 2008; Himler *et al.*, 2011; Zug & Hammerstein, 2015). Beneficial effects of reproductive manipulators are expected when reproductive manipulation is weak in a population (Hoffmann *et al.*, 1998; Fry *et al.*, 2004). Thus, these non-beneficial or even detrimental bacteria in the inducer strain may increase their prevalence in the mite population via reproductive manipulation, but we did not test this in our mites.

### 1.3. Interactions between multiple bacteria in one arthropod host

Bacteria can shape their environment by affecting abiotic factors, such as pH, and as such are true ecosystem engineers (Madigan *et al.*, 2009). Moreover, bacterial strains can utilize and even depend on breakdown products of other strains in a metabolic cross-feeding network (Flint *et al.*, 2007). For instance, bacteria that are unable to use complex carbohydrates depend on the presence of primary degraders (Flint *et al.*, 2007). Furthermore, *in vitro* experiments with human gut bacteria showed that lactate utilizing bacteria grow on starch only in the presence of particular strains of lactate producing *Bifidobacterium adolescentis*, which are able to metabolize starch (Belenguer *et al.*, 2006). Because of their strong influence on community structure, the presence of bacterial strains with keystone function could underlie high variability between bacterial communities. Another common type of interaction between bacterial strains is competition for resources and space both at the intra- and inter-specific level (Hibbing *et al.*, 2010). The outcome of competition may depend on environmental factors that promote or inhibit the growth of particular bacterial strains.

The moth *H. virescens* and the spider mite *T. urticae* harbour multiple bacterial strains (CHAPTERS 2 and 5). Interactions between these strains are expected in the hosts and may in part explain the variability of bacterial communities in *H. virescens* and the combination effects of *Spiroplasma* and *Wolbachia* in *T. urticae* that we

observed. For example, competition could underlie some of the variation in bacterial communities *H. virescens* in relation to diet, life stage and rearing background (CHAPTER 2). Such competition may explain the fact that *H. virescens* larvae, collected in the field and reared in the laboratory for four generations, were colonized with enterococci that were absent in field larvae. Enterococci may thus be strong competitors in larvae under laboratory conditions. Another possibility is that under laboratory conditions different bacterial strains disappeared because of a lack of replenishment of new individuals from the environment. Enterococci could thus have replaced these strains without competition.

In *T. urticae*, the combined presence of *Wolbachia* and *Spiroplasma* affected induced plant responses and mite gene expression. This observation suggests that either the two bacteria affected plant responses and mite gene expression independently, or that the bacteria interacted. Based on sequencing data, we speculated that *Wolbachia* negatively affects the abundance of *Spiroplasma* in the mites (CHAPTER 5). For many maternally transmitted bacteria, colonization of the ovaries is important; in the ovaries co-infecting bacteria may compete for space or nutrients. For instance, *Wolbachia* impeded *Asaia* bacteria from colonizing of reproductive organs in mosquitos (Rossi *et al.*, 2015). In *T. urticae*, *Wolbachia* was shown to reside in the ovaries (Zhao *et al.*, 2013). Possibly, *Spiroplasma* also colonizes the ovaries in the mite, because this bacterium stably infected our mite populations and is likely vertically transmitted. Thus, *Wolbachia* and *Spiroplasma* may compete in the ovaries.

*Wolbachia* not only resides in the ovaries but also in the gnathosoma (i.e., mouth and feeding parts of mites, including salivary glands, amongst others) of *T. urticae* (Zhao *et al.*, 2013). From the gnathosoma, *T. urticae*-associated *Wolbachia* possibly affects plant physiology, e.g., SA and OPDA related plant responses after mite feeding (CHAPTER 5). *Wolbachia* seems to affect plant responses in an opposite way of *Spiroplasma*. Competition for space in the gnathosoma could underlie this result.

## **2. Effect of immune challenge on life history traits**

### **2.1. Cost of immune defence**

Activation and maintenance of the immune system in animals is costly and can trade off against other life history traits such as reproductive traits (Sheldon & Verhulst, 1996). The trade-off between immunity and reproduction can depend on sex and the investment of each sex into offspring (Roff, 2002; Zuk & Stoehr, 2002; Roth *et al.*, 2011; Vincent & Gwynne, 2014). Males and females of *H. virescens* differ in their investment strategy into immunity (CHAPTER 3). Non-challenged males had higher expression levels of immune-related genes than non-challenged females. This suggests that males have higher immune maintenance than females, which implies physiological costs to keep the immune system at a certain level of readiness. Females showed higher immune deployment than males after immune challenge, which implies physiological costs of immune system activation to combat pathogens

(see Schmid-Hempel, 2011). These findings are in line with the result that bacteria-challenged females had lower mating success than unchallenged females, but in males there was no difference between bacteria-injected and non-injected mates. Thus, in *H. virescens*, immune response is negatively correlated to mating success in females but not in males.

In addition to the differences between the sexes in immune deployment and maintenance, the differences between wounding and bacterial challenge were more pronounced in females than in males, suggesting that the female immune system differentiates more specifically between wounding and bacterial challenge than the male system. Together, female immune response may be more cost efficient than the male immune response, because different types of immune responses differ in their costliness: non-specific and constitutive immune responses are assumed to have higher (continuous) energy costs, with a higher risk of self-reactivity, than induced and specific responses (Schmid-Hempel, 2011).

In general, immunity is hypothesized to be more important for females than for males, because males maximize their fitness by increasing mating rate, while females maximize their fitness by increasing longevity (Rolff, 2002). However, in moths longevity is important for both sexes as discussed in CHAPTER 3. As in moths females are sexual signalers to which males respond (Nesbitt *et al.*, 1979; Vetter & Baker, 1983; Tumlinson *et al.*, 1986; Groot *et al.*, 2014), female fitness may not only depend on longevity, but also on access to males. A (cost) efficient immunity may therefore be necessary to uphold a high quality of the sexual signal. To assess how exactly males and females differ in their type of immunity, additional immune responses should be measured, such as phagocytosis (or encapsulation) (i.e., a not delayed, constitutive response with little specificity) and levels of antimicrobial peptides (i.e., a delayed, induced and more specific response) (Mallon *et al.*, 2003; Schmid-Hempel & Ebert, 2003).

The sex pheromone blend of *H. virescens* females seems to be condition dependent, as the blend of bacteria-challenged females was less attractive than that of wounded or unchallenged females (CHAPTER 3). Thus, infection could possibly affect attractiveness and therefore mating success of female *H. virescens* in the field. The costs of the quality of female sex pheromones in moths has scarcely been assessed, but may be high (Harari *et al.*, 2011; Xu *et al.*, 2014). In addition, our result suggests that the quality of the female sex pheromone of this moth could be an honest signal that is costly and indicates to the males the condition of the females (Zahavi, 1975; Hamilton & Zuk, 1982; Zahavi & Zahavi, 1997). Possibly, there is a physiological trade-off between immune defence activation and regulation of biosynthesis of the female sex pheromone of *H. virescens*.

In insects, juvenile hormone (JH) plays a central role in the regulation of development and reproduction and was shown to have an immunosuppressive effect, which may thus mediate the trade-off between immunity and reproduction (Nijhout & Williams, 1974; Wyatt & Davey, 1996; Rolff & Siva-Jothy, 2002;

Rantala *et al.*, 2003). JH could also play a role in the trade-off between attractiveness and immunity, a situation that was encountered in mealworm beetles, in which JH increased the attractiveness of male sex pheromone which coincided with lower phenoloxidase (PO) activity in response to the JH treatment (Rantala *et al.*, 2003). The altered sex pheromone blend in *H. virescens* upon immune system activation in bacteria-challenged females could thus possibly result from reduced JH levels after an immune challenge. Unlike in beetles, female sex pheromone production in moths is regulated by pheromone biosynthesis-activating neuropeptide (PBAN) and not directly by JH (Raina *et al.*, 1989; Jurenka, 1996, 2004; Jurenka & Rafaeli, 2011). In migratory moths, JH seems to regulate the release or synthesis of PBAN (Cusson & McNeil, 1989; Gadenne, 1993; Picimbon *et al.*, 1995), while the role of JH in non-migratory moth species is less clear. In the non-migratory moth *Helicoverpa armigera*, JH primes the sex pheromone gland to respond to PBAN, indicating that JH is involved in the initiation of female sex pheromone biosynthesis (Fan *et al.*, 1999; Rafaeli *et al.*, 2003). Thus, even though the role of JH in an immunity-sex pheromone trade-off may not be as straightforward as in beetles, JH remains to be a candidate for mediating such a trade-off in moths. Since the sex pheromone change in *H. virescens* in response to bacterial challenge concerned pheromone composition and not total amounts of the sex pheromone (CHAPTER 3), we expect that later steps than the initiation of pheromone synthesis are (also) involved in the trade-off.

### *2.2 Behavioural changes in response to immune challenge*

Many organisms have evolved behavioural adaptations in the face of infection (Schmid-Hempel, 2011; de Roode & Lefèvre, 2012). Prior to infection, animals can show infection avoidance behaviour, i.e., spatial and temporal avoidance of contaminated food or contaminated co-specifics such as mates (Christe *et al.*, 1994; Kavaliers & Colwell, 1995; Penn *et al.*, 1998; Hutchings *et al.*, 2001; Alma *et al.*, 2010). We found that *H. virescens* males mated less with bacteria-challenged than with unchallenged females (CHAPTER 3), indicating that males avoid bacteria-challenged females. As an alternative explanation, bacteria-challenged females could have been less apt to mate than unchallenged females. However, we found no difference in calling behaviour (extrusion of the sex pheromone gland to emit pheromone) between bacteria-challenged, wounded and unchallenged females, which makes this alternative explanation less likely. Avoidance of mating with infected mates has been shown in vertebrates (Kavaliers & Colwell, 1995; Penn & Potts, 1998; Penn *et al.*, 1998; Deaton, 2009). However, in experiments testing avoidance of mating with infected mates in invertebrates, infected and uninfected mates did equally well (with exception of infection with reproductive parasites; Vala *et al.*, 2004) (Abbot & Dill, 2001; Luong & Kaya, 2005; Burand & Tan, 2006; de Roode & Lefèvre, 2012). We thus show for the first time that avoidance of infected mates can also occur in invertebrates.

Once infection has been contracted, adaptations may either take the form of curative behaviours, such as grooming and therapeutic medication, or behaviour that leads to an increase in fecundity to maximize fitness in the short lifespan that remains, such as terminal investment (Schmid-Hempel, 2011; de Roode & Lefèvre, 2012; Abbott, 2014). Terminal investment may involve an increase in courtship activity, oviposition rate or parental care (Minchella & Loverde, 1981; Part *et al.*, 1992; Polak & Starmer, 1998; Adamo, 1999; Bonneaud *et al.*, 2004; Creighton *et al.*, 2009). Bacteria-challenged *H. virescens* females had a higher oviposition rate than control females one night after the challenge (CHAPTER 4). Surprisingly, even though immune-challenged females had shorter life spans than control females, the total amount of eggs females produced was not significantly different between the two groups of females. This suggests that bacteria-challenged females may be able to compensate for shortened life span by higher oviposition rate. However, the increase in oviposition rate was linked to reduced oviposition site selectivity between plants that were damaged by *H. virescens* larvae and undamaged plants (CHAPTER 4), which increases the chance that hatching larvae have lower survival and/or develop more slowly than larvae hatching from eggs laid on undamaged plants. The costs incurred by an increased oviposition rate in infected females are thus likely to become apparent in the offspring generation, indicating that not only immune system activation, but also behavioural changes due to infection may be costly (see also Minchella & Loverde, 1981; Minchella, 1985).

The concept of terminal investment implies that the actual strategy after an infection may depend on the acuteness of the threat for survival and thus on the dose and virulence of an infecting agent (Williams, 1966; Clutton-Brock, 1984). Females that are in acute mortal danger may follow a terminal investment strategy and spend all the resources they have into reproduction, while females that are not in acute life danger may invest in immune system activation or curative behaviours and thus survival. Possibly, there is an ‘acuteness threshold’ at which species switch from a curative to a terminal investment strategy.

Finally, the immune response may depend on the mating status of an animal (Rolff & Siva-Jothy, 2002). In several insect species, including *H. virescens*, juvenile hormone levels rise significantly after mating compared to virgin adults (Loher *et al.*, 1983; Couche *et al.*, 1985; Shu *et al.*, 1998; Rolff & Siva-Jothy, 2002). Since the juvenile hormone can have an immunosuppressive effect (Rolff & Siva-Jothy, 2002), it is possible that immune response is stronger in virgin than in mated females in this moth. Consequently, the immune challenge with *S. entomophila* could have been a stronger and more acute life threat for mated females in the oviposition assay (CHAPTER 3) than for the virgin females in the mating assay (CHAPTER 4). This is in line with our finding that immunity is negatively correlated to reproduction, indicating a trade-off (CHAPTER 3), while we found mated females to make a terminal investment after bacterial challenge (CHAPTER 4). Moreover, while all tested virgin females survived for at least three days (as our experiment

ended after three days) (CHAPTER 3), 38.5% of the mated females of the oviposition experiment died after two nights (CHAPTER 4). It will be interesting to determine whether virgin and mated *H. virescens* females differ in their immune system activation.

Together, the data presented in this thesis underline that arthropods may be exposed to and colonized by multiple bacteria during their lifetime and that bacteria may have a wide variety of effects on their arthropod hosts, ranging from beneficial to lethal. These effects may be due to the effect of bacteria on host physiology, which may extend to induced plant responses after herbivore feeding. In addition, immune challenge via bacterial pathogens can activate costly immune defences which may trade off with other traits or may result in terminal investment behaviour by which a shortened lifespan can be compensated.

## REFERENCES

- Abbot, P. & Dill, L.M. 2001. Sexually transmitted parasites and sexual selection in the milkweed leaf beetle, *Labidomera clivicollis*. *Oikos* **92**: 91-100.
- Abbott, J. 2014. Self-medication in insects: current evidence and future perspectives. *Ecol Entomol* **39**: 273-280.
- Adamo, S.A. 1999. Evidence for adaptive changes in egg laying in crickets exposed to bacteria and parasites. *Anim Behav* **57**: 117-124.
- Alba, J.M., Schimmel, B.C.J., Glas, J.J., Ataide, L.M.S., Pappas, M.L., Villarroel, C.A., Schuurink, R.C., Sabelis, M.W. & Kant, M.R. 2015. Spider mites suppress tomato defenses downstream of jasmonate and salicylate independently of hormonal crosstalk. *New Phytol* **205**: 828-840.
- Alma, C.R., Gillespie, D.R., Roitberg, B.D. & Goettel, M.S. 2010. Threat of infection and threat-avoidance behavior in the predator *Dicyphus hesperus* feeding on whitefly nymphs infected with an entomopathogen. *J Insect Behav* **23**: 90-99.
- Anand, A.A.P., Vennison, S.J., Sankar, S.G., Prabhu, D.I.G., Vasan, P.T., Raghuraman, T., Geoffrey, C.J. & Vendan, S.E. 2010. Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin and starch and their impact on digestion. *J Insect Sci* **10**: 107.
- Baumann, P. 2005. Biology of bacteriocyte-associated endosymbionts of plant sap-sucking insects. *Annu Rev Microbiol* **59**: 155-189.
- Belda, E., Pedrola, L., Peretó, J., Martínez-Blanch, J.F., Montagud, A., Navarro, E., Urchueguía, J., Ramón, D., Moya, A. & Porcar, M. 2011. Microbial diversity in the midguts of field and lab-reared populations of the European corn borer *Ostrinia nubilalis*. *PLoS ONE* **6**.
- Belenguer, A., Duncan, S.H., Calder, A.G., Holtrop, G., Louis, P., Lobley, G.E. & Flint, H.J. 2006. Two routes of metabolic cross-feeding between *Bifidobacterium adolescentis* and butyrate-producing anaerobes from the human gut. *Appl Environ Microbiol* **72**: 3593-3599.
- Bonneaud, C., Mazuc, J., Chastel, O., Westerdahl, H. & Sorci, G. 2004. Terminal investment induced by immune challenge and fitness traits associated with major histocompatibility complex in the house sparrow. *Evolution* **58**: 2823-2830.
- Bosch, M., Wright, L.P., Gershenzon, J., Wasternack, C., Hause, B., Schaller, A. & Stintzi, A. 2014. Jasmonic acid and its precursor 12-oxophytodienoic acid control different aspects of constitutive and induced herbivore defenses in tomato. *Plant Physiol* **166**: 396-410.

- Broderick, N.A., Raffa, K.F., Goodman, R.M. & Handelsman, J. 2004. Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. *Appl Environ Microbiol* **70**: 293-300.
- Brownlie, J.C., Cass, B.N., Riegler, M., Witsenburg, J.J., Iturbe-Ormaetxe, I., McGraw, E.A. & O'Neill, S.L. 2009. Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS Pathol* **5**: e1000368.
- Buchner, P. 1965. *Endosymbiosis of animals with plant microorganisms*. Interscience, New York, USA.
- Burand, J.P. & Tan, W. 2006. Mate preference and mating behavior of male *Helicoverpa zea* (Lepidoptera: Noctuidae) infected with the sexually transmitted insect virus Hz-2V. *Ann Entomol Soc Am* **99**: 969-973.
- Christe, P., Oppliger, A. & Richner, H. 1994. Ectoparasite affects choice and use of roost sites in the great tit, *Parus major*. *Anim Behav* **47**: 895-898.
- Chung, S.H., Rosa, C., Scully, E.D., Peiffer, M., Tooker, J.F., Hoover, K., Luthe, D.S. & Felton, G.W. 2013. Herbivore exploits orally secreted bacteria to suppress plant defenses. *Proc Natl Acad Sci USA* **110**: 15728-15733.
- Cleveland, L.R. 1923. Symbiosis between termites and their intestinal protozoa. *Proc Natl Acad Sci USA* **9**: 424.
- Clutton-Brock, T.H. 1984. Reproductive effort and terminal investment in iteroparous animals. *Am Nat* **123**: 212-229.
- Couche, G., Gillott, C., Tobe, S. & Feyereisen, R. 1985. Juvenile hormone biosynthesis during sexual maturation and after mating in the adult male migratory grasshopper, *Melanoplus sanguinipes*. *Can J Zoolog* **63**: 2789-2792.
- Creighton, J.C., Heflin, N.D. & Belk, M.C. 2009. Cost of reproduction, resource quality, and terminal investment in a burying beetle. *Am Nat* **174**: 673-684.
- Cusson, M. & McNeil, J.N. 1989. Involvement of juvenile hormone in the regulation of pheromone release activities in a moth. *Science* **243**: 210-212.
- Darby, A.C., Armstrong, S.D., Bah, G.S., Kaur, G., Hughes, M.A., Kay, S.M., Koldkjær, P., Rainbow, L., Radford, A.D. & Blaxter, M.L. 2012. Analysis of gene expression from the *Wolbachia* genome of a filarial nematode supports both metabolic and defensive roles within the symbiosis. *Genome Res* **22**: 2467-2477.
- de Roode, J.C. & Lefèvre, T. 2012. Behavioral immunity in insects. *Insects* **3**: 789-820.
- Deaton, R. 2009. Effects of a parasitic nematode on male mate choice in a livebearing fish with a coercive mating system (western mosquitofish, *Gambusia affinis*). *Behav Process* **80**: 1-6.
- Dillon, R. & Dillon, V. 2004. The gut bacteria of insects: nonpathogenic interactions. *Annu Rev Entomol* **49**: 71-92.
- Douglas, A. 2009. The microbial dimension in insect nutritional ecology. *Funct Ecol* **23**: 38-47.
- Duron, O., Bouchon, D., Boutin, S., Bellamy, L., Zhou, L., Engelstädter, J. & Hurst, G.D. 2008. The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol* **6**: 27.
- Engel, P. & Moran, N.A. 2013. The gut microbiota of insects—diversity in structure and function. *FEMS Microbiol Rev* **37**: 699-735.
- Fan, Y., Rafaeeli, A., Gileadi, C. & Applebaum, S.W. 1999. Juvenile hormone induction of pheromone gland PBAN-responsiveness in *Helicoverpa armigera* females. *Insect Biochem Mol Biol* **29**: 635-641.
- Flint, H.J., Duncan, S.H., Scott, K.P. & Louis, P. 2007. Interactions and competition within the microbial community of the human colon: links between diet and health. *Environ Microbiol* **9**: 1101-1111.

## General discussion

- Foster, J., Ganatra, M., Kamal, I., Ware, J., Makarova, K., Ivanova, N., Bhattacharyya, A., Kapatral, V., Kumar, S. & Posfai, J. 2005. The *Wolbachia* genome of *Brugia malayi*: endosymbiont evolution within a human pathogenic nematode. *PLoS Biol* **3**: e121.
- Fry, A., Palmer, M. & Rand, D. 2004. Variable fitness effects of *Wolbachia* infection in *Drosophila melanogaster*. *Heredity* **93**: 379-389.
- Fürnkranz, M., Wanek, W., Richter, A., Abell, G., Rasche, F. & Sessitsch, A. 2008. Nitrogen fixation by phyllosphere bacteria associated with higher plants and their colonizing epiphytes of a tropical lowland rainforest of Costa Rica. *ISME J* **2**: 561-570.
- Gadenne, C. 1993. Effects of fenoxycarb, juvenile hormone mimetic, on female sexual behaviour of the black cutworm, *Agrotis ipsilon* (Lepidoptera: Noctuidae). *J Insect Physiol* **39**: 25-29.
- Groot, A.T., Schöfl, G., Inglis, O., Donnerhacke, S., Classen, A., Schmalz, A., Santangelo, R.G., Emerson, J., Gould, F. & Schal, C. 2014. Within-population variability in a moth sex pheromone blend: genetic basis and behavioural consequences. *Proc R Soc B Biol Sci* **281**: 20133054.
- Guerrero, R., Margulis, L. & Berlanga, M. 2013. Symbiogenesis: the holobiont as a unit of evolution. *Int Microbiol* **16**: 133-143.
- Hamilton, W.D. & Zuk, M. 1982. Heritable true fitness and bright birds: a role for parasites? *Science* **218**: 384-387.
- Hammer, T.J., McMillan, W.O. & Fierer, N. 2014. Metamorphosis of a butterfly-associated bacterial community. *PLoS ONE*: e86995.
- Hansen, A.K. & Moran, N.A. 2014. The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol Ecol* **23**: 1473-1496.
- Harari, A.R., Zahavi, T. & Thiéry, D. 2011. Fitness cost of pheromone production in signaling female moths. *Evolution* **65**: 1572-1582.
- Hibbing, M.E., Fuqua, C., Parsek, M.R. & Peterson, S.B. 2010. Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol* **8**: 15-25.
- Himler, A.G., Adachi-Hagimori, T., Bergen, J.E., Kozuch, A., Kelly, S.E., Tabashnik, B.E., Chiel, E., Duckworth, V.E., Dennehy, T.J. & Zchori-Fein, E. 2011. Rapid spread of a bacterial symbiont in an invasive whitefly is driven by fitness benefits and female bias. *Science* **332**: 254-256.
- Hoffmann, A.A., Hercus, M. & Dagher, H. 1998. Population dynamics of the *Wolbachia* infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* **148**: 221-231.
- Hosokawa, T., Koga, R., Kikuchi, Y., Meng, X.-Y. & Fukatsu, T. 2010. *Wolbachia* as a bacteriocyte-associated nutritional mutualist. *Proc Natl Acad Sci USA* **107**: 769-774.
- Howe, G.A., Lightner, J. & Ryan, C. 1996. An octadecanoid pathway mutant (JL5) of tomato is compromised in signaling for defense against insect attack. *Plant Cell* **8**: 2067-2077.
- Hutchings, M.R., Kyriazakis, I. & Gordon, I.J. 2001. Herbivore physiological state affects foraging trade-off decisions between nutrient intake and parasite avoidance. *Ecology* **82**: 1138-1150.
- Inoue, T., Murashima, K., Azuma, J.-I., Sugimoto, A. & Slaytor, M. 1997. Cellulose and xylan utilisation in the lower termite *Reticulitermes speratus*. *J Insect Physiol* **43**: 235-242.
- Jurenka, R. 2004. Insect pheromone biosynthesis. In: *The Chemistry of Pheromones and Other Semiochemicals I*, pp. 97-132. Springer, Berlin.
- Jurenka, R. & Rafeali, A. 2011. Regulatory role of PBAN in sex pheromone biosynthesis of heliothine moths. *Front Endocrinol* **2**: 1-8.
- Jurenka, R.A. 1996. Signal transduction in the stimulation of sex pheromone biosynthesis in moths. *Arch Insect Biochem Physiol* **33**: 245-258.
- Kaiser, W., Huguet, E., Casas, J., Commin, C. & Giron, D. 2010. Plant green-island phenotype induced by leaf-miners is mediated by bacterial symbionts. *Proc R Soc B Biol Sci*: rspb20100214.

- Karamanoli, K., Menkissoglu-Spirodi, U., Bosabalidis, A.M., Vokou, D. & Constantinidou, H.-I.A. 2005. Bacterial colonization of the phyllosphere of nineteen plant species and antimicrobial activity of their leaf secondary metabolites against leaf associated bacteria. *Chemoecology* **15**: 59-67.
- Karban, R. & Baldwin, I.T. 2007. *Induced responses to herbivory*. University of Chicago Press.
- Kavaliers, M. & Colwell, D.D. 1995. Discrimination by female mice between the odours of parasitized and non-parasitized males. *Proc R Soc B Biol Sci* **261**: 31-35.
- Koch, H. & Schmid-Hempel, P. 2011. Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proc Natl Acad Sci USA* **108**: 19288-19292.
- Li, C., Williams, M.M., Loh, Y.-T., Lee, G.I. & Howe, G.A. 2002. Resistance of cultivated tomato to cell content-feeding herbivores is regulated by the octadecanoid-signaling pathway. *Plant Physiol* **130**: 494-503.
- Loher, W., Ruzo, L., Baker, F., Miller, C. & Schooley, D. 1983. Identification of the juvenile hormone from the cricket, *Teleogryllus commodus*, and juvenile hormone titre changes. *J Insect Physiol* **29**: 585-589.
- Luong, L.T. & Kaya, H.K. 2005. Sexually transmitted parasites and host mating behavior in the decorated cricket. *Behav Ecol* **16**: 794-799.
- Madigan, M.T., Martinko, J., Stahl, D. & Clark, D. 2009. *Brock Biology of Microorganisms*. Benjamin Cummings, San Francisco.
- Mallon, E.B., Loosli, R. & Schmid-Hempel, P. 2003. Specific versus nonspecific immune defense in the bumblebee, *Bombus terrestris* L. *Evolution* **57**: 1444-1447.
- Margulis, L. & Chapman, M.J. 1998. Endosymbioses: cyclical and permanent in evolution. *Trends Microbiol* **6**: 342-345.
- Mason, C.J., Couture, J.J. & Raffa, K.F. 2014. Plant-associated bacteria degrade defense chemicals and reduce their adverse effects on an insect defoliator. *Oecol* **175**: 901-910.
- Minchella, D. 1985. Host life-history variation in response to parasitism. *Parasitology* **90**: 205-216.
- Minchella, D.J. & Loverde, P.T. 1981. A cost of increased early reproductive effort in the snail *Biomphalaria glabrata*. *Am Nat* **118**: 876-881.
- Moll, R.M., Romoser, W.S., Modrakowski, M.C., Moncayo, A.C. & Lerdthusnee, K. 2001. Meconial peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae) metamorphosis. *J Med Entomol* **38**: 29-32.
- Moreira, L.A., Iturbe-Ormaetxe, I., Jeffery, J.A., Lu, G., Pyke, A.T., Hedges, L.M., Rocha, B.C., Hall-Mendelin, S., Day, A. & Riegler, M. 2009. A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and *Plasmodium*. *Cell* **139**: 1268-1278.
- Nesbitt, B., Beevor, P., Hall, D. & Lester, R. 1979. Female sex pheromone components of the cotton bollworm, *Heliothis armigera*. *J Insect Physiol* **25**: 535-541.
- Nijhout, H.F. & Williams, C.M. 1974. Control of moulting and metamorphosis in the tobacco hornworm, *Manduca sexta* (L.): cessation of juvenile hormone secretion as a trigger for pupation. *J Exp Biol* **61**: 493-501.
- Oliver, K.M., Degan, P.H., Burke, G.R. & Moran, N.A. 2010. Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. *Annu Rev Entomol* **55**: 247-266.
- Part, T., Gustafsson, L. & Moreno, J. 1992. "Terminal investment" and a sexual conflict in the collared flycatcher (*Ficedula albicollis*). *Am Nat* **140**: 868-882.
- Penn, D. & Potts, W.K. 1998. Chemical signals and parasite-mediated sexual selection. *Trends Ecol Evol* **13**: 391-396.
- Penn, D., Schneider, G., White, K., Slev, P. & Potts, W. 1998. Influenza infection neutralizes the attractiveness of male odour to female mice (*Mus musculus*). *Ethol* **104**: 685-694.

## General discussion

- Picimbon, J.-F., Becard, J.-M., Sreng, L., Clement, J.-L. & Gadenne, C. 1995. Juvenile hormone stimulates pheromonotropic brain factor release in the female black cutworm, *Agrotis ipsilon*. *J Insect Physiol* **41**: 377-382.
- Pieterse, C.M., Van der Does, D., Zamioudis, C., Leon-Reyes, A. & Van Wees, S.C. 2012. Hormonal modulation of plant immunity. *Annu Rev Cell Dev Biol* **28**: 489-521.
- Pinto-Tomas, A., Uribe-Lorio, L., Blanco, J., Fontecha, G., Rodríguez, C., Mora, M., Janzen, D., Chavarria, F., Díaz, J. & Sittenfeld, A. 2007. Enzymatic activities of bacteria isolated from the digestive tract of caterpillars and the pupal content of *Automeris zugana* and *Rothschildia lebeau* (Lepidoptera: Saturniidae). *Rev Biol Trop* **55**: 401-415.
- Polak, M. & Starmer, W.T. 1998. Parasite-induced risk of mortality elevates reproductive effort in male *Drosophila*. *Proc R Soc B Biol Sci* **265**: 2197-2201.
- Rafaeli, A., Zakharova, T., Lapsker, Z. & Jurenka, R. 2003. The identification of an age- and female-specific putative PBAN membrane-receptor protein in pheromone glands of *Helicoverpa armigera*: possible up-regulation by juvenile hormone. *Insect Biochem Mol Biol* **33**: 371-380.
- Raina, A., Jaffe, H., Kempe, T., Keim, P., Blacher, R., Fales, H., Riley, C., Klun, J., Ridgway, R. & Hayes, D. 1989. Identification of a neuropeptide hormone that regulates sex pheromone production in female moths. *Science* **244**: 796-798.
- Rani, A., Sharma, A., Rajagopal, R., Adak, T. & Bhatnagar, R.K. 2009. Bacterial diversity analysis of larvae and adult midgut microflora using culture-dependent and culture-independent methods in lab-reared and field-collected *Anopheles stephensi*-an Asian malarial vector. *BMC Microbiol* **9**: 96.
- Rantala, M.J., Vainikka, A. & Kortet, R. 2003. The role of juvenile hormone in immune function and pheromone production trade-offs: a test of the immunocompetence handicap principle. *Proc R Soc B Biol Sci* **270**: 2257-2261.
- Rohwer, F., Seguritan, V., Azam, F. & Knowlton, N. 2002. Diversity and distribution of coral-associated bacteria. *Mar Ecol Prog Ser* **243**: 1-10.
- Rolff, J. 2002. Bateman's principle and immunity. *Proc R Soc B Biol Sci* **269**: 867-872.
- Rolff, J. & Siva-Jothy, M.T. 2002. Copulation corrupts immunity: a mechanism for a cost of mating in insects. *Proc Natl Acad Sci USA* **99**: 9916-9918.
- Rossi, P., Ricci, I., Cappelli, A., Damiani, C., Ulissi, U., Mancini, M.V., Valzano, M., Capone, A., Epis, S. & Crotti, E. 2015. Mutual exclusion of *Asaia* and *Wolbachia* in the reproductive organs of mosquito vectors. *Parasite Vector* **8**: 278.
- Roth, O., Scharsack, J., Keller, I. & Reusch, T.B. 2011. Bateman's principle and immunity in a sex-role reversed pipefish. *J Evol Biol* **24**: 1410-1420.
- Scalschi, L., Sanmartín, M., Camañes, G., Troncho, P., Sánchez-Serrano, J.J., García-Agustín, P. & Vicedo, B. 2015. Silencing of OPR3 in tomato reveals the role of OPDA in callose deposition during the activation of defense responses against *Botrytis cinerea*. *Plant J* **81**: 304-315.
- Schmid-Hempel, P. & Ebert, D. 2003. On the evolutionary ecology of specific immune defence. *Trends Ecol Evol* **18**: 27-32.
- Schmid-Hempel, P. 2011. *Evolutionary parasitology: the integrated study of infections, immunology, ecology, and genetics*. Oxford University Press, New York.
- Schoonhoven, L.M., Van Loon, J.J. & Dicke, M. 2005. *Insect-Plant Biology, 2nd edn*. Oxford University Press, New York.
- Sheldon, B.C. & Verhulst, S. 1996. Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends Ecol Evol* **11**: 317-321.
- Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y. & Ishikawa, H. 2000. Genome sequence of the endocellular bacterial symbiont of aphids *Buchnera* sp. APS. *Nature* **407**: 81-86.
- Shu, S., Park, Y.I., Ramaswamy, S.B. & Srinivasan, A. 1998. Temporal profiles of juvenile hormone titers and egg production in virgin and mated females of *Heliothis virescens* (Noctuidae). *J Insect Physiol* **44**: 1111-1117.

- Stintzi, A., Weber, H., Reymond, P., Browse, J. & Farmer, E.E. 2001. Plant defense in the absence of jasmonic acid: The role of cyclopentenones. *Proc Natl Acad Sci USA* **98**: 12837-12842.
- Su, Q., Oliver, K.M., Xie, W., Wu, Q., Wang, S. & Zhang, Y. 2015. The whitefly-associated facultative symbiont *Hamiltonella defensa* suppresses induced plant defences in tomato. *Funct Ecol* **29**: 1007-1018.
- Sy, A., Timmers, A.C., Knief, C. & Vorholt, J.A. 2005. Methylo-trophic metabolism is advantageous for *Methylobacterium extorquens* during colonization of *Medicago truncatula* under competitive conditions. *Appl Environ Microbiol* **71**: 7245-7252.
- Teixeira, L., Ferreira, Á. & Ashburner, M. 2008. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol*: e1000002.
- Thaler, J.S., Farag, M.A., Paré, P.W. & Dicke, M. 2002. Jasmonate-deficient plants have reduced direct and indirect defences against herbivores. *Ecol Lett* **5**: 764-774.
- Thaler, J.S., Humphrey, P.T. & Whiteman, N.K. 2012. Evolution of jasmonate and salicylate signal crosstalk. *Trends Plant Sci* **17**: 260-270.
- Tumlinson, J., Mitchell, E., Teal, P., Heath, R. & Mengelkoch, L. 1986. Sex pheromone of fall armyworm, *Spodoptera frugiperda* (JE Smith). *J Chem Ecol* **12**: 1909-1926.
- Vala, F., Egas, M., Breeuwer, J. & Sabelis, M. 2004. *Wolbachia* affects oviposition and mating behaviour of its spider mite host. *J Evol Biol* **17**: 692-700.
- Vetter, R.S. & Baker, T.C. 1983. Behavioral responses of male *Heliothis virescens* in a sustained-flight tunnel to combinations of seven compounds identified from female sex pheromone glands. *J Chem Ecol* **9**: 747-759.
- Vincent, C.M. & Gwynne, D.T. 2014. Sex-biased immunity is driven by relative differences in reproductive investment. *Proc R Soc B Biol Sci* **281**: 20140333.
- Vorholt, J.A. 2012. Microbial life in the phyllosphere. *Nat Rev Microbiol* **10**: 828-840.
- Wallace, R.J. 2004. Antimicrobial properties of plant secondary metabolites. *Proc Nutr Soc* **63**: 621-629.
- Warnecke, F., Luginbühl, P., Ivanova, N., Ghassemian, M., Richardson, T.H., Stege, J.T., Cayouette, M., McHardy, A.C., Djordjevic, G. & Aboushadi, N. 2007. Metagenomic and functional analysis of hindgut microbiota of a wood-feeding higher termite. *Nature* **450**: 560-565.
- Werren, J.H., Baldo, L. & Clark, M.E. 2008. *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* **6**: 741-751.
- Williams, G.C. 1966. Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am Nat* **100**: 687-690.
- Wyatt, G.R. & Davey, K.G. 1996. Cellular and molecular actions of juvenile hormone. II. roles of juvenile hormone in adult insects. *Adv Insect Physiol* **26**: 1-155.
- Xu, J., Huigens, M.E., Orr, D. & Groot, A.T. 2014. Differential response of *Trichogramma* wasps to extreme sex pheromone types of the noctuid moth *Heliothis virescens*. *Ecol Entomol* **39**: 627-636.
- Zahavi, A. 1975. Mate selection—a selection for a handicap. *J Theor Biol* **53**: 205-214.
- Zahavi, A. & Zahavi, A. 1997. *The handicap principle: a missing piece of Darwin's puzzle*. Oxford University Press.
- Zhao, D.-X., Zhang, X.-F., Chen, D.-S., Zhang, Y.-K. & Hong, X.-Y. 2013. *Wolbachia*-host interactions: host mating patterns affect *Wolbachia* density dynamics. *PLoS ONE* **8**: e66373.
- Zilber-Rosenberg, I. & Rosenberg, E. 2008. Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* **32**: 723-735.
- Zug, R. & Hammerstein, P. 2015. Bad guys turned nice? A critical assessment of *Wolbachia* mutualisms in arthropod hosts. *Biol Rev* **90**: 89-111.
- Zuk, M. & Stoehr, A.M. 2002. Immune defense and host life history. *Am Nat* **160**: S9-S22.

# Summary

Bacteria are omnipresent in nature and a multitude of associations between bacteria and higher organisms (plants, animals and fungi) have been described. The effect of bacteria can vary from beneficial to neutral to detrimental as for the fitness of the organism with which the bacteria are associated. In this thesis, I investigated the bacterial communities as well as pathogenic and non-pathogenic effects of bacteria on life history traits of two generalist herbivorous arthropods. In the noctuid moth *Heliothis virescens*, I described the variability of bacterial communities that are associated with this moth in the laboratory and the field (CHAPTER 2). Furthermore, I determined the effect of immune challenge on mating success, female sex pheromone composition and oviposition strategy (CHAPTERS 3 & 4). In the two-spotted spider mite, *Tetranychus urticae*, I investigated the bacterial communities of two of its strains, as well as the effects of *Wolbachia*, *Cardinium* and *Spiroplasma* on mite survival and oviposition, mite gene expression and the induction of tomato responses after mite feeding (CHAPTER 5).

In CHAPTER 2, I describe the variability of bacterial communities that are associated with the moth *H. virescens* that was determined by using bacterial tag-encoded FLX pyrosequencing of 16S rRNA amplicons. Bacterial communities varied between *H. virescens* larvae that had fed on different host plants and between larvae originating from the field and those from laboratory rearing. Notably, while larvae from the laboratory contained an unidentified *Enterococcus* strain in high relative amounts, this bacterial strain was absent in field larvae. Moreover, field larvae contained a more diverse bacterial community than larvae from the laboratory. Bacterial communities also varied between biological replicates in larvae from the field and between moths of different developmental stages, i.e., eggs, larvae and female adults. Overall, the variability of bacterial communities in *H. virescens* suggests that this moth has not evolved a long-term, stable symbiosis with the detected bacteria. Whether bacteria that are transiently present in these moths have a beneficial effect on fitness and may for instance support host plant utilization remains to be investigated.

In CHAPTER 3, I show sexual dimorphism in immune system activation of the moth *H. virescens* in response to an immune challenge with dead cells of the bacterial entomopathogen *Serratia entomophila*. When measuring expression levels of immunity-related genes, females showed higher immune deployment, i.e., immune system activation in response to a pathogenic threat, whereas males had higher immune maintenance, i.e., their immune system was at a higher basic level of immune defence than females. Furthermore, the female immune response seemed to differentiate more between wounding and bacterial challenge, suggesting a more specific and possibly more cost-efficient immune response than the male immune system. Sexual dimorphism in immune response matched the results from mating experiments: bacteria-challenged females had lower mating success than non-

challenged females, while males showed no differences in mating success between bacteria-challenged, wounded or non-challenged males. This suggests that females trade off immunity for mating success, while males do not. Lastly, the sex pheromone composition of bacteria-challenged females differed from that of non-challenged and wounded females, suggesting that females may trade off immunity for sexual signalling, although the differences in female sex pheromone cannot explain the differences in mating success between females. Overall, the observed difference in the type of immunity between males and females adds a new dimension to the research of sexual dimorphism in immunity.

In CHAPTER 4, I demonstrate that immune challenge with dead cells of the bacterial entomopathogen *S. entomophila* affects the oviposition strategy of *H. virescens*. Females increased their oviposition rate in response to bacterial challenge. Notably, bacteria-challenged females had shorter life spans, but did not lay significantly fewer eggs than control females, indicating that bacteria-challenged females can compensate for a shorter life-span. However, bacteria-challenged females were less selective in terms of oviposition site than control females. Bacteria-challenged females laid equal numbers of eggs on plants that were infested with *H. virescens* larvae as on uninfested plants, whereas control females preferred uninfested plants. Since *H. virescens* larvae are cannibalistic and infested plants can be expected to have higher levels of secondary plant metabolites, non-preference of bacteria-challenged females likely has a negative effect on female fitness. The strategy of *H. virescens* females to increase their oviposition rate after an immune-challenge, which may compensate for a shortened life span, might thus come at the cost of reduced growth and survival of their offspring.

In CHAPTER 5 it is shown that various (endo)symbiotic bacteria of *T. urticae*, i.e., *Wolbachia*, *Cardinium* and *Spiroplasma*, affect mite survival and oviposition, and mite gene expression as well as induced responses of tomato, the mite's host plant, after mite feeding. These effects were investigated in two strains of *T. urticae*, one plant defence suppressor strain (DeLier-1) and one plant defence inducer strain (Santpoort-2) which contained different endosymbionts: while the suppressor strain harboured *Wolbachia* (W) and *Spiroplasma* (S), the inducer strain harboured *Cardinium* (C) and *Spiroplasma* (S). By means of antibiotic treatment, different mite groups were produced, containing or not containing W and S or C and S.

In the suppressor strain, the presence of *Wolbachia* was associated with higher survival of the mites, and higher expression of digestion- and detoxification-related genes, compared to mites that did not contain *Wolbachia*. Additionally, mites with *Wolbachia* reduced salicylic acid (SA) levels in tomato leaflets compared to leaflets that were infested with mites that contained only *Spiroplasma*. Further, *Spiroplasma* was associated with reduced JA-precursor [12-oxophytodienoic acid (OPDA)] levels in tomato leaflets, while *Wolbachia* had the opposite effect.

In the inducer strain, mites that contained *Cardinium* and *Spiroplasma* had lower survival, laid lower amounts of eggs and fed less than mites that did not

## Summary

contain the two bacterial strains. Moreover, leaflets that were infested with C+S+ mites had higher expression of SA-marker genes, and leaflets accumulated lower amounts of abscisic acid (ABA), than leaflets infested with mites without these bacteria. Further, C+S+ mites caused rusty/red feeding scars, while C-S- mites caused white feeding scars. Overall, *Wolbachia* seems to be beneficial for the mites, while *Cardinium* is likely a parasite. Even though the role of *Spiroplasma* is less clear, the results suggest that *Spiroplasma* may interact with *Wolbachia* in the suppressor strain, as *Wolbachia* seems to counteract the effects of *Spiroplasma*.

# Samenvatting

Bacteriën zijn alom aanwezig in de natuur en derhalve bestaat er een grote verscheidenheid aan associaties tussen bacteriën en hogere organismen (planten, dieren en schimmels). Effecten van bacteriën kunnen variëren van gunstig tot neutraal en schadelijk voor het organisme waarmee de bacteriën geassocieerd zijn. In dit proefschrift heb ik de samenstelling van de bacteriële gemeenschappen en de pathogene en niet-pathogene effecten van bacteriën op de levensgeschiedenissenmerken van twee generalistische geleedpotigen onderzocht. In de nachtvlinder *Heliothis virescens* heb ik de variabiliteit van de aanwezige bacteriële gemeenschappen in het veld en in het laboratorium in kaart gebracht (HOOFDSTUK 2). Ook heb ik het effect van een immuunuitdaging (d.w.z. een infectie) op het paringssucces en de samenstelling van het seksferomoon in vrouwtjes en hun ovipositie-strategie bepaald (HOOFDSTUKKEN 3 en 4). In twee lijnen van de spintmijt *Tetranychus urticae* heb ik de bacteriële gemeenschappen onderzocht en heb ik bepaald wat de effecten van *Wolbachia*, *Cardinium* en *Spiroplasma* bacteriën zijn op de overleving, ovipositie en gen-expressie van de gastheer en op de inductie van de reactie van tomatenplanten die door mijten zijn aangevreten (HOOFDSTUK 5).

In **HOOFDSTUK 2** beschrijf ik de variabiliteit van bacteriële gemeenschappen die geassocieerd zijn met *H. virescens*; de bacteriën werden moleculair-genetisch getypeerd met behulp van zogenoemde bacteriële tag-gecodeerde FLX pyrosequencing van 16S rRNA amplicons. De bacteriële gemeenschappen verschilden tussen *H. virescens* rupsen die gegeten hadden op verschillende planten en tussen rupsen uit het veld vergeleken met die uit het laboratorium. Verder bleek dat rupsen uit het laboratorium voornamelijk veel van een ongeïdentificeerde *Enterococcus*-soort bevatten, terwijl deze soort afwezig was in rupsen uit het veld. Rupsen uit het veld bevatten ook een meer diverse bacteriële gemeenschap dan rupsen uit het laboratorium. Aangezien de bacteriële gemeenschappen tussen biologische replicas van rupsen uit het veld en tussen verschillende ontwikkelingsstadia (eieren, rupsen en volwassen vrouwtjes) nogal varieerden, lijkt het erop dat in deze nachtvlinder geen stabiele, langdurige symbiose is ontwikkeld met de gevonden bacteria. Of de tijdelijk aanwezige bacteriën een gunstig effect hebben op de fitness van de nachtvlinders, doordat ze bijvoorbeeld betrokken zijn bij het verteren en ontgiften van voedsel zodat de rupsen op planten kunnen leven die in afwezigheid van de bacteriën niet geschikt zouden zijn, zal verder onderzocht moeten worden.

In **HOOFDSTUK 3** toon ik aan dat, wanneer volwassen *H. virescens* geïnjecteerd worden met dode cellen van de bacteriële entomopathogeen *Serratia entomophila*, de motten een sekse-specifieke activatie van het immuun-systeem laten zien. Bij het meten van de expressie-niveaus van immuun-gerelateerde genen vertoonden vrouwtjes een hogere activatie dan mannetjes, terwijl mannetjes een hogere handhaving van het immuunsysteem vertoonden, d.w.z. hun immuunsysteem had een hoger basis

expressie-niveau dan dat van vrouwtjes. Bovendien leek de immunrespons in vrouwtjes meer te differentiëren tussen verwonding op zich (injectie met controlestof) en een bacteriële injectie, hetgeen een specifiekere en mogelijk kosten-efficiëntere immunrespons in vrouwtjes dan in mannetjes suggereert. Dit sekse-specifieke onderscheid in immunrespons kwam overeen met de resultaten van de parings-experimenten: met bacterie geïnjecteerde vrouwtjes hadden een lager paringssucces dan niet-geïnjecteerde vrouwtjes, terwijl het paringssucces van mannetjes niet verschilde tussen bacterieel-geïnjecteerde, gewonde of niet-geïnjecteerde mannetjes. Dit suggereert dat vrouwtjes een balans vertonen tussen immuniteit en seksuele aantrekkingskracht, hoewel de verschillen in paringssucces niet verklaard kunnen worden door verschuivingen in het seksferomoon van het vrouwtje. De gevonden verschillen in immuniteit tussen vrouwtjes en mannetjes voegen een nieuwe dimensie toe aan het onderzoek naar sekse-specifieke immuniteit.

In **HOOFDSTUK 4** toon ik aan dat activatie van het immuunsysteem met dode cellen van de bacteriële entomopathogeen *S. entomophila* het ovipositie-gedrag van *H. virescens* beïnvloedt. Vrouwtjes verhoogden het aantal gelegde eieren na immun-activatie, maar leefden ook korter, zodat ze uiteindelijk dezelfde hoeveelheid eieren legden als niet-geactiveerde vrouwtjes. Dit suggereert dat bacterieel-geïnfecteerde vrouwtjes een kortere levensduur kunnen compenseren door sneller meer eieren te leggen. Dergelijke vrouwtjes waren echter tegelijkertijd minder selectief in hun ovipositiekeuze dan controle-vrouwtjes: de bacterieel-geactiveerde vrouwtjes legden dezelfde hoeveelheid eieren op planten met *H. virescens* rupsen als op planten zonder die rupsen, terwijl controle-vrouwtjes bij voorkeur hun eieren legden op de niet-aangevreten planten. Omdat *H. virescens*-rupsen kannibalistisch zijn, en met rupsen bezette planten waarschijnlijk meer secundaire plantenstoffen bevatten (als verdedigingsreactie van de plant op de insectenvraat), heeft het ontbreken van keuzegedrag van geïnfecteerde vrouwtjes hoogstwaarschijnlijk een negatief effect op de fitness van deze vrouwtjes. De strategie van *H. virescens* vrouwtjes om meer eieren te leggen na een immun-activatie, hetgeen kan compenseren voor een verkorte levensduur, gaat dus waarschijnlijk ten koste van een verminderde groei en overleving van hun nakomelingen.

In **HOOFDSTUK 5** beschrijf ik dat verschillende (endo)symbiotische bacteriën van de mijt *T. urticae*, in het bijzonder *Wolbachia*, *Cardinium* en *Spiroplasma*, de overleving en ovipositie van mijten beïnvloeden, evenals hun gen-expressie patronen en, na mijtenvraat, ook geïnduceerde responsen in tomaat, een waardplant van de mijt. Deze effecten zijn in twee mijten-lijnen onderzocht, een plantenafweer-onderdrukkende lijn (DeLier-1) en een plantenafweer-inducerende lijn (Santpoort-2), lijnen die verschillende endosymbionten bevatten: terwijl de onderdrukkende lijn *Wolbachia* (W) en *Spiroplasma* (S) bevat, herbergt de inducerende lijn *Cardinium* (C) en *Spiroplasma*. Door de mijten te behandelen met antibiotica konden verschillende groepen mijten geproduceerd worden, die wel of niet W en S of C bevatten.

In de onderdrukkende lijn was de aanwezigheid van *Wolbachia* (W+) geassocieerd met hogere overleving van de mijten en een hoger expressieniveau van verterings- en detoxificatie-gerelateerde genen, vergeleken met mijten zonder *Wolbachia* (W-). Ook verlaagden W+ mijten salicylzuur (SA)-niveaus in tomatenbladeren vergeleken met tomatenbladeren die met S+ mijten geïnfecteerd waren. De aanwezigheid van *Spiroplasma* was geassocieerd met gereduceerde niveaus van 12-oxofytodienoisch zuur, een voorloperstof van jasmonzuur, in tomatenbladeren, terwijl *Wolbachia* juist het tegenovergestelde effect hadden.

In de inducerende mijtenlijn hadden mijten met *Cardinium* en *Spiroplasma* (C+S+) een lagere overleving, legden ze minder eieren en vraten ze minder dan mijten die deze bacteriën niet bevatten. Bovendien vertoonden bladeren die geïnfecteerd waren met C+S+ mijten een hogere expressie van SA-merker genen en accumuleerden deze bladeren lagere hoeveelheden abscisinezuur dan bladeren met C-S- mijten. Ook veroorzaakten C+S+ mijten roestige, rode vraat-littekens, terwijl C-S- mijten juist witte vraat-littekens veroorzaakten. Kortom, de aanwezigheid van *Wolbachia* lijkt gunstig te zijn voor de mijten, terwijl *Cardinium* een parasiet lijkt te zijn. Hoewel de rol van *Spiroplasma* minder duidelijk is, suggereren de resultaten dat *Spiroplasma* en *Wolbachia* elkaar beïnvloeden in de onderdrukkende mijtenlijn, aangezien *Wolbachia* de effecten van *Spiroplasma* lijkt tegen te werken.

# Author contributions

## **Variability of bacterial communities in the moth *Heliothis virescens* indicates transient association with the host**

HEIKE STAUDACHER, *Martin Kaltenpoth, Johannes A.J. Breeuwer, Steph B.J. Menken, David G. Heckel & Astrid T. Groot*

HS, MK & ATG planned the experiments, HS conducted experiments, HS analysed the data, HS, MK, JAJB, DGH, SBJM & ATG wrote the manuscript

## **Sex-specific trade-offs between immune response and reproduction in a moth**

*Andrea Barthel\**, HEIKE STAUDACHER\*, *Antje Schmaltz, David G. Heckel & Astrid T. Groot*

AB, HS, DGH & ATG planned the experiments, AB & AS conducted the experiments, HS analysed the data, AB, HS, DGH & ATG wrote the manuscript

*\*Contributed equally to this study*

## **Effects of immune challenge on the oviposition strategy of a noctuid moth**

HEIKE STAUDACHER, *Steph B.J. Menken & Astrid T. Groot*

HS & ATG planned the experiments, HS conducted the experiments and analysed the data, HS, SBJM & ATG wrote the article

## **The bacterial symbionts *Wolbachia*, *Cardinium* and *Spiroplasma* affect gene expression and survival of their spider mite host and impact distinct induced responses in plants**

HEIKE STAUDACHER\*, *Bernardus C.J. Schimmel\**, *Mart M. Lamers, Nicky Wybouw, Astrid T. Groot & Merijn R. Kant*

HS, BCJS, ATG, & MRK planned the experiments, BCJS, HS & MML conducted the experiments, HS & NW analysed the data, HS, BCJS, ATG & MRK wrote the manuscript

*\*Contributed equally to this study*

## Author addresses

*University of Amsterdam, Institute for Biodiversity and Ecosystem Dynamics,  
Science Park 904, 1098 XH Amsterdam, The Netherlands*

Johannes A.J. Breeuwer, Astrid T. Groot, Heike Staudacher, Merijn R. Kant, Steph  
B.J. Menken, Bernardus C.J. Schimmel, Nicky Wybouw

*Max Planck Institute for Chemical Ecology, Hans-Knöll-Str. 8, 07745 Jena,  
Germany*

Andrea Barthel, Astrid T. Groot, David G. Heckel, Martin Kaltenpoth, Antje  
Schmaltz

*Johannes Gutenberg University, Department for Evolutionary Ecology, Johann-  
Joachim-Becher-Weg 13, 55128 Mainz, Germany*

Martin Kaltenpoth

*Erasmus MC, Department of Viroscience, Wytemaweg 80, Ee 1726, 3015 CN  
Rotterdam, The Netherlands*

Mart M. Lamers

# Biography

Heike Staudacher was born on the 30th of January, 1982 in Schramberg, Germany. She obtained the Abitur at the Gymnasium Schramberg in 2001. After the Abitur, Heike studied media informatics for one semester and took a 1-year course in digital film directing at the SAE in Amsterdam. In 2004, she started her studies of Biology at the Free University in Berlin, with paleontology, zoology and genetics as minors and ecology as major subject. She conducted her Diploma thesis at the Max Planck Institute for Chemical Ecology in Jena, Germany, and finished her studies in 2009. In 2010, Heike worked as a field technician on population dynamics of *Aedes aegypti* in Mexico for three months. In 2011, she started her PhD project at the Institute of Biodiversity and Ecosystem Dynamics of the University of Amsterdam with Dr. Astrid Groot as her supervisor and Prof. Dr. Steph Menken as her promotor.

## PUBLICATIONS

- Sato, Y., STAUDACHER, H. & Sabelis, M.W. 2016. Why do males choose heterospecific females in the red spider mite? *Experimental and Applied Acarology* **68**: 21-31.
- Barthel, A.\*, Staudacher, H.\*, Schmaltz, A., Heckel, D. G., & Groot, A.T. 2015. Sex-specific consequences of an induced immune response on reproduction in a moth. *BMC Evolutionary Biology* **15**: 282. (\*equal contribution)
- Melo, J.W.S., Lima, D.B., STAUDACHER, H., Silva, F.R., Gondim Jr, M.G.C. & Sabelis, M.W. 2015. Evidence of *Amblyseius largoensis* and *Euseius alatus* as biological control agent of *Aceria guerreronis*. *Experimental and Applied Acarology* **67**: 411-421.
- STAUDACHER, H., Menken, S.B.J. & Groot, A.T. 2015. Effects of immune challenge on the oviposition strategy of a noctuid moth. *Journal of Evolutionary Biology* **28**: 1568-1577.
- Groot, A.T., STAUDACHER, H., Barthel, A., Inglis, O., Schöfl, G., Santangelo, R., Gebauer-Jung, S., Vogel, H., Emerson, J. & Schal, C. 2013. One quantitative trait locus for intra- and interspecific variation in a sex pheromone. *Molecular Ecology* **22**: 1065-1080.
- Groot, A.T., Claßen, A., STAUDACHER, H., Schal, C. & Heckel, D.G. 2010. Phenotypic plasticity in sexual communication signal of a noctuid moth. *Journal of Evolutionary Biology* **23**: 2731-2738.

# Acknowledgements

First of all, I would like to thank my co-promotor Astrid Groot. Astrid, thanks for all the support that you gave to me in the last years. Thanks for all your ideas, your great enthusiasm for my project and for the freedom to investigate my own ideas. You taught me how important it is to plan well but also to let go, finish and submit things. Your amazing correction speed was particularly helpful with finishing the thesis. I learned from you that openness and courage are very important to move forward. Thanks for everything!

I was lucky to get much additional input from my advisors/ supervisors at the University of Amsterdam and the Max Planck institute for Chemical Ecology in Jena. Steph, I was very happy to have you as my promotor. Thank you for always having an open door for me, for your suggestions about how to proceed and especially for your input on my writing. Martin, I was lucky that you were involved in my thesis from the beginning on. Thank you for giving me a lot of input on my results and plans, especially for your help in investigating bacterial communities, where you supported me in planning, analysing and writing. Thank you also for letting me use your lab facilities in Jena and for the great congress travel to California with your group. Hans, thanks a lot for your input on my work. Your comments often helped me to see where I stand. You always took the time to discuss research plans and results with me and helped me in particular to improve one chapter as a co-author. David, thank you for letting me work in your lab in Jena, for supporting the sequencing of bacterial communities of *Heliothis virescens* and your input on the manuscript that it lead to. Thank you for your ideas and support of the trade-off manuscript of Andrea and me. Merijn, thank you for letting me sneak into your research about spider mite-plant interactions and supporting me and Bart with your suggestions, your investment into our ideas and your help with writing up the chapter. Thanks to the members of my doctorate committee for reading and evaluating my thesis.

Andrea and Bart, we did a lot of work together and I learned a lot from you. Andrea, thank you for your collaboration and for many tips during my PhD. It was great to plan and write our study together and share time with you at the course in Holland and the congress trip to California. Bart, thanks a lot, I am happy that I could do one chapter of my PhD project together with you. We made a good team and I am sure that we will get something great out of our 'side' project.

Thanks to the members of Astrid's group, Michiel, thanks for taking the time to think about ideas and especially discuss statistics with me. Your input helped me a lot to get the statistical part of my thesis done. Dennis, thank you for taking such good care of the moths. Pascaline, thank you, you brightened up my lab and culinary life and helped me a lot with phylogenetic analyses. Rik, thanks for all your tips and help in building things and for many nice dinners together with Sijing. Mart, thanks

for your ideas and motivation, you have helped me a lot with your work for us. Seyed, Masoumeh and Gao Ke, thanks for being such nice colleagues.

Many thanks to the lab technicians, Peter, Betsy and Lin for your good mood, your ideas and support in the molecular lab. Also, many thanks to the infamous greenhouse team, Ludek, Harold and Thijs, for planting and taking care of literally thousands of plants for me. Thank you Jan, for taking great fotos and for being such a fun bass player.

Karen and Rocio thank you for your friendship, your openness, positive energy and the many invaluable hours that we spent together in and outside of UvA in the past years. Debora and Wagner, it was a pleasure to collaborate, eat, chat and play in the snow with you. I am happy that I found such good friends. Joris, thanks for your friendship and many talks and excursions together. Yukie thank you for being such a kind person and for letting me participate in your work. Thanks to my many colleagues at UvA for your company at dinners, coffee breaks, BBQs and other excursions: Juanma, Carlos, Dan, Jie, Paulien, Nina, Sabina, Nicky, George, Maria, Bram, Nazer, Rizwan, Paul, Saioa, Marcus, Maus, Iza, Martijn, Arne, Thomas, Peter R, Firdevs, Josi, Marian, Ciska, Livia, Rachid, Kathrin, Alexandra, Orsi, Claire, Felipe, Ernesto, Estefania, Laura, Cleide, Louise, Anieke, Floor, Hanna. Meine Paranimf Alice, danke, dass du über Alles lachen kannst.

Meine liebe Familie, Mama, Papa, Christian, Fernando und Siglinde, ihr seid immer für mich da, egal was ich mache. Danke für eure unermüdliche Unterstützung. Es ist geschafft!