



UvA-DARE (Digital Academic Repository)

Brain networks of perceptual decision-making: an fMRI ALE meta-analysis

Keuken, M.C.; Müller-Axt, C.; Langner, R.; Eickhoff, S.B.; Forstmann, B.U.; Neumann, J.

DOI

[10.3389/fnhum.2014.00445](https://doi.org/10.3389/fnhum.2014.00445)

Publication date

2014

Document Version

Other version

Published in

Frontiers in Human Neuroscience

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Keuken, M. C., Müller-Axt, C., Langner, R., Eickhoff, S. B., Forstmann, B. U., & Neumann, J. (2014). Brain networks of perceptual decision-making: an fMRI ALE meta-analysis. *Frontiers in Human Neuroscience*, 8, [445]. <https://doi.org/10.3389/fnhum.2014.00445>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



Corrigendum: Brain networks of perceptual decision-making: an fMRI ALE meta-analysis

Max C. Keuken^{1*}, Christa Müller-Axt², Robert Langner^{3,4}, Simon B. Eickhoff^{3,4}, Birte U. Forstmann¹ and Jane Neumann^{2,5}

¹ Faculty of Social and Behavioural Science, Cognitive Science Center Amsterdam, University of Amsterdam, Amsterdam, Netherlands, ² Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany, ³ Institute of Clinical Neuroscience and Medical Psychology, Heinrich Heine University Düsseldorf, Düsseldorf, Germany, ⁴ Research Centre Jülich, Institute of Neuroscience and Medicine (INM-1), Jülich, Germany, ⁵ Leipzig University Medical Center, IFB Adiposity Diseases, Leipzig, Germany

Keywords: decision-making, meta-analysis, fronto-parietal-basal ganglia, corrigendum, GingerALE

A corrigendum on

Brain networks of perceptual decision-making: an fMRI ALE meta-analysis

by Keuken, M. C., Müller-Axt, C., Langner, R., Eickhoff, S. B., Forstmann, B. U., and Neumann, J. (2014). *Front. Hum. Neurosci.* 8:445. doi: 10.3389/fnhum.2014.00445

INTRODUCTION

Recently, Eickhoff et al. (2017) reported that the software version of GingerALE (2.3 <http://brainmap.org/ale/>) which was used for our meta-analysis published in Keuken et al. (2014) contained several severe implementation errors regarding the multiple-comparison correction. These errors resulted in a more liberal statistical threshold than was specified by the authors. With the newest release of GingerALE (2.3.6), these implementation errors should be remedied.

METHODS

In line with the recommendation by Eickhoff et al. (2017) and following the example by Smith and Delgado (2017) we re-analyzed the original data implementing the corrected multiple-comparison correction using GingerALE (2.3.6). All other statistical parameters were identical to those previously reported in Keuken et al. (2014). As expected, when applying a more stringent threshold, the major change in results pertains to a reduction in the number of significant clusters that survived correction for multiple comparisons. Here, we report results of our re-analysis, providing an adjusted original Table 4, indicating which clusters did not survive the correct statistical threshold (see **Table 4**).

RESULTS

For the contrast Task > Control condition, no clusters survived the multiple-comparison correction.

For the contrast Hard > Easy condition, the right pre-supplementary motor area (pre-SMA), right pre-central gyrus, right angular gyrus (hIP3) and the bilateral anterior insula were the only clusters that remained significant after correction. The biggest change was that there was no cluster in the inferior frontal gyrus (IFG), superior frontal gyrus, and occipital lobe.

OPEN ACCESS

Edited by:

Aron K. Barbey,
University of Illinois at
Urbana-Champaign, USA

Reviewed by:

Aron K. Barbey,
University of Illinois at
Urbana-Champaign, USA
Matthew J. Sutterer,
University of Iowa, USA

*Correspondence:

Max C. Keuken
mckeuken@gmail.com

Received: 09 November 2016

Accepted: 09 March 2017

Published: 22 March 2017

Citation:

Keuken MC, Müller-Axt C, Langner R,
Eickhoff SB, Forstmann BU and
Neumann J (2017) Corrigendum:
Brain networks of perceptual
decision-making: an fMRI ALE
meta-analysis.
Front. Hum. Neurosci. 11:139.
doi: 10.3389/fnhum.2017.00139

TABLE 4 | The original Table 4 of Keuken et al. (2014) adjusted for the correct statistical multiple comparison implementation.

Contrast	Region	Volume	x	y	z	ALE($\times 10^3$)
Task > Control (minimum cluster size 304 mm ³)	L pre-SMA		–	–	–	n.s.
	R pre-SMA		–	–	–	n.s.
	R insula; anterior part		–	–	–	n.s.
	R insula; anterior part		–	–	–	n.s.
	L insula; anterior part		–	–	–	n.s.
	R putamen		–	–	–	n.s.
	R inferior parietal lobule (PFop)		–	–	–	n.s.
	L middle frontal gyrus		–	–	–	n.s.
	R posterior cingulate gyrus		–	–	–	n.s.
	R inferior parietal lobule (hIP2)		–	–	–	n.s.
	R anterior occipital sulcus (hOC5)		–	–	–	n.s.
	L inferior frontal gyrus; <i>P. opercularis</i>		–	–	–	n.s.
	Hard > Easy (minimum cluster size 320 mm ³)	L pre-SMA	1,208	2	18	46
R insula; anterior part		680	38	20	0	20.1
R inferior frontal gyrus; <i>P. triangularis</i>		–	–	–	–	n.s.
R precentral gyrus		752	42	4	32	25.4
R angular gyrus; hIP3		32	28	–60	46	14
R superior occipital gyrus; SPL			–	–	–	n.s.
L inferior frontal gyrus; <i>P. opercularis</i>			–	–	–	n.s.
L precentral gyrus			–	–	–	n.s.
L precentral gyrus			–	–	–	n.s.
L insula; anterior part		280	–32	22	4	17.6
R precentral gyrus		128	32	–6	54	15.3
L superior frontal gyrus			–	–	–	n.s.
L superior frontal gyrus			–	–	–	n.s.
L superior parietal lobule (SPL)			–	–	–	n.s.
R inferior parietal lobule (hIP3)			–	–	–	n.s.
L inferior parietal lobule (hIP3)			–	–	–	n.s.
L middle occipital gyrus (hOC3v)			–	–	–	n.s.
R middle occipital gyrus			–	–	–	n.s.
L calcarine gyrus			–	–	–	n.s.
R calcarine gyrus			–	–	–	n.s.
R middle frontal gyrus			–	–	–	n.s.
L superior occipital gyrus		–	–	–	n.s.	
Reward anticipation > Control (minimum cluster size 288 mm ³)	R caudate nucleus	10,048	12	10	–10	33
	L putamen		–12	8	–10	26.5
	L caudate nucleus		–6	2	0	23
	R pallidum		10	4	–2	17.7
	R rectal gyrus		22	12	–16	17.2
	R amygdala		22	2	–20	15
	L amygdala		–14	2	–16	14.3
	L thalamus	1,344	0	–18	10	16
	R substantia nigra	1,192	8	–16	–18	19.6
	L mammillary body		–2	–16	–18	14.1
	R inferior frontal gyrus; <i>P. orbitalis</i>	1,032	36	22	–22	14.4
	R inferior frontal gyrus; <i>P. orbitalis</i>		42	22	–14	13
	R superior medial gyrus	640	6	46	30	15.2
	L frontal orbital cortex	544	–38	14	–16	14.5
	L cerebellum; lobule VII crus II	304	–22	–74	–42	13.8
	L parahippocampal gyrus	72	–22	–26	–12	9.6
	L anterior cingulate gyrus	72	0	42	12	10
	R parahippocampal gyrus	40	42	–46	–2	9.5
R superior medial frontal gyrus	16	6	52	16	9.3	

The structures that did not survive the correct statistical correction are now reported as not significant (n.s.).

For the contrast Reward Anticipation > Control condition there was no change in the reported clusters.

Given that there were no surviving clusters for the Task > Control contrast, the conjunction analysis as reported in the original Table 5 is non-informative.

DISCUSSION

It was surprising that the Task > Control analysis with 11 incorporated experiments resulted in no surviving clusters. The Hard > Easy condition with 13 incorporated experiments, and Reward anticipation > Control condition with 14 incorporated experiments replicate most, if not all, previously found clusters. This did not seem to be driven by the initially reported cluster size as the original Task > Control analysis reported similar cluster volumes as the other two analyses.

The changes in results should also be seen in light of recent recommendations of sample size in coordinate-based meta-analyses (Eickhoff et al., 2016). To have sufficient

power for moderate effects it is recommended to include a larger number of experiments in a given contrast than we included.

CONCLUSION

The original conclusion regarding a task-general network for perceptual decision-making is no longer warranted based on the corrected results in the Task > Control analysis. It thus remains an open question whether the lack of significant convergence is just a matter of limited power or whether there simply is no common network involved across the various included paradigms taxing perceptual decision-making.

To allow others to re-analyze our results and to incorporate additional experiments for sufficient statistical power, we have uploaded the raw input files. The data can be found on (<https://app.box.com/s/v974c7fdo6r1o89v96tuktyw170ol3>).

REFERENCES

- Eickhoff, S. B., Laird, A. R., Fox, P. M., Lancaster, J. L., and Fox, P. T. (2017). Implementation errors in the GingerALE Software: description and recommendations. *Hum. Brain Mapp.* 38, 7–11. doi: 10.1002/hbm.23342
- Eickhoff, S. B., Nichols, T. E., Laird, A. R., Hoffstaedter, F., Amunts, K., Fox, P. T., et al. (2016). Behavior, sensitivity, and power of activation likelihood estimation characterized by massive empirical simulation. *NeuroImage* 137, 70–85. doi: 10.1016/j.neuroimage.2016.04.072
- Keuken, M. C., Müller-Axt, C., Langner, R., Eickhoff, S. B., Forstmann, B. U., and Neumann, J. (2014). Brain networks of perceptual decision-making: an fMRI ALE meta-analysis. *Front. Hum. Neurosci.* 8, 1–14. doi: 10.3389/fnhum.2014.00445
- Smith, D. V., and Delgado, M. R. (2017). Meta-analysis of psychophysiological interactions: revisiting cluster-level thresholding and sample sizes. *Hum. Brain Mapp.* 38, 588–591. doi: 10.1002/hbm.23354

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Keuken, Müller-Axt, Langner, Eickhoff, Forstmann and Neumann. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.