



UvA-DARE (Digital Academic Repository)

Free SepF interferes with recruitment of late cell division proteins

Gao, Y.; Wenzel, M.; Jonker, M.J.; Hamoen, L.W.

Published in:
Scientific Reports

DOI:
[10.1038/s41598-017-17155-x](https://doi.org/10.1038/s41598-017-17155-x)

[Link to publication](#)

Citation for published version (APA):

Gao, Y., Wenzel, M., Jonker, M. J., & Hamoen, L. W. (2017). Free SepF interferes with recruitment of late cell division proteins. *Scientific Reports*, 7, [16928]. <https://doi.org/10.1038/s41598-017-17155-x>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <http://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Supplementary Information

Free SepF interferes with recruitment of late cell division proteins

Yongqiang Gao, Michaela Wenzel, Martijs J. Jonker and Leendert W. Hamoen

Table S1: Effect of SepF overexpressing on WalR regulon

Table S2: Effect of SepF overexpressing on genes involved in fatty acid synthesis

Table S3: Strains and plasmids used in this study

Table S4: Primer sequences used in this study

References

Fig. S1: SepF overexpression

Fig. S2: Aberrant membrane invaginations

Fig. S3: Transmission electron microscopy

Fig. S4: High SepF levels do not induce AccDA expression

Fig. S5: Most Z-rings do not co-localize with membrane invaginations

Fig. S6: Correlation of GFP-Pbp2B and FM5-95 fluorescence signals

Fig. S7: Walk is delocalized from Z-rings

Fig. S8. Effect of SepF overexpression on gene expression

Fig. S9: Suppression of growth and cell division defect by deleting either *ezrA* or *ftsA*

Table S1: Effect of SepF overexpressing on WalR regulon

Genes belonging to the WalR regulon¹⁻⁴ are listed with their fold expression difference (YK240/168) and adjusted p-value. “-” represents down-regulation genes.

Activated genes			
	YK240/wt	p.val	function
<i>cwlO</i>	1.5	0.169	cell wall synthesis, cell elongation
<i>ftsA</i>	-1.0	0.991	formation of Z-ring
<i>ftsZ</i>	-1.0	0.955	formation of Z-ring
<i>lytE</i>	-2.7	0.004	major autolysin, cell elongation and separation
<i>mreBH</i>	1.7	0.178	cell shape determation
<i>sigI</i>	1.3	0.593	control of a class of heat shock genes
<i>rsgI</i>	1.5	0.364	control of SigI activity
<i>tagA</i>	1.1	0.826	biosynthesis of teichoic acid
<i>tagB</i>	1.3	0.626	biosynthesis of teichoic acid
<i>tagD</i>	1.0	0.996	biosynthesis of teichoic acid
<i>tagE</i>	1.1	0.962	biosynthesis of teichoic acid
<i>tagF</i>	1.0	0.982	biosynthesis of teichoic acid
<i>ydjM</i>	3.0	0.071	may be involved in cell wall metabolism
<i>ykvT</i>	-1.1	0.957	hypothetical protein
<i>yocH</i>	-1.3	0.462	cell wall turnover
Repressed genes			
	YK240/wt	p.val	function
<i>iseA</i>	-3.0	0.023	protection against cell envelope stress
<i>pdaC</i>	-2.2	0.073	cell wall modification
<i>wapA</i>	-9.7	0.000	contact-dependent growth inhibition protein
<i>wapI</i>	-10.1	0.000	immunity protein against toxic activity of WapA

Table S2: Effect of SepF overexpressing on genes involved in fatty acid synthesis

Genes belonging to fatty acid synthetic systems were listed with their fold expression variation (YK240/168) and adjusted p-values. “-” represents down-regulated genes.

Genes involving in fatty acid synthesis			
	YK240/wt	p.val	function
<i>accA</i>	-1.02	0.962	acetyl-CoA carboxylase (alpha subunit)
<i>accB</i>	1.01	0.983	acetyl-CoA carboxylase (biotin carboxyl carrier subunit)
<i>accC</i>	1.00	0.995	acetyl-CoA carboxylase (biotin carboxylase subunit)
<i>accD</i>	-1.04	0.889	acetyl-CoA carboxylase (beta subunit)
<i>acpA</i>	-1.08	0.771	acyl carrier protein
<i>acpS</i>	1.14	0.512	acyl-carrier protein synthase, 4-phosphopantetheine transferase
<i>birA</i>	1.11	0.499	regulation of biotin synthesis, addition of biotin to proteins
<i>fabD</i>	1.14	0.746	malonyl CoA-acyl carrier protein transacylase
<i>fabF</i>	1.06	0.892	β -ketoacyl-acyl carrier protein synthase II, involved in the control of membrane fluidity
<i>fabG</i>	1.17	0.628	β -ketoacyl-acyl carrier protein reductase
<i>fabHA</i>	1.06	0.883	β -ketoacyl-acyl carrier protein synthase III, principal condensing enzyme responsible for the initiation of fatty acid synthesis in non-stressed B.subtilis cells
<i>fabHB</i>	1.41	0.517	β -ketoacyl-acyl carrier protein synthase III
<i>fabI</i>	1.11	0.664	enoyl-acyl carrier protein reductase
<i>fabL</i>	-1.01	0.980	enoyl-acyl carrier protein reductase
<i>fapR</i>	1.15	0.742	repressor of fatty acid synthetic genes
<i>plsC</i>	1.09	0.705	acyl-ACP:1-acylglycerolphosphate acyltransferase
<i>plsY</i>	-1.13	0.421	acylphosphate:glycerol-phosphate acyltransferase
<i>plsX</i>	1.18	0.680	acyl-acyl carrier protein (ACP): phosphate acyltransferase, catalyzes the synthesis of the intermediate fatty acyl-phosphate, coordinates membrane synthesis with cell division
<i>ycsD</i>	1.08	0.681	β -hydroxyacyl-ACP dehydratase
<i>ywpD</i>	1.48	0.523	two-component orphan sensor kinase
<i>ywpB</i>	-1.13	0.677	β -hydroxyacyl-ACP dehydratase

Table S3: Strains and plasmids used in this study

All strains were made in the *B. subtilis* 168 wild type background. Antibiotic resistance genes were abbreviated as follows: *kan* (kanamycin), *cat* (chloramphenicol), *erm* (erythromycin), *phleo* (phleomycin), *spec* (spectinomycin), *bla* (ampicillin).

Strains	Relevant features or genotype	Construction or reference
<i>B. Subtilis</i>		
168	<i>trpC2</i>	5
YK240	<i>amyE::cat P_{xyl}-sepF</i>	Yoshi Kawai, unpub.
YK1738	<i>amyE::spec P_{xyl}-accDA</i>	6
GYQ215	<i>amyE::cat P_{xyl}-sepF</i>	168 transformed with YK240
874	<i>ftsZ::spec ftsZ-gfp</i>	Laboratory stock
GYQ298	<i>amyE::cat P_{xyl}-sepF ftsZ::spec ftsZ-gfp</i>	GYQ215 transformed with 874
GYQ73	<i>aprE::erm P_{spac}-gfp-pbpB</i>	This study
GYQ81	<i>aprE::erm P_{spac}-gfp-ftsW</i>	This study
GYQ203	<i>aprE::erm P_{spac}-gfp-ftsL</i>	This study
GYQ72	<i>amyE::cat P_{xyl}-sepF aprE::erm P_{spac}-gfp-pbpB</i>	GYQ73 transformed with GYQ215
GYQ74	<i>amyE::cat P_{xyl}-sepF aprE::erm P_{spac}-gfp-ftsW</i>	GYQ81 transformed with GYQ215
GYQ204	<i>amyE::cat P_{xyl}-sepF aprE::erm P_{spac}-gfp-ftsL</i>	GYQ203 transformed with GYQ215
GYQ135	<i>chr. (pMUTIN4 P_{accDA}-lacZ erm)</i>	This study
GYQ136	<i>chr. (pMUTIN4 P_{accDA}-lacZ erm) amyE::cat P_{xyl}-sepF</i>	GYQ135 transformed with GYQ215
TB74	<i>yycH::erm</i>	Laboratory stock
BKE40380	<i>yycI::erm</i>	BGSC stock
BKE40370	<i>yycJ::erm</i>	BGSC stock
GYQ201	<i>yycH::erm</i>	168 transformed with TB74
GYQ202	<i>yycI::erm</i>	168 transformed with BKE40380
GYQ470	<i>yycJ::erm</i>	168 transformed with BKE40370
GYQ17	<i>amyE::cat P_{xyl}-sepF yycH::erm</i>	GYQ201 transformed with GYQ215
GYQ67	<i>amyE::cat P_{xyl}-sepF yycI::erm</i>	GYQ202 transformed with GYQ215
GYQ471	<i>amyE::cat P_{xyl}-sepF yycJ::erm</i>	GYQ470 transformed with GYQ215
bSS421	<i>amyE::spec P_{rpsD}-gfp</i>	Syvertsson, unpub.
GYQ254	<i>aprE::spec P_{rpsD}-gfp</i>	This study
GYQ257	<i>amyE::cat P_{xyl}-sepF aprE::spec P_{rpsD}-gfp</i>	GYQ254 transformed with GYQ215
GYQ132	<i>chr : (pMUTIN4 P_{ycdF}-lacZ erm)</i>	This study
GYQ133	<i>chr : (pMUTIN4 P_{ycdF}-lacZ erm) amyE::cat P_{xyl}-sepF</i>	GYQ132 transformed with GYQ215
GYQ195	<i>aprE::spec P_{srfAA}-lacZ</i>	This study
GYQ199	<i>amyE::cat P_{xyl}-sepF aprE::spec P_{srfAA}-lacZ</i>	GYQ195 transformed with GYQ215
GYQ217	<i>aprE::spec P_{mtnK}-lacZ</i>	This study
GYQ218	<i>amyE::cat P_{xyl}-sepF aprE::spec P_{mtnK}-lacZ</i>	GYQ217 transformed with GYQ215
GYQ144	<i>aprE::erm P_{spac}-gfp-walk</i>	This study
GYQ139	<i>amyE::cat P_{xyl}-sepF aprE::erm P_{spac}-gfp-walk</i>	GYQ144 transformed with GYQ215
GYQ124	<i>amyE::spec P_{xyl}-gfp-walk</i>	This study
3294	<i>chr::(P_{spac}-pbpB Kan) divIVA::P_{divIVA}-gfp-divIVA cat</i>	7
TNVS87	<i>chr::(P_{spac}-pbpB Kan)</i>	168 transformed with 3294
GYQ174	<i>chr::(P_{spac}-pbpB Kan) amyE::spec P_{xyl}-gfp-walk</i>	GYQ124 transformed with TNVS87
GYQ570	<i>aprE::erm P_{spac}-mcherry-walk</i>	This study
GYQ571	<i>amyE::cat P_{xyl}-sepF ftsZ::spec ftsZ-gfp aprE::erm P_{spac}-mcherry-walk</i>	GYQ298 transformed with GYQ570
GYQ152	<i>aprE::Kan P_{spac}-walR* R204C</i>	This study
GYQ159	<i>amyE::cat P_{xyl}-sepF aprE::Kan P_{spac}-walR* R204C</i>	GYQ152 transformed with GYQ215

TB07	<i>ftsA::erm</i>	Laboratory stock
PG49	<i>ezrA::spec</i>	⁸
TNVS158	<i>ezrA::spec</i>	168 transformed with PG49
TNVS101	<i>aprE::spec Pspac-ftsZ</i>	Saaki, unpub.
GYQ10	<i>amyE::cat P_{xyl}-sepF ftsA::erm</i>	GYQ215 transformed with TB07
GYQ130	<i>amyE::cat P_{xyl}-sepF ezrA::spec</i>	GYQ215 transformed with TNVS158
GYQ77	<i>amyE::cat P_{xyl}-sepF aprE::spec Pspac-ftsZ</i>	TNVS101 transformed with GYQ215
GYQ178	<i>amyE::cat P_{xyl}-sepF-L7D</i>	This study
GYQ179	<i>amyE::cat P_{xyl}-sepF-G109K</i>	This study
GYQ180	<i>amyE::cat P_{xyl}-sepF-F126S</i>	This study
3357	<i>ylmB-H::kan</i>	⁹
GYQ205	<i>amyE::cat P_{xyl}-sepF-L7D sepF::erm</i>	GYQ178 transformed with GYQ134
GYQ206	<i>amyE::cat P_{xyl}-sepF-G109K sepF::erm</i>	GYQ179 transformed with GYQ134
GYQ207	<i>amyE::cat P_{xyl}-sepF-F126S sepF::erm</i>	GYQ180 transformed with GYQ134
GYQ185	<i>amyE::cat P_{xyl}-sepF-F126S ylmB-H::kan</i>	GYQ215 transformed with 3357
GYQ223	<i>amyE::cat P_{xyl}-sepF-F126S ylmB-H::kan yycH::erm</i>	GYQ185 transformed with GYQ201
GYQ224	<i>amyE::cat P_{xyl}-sepF-F126S ylmB-H::kan yycI::erm</i>	GYQ185 transformed with GYQ202
GYQ210	<i>amyE::cat P_{xyl}-sepF-F126S sepF::erm aprE::(Pspac-ftsZ spec)</i>	GYQ207 transformed with TNVS101
PG62	<i>aprE::spec Pspac-yfp-ftsA</i>	
GYQ33	<i>amyE::cat P_{xyl}-sepF aprE::spec Pspac-yfp-ftsA</i>	GYQ215 transformed with PG62
EKB36	<i>zapA:(Cm P_{xyl}-mcherry-zapA)</i>	Koning, unpub.
GYQ29	<i>zapA:(Kan P_{xyl}-mcherry-zapA)</i>	This study
GYQ212	<i>amyE::cat P_{xyl}-sepF aprE::spec Pspac-yfp-ftsA zapA:(Kan P_{xyl}-mcherry-zapA)</i>	GYQ33 transformed with GYQ29
GYQ211	<i>amyE::cat P_{xyl}-sepF ftsZ::spec ftsZ-gfp zapA:(Kan P_{xyl}-mcherry-zapA)</i>	GYQ298 transformed with GYQ29
4057	<i>ezrA::Cm ezrA-gfp</i>	Laboratory stock
GYQ28	<i>ezrA::Kan ezrA-gfp</i>	This study
GYQ30	<i>amyE::cat P_{xyl}-sepF ezrA::Kan ezrA-gfp</i>	GYQ215 transformed with GYQ28

E. coli

Top10		Laboratory stock
Plasmid	Relevant features or genotype	Construction or reference
pAPNC213 <i>Cm</i>	<i>bla, aprE3', Cm, lacI, Pspac, aprE5'</i>	¹⁰
pAPNC213 <i>Erm</i>	<i>bla, aprE3', Erm, lacI, Pspac, aprE5'</i>	¹⁰
pAPNC213 <i>Kan</i>	<i>bla, aprE3', Kan, lacI, Pspac, aprE5'</i>	¹⁰
pMarB	<i>bla, erm Pctc Himar1 Kan (TnYLB-1)</i>	¹¹
pMutin4	<i>bla, erm, lacI, Pspac-lacZ</i>	⁵
pUC19	<i>bla, Plac</i>	¹²
pHJS105	<i>bla, amyE3', spec, P_{xyl}-gfp-MCS, amyE5'</i>	¹³ and H. Strahl
pEKC12	<i>bla, amyE3', spec, P_{xyl}-gfp-pbpB, amyE5'</i>	This study
pEKC13	<i>bla, amyE3', spec, P_{xyl}-gfp-ftsL, amyE5'</i>	This study
pEKC14	<i>bla, amyE3', spec, P_{xyl}-gfp-ftsW, amyE5'</i>	This study
pTNV9	<i>bla, aprE3', erm, lacI, Pspac-gfp, aprE5'</i>	This study
pTNV42	<i>bla, Cm</i>	This study
pTNV60	<i>bla, Cm 3', Kan, Cm 5'</i>	This study
pYQ01	<i>bla, aprE3', erm, lacI, Pspac-gfp-ftsW, aprE5'</i>	This study
pYQ02	<i>bla, aprE3', erm, lacI, Pspac-gfp-ftsL, aprE5'</i>	This study
pYQ03	<i>bla, aprE3', erm, lacI, Pspac-gfp-pbpB, aprE5'</i>	This study
pYQ05	<i>bla, erm, lacI, PydcF-lacZ</i>	This study
pYQ10	<i>bla, amyE3', spec, P_{xyl}-gfp-walk, amyE5'</i>	This study
pYQ11	<i>bla, aprE3', erm, lacI, Pspac-gfp-walk, aprE5'</i>	This study
pYQ13	<i>bla, aprE3', Kan, lacI, Pspac-walR, aprE5'</i>	This study
pYQ14	<i>bla, aprE3', Kan, lacI, Pspac-walR R204C, aprE5'</i>	This study
pYQ40	<i>bla, erm, lacI, PaccDA-lacZ</i>	This study
pYQ47	<i>bla, aprE3', spec, PsrfAA-lacZ, aprE5'</i>	This study
pYQ56	<i>bla, aprE3', spec, PmthK-lacZ, aprE5'</i>	This study
pYQ73	<i>bla, aprE3', spec, PrpsD-gfp, aprE5'</i>	This study
pYQ87	<i>bla, aprE3', spec, lacZ, aprE5'</i>	This study

Table S4: Primer sequences used in this study

Name	Sequence (5'-3')
YQ41	GCGCTCACAATTAGAAAAGGAGATTCCTAGGATGG
YQ42	GGGCTAACGCCTAAATAGTACATAATGGATTTCTT
YQ43	GTAATAATTTAGGCGTTAGCCCAAGCGCATC
YQ44	TCCTTTCTAATTGTGAGCGCTCACAATTCCACA
YQ52	ATGGGGAAGAGAACCGCTTAAG
YQ53	ATGTTTGCAAACGATTCAAAC
YQ72	ACAGCGGAATTGACTCCACATTGTGAAATCTATTGAC
YQ73	CACAATGTGGAGTCAATTCGGCTGTCGATAACA
YQ75	GGCTAACGCCCAATTTCGAGCTCttaGTCCTGTTCTGGGTTTCTCA
YQ76	GAGCTCGAATTCGGCGTTAG
YQ78	ACGTAACAGTCTGCCGGCTTCGTG
YQ79	CACGAAGCCGGCAGACTGTTACGT
YQ94	acgtaagcttACTCAACGTACCTGATATCCGCT
YQ95	TCATggatccTTTAAAATTGCCTTTCAAAGAATC
YQ98	CTTCCAGATAACTGCCGCTACT
YQ99	AAGGCTCAGGAAGCGGCTCAATGAATAAGGTTGGTTTTTTTCGGT
YQ100	TGAGCCGTTCTCTGAGCCTTTGTAGAGCTCATCCATGCCAT
YQ101	AGCTTATCGATACCGTCGACTCACGCTTCATCCCAATCATC
YQ102	acgtAAGCTTCGATGGAAACAGGTGTTGCA
YQ103	tcatggatccATGATTACCTCCCTTTTGTGAA
YQ104	ATGAAGTGGCGAAGTTCACGA
YQ105	AAACTATGCGAGTGAAGACGTAGA
YQ142	tagGCAGGAGGAAAATCAAATGatgGATAAAAAGATCCTTGTAGTAG
YQ143	CATTTTGATTTTCTCCTGCctaGCAGGTCAATTGTGAGCGCTCACAATTCCAC
YQ149	GAGTATGAAAAATAAAGACAAAAACTTTTTCTCAATG
YQ150	CATTGAGAAAAAGTTTTTGTCTTTATTTTTCATACTC
YQ151	GACTTTTTAAGCAAGACCGTTTATGCCA
YQ152	TGGCATAAACGGTCTTGCTTAAAAAGTC
YQ153	TCGGCTCAGATATTTCCCTCTGCACGCCTGAC
YQ154	GTCAGGCGTGCAGAGGGAAATATCTGAGCCGA
YQ214	ATGGAAGTTACTGACGTAAGATTAC
YQ215	ATTTAGTCCGAATAGTCTGGA
YQ216	CTTACGTACGTAACCTCCATCTTGACCACTTACCCATAATTTT
YQ217	AAGCTTGTAGTTAAAGCTTTTTAGACATCTAATGCCCGGTTATTATTATTTTGGAC
YQ218	AAAGCTTTAACTACAAGCTTTTTAGACATCTAATGATATCGAATTCTAGTTCTAGAGC
YQ219	CAAAAGCCTAATTGAGAGAAGTTTCTATAGA
YQ220	CTTCTCTCAATTAGGCTTTTGTAAATTTGGAAAGTTACAC
YQ236	CAGACTATTCGGCACTGAAATGAAAGCCTCATGCCTATTCTTG
YQ237	TCTTACGTACGTAACCTCCATATTGTCATACCTCCCCTAATCT
YQ246	AGACTATTCGGCACTGAAATAGGAAGGGCAAATCATTAAGAGT
YQ247	TCTTACGTACGTAACCTCCATAACCTCCAATTATGTAATTAATTAATATG
YQ450	CTAGGATGGGTACCCTGCAGATGGTCAGCAAGGGAGAGGA
YQ451	CTGCAGGGTACCCATCCTAGGAATC
YQ452	CCGGCTCAGGAAGCGGCTCAATGAATAAGGTTGGTTTTTTTCGGT
YQ453	TGAGCCGTTCTCTGAGCCGGATCCTGAGCCGCTTCTGA
TerS117	TCGACTCTAGAGGATCCCCGGGT

TerS118	CCTGCAGGCATGCAAGCTTGGCGT
TerS125	CAAGCTTGCATGCCTGCAGGATGAACTTTAATAAAAATTGATTTAGACAATTGGA
TerS126	CGGGGATCCTCTAGAGTCGAATAAAAAGCCAGTCATTAGGCCT
TerS135	GGGCGTTAGCCCAAGCGCATCA
TerS136	GGTCAATTGTGAGCGCTCACAATTCCACA
TerS139	GTGAGCGCTCACAATTGACCGGGTACCCTGCAGATGAGCAAAGGA
TerS140	ATGCGCTTGGGCTAACGCCCGCGGCCGCTCTAGAACTAGA
TerS257	TagGACTTCAAAGAGTTTTATGATTTATACCT
TerS258	cTAGTCATCCTTTACAGGAGTCAAATACCA
TerS259	CTCCTGTAAAGGATGACTAgTGGTTTCAAATCGGCTCCGTCGA
TerS260	TAAAACCTTTGAAGTCctAACATCAGAGTATGGACAGTTGCGGA
TerS350	CACCGCCGACATTGCGGTGGCTCCA
TerS351	GCATCAGGGCTGCGGCATCCGGA
TerS352	GGGGCCAATAAACGGATTGTATTGT
TerS353	GCCTCTGCCCTTGCAAATCGGATGCCT
EKP22	GTCGACGGTATCGATAAGCTTGAT
EKP30	GTGGATCCGAAGTCTGGACATTTT
EKP31	TGTCCAGACTTCGGATCCACatgATTCAAATGCCAAAAAA
EKP32	ATCAAGCTTATCGATACCGTCGACTtaATCAGGATTTTTA
EKP33	TGTCCAGACTTCGGATCCACatgAGCAATTTAGCTTACCA
EKP34	AGCTTATCGATACCGTCGACTcaTTCCTGTATGTTTTTCA
EKP38	TGTCCAGACTTCGGATCCACatgTAAAAAAAATGCTAAA
EKP39	AGCTTATCGATACCGTCGACTtaCAGATAAACAGTTTTTT

References

- 1 Salzberg, L. I. *et al.* The WalRK (YycFG) and σ^I RsgI regulators cooperate to control CwlO and LytE expression in exponentially growing and stressed *Bacillus subtilis* cells. *Molecular microbiology* **87**, 180-195 (2013).
- 2 Bisicchia, P. *et al.* The essential YycFG two-component system controls cell wall metabolism in *Bacillus subtilis*. *Molecular microbiology* **65**, 180-200 (2007).
- 3 Howell, A. *et al.* Genes controlled by the essential YycG/YycF two-component system of *Bacillus subtilis* revealed through a novel hybrid regulator approach. *Molecular microbiology* **49**, 1639-1655 (2003).
- 4 Fukuchi, K. *et al.* The essential two-component regulatory system encoded by *yycF* and *yycG* modulates expression of the *ftsAZ* operon in *Bacillus subtilis*. *Microbiology* **146**, 1573-1583 (2000).
- 5 Vagner, V., Dervyn, E. & Ehrlich, S. D. A vector for systematic gene inactivation in *Bacillus subtilis*. *Microbiology* **144**, 3097-3104 (1998).
- 6 Mercier, R., Kawai, Y. & Errington, J. Excess membrane synthesis drives a primitive mode of cell proliferation. *Cell* **152**, 997-1007 (2013).
- 7 Hamoen, L. W. & Errington, J. Polar Targeting of DivIVA in *Bacillus subtilis* Is Not Directly Dependent on FtsZ or PBP2B. *Journal of bacteriology* **185**, 693-697 (2003).
- 8 Gamba, P., Rietkötter, E., Daniel, R. A. & Hamoen, L. W. Tetracycline hypersensitivity of an *ezrA* mutant links GalE and TseB (YpmB) to cell division. *Frontiers in microbiology* **6** (2015).
- 9 Hamoen, L. W., Meile, J. C., de Jong, W., Noirot, P. & Errington, J. SepF, a novel FtsZ-interacting protein required for a late step in cell division. *Molecular microbiology* **59**, 989-999 (2006).
- 10 Morimoto, T. *et al.* Six GTP-binding proteins of the Era/Obg family are essential for cell growth in *Bacillus subtilis*. *Microbiology* **148**, 3539-3552 (2002).
- 11 Le Breton, Y., Mohapatra, N. P. & Haldenwang, W. G. *In vivo* random mutagenesis of *Bacillus subtilis* by use of TnYLB-1, a mariner-based transposon. *Applied and environmental microbiology* **72**, 327-333 (2006).
- 12 Yanisch-Perron, C., Vieira, J. & Messing, J. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mpl8 and pUC19 vectors. *Gene* **33**, 103-119 (1985).
- 13 Lewis, P. J. & Marston, A. L. GFP vectors for controlled expression and dual labelling of protein fusions in *Bacillus subtilis*. *Gene* **227**, 101-109 (1999).




Fig. S1: SepF overexpression

Western blot analysis of SepF levels in SepF-overexpressing strain GYQ215 (*amyE::Pxyl-sepF*) grown in the presence (+) or absence (-) of 1% xylose for 3h. SepF primary antibody was used to detect the protein.




Fig. S2: Aberrant membrane invaginations

Microscopic images of strain GYQ215 (*amyE::Pxyl-sepF*) grown in presence of 1% xylose for 3h. Membranes were fluorescently stained with FM5-95.




Fig. S3: Transmission electron microscopy

Transmission electron microscopy images of GYQ215 cells (*amyE::P_{xyI}-sepF*) grown in LB with 1% xylose to overexpress SepF. Arrows indicate membrane invaginations. Scale bar is 200 nm.




Fig. S4: High SepF levels do not induce AccDA expression

(A) N-SIM image of strain GYQ1738 (*amyE::P_{xyI}-accDA*) grown with 1% xylose and stained with the membrane dye MitoTracker green showing the membrane invaginations. In the lower panel the brightness is reduced to better reveal membrane structures. (B) Effect of SepF overexpression on *accDA* promoter activity. β -galactosidase activities of *P_{accDA}-lacZ* in the SepF overexpression strain (GYQ136, *amyE::P_{xyI}-sepF P_{accDA}-lacZ*) grown in either the absence or presence (SepF overexpression) of 1 % xylose for 2 and 3 h.





Fig. S5: Most Z-rings do not co-localize with membrane invaginations

(A) Microscopic images of strain GYQ298 (*amyE::P_{xyl}-sepF ftsZ::ftsZ-gfp*) grown in the presence of 1% xylose for 3 h to overproduce SepF. Membranes were fluorescently stained with FM5-95. Scale bar is 5 μ m. (B) Analysis of Z-rings that co-localized with membrane invaginations or not. In total 258 Z-rings were counted in 42 cells.




Fig. S6: Correlation of GFP-Pbp2B and FM5-95 fluorescence signals

Strain GYQ72 (*amyE::Pxyl-sepF aprE::P_{spac}-gfp-pbpB*) expressing GFP-Pbp2B) was grown with 1 % xylose for SepF overexpression and 5 mM IPTG for GFP-Pbp2B induction. Membranes were fluorescently stained with FM5-95. Scale bar is 2 μ m. (A) White box with solid line indicates an invagination region selected for quantification of the GFP-Pbp2B and FM5-95 signals, and the box with dotted line shows a normal membrane region selected for quantification of the GFP-Pbp2B and FM5-95 signals. (B) Average ratio of GFP-Pbp2B and FM5-95 signals of 8 invaginations regions and 8 normal membrane regions.




Fig. S7: Walk is delocalized from Z-rings

Microscopic images of strain GYQ571 (*amyE::Pxyl-sepF ftsZ-gfp aprE::Pspac-mCherry-walk*) expressing both FtsZ-GFP and mCherry-WalkK in the presence of 1 % xylose to overexpress SepF. Scale bar is 5 μm .




Fig. S8. Effect of SepF overexpression on gene expression

Transcriptional activity of promoters *PydcF*, *PsrfAA* and *PmtnK* in the absence (-) or presence (+) of extra SepF. SepF overexpression was achieved by growth in the presence of 1 % xylose. Transcription was measured using *lacZ* reporters (strain GYQ133, *amyE::P_{xyl}-sepF PydcF-lacZ*, strain GYQ199, *amyE::P_{xyl}-sepF PsrfA-lacZ*, and strain GYQ218, *amyE::P_{xyl}-sepF PmtnK-lacZ*, respectively).




Fig. S9: Suppression of growth and cell division defect by deleting either *ezrA* or *ftsA*

(A) Growth of strains 168 (wild type), strain GYQ215 (*amyE*::*PxyI-sepF*), GYQ10 (*amyE*::*PxyI-sepF* Δ *ftsA*) and GYQ130 (*amyE*::*PxyI-sepF* Δ *ezrA*) in the presence of 1 % xylose (SepF overexpression). (B) Microscopic images of GYQ10 and GYQ130 after 3 h growth in medium with 1 % xylose. Membranes were stained with FM5-95. Scale bar is 5 μ m.