



## UvA-DARE (Digital Academic Repository)

### VLA observations identify the currently active source in Terzan 5 as the neutron star transient EXO 1745-248

Tremou, E.; Sivakoff, G.; Bahramian, A.; Heinke, C.; Tetarenko, A.; Wijnands, R.; Degenaar, N.; Linares, M.; Miller-Jones, J.; Patruno, A.; Chomiuk, L.; Strader, J.; Altamirano, D.; Maccarone, T.; Sanna, A.

**Publication date**

2015

**Document Version**

Final published version

**Published in**

The astronomer's telegram

[Link to publication](#)

**Citation for published version (APA):**

Tremou, E., Sivakoff, G., Bahramian, A., Heinke, C., Tetarenko, A., Wijnands, R., Degenaar, N., Linares, M., Miller-Jones, J., Patruno, A., Chomiuk, L., Strader, J., Altamirano, D., Maccarone, T., & Sanna, A. (2015). VLA observations identify the currently active source in Terzan 5 as the neutron star transient EXO 1745-248. *The astronomer's telegram*, 7262. <http://www.astronomerstelegam.org/?read=7262>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

# The Astronomer's Telegram

Post | Search | Policies  
Credential | Feeds | Email

24 Dec 2015; 09:17 UT

## Outside

GCN  
IAUCs

## Other

ATel on [Twitter](#) and [Facebook](#)  
ATELstream  
ATel Community Site  
MacOS: [Dashboard Widget](#)

## Your Conference Here

This space for free for  
conference advertising.

[ [Previous](#) | [Next](#) | [ADS](#) ]

## VLA observations identify the currently active source in Terzan 5 as the neutron star transient EXO 1745-248

ATel #7262; *E. Tremou (Michigan State), G. Sivakoff, A. Bahramian, C. Heinke, A. Tetarenko (Alberta), R. Wijnands (Amsterdam), N. Degenaar (Cambridge), M. Linares (IAC), J. Miller-Jones (ICRAR-Curtin), A. Patruno (Leiden), L. Chomiuk, J. Strader (Michigan State), D. Altamirano (Southampton), T. Maccarone (Texas Tech), A. Sanna (UniCA)*  
on 20 Mar 2015; 18:19 UT

*Distributed as an Instant Email Notice Transients*

*Credential Certification: Gregory R Sivakoff (sivakoff@ualberta.ca)*

Subjects: Radio, X-ray, Binary, Globular Cluster, Neutron Star, Transient

Referred to by ATel #: [7430](#), [7470](#)

Tweet

Recommend

1

Following the recent detection of a transient outburst in Terzan 5, and initial Swift X-ray follow-up (ATels [#7240](#), [#7242](#), [#7247](#)), we triggered radio observations with the VLA to identify and classify the source via its radio emission. VLA observations were taken from 10:14 UT to 10:43 UT on 2015 March 19 (MJD=57100.43639) with the array in its relatively-extended B-configuration using two 2048-MHz basebands centred at 9.0 and 11.0 GHz. The total exposure time on source was 26 min. Compared to existing VLA data, a transient source was detected and was consistent with an unresolved point source down to the beam size of 1.75 x 0.54 arcseconds (at a position angle of 146.6 deg) at 10.0 GHz.

We therefore fit a point source model in the image plane, yielding flux densities of 23 $\pm$ 5 microJy/beam averaging the two bands together. This corresponds to a radio luminosity ( $L_R = v L_\nu$ ) of 1.0 $\pm$ 0.2 x 1e28 (d/5.9 kpc)<sup>2</sup> erg/s at 10 GHz. The source was well-detected in the lower 9 GHz baseband (flux density 27 $\pm$ 7 microJy/beam) but only marginally detected in the upper 11 GHz baseband (flux density 16 $\pm$ 8 microJy/beam). This is consistent with a steep spectral slope, but the uncertainties are extremely large and include a flat spectral slope; our measured spectral index (defined by  $S_\nu \propto \nu^\alpha$ ) is -2.7  $\pm$  2.8.

The source position in the combined image was

RA (J2000) = 17:48:05.231  $\pm$  0.008 (267.021794 deg;  $\pm$ 0.11 arcsec)

DEC (J2000) = -24:46:47.73  $\pm$  0.14 (-24.779925 deg;  $\pm$ 0.14 arcsec),

where the errors only indicate the statistical errors from the source fitting procedure. We note that nominal systematic uncertainties are estimated to be of order 10% of the beam size, i.e., 0.18 x 0.05 arcseconds. The source position is within 0.4 arcseconds of the published location of EXO 1745-248 (CXOG1b J174805.2-244647 = CX 3; Heinke et al. 2006), but is 2.5 arcseconds away from the only other close, previously identified faint X-ray source (CXOGb J174805.1-244645 = CX 24; Heinke et al. 2006). We therefore confirm the previously suggested identification based on the X-ray location (ATel [#7247](#)) that EXO 1745-248 is the currently outbursting source in Terzan 5.

## Related

- [7470](#) **Optical identification of the X-ray burster EXO 1745-248 in Terzan 5**
- [7430](#) **A hard-to-soft state transition is going on in neutron star transient EXO 1745-248**
- [7262](#) **VLA observations identify the currently active source in Terzan 5 as the neutron star transient EXO 1745-248**
- [7247](#) **Swift refined location points to the neutron star transient EXO 1745-248 as the source currently active in Terzan 5**
- [7242](#) **Swift/XRT follow up of the X-ray transient in the globular cluster Terzan 5**
- [7240](#) **Swift/BAT detects an outburst from an X-ray transient in the globular cluster Terzan 5**

The increasing Swift/BAT flux suggests that the source remains in the hard spectral state, and that the radio emission is therefore likely to arise from compact jets.

On 2015 March 17 17:41 UT, the X-ray luminosity of the source was  $\sim 1.6e36 ((d/5.9 \text{ kpc})^2)$ ; 0.5-10 keV) erg/s, which given the measured spectrum corresponds to  $\sim 1e36 ((d/5.9 \text{ kpc})^2)$ ; 3-9 keV) erg/s. Assuming that the spectrum has not changed since its first detection, we use the Swift BAT count rates (15-50 keV) for the entire globular cluster to estimate the X-ray luminosities at the time of our radio observations to be  $\sim 2.4e36 ((d/5.9 \text{ kpc})^2)$ ; 0.5-10 keV) erg/s and  $\sim 1.5e36 ((d/5.9 \text{ kpc})^2)$ ; 3-9 keV) erg/s. We compare the radio and X-ray luminosity of this source with the two atoll neutron stars with well established tracks in the  $L_R$ - $L_X$  "fundamental plane" (e.g., Chomiuk et al. 2013), both of which are less radio-luminous than accreting millisecond X-ray pulsars / transitional millisecond pulsars. The current outburst of EXO 1745-248 is more consistent with the neutron star track of 4U 1728-34 (Migliari et al. 2003) than that of Aql X-1 (Tudose et al. 2009). We note that EXO 1745-248 is (about 2.5 - 3 times) less radio luminous than both known neutron star tracks; however, given the non-simultaneous nature of our radio and X-ray measurements, further observations are needed to compare these sources.

We strongly encourage simultaneous multi-wavelength observations at all frequencies.

We thank the NRAO staff for rapid scheduling of these VLA observations.

---

[ **Telegram Index** ]

R. E. Rutledge, Editor-in-Chief

`rrutledge@astronomerstelegam.org`

Derek Fox, Editor

`dfox@astronomerstelegam.org`

Mansi M. Kasliwal, Co-Editor

`mansi@astronomerstelegam.org`